

1-1-2021

Explicit tight frames for simulating a new system of fractional nonlinear partial differential equation model of Alzheimer disease

Mutaz Mohammad
Zayed University

Alexander Trounev
Kuban State Agrarian University

Follow this and additional works at: <https://zuscholars.zu.ac.ae/works>




Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Mohammad, Mutaz and Trounev, Alexander, "Explicit tight frames for simulating a new system of fractional nonlinear partial differential equation model of Alzheimer disease" (2021). *All Works*. 1590.
<https://zuscholars.zu.ac.ae/works/1590>

This Article is brought to you for free and open access by ZU Scholars. It has been accepted for inclusion in All Works by an authorized administrator of ZU Scholars. For more information, please contact Yrjo.Lappalainen@zu.ac.ae, nikesh.narayanan@zu.ac.ae.



Explicit tight frames for simulating a new system of fractional nonlinear partial differential equation model of Alzheimer disease

Mutaz Mohammad^{a,*}, Alexander Trounev^b

^a Zayed University, United Arab Emirates

^b Kuban State Agrarian University, Russia

ARTICLE INFO

Keywords:

Time fractional differential equation
Caputo's fractional derivative
Alzheimer disease
Mathematical modeling
Drug treatment
Tight frames

ABSTRACT

This paper is devoted to develop a new mathematical model for Alzheimer disease based on a system of fractional-order partial differential equations. The system of Alzheimer disease includes neurons, astrocytes, microglia and peripheral macrophages, as well as amyloid β aggregation and hyperphosphorylated tau proteins. We consider the Caputo fractional derivative definition to analyze the formulated system by simulating the effect of drugs that either failed or currently in clinical trials. To simulate the model, we use tight frame (framelet) systems generated using the unitary and oblique extension principle. According to the simulation results, and based on using such new direction of fractional modeling, the progression of Alzheimer disease and its consequences will be slowing down. Which may give clinical insights on intervention measures against the disease and its effective therapies.

Introduction

Alzheimer's disease (AD) is a progressive neurodegenerative disease that causes brain cells to waste away which leads to protein aggregation, where the neurodegeneration are only partially understood. It is known as the most common form of dementia and mortality in the elderly [1,2]. Noticing that, the increasing number of older adults afflicted with AD, is causing many challenges for the global health community. For example, the prevalence of AD worldwide in 2006 was 26.12 million, in 2019 it is estimated that 46.8 million people suffer from AD, and by 2030 the prevalence estimated to be 74.7 million people [3], and approximately it is expected that this number to be increased by 1.2% in 2050 [4] taking into account the huge amount of cost of caring for AD adults afflicted. For example, in the United States of America the cost was approximately \$290 billion in 2019, but the costs extend to many other reasons such as the negative mental and physical health outcomes [5]. Given these circumstances (among others), developing models that describe the dynamics of such disease is essential. Consequently, there is an immediate call for effective and efficient models that be able to provide some insights for effective description of the dynamics of AD including neurons, astrocytes, therapies and many other factors for the disease.

Most recently, many scientists have applied fractional derivatives using wavelets, framelets and many other systems on many scientific

and disease models in order to achieve a better matched with the reality and dynamics of these models rather than using the classical ordinary differential equations [6–15,17–37]. It is well-known that the fractional order options, especially with the Caputo fractional derivative, are more suitable to describe the medical phenomena than the classical and integer order [38–40]. This is also due to the fact that Caputo fractional definition describe the modeling of the non-local issues more efficiently.

The main aim of this work is to extend the AD mathematical model presented in [41], that is initially formulated based on integer-order derivative, by applying the Caputo fractional operator derivative. There are many advantages of using such derivative operator in applications. In future, we intended to study this AD model in other sense such as Atangana-Baleanu and Caputo-Fabrizio and compare their differences with the Caputo derivative in order to investigate which one is more appropriate and reliable for describing such model.

Preliminaries and notations

Let us introduce some definitions and notations of tight frames and fractional calculus needed through the paper. An $L^2(\mathbb{R})$ -function is a function maps \mathbb{R} to itself such that

* Corresponding author.

E-mail address: Mutaz.Mohammad@zu.ac.ae (M. Mohammad).

$$\|f\|_{L^2(\mathbb{R})}^2 = \left(\int_{\mathbb{R}} |f(x)|^2 dx \right),$$

exists and finite.

The Fourier transform of a function $g \in L_1(\mathbb{R})$ (can be extended to $L_2(\mathbb{R})$) is defined to be

$$\mathcal{F}(g)(\omega) = \widehat{g}(\omega) = \int_{\mathbb{R}} g(x) e^{-i\omega x} dx, \omega \in \mathbb{R},$$

and the Fourier series of a sequence $a \in \ell_2(\mathbb{Z})$ is defined by

$$\mathcal{F}(h)(\omega) = \widehat{h}(\omega) = \sum_{k \in \mathbb{Z}} a[k] e^{-i\omega k}, \quad \omega \in \mathbb{R}.$$

Definition 0.1. A sequence of functions $\{g_k \in L^2(\mathbb{R}), k = 1, \dots, \infty\}$ is called a *frame* of $L^2(\mathbb{R})$ if there exist a positive constants T_1, T_2 such that,

$$T_1 \|g\|^2 \leq \sum_{k=1}^{\infty} |\langle g, g_k \rangle|^2 \leq T_2 \|g\|^2, \quad \forall g \in L^2(\mathbb{R}).$$

The constants T_1, T_2 are called frame bounds. A frame in Definition 0.1 is called *Parseval (tight)* if it is possible to have $T_1 = T_2$, and $T_1 = T_2 = 1$, respectively.

In applications, it is recommend to use a non-negative functions to simulate the problem. B-splines are an example of non-negative functions and have many mathematical features such as symmetry and refinability. The B-spline B_m of order m is defined by the following relation,

$$B_m = B_{m-1} * B_1 = \int_{(0,1)} B_{m-1}(\cdot - x) dx, \quad m = 1, \dots, \infty,$$

where

$$B_1 = \chi_{(0,1)},$$

is defined to be the indicator function over the interval $[0, 1]$.

We say a function g has N vanishing moments if

$$\int t^m g(t) dt = 0, \quad \text{for } m = 0, 1, \dots, N - 1.$$

In this paper, we use the B-splines as refinable functions to construct the tight frame systems in order to simulate thenew formulated AD model.

For $g \in L^2(\mathbb{R})$, we define the dilation operator D by $Dg(x) = \sqrt{2}g(2x)$, and the translation operator T by $T_k g(x) = g(x - k)$ for $k \in \mathbb{Z}$. A compactly supported square-integrable function ϕ is called refinable if we have

$$\begin{aligned} \psi_1(x) &= \frac{1}{\sqrt{2}} \left(-|x-1| + |x+1| - 2\left|x + \frac{1}{2}\right| + \text{sgn}\left(\frac{1}{2} - x\right) + 2x \text{sgn}\left(x - \frac{1}{2}\right) \right), \\ \psi_1(x) &= \frac{1}{2} \left(-6|x| - |x+1| + 4\left|x - \frac{1}{2}\right| + 2|2x+1| + x \text{sgn}(1-x) + \text{sgn}(x-1) \right). \end{aligned}$$

$$\phi(x) = 2 \sum_{k \in \mathbb{Z}} a_0[k] \phi(2x - k),$$

where $a_0[k]$ is finitely supported sequence.

Let $\Psi = \{\psi_\ell\}_{\ell=1}^r$ be a set of square integrable functions such that

$$\psi^\ell = 2 \sum_{k \in \mathbb{Z}} b^\ell[k] \phi(2 \cdot - k),$$

where $\{b_\ell[k], k \in \mathbb{Z}\}_{\ell=1}^r$ is a finitely supported sequence. To simulate the non-linear system of partial differential equations PDEs for the AD in this work we need to construct tight frame systems of the form of $X(\Psi)$, where

$$X(\Psi) = \left\{ \psi_{j,k}^\ell = D^j T_k \psi^\ell; \quad j, k \in \mathbb{Z}, \quad \ell = 1, \dots, r \right\}.$$

The system is generated using the well-known unitary and oblique extension principles [42,43].

Theorem 0.2. [[42]] For a given refinable ϕ , the system $X(\Psi)$ is a tight frame for the space $L^2(\mathbb{R})$ if

$$\sum_{\ell=0}^r \sum_{k \in \mathbb{Z}} \overline{b^\ell[k]} b^\ell[k - q] = \delta_{0,q},$$

and

$$\sum_{\ell=0}^r \sum_{k \in \mathbb{Z}} (-1)^{k-q} \overline{b^\ell[k]} b^\ell[k - q] = 0,$$

hold for all $q \in \mathbb{Z}$, where $\delta_{0,q}$ is the Kronecker delta function. The oblique extension principle presented is a generalization of the unitary extension principle and to have such extension, it is require to have some required constraints on $a[k]$. That is, for some trigonometric function $\widehat{\Theta}$, for $\sigma \in \{0, 1\}$,

$$\begin{aligned} \widehat{\Theta}(\cdot) |\widehat{\phi}(0)|^2 &= 1, \\ \sum_{\xi \in E} \widehat{b}^\ell \left(\cdot / 2 + \pi\sigma \right) \overline{\widehat{b}^\ell \left(\xi / 2 \right)} &= \Theta \left(\xi / 2 \right) \delta_\sigma - \widehat{\Theta} \left(\xi \right) \widehat{a} \left(\xi / 2 + \pi\sigma \right). \end{aligned}$$

Note that, according to Definition 0.1 and Theorem 0.2, we have the following tight frame expansion,

$$g = \sum_{\ell=1}^r \sum_{j,k \in \mathbb{Z}} \langle g, \psi_{j,k}^\ell \rangle \psi_{j,k}^\ell. \tag{2.1}$$

Let us now construct some tight frame systems using the above principles. Define the function sgn as

$$\text{sgn}(x) = \frac{x}{|x|}, \quad x \neq 0.$$

Example 0.1. Applying the OEP and considering the B-spline of order two, B_2 , we have

Then, the system $X(\Psi)$ forms a tight frame for $L^2(\mathbb{R})$.

Example 0.2. For the B-spline of order four, B_4 , and again by applying the notion of the OEP we get

$$\begin{aligned} \psi_1(x) &= \frac{1}{24} \left(-4|x-2|^3 + 8|x-1|^3 - 8|x+1|^3 + 4|x+2|^3 + 3|2x+1|^3 - |2x+3|^3 - 3|1-2x|^3 + |3-2x|^3 \right), \\ \psi_2(x) &= \frac{1}{4\sqrt{6}} \left(2|x-2|^3 + 8|x-1|^3 - 20|x|^3 + 8|x+1|^3 + 2|x+2|^3 + |2x+1|^3 - |2x+3|^3 + |1-2x|^3 - |3-2x|^3 \right), \\ \psi_3(x) &= \frac{1}{24} \left(-4|x-2|^3 - 56|x-1|^3 + 56|x+1|^3 + 4|x+2|^3 - 7|2x+1|^3 - 3|2x+3|^3 + 7|1-2x|^3 + 3|3-2x|^3 \right) \\ \psi_4(x) &= \frac{-1}{12} \left(-|x-2|^3 - 28|x-1|^3 - 70|x|^3 - 28|x+1|^3 - |x+2|^3 + 7|2x+1|^3 + |2x+3|^3 + 7|1-2x|^3 + |3-2x|^3 \right). \end{aligned}$$

Then, the system $X(\Psi)$ forms a tight frame for $L^2(\mathbb{R})$.

Definition 0.3. For a real-valued function $u(t)$, suppose that $0 < q < 1$. Let Γ be the Euler gamma function, then:

- The Caputo's fractional order derivative is defined as follows,

$$\mathcal{D}^q u(t) = \frac{1}{\Gamma(1-q)} \int_0^t \frac{u'(x)}{(t-x)^q} dx.$$

- The associated Riemann–Liouville integral fractional operator is given by,

$$\mathcal{I}^q u(t) = \frac{1}{\Gamma(q)} \int_0^t \frac{u(x)}{(t-x)^{1-q}} dx.$$

Caputo fractional order mathematical model of AD

Now, we consider the mathematical model of AD using the Caputo fractional order derivative. The model consists of eighteen nonlinear partial differential equations. It is important to notice that to get a matched dimensions on both sides of the AD system we have to change the dimension of each parameter. Therefore, the order of the equations to q , and the dimension of the left-hand side would be $(time)^{-q}$.

$$\mathcal{D}^q A_\beta^i = \left(\lambda_\beta^i (1 + R) - d_{A_\beta^i} A_\beta^i \right) N / N_0 \tag{3.1}$$

$$\begin{aligned} \mathcal{D}^q A_\beta^0 &= A_\beta^i \left| \frac{\partial N}{\partial t} \right| + \frac{\lambda_N N}{N_0} + \frac{\lambda_A A}{A_0} - \left(d_{A_\beta^0 M} (\widehat{M}_1 + \widehat{M}_2 \theta) + d_{A_\beta^0 M} (M_1 \right. \\ &\quad \left. + M_2 \theta) \right) \frac{A_\beta^0}{A_\beta^0 + \bar{K}_{A_\beta^0}} \end{aligned} \tag{3.2}$$

$$\mathcal{D}^q \tau = \frac{N}{N_0} (\lambda_\tau + \lambda_\tau R - d_\tau \tau) \tag{3.3}$$

$$\mathcal{D}^q F_i = \frac{N}{N_0} (\lambda_F \tau - d_{F_i} F_i) \tag{3.4}$$

$$\mathcal{D}^q F_0 = F_i \left| \frac{\partial N}{\partial t} \right| - f_{F_0} F_0 \tag{3.5}$$

$$\mathcal{D}^q N = \frac{-d_{NF} F_i N}{F_i + K_{F_i}} - \frac{d_{NT} T_\alpha N}{(T_\alpha + K_{T_\alpha})(K_{I_{10}} + \gamma I_{10})} \tag{3.6}$$

$$\mathcal{D}^q A = \lambda_{AA_\beta^0} A_\beta^0 + \lambda_{AT_\alpha} T_\alpha - d_A A. \tag{3.7}$$

$$\mathcal{D}^q N_d = \frac{d_{NF} F_i N}{F_i + K_{F_i}} + \frac{d_{NT} T_\alpha N K_{I_{10}}}{(T_\alpha + K_{T_\alpha})(K_{I_{10}} + \gamma I_{10})} - d_{N_d M} (M_1 + M_2) \frac{N_d}{N_d + \bar{K}_{N_d}} \tag{3.8}$$

$$d_{N_d \widehat{M}} (\widehat{M}_1 + \widehat{M}_2) \frac{N_d}{N_d + \bar{K}_{N_d}} \tag{3.9}$$

$$\mathcal{D}^q A_0 = \lambda_{A_0} A_\beta^0 - d_{A_0} A_0 + D_{A_0} \Delta A_0 \tag{3.10}$$

$$\mathcal{D}^q H = \lambda_H N_d - d_H H + D_H \Delta H \tag{3.11}$$

$$\begin{aligned} \mathcal{D}^q M_1 &= \left(\frac{M_G^0 \lambda_{MF} F_0}{F_0 + K_{F_0}} + \frac{M_G^0 \lambda_{MA} A_0}{A_0 + K_{A_0}} \right) \frac{\beta_{\epsilon_1}}{\beta_{\epsilon_1} + \epsilon_2} - \nabla \cdot (M_1 \nabla H) \\ &\quad + \frac{T_\beta M_1 \lambda_{M_1 T_\beta}}{T_\beta + K_{T_\beta}} - d_{M_1} M_1 \end{aligned} \tag{3.12}$$

$$\begin{aligned} \mathcal{D}^q M_2 &= \left(\frac{M_G^0 \lambda_{MF} F_0}{F_0 + K_{F_0}} + \frac{M_G^0 \lambda_{MA} A_0}{A_0 + K_{A_0}} \right) \frac{\epsilon_2}{\beta_{\epsilon_1} + \epsilon_2} - \nabla \cdot (M_2 \nabla H) \\ &\quad + \frac{T_\beta M_2 \lambda_{M_2 T_\beta}}{T_\beta + K_{T_\beta}} - d_{M_2} M_2 \end{aligned} \tag{3.13}$$

$$\mathcal{D}^q \widehat{M}_1 = \alpha(P) (M_0 - \widehat{M}) \frac{\beta_{\epsilon_1}}{\beta_{\epsilon_1} + \epsilon_2} - \nabla \cdot (\widehat{M}_1 \nabla A_0) + \frac{T_\beta \widehat{M}_1 \lambda_{\widehat{M}_1 T_\beta}}{T_\beta + K_{T_\beta}} - d_{\widehat{M}_1} \widehat{M}_1 \tag{3.14}$$

Table 1
The variables description.

Variable	Description
A_β^i	The amyloid- β within neurons
A_β^0	The extracellular amyloid- β peptides outside neurons
τ	Hyperphosphorylated tau protein
$NFT(F_i)$	The neuronfibrillary tangle within neurons
$NFT(F_0)$	The neuronfibrillary tangle outside neurons
N	Live neurons
A	Astrocytes
$ROS(R)$	Reactive oxygen species
$APP(A_p)$	Amyloid precursor protein
$TNF-\alpha(T_\alpha)$	Tumor necrosis factor alpha
$IL-10(I_{10})$	Interleukin 10
$MG(M_G)$	Microglias
$GSK-3(G)$	Glycogen synthase kinase-type 3
$A\beta O(A_0)$	Amyloid β oligomer (soluble)
$TGF-\beta(T_\beta)$	Transforming growth factor beta
P	MCP-1
M_1	Proinflammatory microglias
\widehat{M}_1	Peripheral proinflammatory macrophages
M_2	Anti-inflammatory microglias
\widehat{M}_2	Peripheral anti-inflammatory macrophages
N_d	Dead neurons
H	High mobility group box 1 (HMGB1)

$$\mathcal{D}^q \widehat{M}_2 = \alpha \left(P \right) \left(M_0 - \widehat{M} \right) \frac{\epsilon_2}{\beta \epsilon_1 + \epsilon_2} - \nabla \cdot \left(\widehat{M}_2 \nabla A_o \right) + \frac{T_\beta \widehat{M}_2 \lambda_{M_2 T_\beta}}{T_\beta + K_{T_\beta}} - d_{M_2} \widehat{M}_2 \tag{3.15}$$

$$\mathcal{D}^q T_\beta = \lambda_{T_\beta M} M_2 + D_{T_\beta} \Delta T_\beta + \lambda_{T_\beta \widehat{M}_2} \widehat{M}_2 - d_{T_\beta} T_\beta \tag{3.16}$$

$$\mathcal{D}^q I_{10} = \lambda_{I_{10} M} M_2 + D_{I_{10}} \Delta T_\beta + \lambda_{I_{10} \widehat{M}_2} \widehat{M}_2 - d_{I_{10}} I_{10} \tag{3.17}$$

$$\mathcal{D}^q T_\alpha = \lambda_{T_\alpha M_1} M_1 + D_{T_\alpha} \Delta T_\alpha + \lambda_{T_\alpha \widehat{M}_1} \widehat{M}_1 - d_{T_\alpha} T_\alpha \tag{3.18}$$

$$\mathcal{D}^q P = \lambda_{PA} A + D_P \Delta P_{\lambda_{PM_2} M_2} - d_P P. \tag{3.19}$$

where

$$\begin{aligned} \epsilon_1 &= \frac{T_\alpha}{T_\alpha + K_\alpha}, \\ \epsilon_2 &= \frac{I_{10}}{I_{10} + K_{I_{10}}}, \\ \widehat{M} &= \widehat{M}_1 + \widehat{M}_2, \\ \alpha \left(P \right) &= \frac{\alpha P}{P + K_P}, \end{aligned}$$

and

$$R \left(t \right) = \begin{cases} 0.01 R_0 t, & 0 \leq t \leq 100 \\ R_0, & t > 100 \end{cases}$$

where the description of all variables is presented in Table 1. The AD model is subject to the periodic boundary conditions:

$$A_\beta^O(0, y, t) = A_\beta^O(1, y, t); \quad A_\beta^O(x, 0, t) = A_\beta^O(x, 1, t); \quad h(0, y, t) = h(1, y, t); \quad h(x, 0, t) = h(x, 1, t);$$

$$T_\beta(0, y, t) = T_\beta(1, y, t); \quad T_\beta(x, 0, t) = T_\beta(x, 1, t); \quad I_{10}(0, y, t) = I_{10}(1, y, t); \quad I_{10}(x, 0, t) = I_{10}(x, 1, t);$$

$$T_\alpha(0, y, t) = T_\alpha(1, y, t); \quad T_\alpha(x, 0, t) = T_\alpha(x, 1, t); \quad P(0, y, t) = P(1, y, t); \quad P(x, 0, t) = P(x, 1, t);$$

and the following initial conditions:

$$\begin{aligned} A_\beta \left(x, y, 0 \right) &= \frac{1}{10^6}; \quad A_\beta^O \left(x, y, 0 \right) = \frac{1}{10^8}; \quad \tau \left(x, y, 0 \right) \\ &= \frac{1.37}{10^{10}}; \quad F_i \left(x, y, 0 \right) = \frac{3.36}{10^{10}}; \quad F_o \left(x, y, 0 \right) = \frac{3.36}{10^{11}}; \end{aligned}$$

$$\begin{aligned} N \left(x, y, 0 \right) &= 0.14; \quad A \left(x, y, 0 \right) = 0.14; \quad M_1 \left(x, y, 0 \right) = 0.02; \quad M_2 \left(x, y, 0 \right) \\ &= 0.02; \quad \widehat{M}_1 \left(x, y, 0 \right) = 0; \end{aligned}$$

$$\begin{aligned} \widehat{M}_2 \left(x, y, 0 \right) &= 0, N_d \left(x, y, 0 \right) = 0; \quad H \left(x, y, 0 \right) = \frac{1.3}{10^{11}}; \quad T_\beta \left(x, y, 0 \right) \\ &= \frac{1}{10^6}; \quad T_\alpha \left(x, y, 0 \right) = \frac{2}{10^5}; \end{aligned}$$

$$I_{10} \left(x, y, 0 \right) = \frac{1}{10^5}; \quad P \left(x, y, 0 \right) = \frac{5}{10^9}; \quad A_o \left(x, y, 0 \right) = 0.$$

The parameters and their values are given by the following:

$$\begin{aligned} D_{A_o} &= (4.32 \times 10^{-2})^q; \quad D_H = (8.11 \times 10^{-2})^q; \quad D_{T_\alpha} \\ &= (6.55 \times 10^{-2})^q; \quad D_{T_\beta} = (6.55 \times 10^{-2})^q; \quad D_{I_{10}} = (6.04 \times 10^{-2})^q; \end{aligned}$$

$$\begin{aligned} D_P &= (1.2 \times 10^{-10})^q; \quad \lambda_{\beta}^i = (9.51 \times 10^{-6})^q; \quad \lambda_N = (8 \times 10^{-9})^q; \quad \lambda_A \\ &= 8 \times 10^{-9}; \quad \lambda_{\tau o} = (8.1 \times 10^{-11})^q; \end{aligned}$$

$$\begin{aligned} \lambda_\tau &= (1.35 \times 10^{-11})^q; \quad \lambda_F = (1.662 \times 10^{-3})^q; \quad \lambda_{A_{T_\alpha}} = (81.54)^q; \quad \lambda_{A_{A_\beta^O}} \\ &= (1.793)^q; \quad \lambda_{A_o} = (5 \times 10^{-2})^q; \end{aligned}$$

$$\begin{aligned} \lambda_H &= (3 \times 10^{-5})^q; \quad \lambda_{MF} = (2 \times 10^{-2})^q; \quad \lambda_{MA} = (2.3 \times 10^{-3})^q; \quad \lambda_{M_1 T_\beta} \\ &= (6 \times 10^{-3})^q; \quad \lambda_{M_1 T_\beta} = (6 \times 10^{-4})^q; \end{aligned}$$

$$\begin{aligned} \lambda_{T_\beta M} &= (1.5 \times 10^{-2})^q; \quad \lambda_{T_\beta \widehat{M}} = (1.5 \times 10^{-2})^q; \quad \lambda_{T_\beta M_1} \\ &= (3 \times 10^{-2})^q; \quad \lambda_{T_\beta M_1} = (3 \times 10^{-2})^q; \end{aligned}$$

$$\begin{aligned} \lambda_{I_{10} M_2} &= (6.67 \times 10^{-3})^q; \quad \lambda_{I_{10} \widehat{M}_2} = (6.67 \times 10^{-3})^q; \quad \lambda_{PA} \\ &= (6.6 \times 10^{-8})^q; \quad \lambda_{PM_2} = (1.32 \times 10^{-7})^q; \quad \theta = (0.9)^q; \end{aligned}$$

$$\begin{aligned} \alpha &= (5)^q; \quad \beta = (10)^q; \quad \gamma = 1; \quad d_{A_\beta^i} = (9.51)^q; \quad d_{A_\beta^O} = (9.51)^q; \quad d_{A_\beta^i M} \\ &= (2 \times 10^{-3})^q; \quad d_{A_\beta^i \widehat{M}} = (10^{-2})^q; \quad d_\tau = (0.77)^q; \end{aligned}$$

$$\begin{aligned} d_{F_i} &= (2.77 \times 10^{-3})^q; \quad d_{F_o} = (2.77 \times 10^{-4})^q; \quad d_N = (1.9 \times 10^{-4})^q; \quad d_{NF} \\ &= (3.4 \times 10^{-4})^q; \end{aligned}$$

$$\begin{aligned} d_{NT} &= (1.7 \times 10^{-4})^q; \quad d_{N_d M} = (0.06)^q; \quad d_{N_d \widehat{M}} = (0.06); \quad d_{AM} \\ &= (1.2 \times 10^{-3})^q; \quad d_{M_1} = (0.015)^q; \end{aligned}$$

$$\begin{aligned} d_{M_2} &= (0.015)^q; \quad d_{M_1} = (0.015)^q; \quad d_{M_2} = (0.015)^q; \quad d_{A_o} \\ &= (0.915)^q; \quad d_H = (58.71)^q; \quad d_{T_\alpha} = (55.45)^q; \end{aligned}$$

$$\begin{aligned} d_{T_\beta} &= (3.33 \times 10^2)^q; \quad d_{I_{10}} = (16.64)^q; \quad d_P = (1.73)^q; \quad R_0 = 6^q; \quad M_0 \\ &= (5 \times 10^{-2})^q; \quad N_0 = (0.14)^q; \quad M_G^0 = (0.047)^q; \end{aligned}$$

$$\begin{aligned} A_0 &= (0.14)^q; \quad \overline{K}_{A_\beta^O} = (7 \times 10^{-3})^q; \quad \overline{K}_{N_d} = (10 - 3)^q; \quad K_{I_{10}} \\ &= (2.5 \times 10^{-6})^q; \quad \overline{K}_{T_\beta} = (2.5 \times 10^{-7})^q; \quad K_M = (0.047)^q; \end{aligned}$$

$$\begin{aligned} K_{M_1} &= (0.047)^q; \quad K_{M_1} = (0.03)^q; \quad K_{M_1} = (0.017)^q; \quad K_{M_1} = (0.04)^q; \quad K_{M_2} \\ &= (0.007)^q; \end{aligned}$$

$$\begin{aligned} K_{F_i} &= (3.36 \times 10^{-10})^q; \quad K_{F_o} = (2.58 \times 10^{-11})^q; \quad K_{A_o} \\ &= (1 \times 10^{-7})^q; \quad K_P = (6 \times 10^{-9})^q; \quad K_{T_\alpha} = (4 \times 10^{-5})^q; \end{aligned}$$

Now by applying the Caputo fractional derivative for each variable of the AD system we get:

$$\begin{aligned} \frac{1}{\Gamma(1-q)} \int_0^t \frac{(A_\beta^i)'}{(t-x)^q} dx &= (\lambda_\beta^i(1+R) - d_{A_\beta^i} A_\beta^i) N / N_0 \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(A_\beta^0)'}{(t-x)^q} dx &= A_\beta^i \left| \frac{\partial N}{\partial t} \right| + \frac{\lambda_N N}{N_0} + \frac{\lambda_A A}{A_0} - \left(d_{A_\beta^0 M} (\widehat{M}_1 + \widehat{M}_2 \theta) + d_{A_\beta^0 M} (M_1 + M_2 \theta) \right) \frac{A_\beta^0}{A_\beta^0 + \overline{K}_{A_\beta^0}} \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{\tau'}{(t-x)^q} dx &= \frac{N}{N_0} (\lambda_{\tau_0} + \lambda_{\tau} R - d_{\tau} \tau) \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(F_i)'}{(t-x)^q} dx &= \frac{N}{N_0} (\lambda_F \tau - d_{F_i} F_i) \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(F_o)'}{(t-x)^q} dx &= F_i \left| \frac{\partial N}{\partial t} \right| - f_{F_o} F_o \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{N'}{(t-x)^q} dx &= \frac{-d_{NF} F_i N}{F_i + K_{F_i}} - \frac{d_{NT} T_\alpha N}{(T_\alpha + K_{T_\alpha})(K_{I_{10}} + \gamma I_{10})} \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{A'}{(t-x)^q} dx &= \lambda_{AA_\beta^0} A_\beta^0 + \lambda_{AT_\alpha} T_\alpha - d_A A, \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(N_d)'}{(t-x)^q} dx &= \frac{d_{NF} F_i N}{F_i + K_{F_i}} + \frac{d_{NT} T_\alpha N K_{I_{10}}}{(T_\alpha + K_{T_\alpha})(K_{I_{10}} + \gamma I_{10})} - d_{N_d M} (M_1 + M_2) \frac{N_d}{N_d + \overline{K}_{N_d}} \\ &\quad d_{N_d M} (\widehat{M}_1 + \widehat{M}_2) \frac{N_d}{N_d + \overline{K}_{N_d}} \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(A_o)'}{(t-x)^q} dx &= \lambda_{A_o} A_\beta^0 - d_{A_o} A_o + D_{A_o} \Delta A_o \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{H'}{(t-x)^q} dx &= \lambda_H N_d - d_H H + D_H \Delta H \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(M_1)'}{(t-x)^q} dx &= \left(\frac{M_G^0 \lambda_{MF} F_o}{F_o + K_{F_o}} + \frac{M_G^0 \lambda_{MA} A_o}{A_o + K_{A_o}} \right) \frac{\beta_{\in 1}}{\beta_{\in 1} + \epsilon_2} - \nabla \cdot (M_1 \nabla H) + \frac{T_\beta M_1 \lambda_{M_1 T_\beta}}{T_\beta + K_{T_\beta}} - d_{M_1} M_1 \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(M_2)'}{(t-x)^q} dx &= \left(\frac{M_G^0 \lambda_{MF} F_o}{F_o + K_{F_o}} + \frac{M_G^0 \lambda_{MA} A_o}{A_o + K_{A_o}} \right) \frac{\epsilon_2}{\beta_{\in 1} + \epsilon_2} - \nabla \cdot (M_2 \nabla H) + \frac{T_\beta M_2 \lambda_{M_2 T_\beta}}{T_\beta + K_{T_\beta}} - d_{M_2} M_2 \\ \frac{1}{\Gamma(1-q)} \int_0^t \frac{(\widehat{M}_1)'}{(t-x)^q} dx &= \alpha(P) \left(M_0 - \widehat{M} \right) \frac{\beta_{\in 1}}{\beta_{\in 1} + \epsilon_2} - \nabla \cdot (\widehat{M}_1 \nabla A_o) + \frac{T_\beta \widehat{M}_1 \lambda_{\widehat{M}_1 T_\beta}}{T_\beta + K_{T_\beta}} - d_{\widehat{M}_1} \widehat{M}_1 \end{aligned}$$

$$\frac{1}{\Gamma(1-q)} \int_0^t \frac{(\widehat{M}_2)'}{(t-x)^q} dx = \alpha(P) \left(M_0 - \widehat{M} \right) \frac{\epsilon_2}{\beta_{\in 1} + \epsilon_2} - \nabla \cdot (\widehat{M}_2 \nabla A_o)$$

$$+ \frac{T_\beta \widehat{M}_2 \lambda_{\widehat{M}_2 T_\beta}}{T_\beta + K_{T_\beta}} - d_{\widehat{M}_2} \widehat{M}_2$$

$$\frac{1}{\Gamma(1-q)} \int_0^t \frac{(T_\beta)'}{(t-x)^q} dx = \lambda_{T_\beta M} M_2 + D_{T_\beta} \Delta T_\beta + \lambda_{T_\beta M} \widehat{M}_2 - d_{T_\beta} T_\beta$$

$$\frac{1}{\Gamma(1-q)} \int_0^t \frac{(I_{10})'}{(t-x)^q} dx = \lambda_{I_{10} M} M_2 + D_{I_{10}} \Delta T_\beta + \lambda_{I_{10} M} \widehat{M}_2 - d_{I_{10}} I_{10}$$

$$\frac{1}{\Gamma(1-q)} \int_0^t \frac{(T_\alpha)'}{(t-x)^q} dx = \lambda_{T_\alpha M_1} M_1 + D_{T_\alpha} \Delta T_\alpha + \lambda_{T_\alpha M_1} \widehat{M}_1 - d_{T_\alpha} T_\alpha$$

$$\frac{1}{\Gamma(1-q)} \int_0^t \frac{P'}{(t-x)^q} dx = \lambda_{PA} A + D_P \Delta P_{\lambda_{PM_2} M_2} - d_P P.$$

Each variable in the AD system will be expanded with respect to t by using the truncated representation \mathcal{S}_n that extracted from Eq. 2.1, where

$$\mathcal{S}_n g = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} \langle g, \Psi_{j,k}^\ell \rangle \Psi_{j,k}^\ell = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_g \Psi_{j,k}^\ell.$$

Thus,

$$A_{\beta}^i \approx \mathcal{S}_n A_{\beta}^i = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{A_{\beta}^i} \Psi_{j,k}^\ell(t); A_{\beta}^0 \approx \mathcal{S}_n A_{\beta}^0 = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{A_{\beta}^0} \Psi_{j,k}^\ell(t),$$

$$\tau \approx \mathcal{S}_n \tau = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{\tau} \Psi_{j,k}^\ell(t); F_i \approx \mathcal{S}_n F_i = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{F_i} \Psi_{j,k}^\ell(t),$$

$$F_o \approx \mathcal{S}_n F_o = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{F_o} \Psi_{j,k}^\ell(t); N' \approx \mathcal{S}_n N = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_N \Psi_{j,k}^\ell(t),$$

$$A' \approx \mathcal{S}_n A = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_A \Psi_{j,k}^\ell(t); N_d' \approx \mathcal{S}_n N_d = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{N_d} \Psi_{j,k}^\ell(t),$$

$$A_o \approx \mathcal{S}_n A_o = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{A_o} \Psi_{j,k}^\ell(t); H' \approx \mathcal{S}_n H = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_H \Psi_{j,k}^\ell(t),$$

$$N_d' \approx \mathcal{S}_n N_d = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{N_d} \Psi_{j,k}^\ell(t); A_o' \approx \mathcal{S}_n A_o = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{A_o} \Psi_{j,k}^\ell(t),$$

$$H' \approx \mathcal{S}_n H = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_H \Psi_{j,k}^\ell(t); M_1 \approx \mathcal{S}_n M_1 = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{M_1} \Psi_{j,k}^\ell(t),$$

$$M_1' \approx \mathcal{S}_n M_1 = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{M_1} \Psi_{j,k}^\ell(t); M_2 \approx \mathcal{S}_n M_2 = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{M_2} \Psi_{j,k}^\ell(t);$$

$$\widehat{M}_1 \approx \mathcal{S}_n \widehat{M}_1 = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{\widehat{M}_1} \Psi_{j,k}^\ell(t); \widehat{M}_2 \approx \mathcal{S}_n \widehat{M}_2 = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{\widehat{M}_2} \Psi_{j,k}^\ell(t),$$

$$T'_\beta \approx \mathcal{S}_n T_\beta = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{T_\beta} \Psi_{j,k}^\ell(t); I_{10} \approx \mathcal{S}_n I_{10} = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{I_{10}} \Psi_{j,k}^\ell(t),$$

$$P' \approx \mathcal{S}_n P = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_P \Psi_{j,k}^\ell(t); T'_a \approx \mathcal{S}_n T_a = \sum_{\ell=1}^r \sum_{j<n} \sum_{k \in \mathbb{Z}} c_{T_a} \Psi_{j,k}^\ell(t).$$

Applying the method proposed in [44] reveals us with the following reduced system

$$A'_\beta(t, x, y) - A'_\beta(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_1(s, A'_\beta(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$A^0_\beta(t, x, y) - A^0_\beta(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_2(s, A^0_\beta(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$\tau(t, x, y) - \tau(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_3(s, \tau(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$F_i(t, x, y) - F_i(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_4(s, F_i(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$F_o(t, x, y) - F_o(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_5(s, F_o(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$N(t, x, y) - N(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_6(s, N(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$A(t, x, y) - A(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_7(s, A(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$N_d(t, x, y) - N_d(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_8(s, N_d(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$A_o(t, x, y) - A_o(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_9(s, A_o(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$H(t, x, y) - H(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{10}(s, H(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$M_1(t, x, y) - M_1(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{11}(s, M_1(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$M_2(t, x, y) - M_2(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{12}(s, M_2(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$\widehat{M}_1(t, x, y) - \widehat{M}_1(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{13}(s, \widehat{M}_1(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$\widehat{M}_2(t, x, y) - \widehat{M}_2(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{14}(s, \widehat{M}_2(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$T_\beta(t, x, y) - T_\beta(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{15}(s, T_\beta(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$I_{10}(t, x, y) - I_{10}(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{16}(s, I_{10}(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$T_a(t, x, y) - T_a(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{17}(s, T_a(s, x, y))}{(t-s)^{1-q}} ds = 0;$$

$$P(t, x, y) - P(0, x, y) - \frac{1}{\Gamma(q)} \int_0^t \frac{\mathcal{L}_{18}(s, P(s, x, y))}{(t-s)^{1-q}} ds = 0.$$

Using collocation technique, we discretize the domain (for t, x , and y) in h step size in order to obtain a sequence of mesh points as follows: for a framelet partial sum of size J , we set

$$M = 2^J, \quad h = 1/2M, \quad x_0 = 0, \quad x_{m+1} = x_m + mh, \quad m = 0, 1, \dots, 2M,$$

where the simulation is concluded within the following domain:

$$\Xi = \{(x, y) \in \mathbb{R}^2 : 0 \leq x, \quad y \leq 1\}.$$

Therefore,

$$A^i_\beta(t_m, x_i, y_j) - A^i_\beta(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_1(s, A^i_\beta(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$A^0_\beta(t_m, x_i, y_j) - A^0_\beta(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_2(s, A^0_\beta(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$\tau(t_m, x_i, y_j) - \tau(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_3(s, \tau(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$F_i(t_m, x_i, y_j) - F_i(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_4(s, F_i(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$F_o(t_m, x_i, y_j) - F_o(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_5(s, F_o(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$N(t_m, x_i, y_j) - N(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_6(s, N(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$A(t_m, x_i, y_j) - A(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_7(s, A(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$N_d(t_m, x_i, y_j) - N_d(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_8(s, N_d(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$A_o(t_m, x_i, y_j) - A_o(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_9(s, A_o(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$H(t_m, x_i, y_j) - H(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{10}(s, H(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$M_1(t_m, x_i, y_j) - M_1(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{11}(s, M_1(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$M_2(t_m, x_i, y_j) - M_2(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{12}(s, M_2(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$\widehat{M}_1(t_m, x_i, y_j) - \widehat{M}_1(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{13}(s, \widehat{M}_1(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$\widehat{M}_2(t_m, x_i, y_j) - \widehat{M}_2(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{14}(s, \widehat{M}_2(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$T_\beta(t_m, x_i, y_j) - T_\beta(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{15}(s, T_\beta(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$I_{10}(t_m, x_i, y_j) - I_{10}(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{16}(s, I_{10}(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$T_a(t_m, x_i, y_j) - T_a(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{17}(s, T_a(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0;$$

$$P(t_m, x_i, y_j) - P(0, x_i, y_j) - \frac{1}{\Gamma(q)} \int_0^{t_m} \frac{\mathcal{L}_{18}(s, P(s, x_i, y_j))}{(t_m-s)^{1-q}} ds = 0.$$

$$\frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_1(s_n, A^i_\beta(s_n, x_i, y_j))}{(t_m-s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_1(s_{n+1}, A^i_\beta(s_{n+1}, x_i, y_j))}{(t_m-s_{n+1})^{1-q}}.$$

Now, by approximating the integrals in the proposed model based on the composite trapezoidal rule, we have,

$$\begin{aligned}
 A_\beta^i(t_m, x_i, y_j) &= A_\beta^i(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_1(s_n, A_\beta^i(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_1(s_{n+1}, A_\beta^i(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 A_\beta^0(t_m, x_i, y_j) &= A_\beta^0(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_2(s_n, A_\beta^0(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_2(s_{n+1}, A_\beta^0(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 \tau(t_m, x_i, y_j) &= \tau(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_3(s_n, \tau(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_3(s_{n+1}, \tau(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 F_i(t_m, x_i, y_j) &= F_i(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_4(s_n, F_i(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_4(s_{n+1}, F_i(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 F_o(t_m, x_i, y_j) &= F_o(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_5(s_n, F_o(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_5(s_{n+1}, F_o(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 N(t_m, x_i, y_j) &= N(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_6(s_n, N(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_6(s_{n+1}, N(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 A(t_m, x_i, y_j) &= A(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_7(s_n, A(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_7(s_{n+1}, A(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 N_d(t_m, x_i, y_j) &= N_d(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_8(s_n, N_d(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_8(s_{n+1}, N_d(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 A_o(t_m, x_i, y_j) &= A_o(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_9(s_n, A_o(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_9(s_{n+1}, A_o(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 H(t_m, x_i, y_j) &= H(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{10}(s_n, H(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{10}(s_{n+1}, H(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 M_1(t_m, x_i, y_j) &= M_1(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{11}(s_n, M_1(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{11}(s_{n+1}, M_1(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}};
 \end{aligned}$$

$$\begin{aligned}
 M_2(t_m, x_i, y_j) &= M_2(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{12}(s_n, M_2(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{12}(s_{n+1}, M_2(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 \widehat{M}_1(t_m, x_i, y_j) &= \widehat{M}_1(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{13}(s_n, \widehat{M}_1(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{13}(s_{n+1}, \widehat{M}_1(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 \widehat{M}_2(t_m, x_i, y_j) &= \widehat{M}_2(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{14}(s_n, \widehat{M}_2(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{14}(s_{n+1}, \widehat{M}_2(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 T_\beta(t_m, x_i, y_j) &= T_\beta(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{15}(s_n, T_\beta(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{15}(s_{n+1}, T_\beta(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 I_{10}(t_m, x_i, y_j) &= I_{10}(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{16}(s_n, I_{10}(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{16}(s_{n+1}, I_{10}(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 T_\alpha(t_m, x_i, y_j) &= T_\alpha(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{17}(s_n, T_\alpha(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{17}(s_{n+1}, T_\alpha(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}; \\
 P(t_m, x_i, y_j) &= P(0, x_i, y_j) + \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{18}(s_n, P(s_n, x_i, y_j))}{(t_m - s_n)^{1-q}} - \frac{1}{\Gamma(q)} \sum_{n=0}^{2m-1} \frac{\mathcal{L}_{18}(s_{n+1}, P(s_{n+1}, x_i, y_j))}{(t_m - s_{n+1})^{1-q}}.
 \end{aligned}$$




Fig. 1. The graphs of generators ψ_1, ψ_2 in Example 0.1.

Thus, the AD system will be reduced to a matrix system that numerically can be easily solved to get the approximate solution for each variable (see Figs. 1 and 2).

By joining the variables and solving the resulting system, we present the simulation results of the PDE system of AD fractional model in Figs. 3–7. The plots represent the average density for all variables of the new model considering different values of fractional orders, namely $q = 0.7, q = 0.8$ and $q = 1$. In Fig. 8 we show the number of neurons for $q = 1, 0.8, 0.7$ depending on A_p^0, T_p and P .

Conclusion

In this paper, we investigated and constructed a new mathematical model of AD, that is initially formulated based on the classical ordinary integer derivative, by extending the base model to a fractional system of PDEs and by involving the Caputo fractional derivative. There are many advantages of using such derivative which resulted in better describing the dynamics of such disease. Thus, the fractional-order model provides better results than the integer-order model.

The new system is simulated by means of framelet approximation/expansion where an explicit form of the framelets (generators) of such method is given explicitly. The generators through out the work is

constructed using important set of non-negative functions, namely the B-splines, and based on the unitary and oblique extension principles. Using such approach is to provide us with an approximated simulation of the new mathematical AD model with high accuracy order, compactly supported framelets, high vanishing moments, and a proper properties of smoothness and symmetries.

We provided several graphical illustrations for all of the simulated variables to visualize their behavior for interpretation. For example, in Figs. 3,4,6,7 the results represent the average density for all variables of the new model using different values of fractional orders. It confirm that the variables just keep a consistent pace and so a steady state behavior with different values of fractional order. In Fig. 8, the results show that neurons are increasingly dying according to the fractional order values.

The obtained results provide us with a fundamental information to further better recognize the dynamics of AD. This may give some insights to researchers in future practical medical trails for many applications.

Compliance with ethics requirements

This article does not contain any studies with human or animal subjects.




Fig. 2. The graphs of generators ψ_1, \dots, ψ_4 in Example 0.2.




Fig. 3. The average density of some variables as indicated under each subfigure for different values of q .





Fig. 4. The average density of some variables as indicated under each subfigure for different values of q .





Fig. 5. The average density of some variables as indicated under each subfigure for different values of q .




Fig. 6. The average density of some variables as indicated under each subfigure for different values of q .







Fig. 7. The average density of some variables as indicated under each subfigure for different values of q .




(a) $T_\beta, q = 1$




(b) $T_\beta, q = 0.7$




(c) $T_\beta, q = 0.8$



(d) $A_\beta^0, q = 1$



(e) $A_\beta^0, q = 0.7$



(f) $A_\beta^0, q = 0.8$

Fig. 8. The number of neurons when $q = 1, 0.8, 0.7$ depending on A_β^0, T_β and P .

CRedit authorship contribution statement

Mutaz Mohammad: Conceptualization, Methodology, Software, Investigation, Supervision, Validation, Writing - review & editing.
Alexander Trounev: Software.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Zhao Q, Tang X. Effects of huperzine A on an acetylcholinesterase isoforms in vitro: comparison with tacrine, donepezil, rivastigmine and physostigmine. *Eur J Pharmacol* 2002;455:101–7.
- [2] Janus C, Westaway D. Transgenic mouse models of Alzheimer's disease. *Physiol Behav* 2001;73(5):873–86.
- [3] Martin Prince A. World Alzheimer Report 2015: the global impact of dementia: an analysis of prevalence, incidence, cost and trends. *Alzheimer's Disease Int* 2015.
- [4] Brookmeyer R. Forecasting the global burden of Alzheimer's disease. *Alzheimer's Dementia* 2007;3(3):186–91.
- [5] Alzheimer's Association, 2019 Alzheimer's disease facts and figures, *Alzheimer's Dementia* 2019; 15: 321–387.
- [6] Baleanu D, Jajarmi A, Asad J. Classical and fractional aspects of two coupled pendulums. *Rom Rep Phys* 2019;71:103.
- [7] Hajipour M, Jajarmi A, Baleanu D, Sun HG. On an accurate discretization of a variable-order fractional reaction–diffusion equation. *Commun Nonlinear Sci* 2018;69:119–33.
- [8] Goullart A, Lazo M, Suarez J. A new parameterization for the concentration flux using the fractional calculus to model the dispersion of contaminants in the planetary boundary layer. *Physica A* 2019;518:38–49.
- [9] Chang A, Sun H, Zhang Y, Zheng C, Min F. Spatial fractional Darcy's law to quantify fluid flow in natural reservoirs. *Physica A* 2019;519:119–26.
- [10] Dubey V, Kumar R, Kumar D. Analytical study of fractional Bratu-type equation arising in electro-spun organic nanofibers elaboration. *Physica A* 2019;521: 762–72.
- [11] Abdulhameed M, Muhammad M, Gital A, Yakubu D, Khan I. Effect of fractional derivatives on transient MHD flow and radiative heat transfer in a micro-parallel channel at high zeta potentials. *Physica A* 2019;519:42–71.
- [12] Altaf K, Atangana A. Dynamics of ebola disease in the framework of different fractional derivatives. *Entropy* 2019;21(3):303.
- [13] Ullah S, Khan M, Gomez-Aguilar JF. Mathematical formulation of hepatitis B virus with optimal control analysis. *Optim Control Appl Methods* 2019;40(4):529–44.
- [14] Abro KA, Khan I, Nisar KS. Use of Atangana-Baleanu fractional derivative in helical flow of a circular pipe. *Fractals* 2020;28:2040049.
- [15] Mohammad M, Lin EB. Gibbs phenomenon in tight framelet expansions. *Commun Nonlinear Sci Numer Simul* 2018;55:84–92.
- [17] Mohammad M. Special B-spline tight framelet and it's applications. *J Adv Math Comput Sci* 2018;29:1–18.
- [18] Mohammad M. On the Gibbs effect based on the quasi-affine dual tight framelets system generated using the mixed oblique extension principle. *Mathematics* 2019; 7.
- [19] Mohammad M, Howari F, Acbas G, Nazzal Y, AlAydarooos F. Wavelets based simulation and visualization approach for unmixing of hyperspectral data. *Int J Earth Environ Sci* 2018;3:152.
- [20] Mohammad M. Biorthogonal-wavelet-based method for numerical solution of Volterra integral equations. *Entropy* 2019;21:1098.
- [21] Mohammad M. A numerical solution of Fredholm integral equations of the second kind based on tight framelets generated by the oblique extension principle. *Symmetry* 2019;11:854.
- [22] Mohammad M, Cattani C. A collocation method via the quasi-affine biorthogonal systems for solving weakly singular type of Volterra-Fredholm integral equations. *Alex Eng J* 2020;59. <https://doi.org/10.1016/j.aej.2020.01.046>.
- [23] Mohammad M. Bi-orthogonal wavelets for investigating Gibbs effects via oblique extension principle, 2020. *J Phys Conf Ser* 2020;1489. <https://doi.org/10.1142/S0218348X20400514>.
- [24] Mohammad M, Cattani C. Applications of bi-framelet systems for solving fractional order differential equations. *Fractals* 2020; 28. doi:10.1142/S0218348X20400514.
- [25] Mohammad M, Trounev A. Implicit Riesz wavelets based-method for solving singular fractional integro-differential equations with applications to hematopoietic stem cell modeling. *Chaos Solitons Fract* 2020;138. <https://doi.org/10.1016/j.chaos.2020.109991>.
- [26] Mohammad M, Trounev A, Cattani C. The dynamics of COVID-19 in the UAE based on fractional derivative modeling using Riesz wavelets simulation. DOI: 10.21203/rs.3.rs-33366/v1.
- [27] Mohammad M, Trounev A, Cattani C. An efficient method based on framelets for solving fractional Volterra integral Eqs. 2020; 22. DOI: 10.21203/rs.3.rs-33366/v1.
- [28] Mohammad M, Trounev A. On the dynamical modeling of Covid-19 involving Atangana-Baleanu fractional derivative and based on Daubechies framelet simulations. *Chaos Solitons Fract* 2020;140. <https://doi.org/10.1016/j.chaos.2020.110171>.
- [29] Mohammad M, Trounev A. Fractional nonlinear Volterra-Fredholm integral equations involving Atangana-Baleanu fractional derivative: framelet applications. *Adv Differ Equ*.
- [30] Ghanbari B, Atangana A. A new application of fractional Atangana-Baleanu derivatives: designing ABC-fractional masks in image processing. *Physica A: Stat Mech Appl* 2020;542:123516.
- [31] Khan MA, Atangana A. Modeling the dynamics of novel coronavirus (2019-nCoV) with fractional derivative 2020; 59(4): 2379–2389.
- [32] Atangana A, Bonyah E, Elsadany A. A fractional order optimal 4D chaotic financial model with Mittag-Leffler law. *Chin J Phys* 2020;65:38–53.
- [33] Atangana A, Aguilar J, Kolade M, Hristov J. Fractional differential and integral operators with non-singular and non-local kernel with application to nonlinear dynamical systems. *Chaos Solitons Fract* 2020;132:109493. <https://doi.org/10.1016/j.chaos.2019.109493>.
- [34] Atangana A, Baleanu D. New fractional derivatives with nonlocal and non-singular kernel: theory and application to heat transfer model. *Therm Sci* 2016;20:763–9.
- [35] Atangana A, Koca I. Chaos in a simple nonlinear system with Atangana-Baleanu derivatives with fractional order. *Chaos Solitons Fract* 2016;89:447–54.
- [36] Atangana A. On the new fractional derivative and application to nonlinear Fisher's reaction-diffusion equation. *Appl Math Comput* 2016;273:948–56.
- [37] Atangana A, Aguilar J. Decolonisation of fractional calculus rules: breaking commutativity and associativity to capture more natural phenomena. *Eur Phys J Plus* 2018;133:166.
- [38] Carvalho A, Pinto C. Non-integer order analysis of the impact of diabetes and resistant strains in a model for TB infection. *Commun Nonlinear Sci Numer Simulat* 2018;61:104–26.
- [39] Sweilam N, AL-Mekhlafi S, Hassan AN. Numerical treatment for solving the fractional two-Group Influenza model, *Progr Fract Differ Appl* 2018; 4: 1–15.
- [40] Kumar S, Ghosh S, Lotayif M, Samet B. A model for describing the velocity of a particle in Brownian motion by Robotnov function based fractional operator. *Alex Eng J* 2020. <https://doi.org/10.1016/j.aej.2020.04.019>.
- [41] Hao W, Friedman A. Mathematical model on Alzheimer's disease. *BMC Syst Biol* 2016;10(108). <https://doi.org/10.1186/s12918-016-0348-2>.
- [42] Daubechies I, Han B, Ron A, Shen Z. Framelets: MRA-based constructions of wavelet frames. *Appl Comput Harmon Anal* 2003;14:1–46.
- [43] Han B. Framelets and wavelets: algorithms, analysis, and applications. In: *Applied and numerical harmonic analysis*. Birkhauser/Springer: Cham; 2017.
- [44] Toufik M, Atangana A. New numerical approximation of fractional derivative with non-local and non-singular kernel: application to chaotic models. *Eur Phys J Plus* 2017;132(10):444.

Further reading

- [16] Mohammad, M, Lin, EB. Gibbs effects using Daubechies and Coiflet tight framelet systems, *contemporary mathematics*. AMS 2018; 706: 271–282.