



Factors controlling the migration phenology of Siberian *Phylloscopus* species

László Bozó¹ · Tibor Csörgő² · Wieland Heim³

Received: 23 March 2020 / Revised: 18 June 2020 / Accepted: 31 July 2020 / Published online: 11 August 2020
© The Author(s) 2020

Abstract

The timing of annual events in migratory species is determined by the position of breeding and wintering grounds, length of the breeding season, the occurrence of their preferred diet or the timing of moult in birds. Previous studies found significant differences in the migration phenology of Asian *Phylloscopus* warbler species, a group of long-distance migratory passerines. However, the factors that influence the observed differences in migration phenology have not been studied yet. Here, we analyze the effect of the above-mentioned factors on the migration timing of seven *Phylloscopus* species at a stopover site in the Russian Far East. We found that spring migration phenology was best explained by moult strategy and the preferred prey size, while autumn migration phenology was linked to the southernmost wintering latitude of the species. Interspecific differences in migration timing were much higher in autumn than in spring. The duration of the species-specific migration periods was also longer in autumn than in spring, most likely caused by higher competition during spring to arrive early at the breeding grounds. Our results contribute to the understanding of migration ecology in songbirds moving along the little-studied East Asian flyway.

Keywords Leaf warbler · Migration strategies · East Asian flyway · Diet

Zusammenfassung

Welche Faktoren bestimmen die Zugphänologie sibirischer Laubsänger?

Der Jahreszyklus ziehender Arten wird von der Lage der Brut- und Überwinterungsgebiete, der Länge der Brutsaison, der Verfügbarkeit bevorzugter Nahrung oder dem Zeitraum der Mauser bestimmt. Vorhergehende Studien fanden signifikante Unterschiede in der Zugphänologie von asiatischen Laubsängern, einer Gruppe von Langstreckenziehern. Die Ursachen für die beobachteten Unterschiede wurden bisher jedoch noch nicht untersucht. Hier analysieren wir den Einfluss der oben genannten Faktoren auf das zeitliche Auftreten von sieben Laubsänger-Arten an einem Zwischenrastplatz im Fernen Osten Russlands. Im Frühjahr konnten die Unterschiede in der Phänologie der Arten am besten mit der Mauserstrategie und der Größe der bevorzugten Nahrung erklärt werden, während die Phänologie des Herbstzuges mit dem südlichsten Breitengrad des Überwinterungsgebietes erklärt werden konnte. Die zeitlichen Unterschiede zwischen den Arten waren im Herbst größer als während des Frühjahrszuges. Die artspezifischen Zugperioden waren im Herbst ebenfalls länger als im Frühjahr. Dies kann mit der Konkurrenz um ein möglichst zeitiges Ankommen in den Brutgebieten erklärt werden. Unsere Ergebnisse tragen zu einem besseren Verständnis der Zugökologie von Singvögeln des wenig untersuchten ostasiatischen Zugweges bei.

Communicated by N. Chernetsov.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10336-020-01805-5>) contains supplementary material, which is available to authorized users.

✉ László Bozó
bozolaszlo91@gmail.com

¹ Department of Systematic Zoology and Ecology,
Eötvös Loránd University, Pázmány Péter sétány 1/C,
1117 Budapest, Hungary

² Department of Anatomy, Cell- and Developmental Biology,
Eötvös Loránd University, Pázmány Péter sétány 1/C,
1117 Budapest, Hungary

³ Institute of Landscape Ecology, University of Münster,
Heisenbergstraße 2, 48149 Münster, Germany

Introduction

Migratory birds connect continents and different bioms during their long journey twice each year (Hahn et al. 2009). An estimated four billion migratory birds, the majority of which are passerines, migrate annually from temperate Eurasia to lower latitudes in Africa and Asia during the boreal winter (Newton 2007).

The phenology of migration relates to repeating cycles of life-history stages, such as migration, moult and breeding (Piersma and Ramenofsky 1998; Piersma and Drent 2003). Moult is costly in terms of energy (Lindström et al. 1993), and there is a strong correlation between the termination of moult and the onset of migration (Pulido and Coppack 2004). Therefore, moult strategy is a key regulator of the autumn migration. Among Palaearctic migratory songbirds, about half of the long-distance migrants moult directly after breeding (postnuptial moult) while the other half moult on the wintering grounds (prenuptial moult) (Barta et al. 2008). The different moult strategies appear to depend on the distance between the breeding and wintering grounds (Ginn and Melville 1983; Jenni and Winkler 2020). Phylogenetical analyses of moult strategies of warblers indicated, that species with longer migration routes tend to moult on the wintering grounds (Svensson and Hedenström 1999).

Another important factor for differences in phenology is the migration distance a species has to cover between breeding and wintering grounds (Marchetti et al. 1995). Species with long migration distances tend to migrate earlier in autumn and later in spring compared to short-distance migrants (Jenni and Kéry 2003).

Besides moult and migration distance, wintering behaviour shapes migration and breeding phenology (Marra et al. 1998). Conditions during winter also affect arrival schedules and the birds' condition in the breeding season, as well as the birds' reproductive success (Verboven and Visser 1998; Marra et al. 1998). The relationship between arrival date on the breeding grounds and the expected reproductive success can strongly influence the migratory behavior of individuals (Weber et al. 1998, 1999). In American Redstarts (*Setophaga ruticilla*), birds wintering in higher-quality habitats were found to depart earlier from the wintering grounds because of their better body conditions (Marra and Holberton 1998; Marra et al. 1998; Studds and Marra 2005). Furthermore, the timing of the onset of autumn migration also depends on the number of broods (Jenni and Kéry 2003) and the length of the breeding season (Ellegren 1990; Fransson 1995).

On top of that, migration phenology might be shaped by the availability of the preferred prey. Katti and Price (2003) found that the size of the preferred prey shapes the distribution of leaf warblers in breeding and wintering grounds in India: smaller species would return earlier to the breeding

grounds than larger taxa because their smaller prey becomes available sooner.

Despite the fact that the East Asian migratory flyway is the largest of all other flyways and holds the highest diversity of migratory songbirds, we have only very limited information about the migration ecology and phenology for most of the species (Yong et al. 2015). Short-term but standardized bird-ringing activities have resulted in a better understanding of feeding habits, habitat use, and migration strategies in several groups of Asian passerines, for example small thrushes (Wang et al. 2006; Maslovsky et al. 2018), and buntings (Nam et al. 2011; Heim et al. 2018). However, in the last two decades the most studied group in Asia were the leaf warblers. We have detailed information on breeding ecology (e.g., Price and Jamdar 1991; Bourski and Forstmeier 2000; Forstmeier et al. 2001a; Forstmeier 2002; Forstmeier and Balsby 2002), habitat use, morphology and foraging strategies (Price 1991; Forstmeier et al. 2001b; Katti and Price 2003; Batova 2011; Bozó et al. 2018a), the sex-specific timing of migration (Bozó and Heim 2016), the effect of weather variables on the migration (Bozó et al. 2018b) and the estimated migration distances (Sander et al. 2017; Bozó et al. 2019) of these species. However, the factors that influence the observed differences in migration phenology have not been studied yet.

In this study, we investigated factors possibly shaping the migration phenology of a set of leaf warbler species at a stop-over site in the Russian Far East. Since the timing of moult determines the migration timing of birds, we assume that (1) species with a postnuptial moult strategy start their migration later in autumn than species exhibiting prenuptial moult. We also tested (2) the effect of the geographical position of species-specific wintering grounds on the migration phenology. We hypothesize that (3) breeding biology influences the migration timing of the study species: species with an early breeding season would migrate earlier in autumn than species with a late breeding season (cf. Jenni and Kéry 2003). Furthermore, we test (4) whether the size of the preferred prey would explain the observed migration phenology.

Material and methods

The study was carried out within the Amur Bird Project (Heim and Smirenski 2013, 2017) at Muraviovka Park along the middle reaches of the Amur River in the Russian Far East. The study site is located 60 km southeast of the city of Blagoveshchensk (49°55'08,27"N, 127°40'19,93"E). Birds were trapped with mist-nets using a constant-effort-site approach in a variety of different habitat types (for details see Heim et al. 2018). We included the following species: Yellow-browed Warbler *Phylloscopus inornatus*, Arctic Warbler *P. borealis*, Two-barred Warbler *P. plumbeitarsus*,

Pale-legged Leaf Warbler *P. tenellipes*, Dusky Warbler *P. fuscatus*, Radde's Warbler *P. schwarzi* and Pallas's Leaf Warbler *P. proregulus*. A total of 9211 trapped individuals of all species combined were available for analysis (Table 1).

Two of the study species, Dusky Warbler and Pale-legged Leaf Warbler, are not only migrants but breed at the study site (Heim 2014). Therefore, mainly in case of the more common Dusky Warbler, it is not possible to separate local and transient individuals, and the median day of occurrence might be biased by local breeding birds. However, both species start to breed from late May-early June and finish breeding in late July in Southeast Russia (Clement 2006a, b), so we assume that the overlap between migration and breeding periods are negligible. Furthermore, the overall number of possible breeders is very small compared to the large numbers of migrating individuals. In addition, we excluded all individuals that were recaptured during the breeding season.

For the analysis of phenology, we included birds captured in mist-nets during spring (25th April to 10th June in 2013, 2015–2017) and autumn (1st August to 20th October in 2013, 2014, 2015 and 2017) migration. We defined the length of the species-specific migration periods as the number of days between the first and the last capture.

We used Kruskal–Wallis tests to analyze differences in the timing of migration between the study species, and paired *t* tests for differences between the length of spring and autumn migration period. To test the impact of different variables on migration timing, we used generalized linear mixed-effect models (GLMMs) within the R package lme4 (Bates et al. 2007). The following factors were used to explain the dependent variable *migration day* for spring and autumn: southernmost wintering latitude, end of the breeding period, moulting strategy (pre-nuptial or post-nuptial) and the average size of the prey. The selected fixed factors were not strongly correlated ($R < 0.5$). *Species* and *year* were fitted as random factors. Significant variables were selected with the help of “backward stepwise model selection”

(Crawley 2013) using the Likelihood-ratio test ($p < 0.05$). Information about the distribution of the study species was gathered from the IUCN range maps (BirdLife International 2019). Data regarding diet and breeding were taken from the following sources: Price (1991); Glutz von Blotzheim and Bauer (1991); Forstmeier and Kessler (2001); Forstmeier et al. (2001b); Katti and Price (2003); del Hoyo et al. (2006); Batova (2011). The average timing of the beginning and end of the nesting period was gathered from the literature for the entire breeding area, as very few data are available from different geographical regions. Mean prey size was calculated by the typical size of the prey based on information from the literature. In many cases, only higher taxa and groups were specified, so these prey sizes represent rough estimates (see supplementary material 2). No detailed information on diet was available for the Pale-legged Leaf Warbler, but based on descriptions in the literature we estimated the average prey size at 10 mm. Two of the seven study species (Arctic Warbler and Two-barred Warbler) have complete pre-nuptial moult, while the other species have complete post-nuptial moult (Svensson 1992; Brazil 2009; Demongin 2016). All data are given in Table 2.

All statistical analyses were carried out using the program R version 3.6.2 (R Core Team 2019).

Results

We found significant differences in the timing of migration among the seven studied leaf warbler species both in spring (Kruskal–Wallis test, $H = 1151$, $p < 0.0001$) and in autumn (Kruskal–Wallis test, $H = 1202$, $p < 0.0001$) (Fig. 1). Spring migration begins with Pallas's Leaf Warbler and Yellow-browed Warbler as the earliest species to arrive at the study site. The species migrating last are Two-barred Warbler and Arctic Warbler. Autumn migration starts in early August for

Table 1 The number of trapped individuals, the median migration dates and the length of the migration periods of the seven study species

Species	Spring		Autumn		Median date		Length of migration period (days)	
	Standard	Non-standard	Standard	Non-standard	Spring	Autumn	Spring	Autumn
<i>P. inornatus</i>	2048	338	1718	603	14 May	7 Sep	116	158
<i>P. borealis</i>	202	38	183	47	27 May	23 Aug	138	160
<i>P. plumbeitarsus</i>	57	8	114	33	31 May	27 Aug	139	161
<i>P. tenellipes</i>	30	11	146	45	19 May	15 Aug	137	158
<i>P. fuscatus</i>	749	154	1328	522	21 May	7 Sep	119	160
<i>P. schwarzi</i>	96	7	191	62	25 May	11 Sep	129	161
<i>P. proregulus</i>	72	11	276	122	22 May	25 Sep	119	157

Numbers are shown separately for birds trapped in standard mist-nets (used in all study years) and for birds trapped in non-standard nets (not used in all study years)

Table 2 Potential factors shaping the migration phenology of East Asian warbler species, including data on distribution, migration distance, breeding biology and diet, start and end of breeding = number of decades from 1st May to 31 August (1–12)

Variable	<i>inornatus</i>	<i>borealis</i>	<i>plumbeitarsus</i>	<i>proregulus</i>	<i>fuscatus</i>	<i>schwarzi</i>	<i>tenellipes</i>
Northernmost breeding latitude	70	72	62	61	66	58	54
Southernmost breeding latitude	40	42	41	43	28	38	37
Northernmost wintering latitude	27	18	24	32	33	23	22
Southernmost wintering latitude	2	– 11	9	18	2	8	3
Migration distance (°latitudes)	40.5	53.5	35	27	29.5	32.5	33
Number of broods	1	1	1	2	1	1	1
Start of breeding	4	4	3	4	3	4	3
End of breeding	10	9	11	9	11	10	9
Clutch size	3	6	6	5	5	4	5
Incubation + nestling phase (days)	25	26	26	26	26	27	22
Winter moult	0	1	1	0	0	0	0
Average prey size (mm)	5.4	10	5.6	5.3	11.8	20.6	10

Furthermore, the moult strategy for the studied species is given (0 = complete summer moult, 1 = complete winter moult)

Pale-legged Leaf Warblers and Arctic Warblers and ends in late September and early October with Pallas's Leaf Warbler as the latest migrating species (Fig. 1). The duration of spring migration was significantly shorter than autumn migration ($t = -3.585$, $p = 0.004$).

Our GLMMs revealed that autumn migration phenology is explained best by the southernmost wintering latitude, while spring migration phenology was explained by moult strategy and prey size (Table 3, supplementary material 1).

Discussion

We found significant differences in the migration phenology among a set of East Asian *Phylloscopus* species. Their migration phenology at the study site was found to be related to moult and the size of their preferred prey in spring, and the position of their southernmost wintering areas in autumn.

While most studies found an effect of moult on the timing of autumn migration (Owen and Krohn 1973; Kjellén 1994; Pérez-Tris et al. 2001; Pulido and Coppack 2004), we found that spring migration was related to moult strategy in our study species. In migratory birds, there is a strong correlation between the termination of moult and the onset of migration (Pulido and Coppack 2004). Summer moult occurs when food has a high peak after breeding, while winter moult occurs if there is a short period of high food availability in summer and a strong winter peak at different locations (Barta et al. 2008). Long-distance migrant species usually moult their flight feathers in the wintering grounds (in the beginning or in the end of the winter season; Salewski et al. 2004), or at stopover sites, including some *Locustella* and *Acrocephalus* species (Hedenström et al. 1993; Neto et al. 2006), as in the short Palearctic summer they have no time left between breeding and the preparation for migration (Barta et al. 2008). As moulting is costly in terms of

Table 3 Effects of southernmost wintering latitude, end of the breeding season, moult strategy and average prey size on the migration phenology of seven warbler species in spring and autumn

Dependent variable	Independent variable	Chi square	p	R^2 marg	R^2 cond
Median migr. day (spring)				0.114	0.139
	Southernmost wintering latitude	2.802	0.094		
	End of breeding	0.127	0.721		
	Moult	9.631	0.002		
Median migr. day (autumn)	Average prey size (mm)	5.182	0.023		
				0.085	0.302
	Southernmost wintering latitude	3.907	0.048		
	End of breeding	0.030	0.862		
	Moult	0.571	0.450		
	Average prey size (mm)	0.002	0.968		

Year and species were fitted as random factors. Categories of variables are the same defined in Table 2. Shown are Chi square, p value as well as marginal and conditional R^2 of GLMMs




Fig. 1 Migration phenology of the seven studied warbler species during spring and autumn. The date is given as the number of the day of the year (114 = 23th April, 162 = 10th June, 200 = 18th July, 290 = 16th October)

energy (Lindström et al. 1993), a pre-breeding moult would negatively affect preparation for migration in species that have to migrate longer distances. Spring migration might be delayed in species moulting in winter, as species replacing flight feathers during winter may delay departure compared to species that moult in summer (Rubolini et al. 2005). This could be the case for Arctic and Two-barred Warblers in our study.

Furthermore, we found an interesting relationship between spring migration phenology and mean prey size. Prey size and abundance have been found to be the main determinants of the spring arrival time of *Phylloscopus* warblers in India as well (Katti and Price 2003). In general, larger-bodied animals are typically able to handle larger prey (McNab 1971), which is true for *Phylloscopus* warblers as well (Price 1991; Gross and Price 2000). In boreal forests, the number of insects related to coniferous trees is much higher than those related to deciduous trees, and insects on coniferous trees appear much earlier than on deciduous trees, as the latter develop leaves later in the spring (Southwood 1961). In the temperate zone, therefore, several species of Passerines (tits, Goldcrests *Regulus regulus*) overwinter in pines, as small insects are available there even in winter (Haraszthy 1998; del Hoyo et al. 2006). The smallest and most typical taiga species (Pallas's Leaf Warbler, Yellow-browed Warbler) migrated earlier in spring than the other species in our study, probably because their small prey becomes available earlier.

Autumn migration phenology was related to the location of the wintering areas in our study. The most southerly wintering study species were the earliest to arrive in autumn, which highlights the role of the distance to the wintering grounds. The longest distances to their wintering grounds are covered by Arctic Warblers and Two-barred Warblers,

which might be the cause for the early autumn migration at the study site. Pallas's Leaf Warblers have a rather short distance to cover between the breeding and wintering areas, which may explain its late migration in autumn (del Hoyo et al. 2006). Consequently, the opposite pattern was found in spring, with species migrating shorter distances arriving earlier than those species with wintering grounds further south. Similar patterns are known from songbirds in other flyway systems (Alerstam 1993). Even within species, northern populations are known to migrate longer distances and to occur later on stop-over sites, e.g., in Eurasian Blackcap *Sylvia atricapilla* (e.g., Lövei et al. 1985), Garden Warbler *Sylvia borin* (Berthold 1988), Common Chiffchaff *P. collybita* (Gaston 1974; Tiainen and Hanski 1985) and Willow Warbler *P. trochilus* (Hedenström and Pettersson 1987).

Despite the fact that departure in autumn depends on the length of the breeding season (Ellegren 1990; Fransson 1995), we found no effect of the end of the breeding period on the timing of migration in our study. This may be due to the fact that the studied species typically have only one brood annually due to the short time available for hatching (del Hoyo et al. 2006), so there might be no significant variation among the species.

Furthermore, we found that migration periods are longer in autumn than in spring for all of our study species. This is most likely caused by the fact, that competition is high to arrive early at the breeding grounds in spring and, therefore, the migration is faster in spring than in autumn (Karlsson et al. 2012; Nilsson et al. 2013). We found the smallest species (Pallas's Leaf Warbler and Yellow-browed Warbler) to arrive earliest in spring. Gaston (1974) hypothesized that small *Phylloscopus* species breed earlier than larger species, and by wintering farther north they can rapidly return to the breeding grounds. This might be true for the species

in our study as well, although no obvious differences in the start of breeding are evident (Table 1). Some early breeding, small species make very long migratory journeys (to Northern Siberia from Southeast Asia) (Gaston 1974), suggesting that wintering far away does not necessarily prevent an early return to breeding grounds (Katti and Price 2003).

Conclusion

We identified moult strategy and the preferred prey size as the most important factors shaping spring phenology, while the position of the wintering areas was related to autumn migration phenology in Siberian leaf warblers. The migration periods of our study species were significantly shorter in spring than in autumn. These results are in line with findings from other flyway systems. Our work contributes to a better understanding of the causes and consequences of the migration of little-known East Asian songbird species.

Acknowledgements Open access funding provided by Eötvös Loránd University. The authors want to thank all volunteers and sponsors of the Amur Bird Project, as well as Sergei M. Smirenski and the staff of Muraviovka Park. We would like to thank Nikolett Olajos for improving the language of the manuscript. The experiments comply with the current laws of Russia.

Author contributions LB and WH designed the project. LB and WH conducted field work and data collection. WH and LB analysed the data and wrote the manuscript. LB, WH and TCS interpreted the results. LB, WH and TCS edited the manuscript. All authors read and approved the last version of the manuscript.

Funding Not applicable.

Code availability Not applicable.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Availability of data and material The datasets analysed during the current study are available from the corresponding author on reasonable request.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alerstam T (1993) Bird migration. Cambridge University Press, Cambridge
- Barta Z, McNamara JM, Houston AI, Weber TP, Hedenström A, Feró O (2008) Optimal moult strategies in migratory birds. *Philos T R Soc B* 363:211–229
- Bates D, Sarkar D, Bates MD, Matrix L (2007) The lme4 package. R package version 2(1):74
- Batova ON (2011) Selection of foraging tactics in leaf warblers (*Phylloscopus*). *Biol Bull* 38:259–265
- Berthold P (1988) The control of migration in European warblers. In: proceedings of XIX. International ornithological congress. University of Ottawa Press, Ottawa
- BirdLife International (2019) IUCN red list for birds. <http://www.birdlife.org>. (Accessed Nov 01 2019)
- Bourski OV, Forstmeier W (2000) Does interspecific competition affect territorial distribution of birds? A long-term study on Siberian *Phylloscopus* warblers. *Oikos* 88:341–350
- Bozó L, Heim W (2016) Sex-specific migration of *Phylloscopus* warblers at a stopover site in Far Eastern Russia. *Ringing Migr* 31:41–46
- Bozó L, Heim W, Csörgő T (2018a) Habitat use by Siberian warbler species at a stopover site in Far Eastern Russia. *Ringing Migr* 33:31–35
- Bozó L, Csörgő T, Heim W (2018b) Weather conditions affect spring and autumn migration of Siberian leaf warblers. *Avian Res* 9:33
- Bozó L, Csörgő T, Anisimov Y (2019) Estimation of flight range of migrant leaf-warblers at Lake Baikal. *Ardeola* 67:101–111
- Brazil M (2009) Birds of East Asia. Christopher Helm, London
- Clement P (2006a) Dusky Warbler (*Phylloscopus fuscatus*). In: del Hoyo J, Elliott A, Christie DA (eds) Handbook of the Birds of the World, vol 11. Lynx Edicions, Barcelona, p 655
- Clement P (2006b) Pale-legged Leaf-warbler (*Phylloscopus tenellipes*). In: del Hoyo J, Elliott A, Christie DA (eds) Handbook of the Birds of the World, vol 11. Lynx Edicions, Barcelona, pp 663–664
- Crawley MJ (2013) The R book, 2nd edn. Wiley, Chichester, U.K.
- del Hoyo J, Elliott A, Christie DA (2006) Handbook of the Birds of the World, vol 11. Lynx Edicions, Barcelona
- Demongin L (2016) Identification guide to birds in the hand. Beauregard-Vendon, France
- Ellegren H (1990) Timing of autumn migration in Bluethroats *Luscinia s. svecica* depends on timing of breeding. *Ornis Fennica* 67:13–17
- Forstmeier W, Kessler A (2001) Morphology and foraging behaviour of Siberian *Phylloscopus* warblers. *J Avian Biol* 32:127–138
- Forstmeier W, Kuijper DP, Leisler B (2001a) Polygyny in the Dusky Warbler, *Phylloscopus fuscatus*: the importance of female qualities. *Anim Behav* 62:1097–1108
- Forstmeier W, Bourski OV, Leisler B (2001b) Habitat choice in *Phylloscopus* warblers: the role of morphology, phylogeny and competition. *Oecologia* 128:566–576
- Forstmeier W (2002) Benefits of early arrival at breeding grounds vary between males. *J Anim Ecol* 71:1–9
- Forstmeier W, Balsby TJ (2002) Why mated Dusky Warblers sing so much: territory guarding and male quality announcement. *Behaviour* 139:89–111
- Fransson T (1995) Timing and speed of migration in North and West European populations of *Sylvia* warblers. *J Avian Biol* 26:39–48
- Gaston AJ (1974) Adaptation in the genus *Phylloscopus*. *Ibis* 116:432–450
- Ginn HB, Melville DS (1983) Moulting in birds. BTO Guide Tring, Britain
- Glutz von Blotzheim UN, Bauer KM (1991) Handbuch der Vögel Mitteleuropas, vol 12. Wiesbaden, AULA-Verlag, Teil II
- Gross SJ, Price TD (2000) Determinants of the northern and southern range limits of a warbler. *J Biogeogr* 27:869–878

- Hahn S, Bauer S, Liechti F (2009) The natural link between Europe and Africa – 2.1 billion birds on migration. *Oikos* 118:624–626
- Haraszthy L (1998) Birds of Hungary. Budapest, Mezőgazda Kiadó (in Hungarian)
- Hedenström A, Petterson J (1987) Migration routes and wintering areas of Willow Warblers *Phylloscopus trochilus* (L) ringed in Fennoscandia. *Ornis Fennica* 64(4):137–143
- Hedenström A, Bensch S, Hasselquist D, Lockwood M, Ottosson U (1993) Migration, stopover and moult of the Great Reed Warbler *Acrocephalus arundinaceus* in Ghana, West Africa. *Ibis* 135:177–180
- Heim W, Smirenski SM (2013) The Amur bird project at Muraviovka Park in Far East Russia. *BirdingASIA* 19:31–33
- Heim W (2014) Birds at Muraviovka Park 2011–2013: Results of the Amur Bird Project. Unpublished technical report
- Heim W, Smirenski SM (2017) The importance of Muraviovka Park/Far East Russia for endangered bird species on regional, national and international scale based on observations from 2011–2016. *Forktail* 33:77–83
- Heim W, Eccard JA, Bairlein F (2018) Migration phenology determines niche use of East Asian buntings (*Emberizidae*) during stopover. *Curr Zool* 64:681–692
- Jenni L, Kéry M (2003) Timing of autumn bird migration under climate change: advances in long-distance migrants, delays in short-distance migrants. *P Roy Soc Lond B Bio* 270:1467–1471
- Jenni L, Winkler R (2020) Moulting and ageing of European Passerines. Second Edition, Helm
- Karlsson H, Nilsson C, Bäckman J, Alerstam T (2012) Nocturnal passerine migrants fly faster in spring than in autumn: a test of the time minimization hypothesis. *Anim Behav* 83:87–93
- Katti M, Price TD (2003) Latitudinal trends in body size among overwintering leaf warblers (genus *Phylloscopus*). *Ecography* 26:69–79
- Kjellén N (1994) Moulting in relation to migration in birds—A review. *Ornis Svec* 4(1):1–24
- Lindström Å, Pearson DJ, Hasselquist D, Hedenström A, Bensch S, Åkesson S (1993) The moult of Barred Warblers *Sylvia nisoria* in Kenya – evidence for a split wing-moult pattern initiated during the birds' first winter. *Ibis* 135:403–409
- Lövei GL, Scebbba S, Milone M (1985) Migration and wintering of the Blackcap *Sylvia atricapilla* on a Mediterranean island. *Ringling Migr* 6(1):39–44
- Marchetti K, Price T, Richman A (1995) Correlates of wing morphology with foraging behaviour and migration distance in the genus *Phylloscopus*. *J Avian Biol* 26:177–181
- Marra PP, Hobson KA, Holmes RT (1998) Linking winter and summer events in a migratory bird by using stable-carbon isotopes. *Science* 282:1884–1886
- Marra PP, Holberton RL (1998) Corticosterone levels as indicators of habitat quality: effects of habitat segregation in a migratory bird during the non-breeding season. *Oecologia* 116:284–292
- Maslovsky KS, Valchuk OP, Leliuchina EV (2018) Differential migration and dynamic state of the state of the trasnit population of Siberian Rubythroat in Southern Primorye: analysis of long-term banding data in the Litovka River valley. *Bull Far East Br Russ Acad Sci* 2:19–28 (in Russian)
- McNab BK (1971) On the ecological significance of Bergmann's rule. *Ecology* 52:845–854
- Nam HY, Choi CY, Park JG, Hong GP, Won IJ, Kim SJ, Bing GC, Chae HY (2011) Protandrous migration and variation in morphological characters in *Emberiza* buntings at an East Asian stopover site. *Ibis* 153:494–501
- Neto JM, Newton J, Gosler AG, Perrins CM (2006) Using stable isotope analysis to determine the winter moult extent in migratory birds: the complex moult of Savi's Warblers *Locustella luscinioides*. *J Avian Biol* 37:117–124
- Newton I (2007) Weather-related mass-mortality events in migrants. *Ibis* 149:453–467
- Nilsson C, Klaassen RH, Alerstam T (2013) Differences in speed and duration of bird migration between spring and autumn. *Am Nat* 181:837–845
- Owen RB, Krohn WB (1973) Molt patterns and weight changes of the American Woodcock. *Wilson Bull* 85:31–41
- Pérez-Tris J, de la Puente J, Pinilla J, Bermejo A (2001) Body moult and autumn migration in the Barn Swallow *Hirundo rustica*: is there a cost of moulting late? In: *Ann Zool Fennici*. Finnish Zoological and Botanical Publishing Board, Helsinki
- Piersma T, Ramenofsky M (1998) Long-term decreases of corticosterone in captive migrant shorebirds that maintain seasonal mass and moult cycles. *J Avian Biol* 29:97–104
- Piersma T, Drent J (2003) Phenotypic flexibility and the evolution of organismal design. *Trends Ecol Evol* 18:228–233
- Price T (1991) Morphology and ecology of breeding warblers along an altitudinal gradient in Kashmir, India. *J Anim Ecol* 60:643–664
- Price T, Jamdar N (1991) Breeding of eight sympatric species of *Phylloscopus* warblers in Kashmir. *J Bom Nat Hist Soc* 88:242–255
- Pulido F, Coppack T (2004) Correlation between timing of juvenile moult and onset of migration in the Blackcap, *Sylvia atricapilla*. *Anim Behav* 68:167–173
- Core Team R (2019) R: A language and environment for statistical computing. Austria, R Foundation for Statistical Computing, Vienna
- Rubolini D, Spina F, Saino N (2005) Correlates of timing of spring migration in birds: a comparative study of trans-Saharan migrants. *Biol J Linn Soc* 85(2):199–210
- Salewski V, Altwegg R, Erni B, Falk KH, Bairlein F, Bernd L (2004) Moulting of three Palaearctic migrants in their West African winter quarters. *J Ornithol* 145:109–116
- Sander MM, Eccard JA, Heim W (2017) Flight range estimation of migrant Yellow-browed Warblers *Phylloscopus inornatus* on the East Asian Flyway. *Bird Study* 64:569–572
- Southwood TRE (1961) The number of species of insect associated with various trees. *J Anim Ecol* 30:1–8
- Studds CE, Marra PP (2005) Nonbreeding habitat occupancy and population processes: an upgrade experiment with a migratory bird. *Ecology* 86:2380–2385
- Svensson L (1992) Identification guide to European passerines. Svensson, Stockholm
- Svensson E, Hedenström A (1999) A phylogenetic analysis of the evolution of moult strategies in Western Palearctic warblers (Aves: Sylviidae). *Biol J Linn Soc* 67:263–276
- Tiainen J, Hanski IK (1985) Wing shape variation of Finnish and central European Willow Warblers *Phylloscopus trochilus* and Chiffchaffs *P. collybita*. *Ibis* 127(3):365–371
- Yong DL, Liu Y, Low BW, Española CP, Choi CY, Kawakami K (2015) Migratory songbirds in the East Asian-Australasian Flyway: a review from a conservation perspective. *Bird Conserv Int* 25:1–37
- Verboven N, Visser ME (1998) Seasonal variation in local recruitment of great tits: the importance of being early. *Oikos* 81:511–524
- Wang Y, Chang JC, Moore FR, Su L, Cui L, Yang X (2006) Stopover ecology of Red-flanked Bush Robin (*Tarsiger cyanurus*) at Maoershan, Northeast China. *Acta Ecol Sinica* 26:638–646 (In Chinese)
- Weber TP, Ens BJ, Houston AI (1998) Optimal avian migration: a dynamic model of fuel stores and site use. *Evol Ecol* 12:377–401
- Weber TP, Houston AI, Ens BJ (1999) Consequences of habitat loss at migratory stopover sites: a theoretical investigation. *J Avian Biol* 30:416–426