

GENERAL RELATIVITY AND THE AKSZ CONSTRUCTION

G. CANEPA, A. S. CATTANEO, AND M. SCHIAVINA

ABSTRACT. In this note the AKSZ construction is applied to the BFV description of the reduced phase space of the Einstein–Hilbert and of the Palatini–Cartan theories in every space-time dimension greater than two. In the former case one obtains a BV theory for the first-order formulation of Einstein–Hilbert theory, in the latter a BV theory for Palatini–Cartan theory with a partial implementation of the torsion-free condition already on the space of fields. All theories described here are BV versions of the same classical system on cylinders. The AKSZ implementations we present have the advantage of yielding a compatible BV-BFV description, which is the required starting point for a quantization in presence of a boundary.

CONTENTS

Introduction	1
Relevance and outlook	5
Acknowledgements	6
1. Background	6
1.1. The Batalin–Vilkovisky formalism	6
1.2. The AKSZ construction	8
1.3. One-dimensional AKSZ construction	10
2. BFV theories of gravity	11
2.1. BFV Einstein–Hilbert Theory	11
2.2. BFV Palatini–Cartan theory	12
3. AKSZ EH	15
3.1. Pushforward	18
3.2. Reconstruction of Einstein–Hilbert theory	20
4. AKSZ PC	26
4.1. Comparison of BV data for PC theory	28
4.2. An interpretation of the restricted theory	32
4.3. Three dimensional case	34
Appendix A. Lengthy calculations	34
References	39

INTRODUCTION

A Lagrangian field theory F on a cylinder $\Sigma \times I$, where I is a “time” interval, can be given a corresponding Hamiltonian description in terms of a symplectic manifold (the phase space) of the possible initial conditions on Σ and a Hamiltonian that

This research was (partly) supported by the NCCR SwissMAP, funded by the Swiss National Science Foundation. G.C. and A.S.C. acknowledge partial support of SNF Grant No. 200020_192080. M.S. acknowledges partial support from Swiss National Science Foundation grants P2ZHP2_164999 and P300P2_177862. Part of the research has been carried out while M.S. was a visiting scientist at Max Planck Institute for Mathematics, Bonn, funded by the Max Planck Foundation.

describes the time evolution. If the Lagrangian is degenerate, its Euler–Lagrange equations yield, in addition to time evolution, some constraints that have to be taken into account when specifying the initial conditions. The true phase space, called the “reduced phase space”, is typically described as the symplectic reduction of the coisotropic submanifold defined by the constraints (hence the name).

This reduction is often singular, and one possible description is by means of a cohomological resolution: one introduces a complex whose cohomology is the algebra of functions of the reduced phase space. In addition, one wants this resolution to feature also the symplectic/Poisson nature of the phase space, and a solution to this problem is provided by the Batalin–Fradkin–Vilkovisky (BFV) formalism [BF83] (see also [Sch08; Sch09; Sta97]). We denote by \mathfrak{F}^∂ the collection of data associated to the reduced phase space of a Lagrangian theory F , as a BFV theory (Definition 3).

On the other hand, a flexible way to deal with a degenerate Lagrangian is the Batalin–Vilkovisky (BV) formalism [BV81], which allows a cohomological resolution of the space of solutions to the Euler–Lagrange equations modulo symmetries but is also the starting point for perturbative quantization. We denote with \mathfrak{F} the BV data associated to a classical Lagrangian field theory F , as a BV theory (Definition 1).

To quantize a Lagrangian field theory F on a cylinder $\Sigma \times I$, one needs a good relation between its associated BV and BFV data \mathfrak{F} and \mathfrak{F}^∂ . In [CMR11] an explicit procedure was introduced to construct what in [CMR14] is called a BV-BFV theory (Definition 5), associating to the BV data \mathfrak{F} certain BFV data denoted by $BFV(\mathfrak{F})$ in a way suitable for quantization [CMR18] — under some regularity assumptions. In regular cases it relates \mathfrak{F} and \mathfrak{F}^∂ , so that that $BFV(\mathfrak{F}) = \mathfrak{F}^\partial$.

While it is true that both \mathfrak{F} and \mathfrak{F}^∂ depend on F , the relation $BFV(\mathfrak{F}) = \mathfrak{F}^\partial$ is not guaranteed, and it is a necessary requirement for BV quantisation with boundary [CMR18]. This relation turns out to hold for a large variety of field theories, including general relativity (GR) in the Einstein–Hilbert (EH) formulation in any space–time dimension greater than 2 [CS16]. However, the procedure notably fails in the case of GR in the Palatini–Cartan (PC) formulation¹ in 3 + 1 dimensions [CS19b], as the construction of $BFV(\mathfrak{F})$ is obstructed. However, \mathfrak{F}^∂ exists and has been presented in [CCS20].

Conversely, given a BFV theory \mathfrak{F}^∂ associated to a manifold Σ , there is a standard way² to produce a BV theory on $\Sigma \times I$ by means of a construction due to Alexandrov, Kontsevich, Schwarz and Zaboronski (AKSZ [Ale+97]). The resulting BV theory, which we temporarily denote here by³ $AKSZ(\mathfrak{F}^\partial)$, satisfies automatically the regularity assumptions required by the BV-BFV formalism, and we also have $BFV(AKSZ(\mathfrak{F}^\partial)) = \mathfrak{F}^\partial$.

On the other hand, in general $AKSZ(BFV(\mathfrak{F}))$ will not be the same as \mathfrak{F} . In fact, the AKSZ construction produces a theory that is invariant under reparametrization of I , which is certainly different from \mathfrak{F} if the latter does not enjoy this invariance. In this case $AKSZ(BFV(\mathfrak{F}))$ is a version of \mathfrak{F} with “frozen time” and may be used to describe a change in the polarization chosen for the quantization of the reduced phase space (see [CMR18, Remark 2.38]). If \mathfrak{F} is reparametrization invariant — e.g. a topological field theory or GR — we may wonder whether $AKSZ(BFV(\mathfrak{F}))$

¹There appears to be no uniform consensus on the nomenclature to best attribute and label the theory that will be described in Sections 2.2 and 4. We discuss our choice “Palatini–Cartan” in [CS19c] (see also references therein). Other choices include the names of Einstein, Weyl, Sciama and Kibble, in various combinations.

²To the best of our knowledge, the first explicit application of the AKSZ construction to a BFV target to produce a BV structure in one dimension goes back to [GD00].

³This construction is clarified in Theorem 14, and $AKSZ(\mathfrak{F}^\partial)$ will be denoted $\mathfrak{F}^{AKSZ}(I; \mathfrak{F}^\partial)$.

and \mathfrak{F} are somehow related. In the case of AKSZ topological field theories, it turns out that $AKSZ(BFV(\mathfrak{F}))$ and \mathfrak{F} are actually the same. For more general reparametrization invariant theories we might expect the two to be equivalent, in one of the possible ways presented below.

A BV theory \mathfrak{F} is essentially composed of a (-1)-symplectic manifold (\mathcal{F}, ϖ) and an action functional S over it. We say that \mathfrak{F}_1 and \mathfrak{F}_2 are strongly BV-equivalent if there is a symplectomorphism $\phi: (\mathcal{F}_1, \varpi_1) \rightarrow (\mathcal{F}_2, \varpi_2)$ that relates their action functionals, i.e. $S_1 = \phi^* S_2$. This in particular implies that their BV cohomology groups are isomorphic. A nontrivial example of strong BV-equivalence is the one between PC and BF theory in 3 space–time dimensions [CS19a; CSS18].

If \mathfrak{F}_2 is obtained from \mathfrak{F}_1 by a partial integration of the fields⁴ (with some partial gauge fixing), we say that \mathfrak{F}_2 is an effective theory for \mathfrak{F}_1 . We say that two BV theories are effectively BV-equivalent if one is (strongly BV-equivalent to) an effective theory for the other. Typical cases for this are Wilson renormalization or the passage to a second-order theory from its associated first-order formulation. Another important example is given by elimination of so-called auxiliary fields. In that case, one can argue that effective equivalence also preserves the BV cohomology [BBH95; Hen90] (see Remark 12).

A third case is when the theories \mathfrak{F}_1 and \mathfrak{F}_2 have the same space of classical solutions modulo symmetries. We speak in this case of classical equivalence. A typical case of classical equivalence is that between EH and PC. Observe that this is equivalent to just asking that the degree-zero BV cohomologies of the two theories coincide, making this kind of equivalence weaker.

In this paper we study this question for EH and PC models of gravity in any space–time dimension greater than 2, assuming that the metric encoded in the BFV data \mathfrak{F}^∂ is nondegenerate (i.e. assuming that the manifold Σ is either spacelike or timelike but not lightlike). In the case of EH, we show that \mathfrak{F} and $AKSZ(BFV(\mathfrak{F}))$ are effectively equivalent, with the former being actually the first-order formulation of the latter.

In the case of PC in three dimensions, where $BFV(\mathfrak{F}) = \mathfrak{F}^\partial$ holds, we show that $AKSZ(BFV(\mathfrak{F}))$ and \mathfrak{F} are strongly BV equivalent, which is not unexpected, since PC is strongly BV equivalent to BF theory [CS19a; CSS18], and the latter is a topological AKSZ theory. Instead, for higher dimensional PC theory we show that $AKSZ(\mathfrak{F}^\partial)$ and \mathfrak{F} are classically equivalent⁵, with \mathfrak{F}^∂ the BFV data constructed from the reduced phase space of PC theory [CCS20; CS19c]. This case is particularly interesting because the BV-BFV construction for PC is obstructed in dimension 4 (and presumably higher). The data $AKSZ(\mathfrak{F}^\partial)$ resulting from the AKSZ construction is a new BV theory defined on cylinders that is still classically equivalent to EH, but also compatible with the BV-BFV formalism (by construction via the AKSZ procedure). Classically, it is simply PC on a smaller space of fields, where part of the torsion-free condition is imposed a priori instead of through the Euler–Lagrange equations.

Our result addresses the problem presented in [CS19b], where it was pointed out that PC theory in dimension greater than three must be complemented with requirements on field configurations at the boundary in order to induce a well-defined BV-BFV structure. One possible way to construct a BV-BFV structure for PC theory is to assume vector fields generating diffeomorphisms transversal to the boundary to vanish at the boundary. Denote by $\underline{\mathfrak{F}}$ the resulting BV theory. In [CS19b, Section 5, Remark 34] this was shown to be insufficient to describe

⁴This is more appropriately called BV-pushforward or BV fiber integral, see [CMR18, Section 2.2.2].

⁵On an open subset of the moduli space of solutions.

the full reduced phase space of GR, as the Hamiltonian constraint is lost in the process: this means that $BV(\mathfrak{F}) \neq \mathfrak{F}^\partial$. Alternatively one may require certain components of the Lorentz connection to vanish on the boundary, although this condition is not natural for general manifolds. One way of reading our paper is to make these conditions natural on cylindrical manifolds, in the sense that we present a version of PC theory, with the compatibility requirements already implemented. In fact, the resulting AKSZ theory has the same equations of motion and the same symmetries, but the AKSZ procedure restricts the moduli space of solutions to an open subset. This is akin to restricting to globally hyperbolic solutions. One can think of the extra required conditions as imposing part of the equations of motion that fix ω to be the Levi-Civita connection for the metric induced by a tetrad e . This is discussed in Section 4.2.

Let us stress that having a well-defined BV-BFV structure is a necessary requirement for the quantisation of BV theories with boundary [CMR18]. The fact that the boundary-compatible AKSZ version of PC theory is (possibly) only classically equivalent to the original PC formulation reinforces the idea that care must be placed when attempting BV quantisation of the latter.

A related approach is the ‘parent formulation’ by Barnich and Grigoriev [BG11; Gri11] which derives an AKSZ construction of the BV theory from the jet space formalism (trivariational complex). What is crucially different in our construction is that we consider, as a target, a symplectic description of the classical boundary states. This involves a careful symplectic reduction of the naively associated boundary spaces⁶. The result of our construction is not only a BV reformulation of the original bulk theory, but a reformulation that is compatible with the boundary as a 1-extended BV-BFV theory (see Definition 5), which is the starting point for quantum (or at least semiclassical) considerations for a theory with boundary [CMR18].

For the same reason, unlike the presymplectic AKSZ formulation presented by Grigoriev et Alkalaev in [AG14] and [Gri16], our BV-BFV description of PC gravity is based on a symplectic structure, which is essential for quantization. This does not arise directly from a reduction of the natural presymplectic BFV structure derived from BV in the bulk, which is impossible for $N \geq 4$ as shown in [CS19b], but it is the symplectic BFV structure [CCS20] that resolves the reduced phase space of the theory.

Finally, note that in this paper we consider two separate applications of the AKSZ “reconstruction” of a parametrization-invariant bulk BV theory from its boundary BFV structure, respectively for two formulations of GR (EH and PC). We do not discuss the equivalence between EH and PC, but we investigate the appropriate BV equivalence between each formulation and its own AKSZ “reconstruction.”

These considerations do not exclude, however, some deeper connection between our construction and the ones mentioned above, which are definitely worth exploring.

The paper is organised as follow. In Sections 1.1 and 1.2 we will outline the BV-BFV and AKSZ constructions, while Section 2 is a brief review of the construction of the BFV data for Einstein–Hilbert and Palatini–Cartan theories of gravity, as presented respectively in [CS16] and [CCS20].

Finally, in Sections 3 and 4 we will apply the AKSZ construction to the BFV data of EH and PC gravity, respectively, and compare it with the BV data for the two formulations as presented in [CS16] and [CS19b].

⁶This association is the natural restriction of fields and normal jets to the boundary, see [CMR14].

Relevance and outlook. This work is intended as a first reaping, as a result of a few years of sowing, in a program directed at an analysis of classical General Relativity seen through the lens of the BV formalism with boundary, an attempt at formalising its quantisation within the BV-BFV formalism [CMR18]. The program was initiated in [CS16; CS19b; Sch15], where a few inconsistencies in the behaviour of GR in the presence of boundaries in dimension 4 were detected, and was later extended in [CCS20; CS19a; CSS18], where the comparison with the three dimensional analogue was made.

This series of works is motivated by the obstruction encountered in defining the BV-BFV data for Palatini–Cartan gravity, a requirement for the BV quantisation program with boundary, which has otherwise provided very reliable and flexible (see [CMR18; CMR20] for the quantisation of BF theory, [IM19] for Yang–Mills theory in dimension 2, [CMW17; CMW19a] for split Chern–Simons theory and [CMW19b] for a general approach to a class of AKSZ models, including the Poisson sigma model). No obstruction to BV quantisation with boundary is otherwise present for Einstein–Hilbert theory, and this discrepancy points at the fact that classical equivalence of field theories might be too coarse a classification to have bearing on the respective quantum theories.

The results contained in this paper close the circle, so to speak, in the comparison of classical BV general relativity with boundary, between EH and PC formulations. As a matter of fact, while the AKSZ construction for EH theory is effectively equivalent to the BV theory analysed in [CS16], this is not the case in PC theory analysed in [CS19b] (it is only included within). This fact, together with the equivalence of the reduced phase spaces for EH and PC theories [CCS20; CS19c], can be interpreted as a confirmation that BV Palatini–Cartan theory *must* be supplemented with additional requirements on fields, or otherwise restricted, in order to be viable for BV quantisation. The requirements we find, summarised by Definition 52 are conditions on the Lorentz connection and its conjugate variable, which effectively restrict the space of fields. We find that these conditions are somewhat natural on cylinders.

The very ultimate goal of the construction presented in this paper is the grail of quantisation of gravity. We do not attempt doing it here. What we present is the preliminary setting for a perturbative quantisation on cylinders resulting in the quantum evolution operator from the initial to the final quantum space of states. Due to the degeneracy of the actions (related to gauge invariance), one needs a formalism that allows imposing gauge fixings and checking that the results are independent thereof, up to equivalences that are under control. In the absence of free boundaries (i.e., in the computation of partition functions and expectation values), there are several good methods to do this, including BRST and BV. In the presence of boundary, the best developed method is a compatible combination of BV in the bulk and BFV on the boundary. The compatibility is the main issue here, and this paper discusses it in the context of GR theories.

Performing the actual quantisation, which is far beyond the scope of this paper, implies choosing a polarization on the boundary and a gauge fixing in the bulk, computing the resulting propagators, regularizing the theory, and performing renormalisation in a compatible way with the BV-BFV data (the quantum master equation, and its version with boundary [CMR18]). In the case of gravity, one of course expects an infinite number of independent counterterms to be taken care of. Clever or miraculous ways to keep them under control are the same issue as in other treatments (without boundary): we do not claim to have a better recipe for this issue, but just to have a method to incorporate free boundaries. A minimal way to proceed, as, e.g., in [BFR16], is to allow for infinite counterterms (which is

algebraically possible and allows for the construction of families of effective theories, even though the predictivity at all energies is missing).

Naturally, since the outlook of this extended program is that of addressing quantisation of General Relativity (with boundary), we wish to stress that without the observations produced in this preliminary phase, an early attempt at directly quantising PC theory might have been thwarted by the very obstructions highlighted by our investigations.

In this sense, we believe the correct preparation of a field theory for its perturbative quantisation to be of crucial importance to drive the scientific effort towards sensible questions, and divert it when evidence is presented of a potential roadblock ahead. This should be of particular interest for the scientific community heavily involved with the study of Palatini–Cartan theory as a fundamental building block for a quantum theory of gravity.

Acknowledgements. We thank G. Barnich and M. Grigoriev for useful discussions on both content and context, as well as the anonymous referees, whose comments have helped improve our manuscript. M.S. would also like to thank C. Blohmann and A. Weinstein for numerous scientific interactions relevant to this paper.

1. BACKGROUND

One of the goals of this paper is the construction of a BV theory on a cylindrical manifold $\Sigma \times I$ by means of the AKSZ construction, with target a BFV theory associated to Σ . In this section we introduce the basic definition of the BV(-BFV) and AKSZ formalisms, together with the relevant notions of equivalence that will allow us to compare theories. We refer to [BF83; BV77; BV81; CMR11; CMR14] for a more detailed introduction and more insight in the meaning and the motivations for the following definitions and theorems. For an introduction of the BV (-BFV) formalism and gravity see [CS19a; CS17; CS19c]. Other versions and interpretations of the BV formalism for gravity can be found in [BFR16].

1.1. The Batalin–Vilkovisky formalism.

Definition 1. A BV theory is a quadruple $\mathfrak{F} = (\mathcal{F}, S, \varpi, Q)$ where \mathcal{F} is a graded manifold (the *space of BV fields*) endowed with a degree -1 symplectic form ϖ , $S: \mathcal{F} \rightarrow \mathbb{R}$ is a degree 0 functional (the *BV action*) and Q is the (odd) Hamiltonian vector field of S with respect to ϖ satisfying $[Q, Q] = 0$.

Remark 2. Since Q is the Hamiltonian vector field of S , i.e. $\iota_Q \varpi = \delta S$ where δ is the de Rham differential on \mathcal{F} and ι_Q is the contraction w.r.t. Q , we can rewrite the equation $[Q, Q] = 0$ as $(S, S) = 0$ where (\cdot, \cdot) denotes the Poisson bracket defined by ϖ . The latter equation is called the Classical Master Equation (CME).

Definition 3. An exact BFV theory is a quadruple $\mathfrak{F}^\partial = (\mathcal{F}^\partial, S^\partial, \varpi^\partial, Q^\partial)$ where \mathcal{F}^∂ is a graded manifold (the *space of boundary fields*) endowed with a degree-0 exact symplectic form $\varpi^\partial = \delta\alpha^\partial$, $S^\partial: \mathcal{F}^\partial \rightarrow \mathbb{R}$ is a degree 1 functional and Q^∂ is the Hamiltonian vector field of S^∂ with respect to ϖ^∂ such that $[Q^\partial, Q^\partial] = 0$.

Remark 4. Typical examples of BV and BFV theories are modeled on sections of bundles over differentiable manifolds, possibly with boundary, with $\varpi^{(\partial)}$, $S^{(\partial)}$ and $Q^{(\partial)}$ respectively a local two-form, functional and vector field. Throughout the paper, when specifying BV theories, we will assume that the equations $\iota_Q \varpi = \delta S$ and $(S, S) = 0$ are satisfied only up to boundary terms. The failure of said equations will be controlled by the data of a BV-BFV theory, as follows. It is often convenient, in this scenario, to define the slightly more general concept of a

relaxed BV theory, i.e. data $\mathfrak{F} = (\mathcal{F}, S, \varpi, Q)$ as in Definition 3, but without the requirement that Q be the Hamiltonian vector field of S . If we are given a BV theory on a closed manifold without boundary, we can consider the same local data as a relaxed BV theory on a manifold with boundary.

Definition 5 ([CMR14]). A (relaxed) BV theory $\mathfrak{F} = (\mathcal{F}, S, \varpi, Q)$ is said to be 1-extended to the BFV theory $\mathfrak{F}^\partial = (\mathcal{F}^\partial, S^\partial, \varpi^\partial, Q^\partial)$ if there exists a surjective submersion $\pi : \mathcal{F} \rightarrow \mathcal{F}^\partial$, such that the following compatibility relation is satisfied:

$$\iota_Q \varpi = \delta S + \pi^* \alpha^\partial \quad (1a)$$

The data $\mathfrak{F}^{\uparrow 1} = (\mathfrak{F}, \mathfrak{F}^\partial, \pi)$ will be called 1-extended BV-BFV theory.

Remark 6. Notice that, from the data above, the following relation follows:

$$\iota_Q \iota_Q \varpi = 2\pi^* S^\partial. \quad (1b)$$

The following definitions compare two different BV (or BFV) theories.

Definition 7. Two B(F)V theories $\mathfrak{F}_1^{(\partial)}$ and $\mathfrak{F}_2^{(\partial)}$ are said to be strongly B(F)V-equivalent if there exists a symplectomorphism

$$\Phi : (\mathcal{F}_1^{(\partial)}, \varpi_1^{(\partial)}) \rightarrow (\mathcal{F}_2^{(\partial)}, \varpi_2^{(\partial)})$$

preserving the BV action: $\Phi^* S_2^{(\partial)} = S_1^{(\partial)}$. The map Φ is called a strong B(F)V-equivalence.

Definition 8. Let \mathfrak{F}_1 and \mathfrak{F}_2 be two (relaxed) BV theories. A (relaxed) BV-inclusion $\mathfrak{J} : \mathfrak{F}_1 \rightarrow \mathfrak{F}_2$ is an inclusion of (super)manifolds $\iota : \mathcal{F}_1 \rightarrow \mathcal{F}_2$ such that $\varpi_1 = \iota^* \varpi_2$ and $\iota^* Q_1 = Q_2 \iota^*$. If the two theories are relaxed we will additionally require $\iota^* S_2 = S_1$. In this case we say that \mathfrak{F}_1 is a BV-subspace⁷ of \mathfrak{F}_2 .

Remark 9. Naturally, if Q_1 and Q_2 are the Hamiltonian vector fields of S_1 and S_2 respectively, the condition $\iota^* S_2 = S_1$ is equivalent to the condition $\iota^* Q_2 = Q_1 \iota^*$, up to a constant.

Proposition 10. *The composition of a strong BV equivalence and a BV inclusion is in turn a BV inclusion.*

Proof. The map $\Phi \circ \iota$ satisfies trivially the properties of a BV inclusion. \square

A notion that we will need to compare theories is that of BV-pushforward. This notion is usually phrased at the quantum level [CMR18; Mne17], where the additional data of a BV Laplacian needs to be provided. However here we are interested mainly in its classical counterpart. The basic setting is the same, although we consider the following simplifying assumptions. Suppose that we have a splitting of a graded symplectic manifold (\mathcal{F}, ϖ) so that $\mathcal{F} = \mathcal{F}' \times \mathcal{F}''$, with $\varpi = \varpi' + \varpi''$, and let \mathcal{L} be a Lagrangian submanifold of $(\mathcal{F}'', \varpi'')$ endowed with a half-density μ on \mathcal{F}'' , which thus defines by restriction a density $\mu_{\mathcal{L}}$ on \mathcal{L} . Denote coordinates (z', z'') respectively in $\mathcal{F}', \mathcal{F}''$, and let $z'' \in \{x, x^\dagger\}$ be Darboux adapted coordinates such that x parametrises \mathcal{L} and x^\dagger are transversal.

Definition 11. We define the Batalin–Vilkovisky–Legendre transform of a functional $S \in C^\infty(\mathcal{F})$, with respect to the Lagrangian $\mathcal{L} \subset \mathcal{F}''$, as $S_{\text{BVL}} \in C^\infty(\mathcal{F}')$:

$$S_{\text{BVL}} = S(z, x_0, x^\dagger = 0) \quad (2)$$

where x_0 is a critical point for S (assumed unique):

$$\left. \frac{\delta S}{\delta x} \right|_{x^\dagger=0} (x_0) = 0.$$

⁷In the math literature, a map with this compatibility between the symplectic structures and the cohomological vector fields is known as a morphism of dg symplectic manifolds.

Starting from a BV theory on (\mathcal{F}, ϖ) we build a theory on (\mathcal{F}', ϖ') by means of a gauge-fixed fiber integral along \mathcal{F}'' , with gauge-fixing Lagrangian \mathcal{L} . In other words, if S denotes a BV action on \mathcal{F} we consider the effective result of the BV pushforward (fiber integral) to be

$$\exp\left(\frac{i}{\hbar} S_{\text{eff}}\right) := \int_{\mathcal{L} \subset \mathcal{F}''} \exp\left(\frac{i}{\hbar} S\right) \Big|_{\mathcal{L}} \mu_{\mathcal{L}} \quad (3)$$

where the integral is defined perturbatively as a power series in \hbar . Note that S_{BVL} is the dominant term of S_{eff} . When S depends only quadratically on the variables on \mathcal{L} , the only correction is $i\hbar/2$ times the logarithm of the determinant of the quadratic form.

Remark 12. Let us comment briefly on the notion of equivalence of theories in the BV formalism. When the moduli spaces of solutions of the equations of motion for two theories coincide, the theories are said to be classically equivalent. A finer notion of equivalence requires that the BV cohomologies be isomorphic⁸, and it allows for an extension to the case with boundary. One looks at the bicomplex given by the BV operator and the de Rham differential $(\Omega^{\bullet, \bullet}(\mathcal{F}_i, M), Q_i - d)$, and equivalence in this sense requires two theories to have quasi-isomorphic local de Rham/BV complexes. The typical argument for equivalence involves the elimination of so-called generalised auxiliary fields and trivial pairs [BBH95; Hen90]. This notion, however, is suboptimal because it relies on homological perturbation theory, which potentially could output an infinite tower of ghosts and antighosts in the process of constructing an equivalent theory. In other words, that approach — albeit somewhat well-established — does not provide a direct answer to the question of whether two *given* BV theories (in the form of two Hamiltonian dg-manifolds) are equivalent, provided their degree-0 sectors are classically equivalent. For this, instead, a spectral sequence argument would be more appropriate.⁹

On the other hand, one could phrase equivalence in the BV-BFV formalism, requiring existence of the BV-BFV structure and equivalence of the respective BV and BFV cohomologies (an explicit example of this, is the strong equivalence of General Relativity and BF theory in three dimensions [CS19a]). This existence requirement might become an obstruction to BV-BFV equivalence¹⁰. For our purposes, then, even assuming that by removal of generalised auxiliary fields one may prove some BV equivalence between Palatini–Cartan and Einstein–Hilbert theories, the results of [CS19b] and of the present paper show a discrepancy of the theories in the BV-BFV sense.

1.2. The AKSZ construction. Let X be a graded manifold and N an ordinary manifold and let μ_N be the canonical Berezinian¹¹ on $T[1]N$.

Definition 13 (Transgression map). Consider the map

$$\mathfrak{T}_N^{(\bullet)} : \Omega^{\bullet}(X) \longrightarrow \Omega^{\bullet}(\text{Map}(T[1]N, X)) \quad (4)$$

⁸Classical equivalence is the less restrictive requirement that only their 0-cohomologies be isomorphic.

⁹In an optimal scenario, to auxiliary fields one can associate a subcomplex with trivial cohomology. More generally, one has a filtration of the original BV complex, so that the associated spectral sequence converges. However, this is in general quite hard to prove.

¹⁰Observe that this is essentially the strictification requirement of the presymplectic BFV data, see Remark 29.

¹¹Recall that a function on $T[1]N$ is the same as a differential form on N . Integrating a function on $T[1]N$ against the canonical Berezinian μ_N is by definition the same as integrating the corresponding differential form on N , which we assume to be oriented.

defined by $\mathfrak{T}_N^{(\bullet)} := p_* \text{ev}^*$, where

$$\begin{array}{ccc} \text{Map}(T[1]N, X) \times T[1]N & \xrightarrow{\text{ev}} & X \\ p \downarrow & & \\ \text{Map}(T[1]N, X) & & \end{array} \quad (5)$$

and we set $p_* = \int_N \mu_N$. We will call $\mathfrak{T}_N^{(\bullet)}$ the transgression map, and its evaluation a transgression.

We endow the graded manifold X with a function S of degree n and parity $n \bmod 2$, together with a one-form α of degree $n - 1$ and parity $n - 1 \bmod 2$, such that $\varpi = d\alpha$ is nondegenerate and $\{S, S\} = 0$ with respect to the Poisson structure defined by ϖ . Then we say that X has a Hamiltonian dg-manifold structure, with differential $\{S, \cdot\}$.

Observing that the de Rham differential d_N on N can be seen as a degree 1 vector field on $\text{Map}(T[1]N, X)$ we have

Theorem 14 ([Ale+97]). *Let (X, S, α) a dg-manifold as described above. Consider the data*

$$\mathfrak{F}^{\text{AKSZ}}(N; X, S, \alpha) := (\mathcal{F}^{\text{AKSZ}}, S^{\text{AKSZ}}, \Omega^{\text{AKSZ}}, Q^{\text{AKSZ}}) \quad (6)$$

with $\mathcal{F}^{\text{AKSZ}} = \text{Map}(T[1]N, X)$, $\Omega^{\text{AKSZ}} := \mathfrak{T}_N^{(2)}(\varpi)$, the functional $S^{\text{AKSZ}}: \mathcal{F}^{\text{AKSZ}} \rightarrow \mathbb{R}$,

$$S^{\text{AKSZ}} := \mathfrak{T}_N^{(0)}(S) + \iota_{d_N} \mathfrak{T}_N^{(1)}(\alpha). \quad (7)$$

and the cohomological vector field Q^{AKSZ} such that $\iota_{Q^{\text{AKSZ}}} \Omega^{\text{AKSZ}} = \delta S^{\text{AKSZ}}$. Then, $\mathfrak{F}^{\text{AKSZ}}(N; X, S, \alpha)$ defines a BV theory.

We will call $\mathcal{F}^{\text{AKSZ}} := \text{Map}(T[1]N, X)$ the AKSZ space of fields. Introducing Darboux coordinates $\{p_i, q^i\}$ in X so that $\alpha = p_i dq^i$, the space of AKSZ fields is composed of inhomogeneous differential forms $\mathfrak{P}, \mathfrak{Q}$ on N . Then, if we consider X to be the space of sections of a bundle $E \rightarrow \Sigma$, that is to say $X = T^*[n - 1]C^\infty(\Sigma, E)$, we can write

$$\Omega^{\text{AKSZ}} = \int_{\Sigma \times N} [(\delta \mathfrak{P}, \delta \mathfrak{Q})]^{\text{top}} \equiv \int_{\Sigma \times N} [\delta \mathfrak{P}_i \delta \mathfrak{Q}^i]^{\text{top}} \quad (8)$$

where we have denoted by δ the deRham differential on spaces of maps and C^∞ -sections, and top denotes the top-form parts of the inhomogeneous differential forms within brackets. We will drop the superscript top in what follows.

Consider this elementary fact:

Lemma 15. *Let A, B, C be graded manifolds, $\phi: B \rightarrow C$ an isomorphism of graded manifolds, and μ_A a measure on A . Consider the diagram*

$$\begin{array}{ccc} A \times B & \xrightarrow{\text{id} \times \phi} & A \times C \\ \pi_B \downarrow & & \downarrow \pi_C \\ B & \xrightarrow{\phi} & C \end{array} \quad (9)$$

Then, setting $\pi_{B*} = \int \mu_A \cdot$ and $\pi_{C*} = \int \mu_A \cdot$, we have $\phi^* \circ \pi_{C*} = \pi_{B*} \circ (\text{id} \times \phi)^*$.

Theorem 16. *Let (X, S_X, α_X) and (Y, S_Y, α_Y) be equivalent Hamiltonian dg-manifolds, i.e. there exists a diffeomorphism $\phi: X \rightarrow Y$ such that $\varpi_X = \phi^* \varpi_Y$, and $S_X = \phi^* S_Y$. Then $\mathfrak{F}^{\text{AKSZ}}(N; X, S_X, \alpha_X)$ and $\mathfrak{F}^{\text{AKSZ}}(N; Y, S_Y, \alpha_Y)$ are strongly equivalent BV(-BFV) theories for every manifold N .*

Proof. $\phi: X \rightarrow Y$ induces an isomorphism

$$\tilde{\phi}: \text{Maps}(T[1]N, X) \rightarrow \text{Maps}(T[1]N, Y)$$

by precomposing maps with ϕ^* or ϕ^{-1*} . Then, we can apply Lemma 15 with $B = \text{Maps}(T[1]N, X)$, $C = \text{Maps}(T[1]N, Y)$ and $A = T[1]N$. \square

1.3. One-dimensional AKSZ construction. Let $I \subset \mathbb{R}$ be an interval, and \mathfrak{F}^∂ an exact BFV theory. We can construct a BV theory by applying Theorem 14 on the Hamiltonian dg-manifold underlying an exact BFV theory:

$$\mathfrak{F}^{\text{AKSZ}}(I; \mathfrak{F}^\partial) := \mathfrak{F}^{\text{AKSZ}}(I; \mathcal{F}^\partial, S^\partial, \alpha^\partial).$$

The resulting space of fields reads

$$\mathcal{F}^{\text{AKSZ}} = \text{Map}(T[1]I, \mathcal{F}^\partial).$$

Since the target space \mathcal{F}^∂ is (locally) a graded vector space, we identify the space of AKSZ fields with

$$\mathcal{F}^{\text{AKSZ}} = \Omega^\bullet(I) \otimes \mathcal{F}^\partial.$$

In particular, when \mathcal{F}^∂ is modeled on sections of some bundle over a $(N-1)$ -dimensional manifold Σ , we can view $\mathcal{F}^{\text{AKSZ}}$ as the space of sections of some (graded) bundle over $\Sigma \times I$. The space $\Omega^\bullet(I)$ splits into:

$$\Omega^\bullet(I) = C^\infty(I) \oplus \Omega^1(I)[-1]$$

hence, to each field in \mathcal{F}^∂ we associate two new fields. For simplicity we denote the field in $C^\infty(I) \otimes \mathcal{F}^\partial$ with the same letter as the old one, and use another letter for the one in $\Omega^1[-1](I) \otimes \mathcal{F}^\partial$.

Proposition 17 ([CMR18] and [CMR14]). *Let $\mathfrak{F}^\partial(\Sigma) = (\mathcal{F}^\partial(\Sigma), S^\partial(\Sigma), \varpi^\partial(\Sigma), Q^\partial(\Sigma))$ be an exact BFV theory, with $\mathcal{F}^\partial(\Sigma) := \Gamma(E \rightarrow \Sigma)$, and $\varpi^\partial(\Sigma) = \delta\alpha^\partial(\Sigma)$. Then, if $I := [0, 1]$ we have that $\mathfrak{F}^{\text{AKSZ}}(I; \mathfrak{F}^\partial(\Sigma))$ is a 1-extended BV-BFV theory over $\mathfrak{F}^\partial(\Sigma)$ (see Definition 5).*

Proof. Theorem 14 tells us that $\mathfrak{F}^{\text{AKSZ}}(I; \mathfrak{F}^\partial(\Sigma))$ is a BV theory (up to boundary terms). If we parametrise fields in $\mathcal{F}^{\text{AKSZ}}$ as

$$\mathfrak{P} = p(t) + q^\dagger(t)dt \quad \mathfrak{Q} = q(t) + p^\dagger(t)dt$$

we get

$$\Omega^{\text{AKSZ}} = \int_{\Sigma \times I} \langle \delta\mathfrak{P}, \delta\mathfrak{Q} \rangle = \int_{\Sigma \times I} \left\{ \langle \delta p, \delta p^\dagger \rangle + (-1)^{|q|+1} \langle \delta q, \delta q^\dagger \rangle \right\} dt$$

and

$$S^{\text{AKSZ}} = \int_{\Sigma \times I} \langle p, d_I q \rangle + [\mathfrak{I}_I^{(0)}(S^\partial(\Sigma))]^{\text{top}}.$$

The transgressed integrand needs to be first-order in dt , which leaves us with

$$[\mathfrak{I}_I^{(0)}(S^\partial(\Sigma))]^{\text{top}} \equiv S^\partial(\Sigma)[p + q^\dagger dt, q + p^\dagger dt] = \frac{\delta S^\partial(\Sigma)}{\delta p}(q^\dagger dt) + \frac{\delta S^\partial(\Sigma)}{\delta q}(p^\dagger dt)$$

Then Q^{AKSZ} splits in a transversal part plus a *tangential* one: $Q^{\text{AKSZ}} = Q^T + \hat{Q}$, where $Q^T q^\dagger = -\dot{p}$ and $Q^T p^\dagger = \dot{q}$ is essentially just deRham differential on I , and \hat{Q} is easily obtained:

$$\begin{aligned} \hat{Q}p &= \frac{\delta S^\partial(\Sigma)}{\delta q} \equiv Q^\partial p & \hat{Q}q &= \frac{\delta S^\partial(\Sigma)}{\delta p} \equiv Q^\partial q \\ \hat{Q}p^\dagger &= \frac{\delta^2 S^\partial(\Sigma)}{\delta q \delta p}(p^\dagger) + \frac{\delta^2 S^\partial(\Sigma)}{\delta p^2}(q^\dagger) & \hat{Q}q^\dagger &= \frac{\delta^2 S^\partial(\Sigma)}{\delta p \delta q}(q^\dagger) + \frac{\delta^2 S^\partial(\Sigma)}{\delta q^2}(p^\dagger). \end{aligned}$$

The boundary terms are easily seen in the given local chart, in fact:

$$\iota_{Q^{\text{AKSZ}}}\Omega^{\text{AKSZ}} = \delta S^{\text{AKSZ}} + \check{\alpha}$$

but $\check{\alpha}$ only sees contributions from $\langle p, dq \rangle$ and, up to sign, we get $\check{\alpha} = \alpha^\partial(\Sigma)$, with $\delta\alpha^\partial(\Sigma) = \varpi_\Sigma^\partial$. Then, the projection of Q^{AKSZ} along the natural projection map from $\mathcal{F}^{\text{AKSZ}}$ to the space of boundary fields, which coincides with $\mathcal{F}^\partial(\Sigma)$, is precisely $Q^\partial(\Sigma)$, concluding the argument. \square

Remark 18. A similar statement to Proposition 17 is presented in Henneaux–Bunster [HT92, Theorem 18.4.5], where one identifies the output of the above AKSZ construction with the (first-order) BV theory obtained by embedding in the BV formalism the generalised Hamiltonian formulation of a given field theory (see also [DGH90]). An analogous construction, already in the context of AKSZ theories, was presented in [BG05; GD00]. The added observation of Proposition 17 is the compatibility between BV for the bulk and BFV on the boundary, viz., what we call a 1-extended BV-BFV theory in Definition 5.

We would like to show that this construction behaves well under equivalences of the relevant BFV data.

Corollary 19 (Theorem 16). *Let \mathfrak{F}_1^∂ and \mathfrak{F}_2^∂ be two strongly BFV-equivalent (exact) theories, then $\mathfrak{F}^{\text{AKSZ}}(I; \mathfrak{F}_1^\partial)$ and $\mathfrak{F}^{\text{AKSZ}}(I; \mathfrak{F}_2^\partial)$ are strongly BV-equivalent.*

Proof. A strong BV equivalence induces an isomorphism of the underlying dg-manifolds. \square

2. BFV THEORIES OF GRAVITY

2.1. BFV Einstein–Hilbert Theory. The BFV theory for GR in the Einstein–Hilbert formalism (as described in [CS16]) associates to any $(N - 1)$ -dimensional (pseudo-)Riemannian¹² manifold Σ the graded 0-symplectic manifold

$$\mathcal{F}_{EH}^\partial(\Sigma) = T^* \underbrace{(S_{nd}^2(T\Sigma) \times \mathfrak{X}[1](\Sigma) \times C^\infty[1](\Sigma))}_{\{\gamma, \xi^\partial, \xi^n\}}, \quad (10)$$

where $S_{nd}^2(\Sigma)$ denotes the space of nondegenerate symmetric tensor fields of rank two, with canonical exact symplectic structure:

$$\varpi_{EH}^\partial(\Sigma) = \delta\alpha_{EH}^\partial(\Sigma) = \delta \int_\Sigma \langle \Pi, \delta\gamma \rangle + \langle \varphi_\partial, \delta\xi^\partial \rangle + \langle \varphi_n, \delta\xi^n \rangle, \quad (11)$$

and $\{\Pi, \varphi_\partial, \varphi_n\}$ denote variables in the cotangent fiber, dual to $\{\gamma, \xi^\partial, \xi^n\}$ respectively, i.e.

$$\begin{aligned} \Pi &\in S^2(T^*\Sigma) \otimes \text{Dens}(\Sigma), \\ \varphi_\partial &\in \Omega^1(\Sigma) \otimes \text{Dens}(\Sigma), \\ \varphi_n &\in C^\infty(\Sigma) \otimes \text{Dens}(\Sigma). \end{aligned}$$

Remark 20. The components $(\gamma)^{ab}$ of a $\gamma \in S_{nd}^2(T\Sigma)$ can be thought of as the inverse of a (pseudo-)Riemannian metric on Σ , which we denote by γ^{-1} . With a slight abuse of notation¹³ we will denote by $\sqrt{\gamma} = \sqrt{\det(\gamma_{ab})}$ the square root of the

¹²In this paper we will mostly focus on the case where Σ is a Riemannian manifold, seen as a spacelike boundary of a cylinder $\Sigma \times \mathbb{R}$. Generalisations of this to the timelike case are straightforward. The relevant BFV data can be found in [CS16].

¹³This is not really problematic, since its variation reads $\delta\sqrt{\gamma} = \frac{1}{2}\sqrt{\gamma}\gamma^{ab}\delta\gamma_{ab} = -\frac{1}{2}\sqrt{\gamma}\gamma_{ab}\delta\gamma^{ab}$ and we can use either formula according to our needs. If we wanted to use the correct notation we should simply replace $\sqrt{\gamma}$ with its reciprocal, in formula (13).

determinant of the metric on Σ that we denote by γ^{-1} everywhere else. In other words, $\sqrt{\gamma}$ is the usual density associated to a metric, i.e. $\sqrt{\gamma}d^{N-1}x$ is a volume form on Σ . Observe that all fields in the fibres of the cotangent bundle (10) are sections of the respective dual bundles, tensored with densities. The conjugate field to γ is a section of the second symmetric tensor power of the cotangent bundle of Σ tensored with densities on Σ , i.e. $\Pi \in S^2(T^*\Sigma) \otimes \text{Dens}(\Sigma)$ is of the form $\Pi = \sqrt{\gamma}\pi$, for $\pi \in S^2(T^*\Sigma)$. A similar decomposition holds for $\varphi_\partial, \varphi_n$.

In addition to $\mathcal{F}_{EH}^\partial(\Sigma)$ and $\varpi_{EH}^\partial(\Sigma)$, the BV-BFV procedure outputs a functional of degree 1 on $\mathcal{F}_{EH}^\partial(\Sigma)$, called BFV action. It is given by the local expression

$$S_{EH}^\partial(\Sigma) = \int_{\Sigma} \left\{ H_n \xi^n + \langle \Pi, L_{\xi^\partial} \gamma \rangle + \varphi_n L_{\xi^\partial} \xi^n - \gamma(\varphi_\partial, d\xi^n) \xi^n + \left\langle \varphi_\partial, \frac{1}{2} [\xi^\partial, \xi^\partial] \right\rangle \right\} \quad (12)$$

where we have denoted the *Hamiltonian constraint density* by

$$H_n(\gamma, \Pi) = \left(\frac{1}{\sqrt{\gamma}} \left(\text{Tr}_\gamma[\Pi^2] - \frac{1}{d-1} \text{Tr}_\gamma[\Pi]^2 \right) + \sqrt{\gamma} (R^\partial - 2\Lambda) \right) \quad (13)$$

with R^∂ is the trace of the Ricci tensor with respect to the metric γ^{-1} , $\Lambda \in \mathbb{R}$ is the cosmological constant, $\text{Tr}_\gamma[\Pi^2] = \gamma^{ab}\gamma^{cd}\Pi_{bc}\Pi_{ad}$ and $\text{Tr}_\gamma\Pi$ is the density $\gamma^{ab}\Pi_{ab}$. Observe that we can also denote the *momentum constraint density* as the density-valued one-form

$$H_\partial: \mathfrak{X}(\Sigma) \rightarrow \text{Dens}(\Sigma) \quad H_\partial(X) = \langle \Pi, L_X \gamma \rangle \quad (14)$$

for $X \in \mathfrak{X}(\Sigma)$.

Remark 21. One can integrate the density of equation (13) against a function $\lambda \in C^\infty(\Sigma)$, or integrate the density in (14) to get local functionals on fields. Then λ and X play the role of Lagrange multipliers, to enforce the so-called Hamiltonian and momentum constraints.

The Hamiltonian vector field $Q_{EH}^\partial(\Sigma)$ of $S_{EH}^\partial(\Sigma)$ with respect to $\varpi_{EH}^\partial(\Sigma)$ is a differential on $C^\infty(\mathcal{F}_{EH}^\partial(\Sigma))$, the BFV differential, and its cohomology in degree zero is the reduced phase space defined by the constraints $\{H_n, H_\partial\}$.

Definition 22. We define BFV Einstein–Hilbert theory associated to be the assignment

$$\Sigma \rightsquigarrow \mathfrak{F}_{EH}^\partial(\Sigma) = (\mathcal{F}_{EH}^\partial(\Sigma), S_{EH}^\partial(\Sigma), \varpi_{EH}^\partial(\Sigma), Q_{EH}^\partial(\Sigma)). \quad (15)$$

2.2. BFV Palatini–Cartan theory. Let Σ be an $(N-1)$ -dimensional compact and orientable¹⁴ smooth manifold and let $P \rightarrow \Sigma$ be an $SO(N-1, 1)$ -principal bundle. Let also \mathcal{V} be the associated vector bundle where each fibre is isomorphic to (V, η) , an N -dimensional vector space with a pseudo-Riemannian inner product η on it. We further identify $\mathfrak{so}(N-1, 1) \cong \bigwedge^2 \mathcal{V}$ using η .

Furthermore we use the following notation:

$$\Omega_\partial^{i,j} := \Omega^i \left(\Sigma, \bigwedge^j \mathcal{V} \right).$$

The BFV data for PC theory has been described in [CCS20] for $N \geq 4$ and in [CS19a] for $N = 3$, the following discussion will be nontrivial for $N \geq 4$, see Remark 28. The classical fields of the theory are then $e \in \Omega_{nd}^1(\Sigma, \mathcal{V})$ — i.e. $\Omega_\partial^{1,1}$ plus the nondegeneracy condition that the induced morphism $T\Sigma \rightarrow \mathcal{V}$ should be injective

¹⁴For simplicity we orient Σ and V , but the formalism generalizes to nonorientable Σ as well, see [CCS20].

— and the equivalence class of a connection $\omega \in \mathcal{A}(\Sigma)$ under the e -dependent relation $\omega \sim \omega + v$ for v such that $W_{e^{N-3}}^{1,1}(v) = 0$, where

$$W_{e^{N-3}}^{1,1} : \Omega_{\partial}^{1,1} \rightarrow \Omega_{\partial}^{2,2}, \quad W_{e^{N-3}}^{1,1}(v) = e^{N-3} \wedge v.$$

We denote this equivalence class and the quotient space it belongs to by $[\omega] \in \mathcal{A}^{\text{red}}(\Sigma)$. We further assume that the boundary metric

$$g_{ij}^{\partial} := \eta(e_i, e_j) \quad (16)$$

is nondegenerate.

The symplectic manifold F_{PC}^{∂} of (degree-0) boundary fields for PC theory is then the total space of the fibre bundle $F_{PC}^{\partial} \rightarrow \Omega_{\text{nd}}^1(\Sigma, \mathcal{V})$, with fiber \mathcal{A}^{red} . The manifold F_{PC}^{∂} is obtained as the symplectic reduction of the naive boundary two-form $\tilde{\omega} = \int e^{N-3} \delta e \delta \omega$, which is pre-symplectic since $\ker(\tilde{\omega}) \simeq \ker(W_{e^{N-3}}^{1,2}) \neq \{0\}$, as described in [CS19c]. Instead of working with equivalence classes of connections, it is convenient to fix a nonvanishing section $\epsilon_n \in \Gamma(\mathcal{V})$ and enforce a condition called *structural constraint*, which was introduced in [CCS20]:

$$(N-3)\epsilon_n e^{N-4} d_{\omega} e \in \text{Im} W_{e^{N-3}}^{1,1}, \quad (17)$$

in the space of boundary tetrads and connections. In order to do this one restricts to tetrads e that are linearly independent from ϵ_n . In general this implies working in patches over the space of the e -fields. However, if g^{∂} is space-like, we may choose once and for all ϵ_n to be time-like, which provides a global choice on the space of the e -fields.

Remark 23. Considering the boundary by itself, the constraint (17) is one of the possible ways to fix the the transformations in the kernel of the presymplectic form. If we take the bulk as well into account, it assumes a more fundamental role: it is the necessary and sufficient condition for the transversal equations of motions to be solvable. Indeed, the (bulk) equations of motion split into a *tangential* equation

$$e^{N-3} d_{\omega} e = 0$$

and a transversal one

$$(N-3)e_n e^{N-4} d_{\omega} e = e^{N-3} (d_{\omega} e)_n,$$

which tells us that $e_n e^{N-4} d_{\omega} e$ must be in the image of $W_{e^{N-3}}^{1,1}$. Then, upon using the tangential equation, we can replace e_n with some fixed ϵ_n . See Section 4.2.

Denoting by $S \subset \Omega_{\text{nd}}^1(\Sigma, \mathcal{V}) \times \mathcal{A}(\Sigma)$ the submanifold defined by Equation (17), we have:

Proposition 24 ([CCS20; CS19c]). *There exists a symplectomorphism*

$$F_{PC}^{\partial}(\Sigma) \rightarrow S.$$

Effectively, then, one can work on S . The main advantage of this explicit description of the symplectic space of boundary fields is that it allows to explicitly compute the *symplectic* BFV data for PC theory (see Remark 29).

In order to write down the BFV data it is sufficient to fix the equivalence class $[\omega] \in \mathcal{A}^{\text{red}}(\Sigma)$ using (17) as done in [CCS20], however, when extending F_{PC}^{∂} to a graded manifold we can choose to modify the structural constraint by adding terms in the ghosts and antifields. This will turn out to be convenient in what follows.

Definition 25. Consider the graded manifold

$$\check{\mathcal{F}}_{PC}(\Sigma) := \Omega_{\text{nd}}^{1,1}(\Sigma, \mathcal{V}) \times \mathcal{A}(\Sigma) \times T^* \left(\Omega_{\partial}^{0,2}[1] \oplus \mathfrak{X}[1](\Sigma) \oplus C^{\infty}[1](\Sigma) \right), \quad (18)$$

where we denote fields by

- $e \in \Omega_{\text{nd}}^{1,1}(\Sigma, \mathcal{V})$ and $\omega \in \mathcal{A}(\Sigma)$ in degree zero,
- $c \in \Omega_{\partial}^{0,2}[1]$, $\xi \in \mathfrak{X}[1](\Sigma)$ and $\lambda \in \Omega^{0,0}[1]$ in degree one,
- $c^\dagger \in \Omega_{\partial}^{3,2}[-1]$, $\lambda^\dagger \in \Omega_{\partial}^{3,4}[-1]$ and $\xi^\dagger \in \Omega_{\partial}^{1,0}[-1] \otimes \Omega_{\partial}^{3,4}$ in degree minus one,

together with a fixed section $\epsilon_n \in \Gamma(M, \mathcal{V})$, completing the image of e to a basis of \mathcal{V} . We define the *BFV structural constraint* to be condition¹⁵:

$$(N-3)\epsilon_n e^{N-4} d_\omega e + (L_\xi^\omega \epsilon_n - [c, \epsilon_n])^{(a)} c_a^\dagger \in \text{Im} W_{e^{N-3}}^{1,1} \quad (19)$$

where $\{(a), (n)\}$ denote components with respect to a basis $\{e_a, \epsilon_n\}$. We define the *space of BFV fields* $\mathcal{F}_{PC}^\partial(\Sigma)$ to be the space of solutions of the BFV structural constraints within $\tilde{\mathcal{F}}_{PC}(\Sigma)$.

In order to have a better expression of the BFV structure, following [CCS20, Section 5.2], we introduce the field $y^\dagger \in \Omega_{\partial}^{3,3}[-1]$ such that the original fields λ^\dagger and ξ^\dagger are recovered through $\epsilon_n y^\dagger = -\lambda^\dagger$ and $e_a y^\dagger = -\xi_a^\dagger$. This also allows us to write a single expression for all $N \geq 3$.

To complete the definition of the BFV data for Palatini–Cartan theory we consider a degree 1 functional and a symplectic form¹⁶ given, respectively, by:

$$\begin{aligned} S_{PC}^\partial(\Sigma) &= \int_{\Sigma} c e^{N-3} d_\omega e + \iota_\xi e e^{N-3} F_\omega + \epsilon_n \lambda e^{N-3} F_\omega + \frac{1}{2} [c, c] c^\dagger - L_\xi^\omega c c^\dagger \\ &\quad + \frac{1}{2} \iota_\xi \iota_\xi F_\omega c^\dagger - [c, \epsilon_n \lambda] y^\dagger + L_\xi^\omega (\epsilon_n \lambda) y^\dagger + \frac{1}{2} \iota_{[\xi, \xi]} e y^\dagger, \end{aligned} \quad (20)$$

$$\varpi_{PC}^\partial(\Sigma) = \int_{\Sigma} e^{N-3} \delta e \delta \omega + \delta c \delta c^\dagger + \delta \omega \delta (\iota_\xi c^\dagger) - \delta \lambda \epsilon_n \delta y^\dagger + \iota_{\delta \xi} \delta (e y^\dagger). \quad (21)$$

Note that each term of the integrals belongs to $\Omega^{N-1, N}$, which can be canonically identified, using $\sqrt{|\det \eta|}$, with the space of densities on Σ . A detailed explanation can be found in [CCS20]. However, we will not write down the factor $\sqrt{|\det \eta|}$ explicitly.

Definition 26. We define BFV Palatini–Cartan theory to be the assignment

$$\Sigma \rightsquigarrow \mathfrak{F}_{PC}^\partial(\Sigma) = (\mathcal{F}_{PC}^\partial(\Sigma), S_{PC}^\partial(\Sigma), \varpi_{PC}^\partial(\Sigma), Q_{PC}^\partial(\Sigma)). \quad (22)$$

with $Q_{PC}^\partial(\Sigma)$ the Hamiltonian vector field of $S_{PC}^\partial(\Sigma)$ with respect to $\varpi_{PC}^\partial(\Sigma)$.

Remark 27. Notice that the data presented in Definition 26 are equivalent to the BFV data presented in [CCS20]. The cohomological vector field of a function on (the presymplectic manifold) $\tilde{\mathcal{F}}_{PC}(\Sigma)$ is uniquely fixed by the tangency to the BFV structural constraint (19). In [CCS20] the equivalent choice of the constraint (17) is made, and the resulting Q 's differ along ω by a term in the kernel of $W_{e^{N-3}}^{1,2}$.¹⁷ We make this choice in this paper because it makes it easier to compare the AKSZ construction for the constrained space $\mathcal{F}_{PC}^\partial(\Sigma)$ with a constrained version of PC, see Remark 28.

¹⁵This BFV structural constraint differs from the classical one of equation (17) by a term depending on ghosts and their antifields, so it reduces to the classical structural constraint if we restrict it to the classical space of boundary fields (the body of $\tilde{\mathcal{F}}_{PC}(\Sigma)$).

¹⁶This version of the BFV data features a particularly simple action functional, at the price of not expressing the symplectic form in its Darboux form. An alternative can be found in [CCS20].

¹⁷Also note that this term vanishes on the body of $\tilde{\mathcal{F}}_{PC}$ when imposing the classical constraints, i.e., the expressions paired to the ghosts in the first three terms of (20). The two BFV constructions, by (17) or by (19), describe the characteristic distribution of the coisotropic submanifold determined by the constraints. They extend this distribution in a different way outside of it, but this is irrelevant.

Remark 28. In [CS19b] two of the authors showed that the natural BV data for PC theory (in dimension $N \geq 4$) cannot be extended to a BV-BFV theory (Definition 5). Since the degree-0 part of our AKSZ target will be the constrained space $\mathcal{F}_{PC}^\partial(\Sigma)$, the extension to the cylinder¹⁸ of Equation (19) provides an explicit realisation of (one of) the conditions that must be imposed on the bulk BV fields, in order to have a 1-extended BV-BFV theory. For $N = 3$, the additional condition imposed by (17) is void, explaining why 3d PC theory can be (fully) extended without additional requirements on the fields [CS19a]. We will comment further on this in Section 4.2.

Remark 29. The main difficulty in constructing BFV data for PC theory comes from the requirement that $(\mathcal{F}_{PC}^\partial, \varpi_{PC}^\partial)$ be a symplectic manifold, as opposed to pre-symplectic (cf. with [AG14] and [Gri16]). We stress that this requirement is essential for quantisation. To the best of our knowledge, a complete description of the *symplectic* BFV structure for PC gravity was not available before [CCS20]. The complexity of the symplectic reduction arising in the classical description of the degree-0 boundary structure [CCS20; CS19c], as well as the obstruction in the BV-BFV induction procedure [CS19b], are relevant features peculiar to this formulation of gravity.

Remark 30. The conventions we choose for the fields in (20) and in (21) differ from those in [CS19a, Proposition 21]. In order to make contact between the formulas one has to perform the following change of variables:

$$\begin{aligned} e^\dagger &\mapsto y^\dagger & \omega^\dagger &\mapsto c^\dagger \\ c &\mapsto -c & \xi &\mapsto -\xi & \xi^n &\mapsto -\lambda. \end{aligned}$$

In the case $N \geq 4$ some signs differ from the ones in [CCS20] due to a sign convention for λ .

3. AKSZ EH

We explore here the idea of reconstructing the $(d+1)$ -dimensional BV extension of Einstein–Hilbert theory by means of the AKSZ construction, with target the BFV data for Einstein–Hilbert theory (as presented in Section 2.1, based on [CS16]). In order to do this one looks at the space $\mathcal{F}_{EH}^{\text{AKSZ}} := \text{Maps}(T[1]I, \mathcal{F}_{EH}^\partial(\Sigma))$, with I an interval. In a chart, to consider the transgression map of Equation (13) means to look at fields composed of a 0-form and a 1-form on the interval I , with values in $\mathcal{F}_{EH}^\partial(\Sigma)$ and fixed total degree. For the case at hand we will then have

$$\mathfrak{G} = \gamma(t) + \Pi^\dagger(t)dt \quad \mathfrak{Z}^n = \xi^n(t) + \eta(t)dt \quad \mathfrak{Z}^a = \xi^a(t) + \beta^a(t)dt \quad (23a)$$

$$\mathfrak{P} = \Pi(t) + \gamma^\dagger(t)dt \quad \mathfrak{H}_n = \varphi_n(t) + \xi_n^\dagger(t)dt \quad \mathfrak{H}_a = \varphi_a(t) + \xi_a^\dagger(t)dt \quad (23b)$$

where, for all $t \in I$, we parametrise $\mathcal{F}_{EH}^{\text{AKSZ}}$ with fields¹⁹

$$\begin{aligned} \gamma(t) &\in \text{Map}(I, S_{nd}^2(T\Sigma)), & \gamma^\dagger(t) &\in \text{Map}(I, S^2[-1](T^*\Sigma)), \\ \Pi(t) &\in \text{Map}(I, S^2(T^*\Sigma)), & \Pi^\dagger(t) &\in \text{Map}(I, S[-1]^2(T\Sigma)), \\ \eta(t), \beta^a(t) &\in \text{Map}(I, C^\infty(\Sigma)), & \varphi_n(t), \varphi_a(t) &\in \text{Map}(I, \text{Dens}[-1](\Sigma)), \\ \xi^n(t), \xi^a(t) &\in \text{Map}(I, C^\infty[1](\Sigma)), & \xi_n^\dagger(t), \xi_a^\dagger(t) &\in \text{Map}(I, \text{Dens}[-2](\Sigma)), \end{aligned}$$

where we required $\gamma(t)$ to be nondegenerate for all $t \in I$. Now, observe that η, ξ^n are functions on Σ whereas β^a, ξ^a can be considered as the components of vector (fields) tangent to Σ , which we will denote by β and ξ^∂ . Similarly, we can promote φ_a and

¹⁸Equation (19) can be extended to the space of fields over a cylinder. We consider this point of view in Definition 58.

¹⁹The motivation for this particular choice of notation will be manifest very soon.

ξ_a^\dagger into Σ -density valued one forms, which we will denote by $\varphi_\partial, \xi_\partial^\dagger$. For simplicity of notation we will often use a unified index $\rho \in \{n, a\}$, so that $\varphi_\rho \in \{\varphi_n, \varphi_a\}$ and $\xi_\rho^\dagger \in \{\xi_n^\dagger, \xi_a^\dagger\}$.

Remark 31. Notice once again that we are using (nondegenerate) sections of $S^2(TM)$ instead of actual metrics. Occasionally we will need to raise/lower indices using γ and its “inverse” which we will denote by γ^{-1} . See Remark 20.

In what follows it will be useful to denote the *Kinetic* part of the Hamiltonian functional (Equation (13)) as

$$\mathcal{K} := \frac{1}{\sqrt{\gamma}} \left(\text{Tr}_\gamma[\Pi^2] - \frac{1}{d-1} \text{Tr}_\gamma[\Pi]^2 \right), \quad (24)$$

and the *cosmological Einstein tensor* with respect to a metric γ^{-1} with cosmological constant Λ will be

$$\mathbf{G}[\gamma, \Lambda] = R[\gamma] + \left(\Lambda - \frac{1}{2} \text{Tr}_\gamma R[\gamma] \right) \gamma^{-1}, \quad (25)$$

where $R[\gamma]$ is the Ricci–Riemann tensor of γ . We also introduce a tensor-valued second order operator \mathbf{D}_γ that on functions $\phi \in C^\infty(\Sigma)$ acts as

$$\mathbf{D}_\gamma \phi = \gamma^{-1} \Delta^\partial \phi - \nabla^\partial \odot \nabla^\partial \phi \quad (26)$$

where ∇^∂ denotes the Levi-Civita connection of γ , and we denoted by $\Delta^\partial = \nabla^\partial \cdot \nabla^\partial$ the Laplace-Beltrami operator. In a coordinate chart this reads:

$$[\mathbf{D}_\gamma \phi]_{ab} = \gamma_{ab} \gamma^{cd} \nabla_c^\partial \nabla_d^\partial \phi - \nabla_a^\partial \nabla_b^\partial \phi$$

In what follows we will also use the metric gradient, i.e. the vector field $\text{grad}_\gamma \phi = \gamma(d\phi, \cdot)$. Since the covariant derivative ∇^∂ will not explicitly appear in what follows, we shall employ the symbol ∇ to introduce a shorthand notation for the metric gradient:

$$\nabla_\gamma \phi \equiv \frac{1}{2} \text{grad}_\gamma \phi.$$

Then, it is a matter of a straightforward calculation to show that

$$\mathbf{D}_\gamma \phi = \frac{1}{2} \left(\gamma^{-1} \text{Tr}[\mathcal{L}_{\nabla_\gamma \phi} \gamma] - \mathcal{L}_{\nabla_\gamma \phi} \gamma^{-1} \right).$$

With this in mind we can state the following:

Theorem 32. *The AKSZ data $\mathfrak{F}_{EH}^{\text{AKSZ}}(I; \mathfrak{F}_{EH}^\partial(\Sigma))$ are given by the (-1) -shifted symplectic manifold*

$$\begin{aligned} \mathcal{F}_{EH}^{\text{AKSZ}} &\simeq T^*[-1] \left(\text{Map}(I, S_{nd}^2(T\Sigma) \times S^2(T^*\Sigma) \times C^\infty(\Sigma) \times \mathfrak{X}(\Sigma) \times \mathfrak{X}[1](\Sigma) \times C^\infty[1](\Sigma)) \right) \\ \Omega_{EH}^{\text{AKSZ}} &= \int_{\Sigma \times I} \left\{ -\langle \delta\gamma, \delta\gamma^\dagger \rangle + \langle \delta\Pi, \delta\Pi^\dagger \rangle + \delta\xi^\rho \delta\xi_\rho^\dagger + \delta\eta \delta\varphi_n + \delta\beta^a \delta\varphi_a \right\} dt \end{aligned} \quad (27)$$

and the AKSZ action functional :

$$S_{EH}^{\text{AKSZ}} = \int_{\Sigma \times I} \left\{ \langle \Pi, \dot{\gamma} \rangle - \langle \varphi_\rho, \dot{\xi}^\rho \rangle + H_n \eta + H_\partial(\beta) - \langle \gamma^\dagger, L_{\xi^\partial} \gamma \rangle + \langle \Pi^\dagger, L_{\xi^\partial} \Pi \rangle \right\} dt \quad (28a)$$

$$- \left(\frac{\delta \mathcal{K}}{\delta \gamma}(\Pi^\dagger) + \frac{\delta \mathcal{K}}{\delta \Pi}(\gamma^\dagger) \right) \xi^n - \sqrt{\gamma} \langle \Pi^\dagger, \mathbf{G}[\gamma, \lambda] \xi^n + \mathbf{D}_\gamma(\xi^n) \rangle \quad (28b)$$

$$+ \langle \varphi_\partial, \nabla_\gamma \eta \xi^n - \eta \nabla_\gamma \xi^n + L_{\xi^\partial} \beta \rangle - \varphi_n L_\beta \xi^n + \varphi_n L_{\xi^\partial} \eta \quad (28c)$$

$$+ \left\langle \xi_\partial^\dagger, \frac{1}{2} [\xi^\partial, \xi^\partial] + \xi^n \nabla_\gamma \xi^n \right\rangle + \Pi^\dagger(\varphi_\partial, d\xi^n) \xi^n + \xi_n^\dagger L_{\xi^\partial} \xi^n \Big\} dt, \quad (28d)$$

together with its Hamiltonian vector field Q_{EH}^{AKSZ} .

Proof. The prescription of Theorem 14, suggests that to construct the data in $\mathfrak{F}^{\text{AKSZ}}(I; \mathfrak{F}_{EH}^\partial)$ we need to compute

$$\Omega_{EH}^{\text{AKSZ}} = \mathfrak{T}_I^{(2)} \varpi_{EH}^\partial(\Sigma) = \int_{\Sigma \times I} \langle \delta \mathfrak{P}, \delta \mathfrak{G} \rangle + \langle \delta \mathfrak{H}_\rho, \delta \mathfrak{Z}^\rho \rangle. \quad (29)$$

By selecting the top-form part of the integrand and observing that $|dt| = 1$ we get

$$\Omega_{EH}^{\text{AKSZ}} = \int_{\Sigma \times I} \{ -\langle \delta \gamma^\dagger, \delta \gamma \rangle + \langle \delta \Pi, \delta \Pi^\dagger \rangle + \delta \xi_\rho^\dagger \delta \xi^\rho + \delta \varphi_n \delta \eta + \delta \varphi_a \delta \beta^a \} dt \quad (30)$$

where the sign comes from $\langle \delta(\gamma^\dagger dt), \delta \gamma \rangle = -\langle \delta \gamma^\dagger, \delta \gamma \rangle dt$, since $|\delta \gamma| = 1$, while $\delta \xi_\rho^\dagger dt \delta \xi^\rho = \delta \xi_\rho^\dagger \delta \xi^\rho dt$, since $|\delta \xi^\rho| = 2$. Ω^{AKSZ} is a (-1) -symplectic structure on $\text{Maps}(T[1]I, \mathcal{F}_{EH}^\partial(\Sigma))$, a BV 2-form.

Now, from $\alpha_{EH}^\partial(\Sigma)$ we can construct a degree-0 functional on $\mathcal{F}_{EH}^{\text{AKSZ}}$ by first applying the transgression map, which yields the 1-form

$$\mathfrak{T}_I^{(1)} \alpha^\partial \in \Omega^1(\text{Maps}(T[1]I, \mathcal{F}_{EH}^\partial(\Sigma))),$$

and then contracting it with the de Rham differential on I seen as an odd cohomological vector field d_I . In a local chart this is tantamount to replacing $\delta \rightsquigarrow d_I := dt \frac{d}{dt}$, so that

$$\iota_{d_I} \mathfrak{T}_I^{(1)} \alpha_{EH}^\partial(\Sigma) = \int_{\Sigma \times I} \{ \langle \Pi, \dot{\gamma} \rangle - \langle \varphi_\rho, \dot{\xi}^\rho \rangle \} dt. \quad (31)$$

where the sign comes from the fact that $\langle \varphi_\rho dt, \dot{\xi}^\rho \rangle = -\langle \varphi_\rho, \dot{\xi}^\rho \rangle dt$. Finally, we want to compute the AKSZ action functional

$$S_{EH}^{\text{AKSZ}} := \mathfrak{T}_I^{(0)}(S_{EH}^\partial(\Sigma)) + \iota_{d_I} \mathfrak{T}_I^{(1)}(\alpha_{EH}^\partial(\Sigma)).$$

This calculation is completely analogous to the previous ones, and it is mostly straightforward. One needs to pay attention to the signs, so it is worthwhile to stress that

$$\Pi^\dagger dt (\varphi_\partial, d\xi^n) \xi^n = [\Pi^\dagger]^{ab} dt \varphi_a \partial_b \xi^n \xi^n = [\Pi^\dagger]^{ab} \varphi_a \partial_b \xi^n \xi^n dt = \Pi^\dagger (\varphi_\partial, d\xi^n) \xi^n dt$$

while

$$-\gamma (\xi_\partial^\dagger dt, d\xi^n) \xi^n - \gamma (\varphi_\partial, d\eta dt \xi^n + d\xi^n \eta dt) = \left(\langle \xi_\partial^\dagger, \xi^n \nabla_\gamma \xi^n \rangle + \langle \varphi_\partial, \nabla_\gamma \eta \xi^n - \eta \nabla_\gamma \xi^n \rangle \right) dt$$

Finally, at first order in dt ,

$$H_n(\gamma + \Pi^\dagger dt, \Pi + \gamma^\dagger) \xi^n = \frac{\delta(H_n \xi^n)}{\delta \gamma} (\Pi^\dagger dt) + \frac{\delta(H_n \xi^n)}{\delta \Pi} (\gamma^\dagger dt),$$

and $dt \xi^n = -\xi^n dt$. We write the formulas above as derivatives of the functional $H_n \xi^n$ to stress that total derivatives will appear, due to the term R^∂ in H_n . Recalling the expression for H_n of equation (13) and the definition of \mathcal{K} , $\mathbf{G}[\gamma, \Lambda]$ and \mathbf{D}_γ of Equations (24), (25) and (26), the variation of $H_n \xi^n$ with respect to γ yields

$$\frac{\delta(H_n \xi^n)}{\delta \gamma} = \frac{\delta \mathcal{K}}{\delta \gamma} \xi^n + \frac{\delta(\sqrt{\gamma} R^\partial \xi^n)}{\delta \gamma} = \frac{\delta \mathcal{K}}{\delta \gamma} \xi^n + \sqrt{\gamma} (\mathbf{G}[\gamma, \Lambda] \xi^n + \mathbf{D}_\gamma(\xi^n)) + d(\dots).$$

The total derivative term is exact with respect to the tangent differential d_Σ . It can be discarded, provided Σ has no boundary (which we are assuming throughout), so:

$$\begin{aligned} H_n(\gamma + \Pi^\dagger dt, \Pi + \gamma^\dagger) \xi^n = \\ - \left(\frac{\delta \mathcal{K}}{\delta \Pi} (\gamma^\dagger) + \frac{\delta \mathcal{K}}{\delta \gamma} (\Pi^\dagger) \right) \xi^n dt - \sqrt{\gamma} \langle \Pi^\dagger, \mathbf{G}[\gamma, \Lambda] \xi^n + \mathbf{D}_\gamma(\xi^n) \rangle dt. \end{aligned} \quad (32)$$

□

Remark 33. In order to compute the cohomological vector field Q_{EH}^{AKSZ} we enforce the Hamiltonian condition $\iota_{Q_{EH}^{\text{AKSZ}}}\Omega_{EH}^{\text{AKSZ}} = \delta S_{EH}^{\text{AKSZ}}$ dropping all possible boundary terms. It reads (we omit the expression for $Q_{EH}^{\text{AKSZ}}\xi^\dagger$ and $Q_{EH}^{\text{AKSZ}}\varphi$):

$$Q_{EH}^{\text{AKSZ}}\gamma = \frac{\delta H_n}{\delta \Pi}\xi^n + L_{\xi^\partial}\gamma \quad (33a)$$

$$Q_{EH}^{\text{AKSZ}}\Pi = -\frac{\delta \mathcal{K}}{\delta \gamma}\xi^n - \sqrt{\gamma}(\mathbf{G}[\gamma, \Lambda]\xi^n + \mathbf{D}_\gamma(\xi^n)) + L_{\xi^\partial}\Pi + \varphi \odot d\xi^n \xi^n \quad (33b)$$

$$Q_{EH}^{\text{AKSZ}}\eta = -\dot{\xi}^n + L_{\xi^\partial}\eta - L_\beta\xi^n \quad (33c)$$

$$Q_{EH}^{\text{AKSZ}}\beta = -\dot{\xi}^\partial + L_{\xi^\partial}\beta + \nabla_\gamma\eta\xi^n - \eta\nabla_\gamma\xi^n - \nabla_{\Pi^\dagger}\xi^n\xi^n \quad (33d)$$

$$Q_{EH}^{\text{AKSZ}}\xi^\partial = \frac{1}{2}[\xi^\partial, \xi^\partial] + \xi^n\nabla_\gamma\xi^n \quad (33e)$$

$$Q_{EH}^{\text{AKSZ}}\xi^n = L_{\xi^\partial}\xi^n \quad (33f)$$

$$Q_{EH}^{\text{AKSZ}}\gamma^\dagger = \dot{\Pi} + \frac{\delta \mathcal{K}}{\delta \gamma}\eta + \sqrt{\gamma}(\mathbf{G}[\gamma, \Lambda]\eta + \mathbf{D}_\gamma(\eta)) + L_\beta\Pi + L_{\xi^\partial}\gamma^\dagger \quad (33g)$$

$$+ \xi_\partial^\dagger \odot d\xi^n \xi^n - \varphi_\partial \odot d\eta\xi^n + \eta\varphi_\partial \odot d\xi^n \quad (33h)$$

$$+ \left[\frac{\delta^2 \mathcal{K}}{\delta \gamma^2}(\Pi^\dagger) + \frac{\delta^2 H_n}{\delta \gamma \delta \Pi}(\gamma^\dagger) \right] \xi^n - \frac{1}{2}\gamma^{-1}\langle \Pi^\dagger, \mathbf{G}[\gamma, \lambda]\xi^n + \mathbf{D}_\gamma(\xi^n) \rangle \quad (33i)$$

$$- \sqrt{\gamma} \left\langle \Pi^\dagger, \frac{\delta \mathbf{G}[\gamma, \lambda]}{\delta \gamma}\xi^n + \frac{\delta \mathbf{D}_\gamma(\xi^n)}{\delta \gamma} \right\rangle \quad (33j)$$

$$Q_{EH}^{\text{AKSZ}}\Pi^\dagger = \dot{\gamma} + \frac{\delta \mathcal{K}}{\delta \Pi}\eta + L_\beta\gamma + L_{\xi^\partial}\Pi^\dagger - \left[\frac{\delta^2 \mathcal{K}}{\delta \Pi^2}(\gamma^\dagger) + \frac{\delta^2 \mathcal{K}}{\delta \gamma \delta \Pi}(\Pi^\dagger) \right] \xi^n. \quad (33k)$$

Remark 34. We notice that the term $\sqrt{\gamma}\mathbf{D}_\gamma(\cdot)$ is the contribution to the field equations for a metric due to the presence of a Brans–Dicke “dilaton” field, whose role is played by the ghost ξ^n in the BVF action $S_{EH}^\partial(\Sigma)$ and by η in S_{EH}^{AKSZ} .

3.1. Pushforward. We would like to compute the BV pushforward of $\mathfrak{F}_{EH}^{\text{AKSZ}}$ along the symplectic submanifold $(\Pi, \Pi^\dagger) \in \mathcal{F}'' = T^*[-1]\text{Map}(I, S^2(T^*\Sigma)) \subset \mathcal{F}_{EH}^{\text{AKSZ}}$.

Remark 35. This is the same as evaluating S_{eff} from equation (3). Since S_{EH}^{AKSZ} is only quadratic in Π , the calculation reduces to computing the Batalin–Vilkovisky–Legendre transform S_{BVL} of S_{EH}^{AKSZ} with respect to \mathcal{L} , as in Definition 11, plus a correction in the integration measure for the remaining (second-order) effective theory. Note that Equation 2 is equivalent to setting to zero the r.h.s. of Equation (33k), together with $\Pi^\dagger = 0$.

Recall that we are assuming $\gamma(t)$ to be a nondegenerate section of $S^2(T\Sigma)$ for every t , i.e. it represents the inverse of a metric, and dually $\Pi(t) \in S^2(T^*\Sigma)$. We can use γ and its inverse (denoted γ^b) to raise/lower indices: explicitly, if $\gamma = \gamma^{ab}\partial_a \odot \partial_b$, we have $\gamma^b = \gamma_{ab}dx^a \odot dx^b$, with $\gamma^{ab}\gamma_{bc} = \delta_c^a$. Then, for $X \in S^2(T^*\Sigma)$, $Y \in S^2(T\Sigma)$ we define $(X^\sharp)^{ab} := \gamma^{ac}\gamma^{bd}X_{bc}$ and $(Y^b)_{ab} = \gamma_{ac}\gamma_{bd}Y^{cd}$.

Definition 36. Consider the space of fields $\mathcal{F}_R(\Sigma \times I) \subseteq \mathcal{F}_{EH}^{\text{AKSZ}}$ as

$$\mathcal{F}_R(\Sigma \times I) := T^*[-1] \left(\text{Map} \left(I, S_{nd}^2(T\Sigma) \times T[1](C^\infty(\Sigma) \times \mathfrak{X}(\Sigma)) \right) \right) \quad (34)$$

parametrised by $(\gamma, \eta, \beta, \xi^n, \xi^\partial, \varphi_n, \varphi_\partial, \xi_n^\dagger, \xi_\partial^\dagger)$, with $\iota_{EH}: \mathcal{F}_R(\Sigma \times I) \rightarrow \mathcal{F}_{EH}^{\text{AKSZ}}$ the inclusion map.

Theorem 37. *The BV-pushforward of $\mathfrak{F}_{EH}^{\text{AKSZ}}(I; \mathfrak{F}_{EH}^\partial(\Sigma))$ with respect to the Lagrangian submanifold $\mathcal{L} = \{(\Pi, \Pi^\dagger) \in \mathcal{F}'' \mid \Pi^\dagger = 0\}$ is the BV theory given by*

$$\mathfrak{F}_R(\Sigma \times I) := (\mathcal{F}_R(\Sigma \times I), S_R(\Sigma \times I), \varpi_R(\Sigma \times I))$$

where

$$\begin{aligned}
S_R(\Sigma \times I) &= \int_{\mathbb{R}} dt \int_{\Sigma} -\eta\sqrt{\gamma} [(\langle K^\sharp, K \rangle - \text{Tr}(K)^2) + R^\partial - 2\Lambda] \\
&\quad - \langle \gamma^\dagger, L_{\xi^\partial} \gamma \rangle - 2\langle K^\sharp, \gamma^\dagger \rangle \xi^n - \langle \varphi_\rho, \dot{\xi}^\rho \rangle \\
&\quad + \langle \varphi_\partial, \nabla_\gamma \eta \xi^n - \eta \nabla_\gamma \xi^n + L_{\xi^\partial} \beta \rangle + \varphi_n (-L_\beta \xi^n + L_{\xi^\partial} \eta) \\
&\quad + \left\langle \xi_\partial^\dagger, \xi^n \nabla_\gamma \xi^n + \frac{1}{2} [\xi^\partial, \xi^\partial] \right\rangle + \xi_n^\dagger L_{\xi^\partial} \xi^n
\end{aligned} \tag{35}$$

with $K := \frac{\eta^{-1}}{2} (\dot{\gamma} + L_\beta \gamma)^\flat$, and

$$\varpi_R(\Sigma \times I) = \iota_{EH}^* \Omega_{EH}^{\text{AKSZ}}. \tag{36}$$

Proof. As discussed in Remark 35, we are interested in finding the effective action one obtains by means of the perturbative expansion of the integral

$$\exp\left(\frac{i}{\hbar} S_{\text{eff}}\right) := \int_{\mathcal{L} \subset \mathcal{F}''} \exp\left(\frac{i}{\hbar} S_{EH}^{\text{AKSZ}}\right) \tag{37}$$

because $S_{EH}^{\text{AKSZ}}|_{\mathcal{L}}$ is quadratic in Π , through the term $\mathcal{K}(\Pi)\eta$. Observing that

$$\frac{\delta \mathcal{K}}{\delta \Pi} = \frac{2}{\sqrt{\gamma}} \left(\Pi^\sharp - \frac{\gamma}{d-1} \text{Tr}(\Pi) \right) \tag{38}$$

$$\frac{\delta^2 \mathcal{K}}{\delta \Pi^2}(\gamma^\dagger) = \frac{2}{\sqrt{\gamma}} \left((\gamma^\dagger)^\sharp - \frac{\gamma}{d-1} \text{Tr}(\gamma^\dagger) \right), \tag{39}$$

we have that Equation (33k) reads

$$\frac{2}{\sqrt{\gamma}} \left(\Pi^\sharp - \frac{\gamma}{d-1} \text{Tr}(\Pi) \right) = -\eta^{-1} \left(\dot{\gamma} + L_\beta \gamma - \frac{2}{\sqrt{\gamma}} \left((\gamma^\dagger)^\sharp - \frac{\gamma}{d-1} \text{Tr}(\gamma^\dagger) \right) \xi^n \right) + F(\Pi^\dagger) \tag{40}$$

where $\text{Tr}(X) = \gamma^{ab} X_{ab}$. We will use the symbol \approx to denote the enforcing of Equation (33k) and of $\Pi^\dagger = 0$. Then, requiring $\Pi^\dagger = 0$ and defining

$$K := \frac{\eta^{-1}}{2} (\dot{\gamma} + L_\beta \gamma)^\flat, \tag{41}$$

we obtain that

$$\Pi \approx -\sqrt{\gamma} \left(K - \text{Tr}(K) \gamma^\flat \right) + \eta^{-1} \gamma^\dagger \xi^n. \tag{42}$$

It is easy to compute now

$$H_n \approx \sqrt{\gamma} [(\langle K^\sharp, K \rangle - \text{Tr}(K)^2) + R^\partial - 2\Lambda] - 2\eta^{-1} \langle K^\sharp, \gamma^\dagger \rangle \xi^n \tag{43}$$

which, together with

$$\langle \Pi, \dot{\gamma} + L_\beta \gamma \rangle \approx -2\sqrt{\gamma} [(\langle K^\sharp, K \rangle - \text{Tr}(K)^2) + R^\partial - 2\Lambda] + 2\eta^{-1} \langle K^\sharp, \gamma^\dagger \rangle \xi^n \tag{44}$$

and

$$-\frac{\delta H_n}{\delta \Pi}(\gamma^\dagger) \xi^n \approx +2 \langle K^\sharp, \gamma^\dagger \rangle \xi^n, \tag{45}$$

yields

$$\begin{aligned}
S_{EH}^{\text{AKSZ}} &\approx \int_{\mathbb{R}} dt \int_{\Sigma} -\eta\sqrt{\gamma} [(\langle K^\sharp, K \rangle - \text{Tr}(K)^2) + R^\partial - 2\Lambda] \\
&\quad - \langle \gamma^\dagger, L_{\xi^\partial} \gamma \rangle + 2\langle K^\sharp, \gamma^\dagger \rangle \xi^n - \langle \varphi_\rho, \dot{\xi}^\rho \rangle \\
&\quad + \langle \varphi_\partial, \nabla_\gamma \eta \xi^n - \eta \nabla_\gamma \xi^n + L_{\xi^\partial} \beta \rangle + \varphi_n (-L_\beta \xi^n + L_{\xi^\partial} \eta) \\
&\quad + \left\langle \xi_\partial^\dagger, \xi^n \nabla_\gamma \xi^n + \frac{1}{2} [\xi^\partial, \xi^\partial] \right\rangle + \xi_n^\dagger L_{\xi^\partial} \xi^n =: S_R(\Sigma \times I).
\end{aligned} \tag{46}$$

So, formula (46) shows that $S_{\text{eff}} = S_R(\Sigma \times I) + O(\hbar)$. The \hbar correction is the (logarithm of the) determinant of the operator defining the quadratic form \mathcal{K} , i.e. the determinant of the deWitt super metric²⁰ [DeW67]

$$W_\gamma^{ijkl} = \frac{1}{\sqrt{\gamma}} \left(\gamma^{ik} \gamma^{jl} - \frac{1}{d-1} \gamma^{ij} \gamma^{kl} \right),$$

or, more invariantly

$$W_\gamma(\Pi, \Pi) = \frac{1}{\sqrt{\gamma}} \left(\langle \Pi^\sharp, \Pi \rangle - \frac{1}{d-1} \text{Tr}_\gamma[\Pi]^2 \right), \quad (47)$$

and it will have the effect of correcting the overall measure on the residual BV space of fields $\mathcal{F}_R(\Sigma \times I)$. \square

Remark 38. Up to boundary, we can compute $Q_R(\Sigma \times I)$ (denoted hereinafter by Q_R) to be:

$$Q_R \gamma = L_{\xi^\partial} \gamma - \eta^{-1} (\dot{\gamma} + L_\beta \gamma) \xi^n \quad (48a)$$

$$Q_R \eta = -L_\beta \xi^n + L_{\xi^\partial} \eta \quad (48b)$$

$$Q_R \beta = \nabla_\gamma \eta \xi^n - \eta \nabla_\gamma \xi^n + L_{\xi^\partial} \beta \quad (48c)$$

$$Q_R \xi^n = L_{\xi^\partial} \xi^n \quad (48d)$$

$$Q_R \xi^\partial = \xi^n \nabla_\gamma \xi^n + \frac{1}{2} [\xi^\partial, \xi^\partial] \quad (48e)$$

and similarly for antifields.

3.2. Reconstruction of Einstein–Hilbert theory. In this section we wish to show that the BV pushforward of the AKSZ theory constructed in Section 3.1 is strongly equivalent to Einstein–Hilbert theory in the BV formalism.

To do this, we begin by considering the following definitions:

$$\tilde{\xi} = -\eta^{-1} \xi^n (\partial_t + \beta) + \xi^\partial \quad (49a)$$

$$\tilde{g} = -\eta^{-2} \partial_t \odot \partial_t - 2\eta^{-2} \beta \odot \partial_t + \gamma - \eta^{-2} \beta \odot \beta \quad (49b)$$

Lemma 39. *We have the following relations*

$$\frac{1}{2} [\tilde{\xi}, \tilde{\xi}] = Q_R \tilde{\xi}, \quad (50a)$$

$$L_{\tilde{\xi}} \tilde{g} = Q_R \tilde{g}. \quad (50b)$$

Proof. It is a straightforward calculation to show

$$\begin{aligned} Q_R \tilde{\xi} &= \eta^{-2} (Q_R \eta) \xi^n (\partial_t + \beta) + \eta^{-1} Q_R \xi^n (\partial_t + \beta) + Q_R \xi^\partial + \eta^{-1} \xi^n Q_R \beta \\ &= \left(-\eta^{-2} \dot{\xi}^n \xi^n - \eta^{-2} L_\beta \xi^n \xi^n + L_{\xi^\partial} (-\eta^{-1} \xi^n) \right) (\partial_t + \beta) \\ &\quad - \eta^{-1} \xi^n \dot{\xi}^\partial + \eta^{-1} \xi^n L_{\xi^\partial} \beta + \frac{1}{2} [\xi^\partial, \xi^\partial]. \end{aligned}$$

Observe that the “algebroid term” (see Remark 40, below) $\xi^n \nabla_\gamma \xi^n$ in $Q_R \xi^\partial$ cancels out with part of $\eta^{-1} \xi^n Q_R \beta$. On the other hand this coincides with

$$\begin{aligned} \frac{1}{2} [\tilde{\xi}, \tilde{\xi}] &= \frac{1}{2} [-\eta^{-1} \xi^n (\partial_t + \beta) + \xi^\partial, -\eta^{-1} \xi^n (\partial_t + \beta) + \xi^\partial] \\ &= (\eta^{-2} \xi^n (\partial_t + \beta) \xi^n + L_{\xi^\partial} (-\eta^{-1} \xi^n)) (\partial_t + \beta) \\ &\quad - \eta^{-1} \xi^n \dot{\xi}^\partial + \eta^{-1} \xi^n L_{\xi^\partial} \beta + \frac{1}{2} [\xi^\partial, \xi^\partial], \end{aligned}$$

²⁰To be precise, W_γ is the inverse of the metric introduced by deWitt, due to our choice of working with inverse metrics γ .

proving the first claim. We compute

$$\begin{aligned}
L_{\tilde{\xi}}\tilde{g} &= -2\eta^{-3}\dot{\xi}^n\partial_t\partial_t + 2\eta^{-3}L_{\xi^\partial}\eta\partial_t\partial_t + 2\eta^{-2}\dot{\xi}\partial_t - 2\eta^{-4}\xi^n L_\beta\eta\partial_t\partial_t \\
&\quad - 2\eta^{-2}\partial_t(\eta^{-1}\xi^n\beta)\partial_t - 4\eta^{-4}\xi^n\partial_t\eta\beta\partial_t + 4\eta^{-3}L_{\xi^\partial}\eta\beta\partial_t - 4\eta^{-4}L_\beta\eta\xi^n\beta\partial_t \\
&\quad - 2\eta^{-2}L_{\xi^\partial}\beta\partial_t - 2\eta^{-2}L_\beta(\eta^{-1}\xi^n)\partial_t\partial_t + 2\eta^{-2}\partial_t(\xi^\partial - \eta^{-1}\xi^n\beta)\beta \\
&\quad - 2\eta^{-2}L_\beta(\eta^{-1}\xi^n)\beta\partial_t - 2\eta^{-2}\partial_t(\eta^{-1}\xi^n)\beta\partial_t + 2\eta^{-2}\xi^n\dot{\beta}\partial_t - \eta^{-1}\xi^n\dot{\gamma}^{ab}\partial_a\partial_b \\
&\quad + L_{\xi^\partial}(\gamma^{ab}\partial_a\partial_b) - \eta^{-1}\xi^n L_\beta(\gamma^{ab})\partial_a\partial_b + 2\nabla_\gamma(\eta^{-1}\xi^n)\partial_t + 2\nabla_\gamma(\eta^{-1}\xi^n\beta)\partial_t \\
&\quad - 2\eta^{-4}\xi^n\dot{\eta}\beta\beta + 2\eta^{-3}L_\xi\eta\beta\beta - 2\eta^{-4}\xi^n L_\beta\eta\beta\beta + 2\eta^{-3}\xi^n\dot{\beta}\beta - \eta^{-2}L_{\xi^\partial}(\beta\beta) \\
&\quad - 2\eta^{-2}L_\beta(\eta^{-1}\xi^n)(\beta\partial_t + \beta\beta)
\end{aligned}$$

where we recall that expressions like $L_{\xi^\partial}(\beta)$ denote the Lie derivative of the vector field $\beta = \beta^a\partial_a$ along ξ^∂ . On the other hand we have

$$\begin{aligned}
Q_R(-\eta^{-2})\partial_t\partial_t &= \left(-2\eta^{-3}\dot{\xi}^n + 2\eta^{-3}L_{\xi^\partial}\eta - 2\eta^{-3}L_\beta\xi^n\right)\partial_t\partial_t \\
Q_R(-2\eta^{-2}\beta)\partial_t &= \left(-4\eta^{-3}\dot{\xi}^n\beta + 4\eta^{-3}L_{\xi^\partial}\eta - 4\eta^{-3}L_\beta\xi^n\beta\right)\partial_t \\
&\quad + \left(2\eta^{-2}\dot{\xi}^\partial - 2\eta^{-2}L_{\xi^\partial}\beta - 2\eta^{-2}\nabla_\gamma\eta\xi^n + 2\eta^{-1}\nabla_\gamma\xi^n\right)\partial_t \\
Q_R(\gamma^{ab})\partial_a\partial_b &= -\eta^{-1}\xi^n\dot{\gamma} - \eta^{-1}\xi^n L_\beta\gamma + L_{\xi^\partial}\gamma \\
Q_R(-\eta^{-2}\beta\beta) &= -2\eta^{-3}\dot{\xi}^n\beta\beta + 2\eta^{-3}L_{\xi^\partial}\eta\beta\beta - 2\eta^{-3}L_\beta\xi^n\beta\beta \\
&\quad + 2\eta^{-2}\beta\dot{\xi}^\partial - 2\eta^{-2}L_{\xi^\partial}\beta\beta - 2\eta^{-2}(\nabla_\gamma\eta\xi^n - \eta\nabla_\gamma\xi^n)\beta
\end{aligned}$$

And it is a matter of a straightforward, but lengthy computation to show that the two expressions coincide. Indeed, subtracting one from the other we obtain

$$\begin{aligned}
L_{\tilde{\xi}}\tilde{g} - Q_R\tilde{g} &= 2\eta^{-3}(-\eta^{-1}\xi^n(\dot{\eta} + L_\beta(\eta)))\beta\beta + 2\eta^{-3}\xi^n\dot{\beta}\beta - 2\eta^{-2}L_\beta(\eta^{-1})\xi^n(\beta\partial_t + \beta\beta) \\
&\quad - 2\eta^{-4}\xi^n L_\beta(\eta)\partial_t^2 + 2\eta^{-4}\dot{\eta}\xi^n\beta\partial_t - 2\eta^{-3}\xi^n\dot{\beta}\xi^n\partial_t - 4\eta^{-4}\xi^n(\dot{\eta} + L_\beta\eta)\beta\partial_t \\
&\quad + 2\eta^{-3}\xi^n\dot{\beta}\partial_t + 2\eta^{-4}L_\beta\eta(\partial_t^2 + \beta\partial_t) + 2\eta^{-4}\dot{\eta}\xi^n(\beta\partial_t + \beta\beta) - 2\eta^{-3}\xi^n\dot{\beta}\beta \equiv 0 \quad (51)
\end{aligned}$$

□

Remark 40. Using Lemma 39 we wish to interpret (49) as a map of Lie algebroids. Consider the (trivial) vector bundle over

$$\text{Map}(I, S_{nd}^2(T\Sigma) \times C^\infty(\Sigma) \times \mathfrak{X}(\Sigma)) \simeq \mathcal{PR}^\Sigma(\Sigma \times I),$$

where $\mathcal{PR}^\Sigma(\Sigma \times I)$ denotes pseudo-Riemannian metrics on $\Sigma \times I$ such that their restriction to Σ is nondegenerate, with fibre

$$\text{Map}(I, C^\infty(\Sigma) \times \mathfrak{X}(\Sigma)) \simeq \mathfrak{X}(\Sigma \times I).$$

We consider two different Lie algebroid structures on this vector bundle. One is the action algebroid with bracket given by the bracket of $(d+1)$ -vector fields, and anchor given by Lie derivatives on metrics. The other algebroid structure is given by formulas (48), with (48a), (48b) and (48c) defining the anchor map, and (48d) and (48e) specifying the bracket of sections. Observe that the morphism of algebroids (49) does not preserve constant sections, as the splitting of a generic vector field $\tilde{\xi}$ depends on the so-called *lapse* η and *shift* β , which are coordinates on the base of the fibre bundle. The latter algebroid encodes the algebraic relations of the constraints of Einstein–Hilbert theory²¹, and was carefully studied by other means in [BFW13]. It was also mentioned as a motivating example for the notion

²¹We stress that, as it is, the structure one can extract from the BFV differential Q^∂ is that of a curved L_∞ algebroid, due to the dependency on fields of negative degree. We thank A. Weinstein, C. Blohmann and N. L. Delgado for enlightening discussions on this matter.

of Hamiltonian Lie Algebroid, introduced in [BW18]. It is an interesting question to check whether this construction satisfies the Hamiltonian requirements for an algebroid.

To proceed, we need to recall the BV data associated with Einstein–Hilbert theory, in the ADM formalism. Given a pseudo-Riemannian (inverse) metric \tilde{g} on a manifold M , we can perform a $d + 1$ decomposition and rewrite it as²²

$$\tilde{g}^{\mu\nu} = \begin{pmatrix} -\eta^{-2} & -\eta^{-2}\beta^b \\ -\eta^{-2}\beta^a & \gamma^{ab} - \eta^{-2}\beta^a\beta^b \end{pmatrix}$$

In the case where M has a boundary, we can define the second fundamental form of the boundary submanifold K_{ab} and its trace K by means of the boundary covariant derivative ∇^∂ (the Levi-Civita connection of γ) as follows

$$K_{ab} = \frac{1}{2}\eta^{-1}(2\nabla_{(a}^\partial\beta_{b)} + \partial_t\gamma_{ab}) \quad K = \gamma^{ab}K_{ab} \quad (52)$$

where t denotes a coordinate transverse to the boundary ∂M . Finally, notice that

$$(L_\beta\gamma)^{cd}\gamma_{ac}\gamma_{bd} = -2\nabla_{(a}^\partial\beta_{b)} \quad (\dot{\gamma})^{cd}\gamma_{ac}\gamma_{bd} = -\partial_t\gamma_{ab}.$$

Definition 41. Let $(\mathcal{F}_{EH}(M), \Omega_{EH}(M))$ be the symplectic manifold

$$\mathcal{F}_{EH}(M) := T^*[-1] \left(\mathcal{PR}^{\partial M}(M) \times \mathfrak{X}[1](M) \right)$$

with its canonical symplectic structure, and $\mathcal{PR}^{\partial M}(M)$ denotes pseudo-Riemannian metrics on M such that their restriction to ∂M is nondegenerate. Consider the functional

$$S_{EH}(M) = \int_M \left\{ -\eta\sqrt{\gamma}(\epsilon(K_{ab}K^{ab} - K^2) + R^\partial - 2\Lambda) \right\} + \tilde{g}^\dagger L_{\tilde{\xi}}\tilde{g} + \frac{1}{2}\iota_{[\tilde{\xi}, \tilde{\xi}]} \tilde{\xi}^\dagger \quad (53)$$

and denote by $Q_{EH}(M)$ the Hamiltonian vector field of $S_{EH}(M)$, up to boundary terms. Then, the assignment of the tuple

$$\mathfrak{F}_{EH} = (\mathcal{F}_{EH}(M), S_{EH}(M), \Omega_{EH}(M), Q_{EH}(M))$$

to every $(d+1)$ -dimensional manifold M that admits a Lorentzian structure will be called Einstein–Hilbert theory in the BV formalism.

Remark 42. The sign convention used above is obtained from the standard ADM decomposition by redefining $(\eta, \beta) \rightarrow (-\eta, -\beta)$. This matches our conventions below. This change is due to the choice of using inverse metrics for the first order formulation, instead of metrics (in fact $\Pi_{ab}\partial_t\gamma^{ab} = -\Pi^{ab}\partial_t\gamma_{ab}$).

Theorem 43. *Einstein–Hilbert theory in the BV formalism $\mathfrak{F}_{EH}(\Sigma \times I)$ is strongly equivalent to $\mathfrak{F}_R(\Sigma \times I)$. Explicitly, the isomorphism of the underlying symplectic*

²²In this paper we will assume that the manifold M has a global product structure $M = \Sigma \times \mathbb{R}$, and the induced metric on Σ will be Riemannian, i.e. the leaves Σ_t are spacelike submanifolds of M . It is straightforward to generalise this to the *timelike* case. The relevant formulas for EH theory in the BV-BFV formalism have been given in [CS16].

dg-manifolds reads:

$$\tilde{g} = -\eta^{-2}\partial_t\partial_t - 2\eta^{-2}\beta\partial_t + \gamma - \eta^{-2}\beta\beta \quad (54a)$$

$$\tilde{\xi} = -\eta^{-1}\xi^n\partial_t + \xi^\partial - \eta^{-1}\xi^n\beta \quad (54b)$$

$$\tilde{\xi}^\dagger = \xi_\partial^\dagger - \left(\eta\xi_n^\dagger + \iota_\beta\xi_\partial^\dagger\right) dt \quad (54c)$$

$$\begin{aligned} \tilde{g}^\dagger &= \left(\frac{1}{2}\eta^3\varphi_n - \eta^2\varphi_a\beta^a - \gamma_{ab}^\dagger\beta^a\beta^b + \eta\beta^a\xi_a^\dagger\xi^n + \frac{1}{2}\eta^{-1}\xi_n^\dagger\xi^n\right) dt^2 \\ &+ \left(\frac{1}{2}\eta^2\varphi_a + \gamma_{ab}^\dagger\beta^b - \frac{1}{2}\eta\xi_a^\dagger\xi^n\right) dx^a dt - \gamma_{ab}^\dagger dx^a dx^b \end{aligned} \quad (54d)$$

with inverse:

$$\eta = [-\tilde{g}^{tt}]^{-\frac{1}{2}} \quad (55a)$$

$$\beta^a = -[-\tilde{g}^{tt}]^{-1}\tilde{g}^{ta} \quad (55b)$$

$$\gamma^{ab} = [-\tilde{g}^{tt}]^{-1}\tilde{g}^{ta}\tilde{g}^{tb} \quad (55c)$$

$$\xi^n = -[-\tilde{g}^{tt}]^{-\frac{1}{2}}\tilde{\xi}^t \quad (55d)$$

$$\xi^a = \tilde{\xi}^a + [-\tilde{g}^{tt}]\tilde{g}^{ta}\tilde{\xi}^t \quad (55e)$$

$$\gamma_{ab}^\dagger = \tilde{g}_{ab}^\dagger \quad (55f)$$

$$\varphi_a = 2[-\tilde{g}^{tt}]\tilde{g}_{at}^\dagger + 2\tilde{g}_{ab}^\dagger\tilde{g}^{tb} + \tilde{\xi}_a^\dagger\tilde{\xi}^t \quad (55g)$$

$$\varphi_n = 2[-\tilde{g}^{tt}]^{\frac{3}{2}}\tilde{g}_{tt}^\dagger - 4[-\tilde{g}^{tt}]^{\frac{1}{2}}\tilde{g}_{ta}^\dagger\tilde{g}^{ta} + 2[-\tilde{g}^{tt}]^{-\frac{1}{2}}\tilde{g}_{ab}^\dagger\tilde{g}^{ta}\tilde{g}^{tb} \quad (55h)$$

$$+ [-\tilde{g}^{tt}]^{\frac{1}{2}}\tilde{\xi}_n^\dagger\tilde{\xi}^t - [-\tilde{g}^{tt}]^{-\frac{1}{2}}\tilde{g}^{ta}\tilde{\xi}_a^\dagger\tilde{\xi}^a \quad (55i)$$

$$\xi_n^\dagger = -[-\tilde{g}^{tt}]^{\frac{1}{2}}\tilde{\xi}_n^\dagger + [-\tilde{g}^{tt}]^{-\frac{1}{2}}\tilde{\xi}_a^\dagger\tilde{g}^{ta} \quad (55j)$$

$$\xi_a^\dagger = \tilde{\xi}_a^\dagger \quad (55k)$$

Proof. We begin observing that the definitions of K in (41) and K in (52) coincide up to sign, after identifying \tilde{g} with the expression of Equation (49b). Since the expression $S_{ADM} := -\eta\sqrt{\gamma}(\epsilon(K_{ab}K^{ab} - K^2) + R^\partial - 2\Lambda)$ is quadratic in K , we conclude that the degree-zero part of (53) and (46) coincide. This means that the two theories are classically equivalent²³, and (49b) is the map between second-order and first-order Einstein–Hilbert theory.

We endeavour now to find the explicit expression for \tilde{g}^\dagger and $\tilde{\xi}^\dagger$ so that

$$\phi^*(\langle\tilde{g}^\dagger, \delta\tilde{g}\rangle + \langle\tilde{\xi}^\dagger, \delta\tilde{\xi}\rangle) = -\langle\delta\gamma^\dagger, \delta\gamma\rangle + \langle\Pi, \delta\Pi^\dagger\rangle + \xi_\rho^\dagger\delta\xi^\rho + \varphi_n\delta\eta + \varphi_a\delta\beta^a.$$

It is straightforward to compute

$$\begin{aligned} \phi^*(\langle\tilde{g}^\dagger, \delta\tilde{g}\rangle + \langle\tilde{\xi}^\dagger, \delta\tilde{\xi}\rangle) &= -\left[(\phi^*\tilde{\xi}^\dagger)_n\eta^{-1} + (\phi^*\tilde{\xi}^\dagger)_a\eta^{-1}\beta^a\right]\delta\xi^n + (\phi^*\tilde{\xi}^\dagger)_a\delta\xi^a \\ &+ \left[2(\phi^*\tilde{g}^\dagger)_{tt}\eta^{-3} + 4(\phi^*\tilde{g}^\dagger)_{at}\eta^{-3}\beta^a + 2(\phi^*\tilde{g}^\dagger)_{ab}\eta^{-3}\beta^a\beta^b\right]\delta\eta \\ &- \left[(\phi^*\tilde{\xi}^\dagger)_t\eta^{-2}\xi^n - (\phi^*\tilde{\xi}^\dagger)_a\eta^{-2}\beta^a\xi^n\right]\delta\eta + (\phi^*\tilde{g}^\dagger)_{ab}\delta\gamma^{ab} \\ &+ \left[2(\phi^*\tilde{g}^\dagger)_{at}\eta^{-2} - 2(\phi^*\tilde{g}^\dagger)_{ab}\eta^{-2}\beta^b + (\phi^*\tilde{\xi}^\dagger)_a\eta^{-1}\xi^n\right]\delta\beta^a \end{aligned} \quad (56)$$

²³Strictly speaking this is only true on an open subset of the moduli space of solutions, due to the nondegeneracy condition on $\gamma(t)$ enforced on the whole cylinder by the AKSZ construction, while Definition 53 only requires the nondegeneracy of $\tilde{g}|_{\partial M}$.

which leaves us with the intermediate expression:

$$\xi_a^\dagger = (\phi^* \tilde{\xi}^\dagger)_a \quad (57a)$$

$$\xi_n^\dagger = - \left[(\phi^* \tilde{\xi}^\dagger)_t \eta^{-1} + (\phi^* \tilde{\xi}^\dagger)_a \eta^{-1} \beta^a \right] \quad (57b)$$

$$\gamma_{ab}^\dagger = -(\phi^* \tilde{g}^\dagger)_{ab} \quad (57c)$$

$$\varphi_a = -2(\phi^* \tilde{g}^\dagger)_{at} \eta^{-2} - 2(\phi^* \tilde{g}^\dagger)_{ab} \eta^{-2} \beta^b + (\phi^* \tilde{\xi}^\dagger)_a \eta^{-1} \xi^n \quad (57d)$$

$$\begin{aligned} \varphi_n &= 2(\phi^* \tilde{g}^\dagger)_{tt} \eta^{-3} + 4(\phi^* \tilde{g}^\dagger)_{at} \eta^{-3} \beta^a + 2(\phi^* \tilde{g}^\dagger)_{ab} \eta^{-3} \beta^a \beta^b \\ &\quad - (\phi^* \tilde{\xi}^\dagger)_t \eta^{-2} \xi^n - (\phi^* \tilde{\xi}^\dagger)_a \eta^{-2} \beta^a \xi^n \end{aligned} \quad (57e)$$

Starting from the top and solving downwards, we easily get

$$(\phi^* \tilde{\xi}^\dagger)_a = \xi_a^\dagger \quad (58)$$

$$(\phi^* \tilde{\xi}^\dagger)_n = -\eta \xi_n^\dagger - \xi_a^\dagger \beta^a \quad (59)$$

$$(\phi^* \tilde{g}^\dagger)_{ab} = -\gamma_{ab}^\dagger \quad (60)$$

$$(\phi^* \tilde{g}^\dagger)_{at} = -\frac{1}{2} \eta^2 \varphi_a + \gamma_{ab}^\dagger \beta^b + \frac{1}{2} \eta \xi_a^\dagger \xi^n \quad (61)$$

$$(\phi^* \tilde{g}^\dagger)_{tt} = \frac{1}{2} \eta^3 \varphi_n + \eta^2 \varphi_a \beta^a - \gamma_{ab}^\dagger \beta^a \beta^b - \eta \xi_a^\dagger \beta^a \xi^n - \frac{1}{2} \eta \xi_n^\dagger \xi^n \quad (62)$$

Alternatively, from (57), observing that the assignment (49) can be inverted to yield

$$\phi^{-1*} \eta = [-\tilde{g}^{tt}]^{-\frac{1}{2}}, \quad \phi^{-1*} \beta^a = -[-\tilde{g}^{tt}]^{-1} \tilde{g}^{ta}, \quad \phi^{-1*} \gamma^{ab} = [-\tilde{g}^{tt}]^{-1} \tilde{g}^{ta} \tilde{g}^{tb}$$

together with

$$\phi^{-1*} \xi^n = -[-\tilde{g}^{tt}]^{-\frac{1}{2}} \tilde{\xi}^t; \quad \phi^{-1*} \xi^a = \tilde{\xi}^a + [-\tilde{g}^{tt}]^{-1} \tilde{g}^{ta} \tilde{\xi}^t$$

we can similarly obtain the inverse:

$$\phi^{-1*} \eta = [-\tilde{g}^{tt}]^{-\frac{1}{2}}$$

$$\phi^{-1*} \beta^a = -[-\tilde{g}^{tt}]^{-1} \tilde{g}^{ta}$$

$$\phi^{-1*} \gamma^{ab} = [-\tilde{g}^{tt}]^{-1} \tilde{g}^{ta} \tilde{g}^{tb}$$

$$\phi^{-1*} \xi^n = -[-\tilde{g}^{tt}]^{-\frac{1}{2}} \tilde{\xi}^t$$

$$\phi^{-1*} \xi^a = \tilde{\xi}^a + [-\tilde{g}^{tt}] \tilde{g}^{ta} \tilde{\xi}^t$$

$$\phi^{-1*} \gamma_{ab}^\dagger = -\tilde{g}_{ab}^\dagger$$

$$\phi^{-1*} \xi_a^\dagger = \tilde{\xi}_a^\dagger$$

$$\phi^{-1*} \xi_n^\dagger = -\tilde{\xi}_n^\dagger [-\tilde{g}^{tt}]^{\frac{1}{2}} + \tilde{\xi}_a^\dagger [-\tilde{g}^{tt}]^{-\frac{1}{2}} \tilde{g}^{ta}$$

$$\phi^{-1*} \varphi_a = -2\tilde{g}_{at}^\dagger [-\tilde{g}^{tt}] + 2\tilde{g}_{ab}^\dagger \tilde{g}^{tb} - \tilde{\xi}_a^\dagger \tilde{\xi}^t$$

$$\begin{aligned} \phi^{-1*} \varphi_n &= 2\tilde{g}_{tt}^\dagger [-\tilde{g}^{tt}]^{\frac{3}{2}} - 4\tilde{g}_{ta}^\dagger [-\tilde{g}^{tt}]^{\frac{1}{2}} \tilde{g}^{ta} + 2\tilde{g}_{ab}^\dagger [-\tilde{g}^{tt}]^{-\frac{1}{2}} \tilde{g}^{ta} \tilde{g}^{tb} \\ &\quad + [-\tilde{g}^{tt}]^{\frac{1}{2}} \tilde{\xi}_t^\dagger \tilde{\xi}^t - [-\tilde{g}^{tt}]^{-\frac{1}{2}} \tilde{g}^{ta} \tilde{\xi}_a^\dagger \tilde{\xi}^t. \end{aligned}$$

Now, using again the intermediate expressions (57) let us consider the following terms, coming from Equation (46):

$$\begin{aligned} \xi_n^\dagger L_{\xi^\partial} \xi^n &= -(\phi^* \tilde{\xi}^\dagger)_t \eta^{-1} L_{\xi^\partial} \xi^n - (\phi^* \tilde{\xi}^\dagger)_a \eta^{-1} \beta^a L_{\xi^\partial} \xi^n \\ \langle \xi_\partial^\dagger, (\xi^n \nabla_\gamma \xi^n + \frac{1}{2} [\xi^\partial, \xi^\partial]) \rangle &= \langle (\phi^* \tilde{\xi}^\dagger)_\partial, (\xi^n \nabla_\gamma \xi^n + \frac{1}{2} [\xi^\partial, \xi^\partial]) \rangle \end{aligned}$$

$$\begin{aligned}
& \varphi_n \left(L_{\xi^\partial} \eta - L_\beta \xi^n - \dot{\xi}^n \right) \\
&= \eta^{-3} \left[2(\phi^* \tilde{g}^\dagger)_{tt} + 4(\phi^* \tilde{g}^\dagger)_{at} \beta^a + (\phi^* \tilde{g}^\dagger)_{ab} \beta^a \beta^b \right] \left(L_{\xi^\partial} \eta - L_\beta \xi^n - \dot{\xi}^n \right) \\
&\quad - \eta^{-2} \left[(\phi^* \tilde{\xi}^\dagger)_n \xi^n + (\phi^* \tilde{\xi}^\dagger)_a \beta^a \xi^n \right] \left(L_{\xi^\partial} \eta - L_\beta \xi^n - \dot{\xi}^n \right) \\
& \langle \varphi_\partial, (\nabla_\gamma \xi^n - \eta \nabla_\gamma \xi^n + L_{\xi^\partial} \beta) \rangle \\
&= -2\eta^{-2} \left((\phi^* \tilde{g}^\dagger)_{at} + (\phi^* \tilde{g}^\dagger)_{ab} \beta^b \right) \left((\nabla_\gamma \eta)^a \xi^n - \eta (\nabla_\gamma \xi^n)^a + (L_{\xi^\partial} \beta)^a - \dot{\xi}^a \right) \\
&\quad + (\phi^* \tilde{\xi}^\dagger)_a \eta^{-1} \xi^n \left((L_{\xi^\partial} \beta)^a - \eta (\nabla_\gamma \xi^n)^a - \dot{\xi}^a \right) \\
& \langle \gamma^\dagger, \eta^{-1} (\dot{\gamma} + L_\beta \gamma) \xi^n - L_{\xi^\partial} \gamma \rangle = -(\phi^* \tilde{g}^\dagger)_{ab} \left(\eta^{-1} (\dot{\gamma} + L_\beta \gamma) \xi^n - L_{\xi^\partial} \gamma \right)^{ab}.
\end{aligned}$$

Then, summing all terms on the left hand side and factoring $(\phi^* \tilde{\xi}^\dagger)_t$, $(\phi^* \tilde{\xi}^\dagger)_a$ and $(\phi^* \tilde{g}^\dagger)$, we obtain

$$\begin{aligned}
& (\phi^* \tilde{\xi}^\dagger)_t \left[-L_{\xi^\partial} (\eta^{-1} \xi^n) + \eta^{-2} \xi^n \left(L_\beta \xi^n + \dot{\xi}^n \right) \right] \\
&+ (\phi^* \tilde{g}^\dagger)_{tt} \left[2\eta^{-3} \left(L_{\xi^\partial} \eta - L_\beta \xi^n - \dot{\xi}^n \right) \right] + \left\langle (\phi^* \tilde{\xi}^\dagger)_\partial, \frac{1}{2} [\xi^\partial, \xi^\partial] \right\rangle \\
&+ \left\langle (\phi^* \tilde{\xi}^\dagger)_\partial, -L_{\xi^\partial} (\eta^{-1} \xi^n) \beta + \eta^{-1} L_{\xi^\partial} \beta \xi^n + \eta^{-2} \beta \xi^n L_\beta \xi^n + \eta^{-2} \beta \xi^n \dot{\xi}^n - \eta^{-1} \xi^n \dot{\xi}^\partial \right\rangle \\
&+ (\phi^* \tilde{g}^\dagger)_{ab} \left[-\eta^{-1} \dot{\gamma}^{ab} \xi^n - \eta^{-1} (L_\beta \gamma)^{ab} \xi^n + (L_{\xi^\partial} \gamma)^{ab} + 4\eta^{-3} \beta^a \left(L_{\xi^\partial} \eta - L_\beta \xi^n - \dot{\xi}^n \right) \right] \\
&+ (\phi^* \tilde{g}^\dagger)_{ab} \left[-2\eta^{-2} \left((\nabla_\gamma \eta)^a \xi^n - \eta (\nabla_\gamma \xi^n)^a + (L_{\xi^\partial} \beta)^a - \dot{\xi}^a \right) \right] \\
&+ (\phi^* \tilde{g}^\dagger)_{ab} \left[2\eta^{-3} \beta^a \beta^b \left(L_{\xi^\partial} \eta - L_\beta \xi^n - \dot{\xi}^n \right) \right] \\
&+ (\phi^* \tilde{g}^\dagger)_{ab} \left[-2\eta^{-2} \beta^b \left((\nabla_\gamma \eta)^a \xi^n - \eta (\nabla_\gamma \xi^n)^a + (L_{\xi^\partial} \beta)^a - \dot{\xi}^a \right) \right]
\end{aligned}$$

Which, using Lemma 39, can be shown to be

$$\phi^* \left(\tilde{g}^\dagger L_{\tilde{\xi}} \tilde{g} + \iota_{[\tilde{\xi}, \tilde{\xi}]} \tilde{\xi}^\dagger \right) \quad (63)$$

leading to

$$\phi^* S_{EH}(\Sigma \times I) = S_R(\Sigma \times I). \quad (64)$$

□

Remark 44. We would like to stress here that the results in this section are a “strictification” of the general construction of a solution of the classical master equation for the extended Hamiltonian, as presented by Henneaux and Bunster in [HT92, Theorem 18.8]. Indeed, the Hamiltonian analysis for a field theory relies on a (possibly) non-reduced version of the strict BFV data we consider, where strict indicates that we require all spaces of fields to be smooth symplectic manifolds. The AKSZ construction yields a BV theory (Theorem 32) which is effectively equivalent to the natural BV extension of Einstein–Hilbert theory (Theorems 37 and 43). It could be argued that this effective equivalence preserves the BV cohomology [BBH95; DGH90; Hen90]. However, note that the quantisation procedure outlined in [CMR18] does indeed require the strict version of a BV-BFV structure²⁴, and its existence is not to be taken for granted, as was shown in [CS19b] and [CS17].

²⁴See [MSW19] for the comparison between strict and lax BV-BFV structures.

4. AKSZ PC

Following the construction outlined in Section 1.3, starting from the BFV theory of Palatini–Cartan gravity (see Section 2.2), we can construct the AKSZ space of fields $\mathcal{F}_{PC}^{\text{AKSZ}}$. We will use the following notation:

$$\mathbf{e} = e + f^\dagger \qquad \mathbf{w} = \omega + u^\dagger \qquad (65a)$$

$$\mathbf{c} = c + w \qquad \mathbf{x} = \xi + z \qquad (65b)$$

$$\mathbf{l} = \lambda + \mu \qquad \mathbf{c}^\dagger = k^\dagger + c^\dagger \qquad (65c)$$

$$\mathbf{\eta}^\dagger = e^\dagger + y^\dagger \qquad (65d)$$

where

$$\begin{aligned} e &\in C^\infty(I) \otimes \Omega_{nd}^1(\Sigma, \mathcal{V}) & f^\dagger &\in \Omega^1[-1](I) \otimes \Omega^1(\Sigma, \mathcal{V}) \\ \omega &\in C^\infty(I) \otimes \mathcal{A}(\Sigma) & u^\dagger &\in \Omega^1[-1](I) \otimes \mathcal{A}(\Sigma) \\ c &\in \Omega^0[1](I \times \Sigma, \wedge^2 \mathcal{V}) & w &\in \Omega^1[-1](I) \otimes \Omega^0[1](\Sigma, \wedge^2 \mathcal{V}) \\ \xi &\in C^\infty(I) \otimes \mathfrak{X}[1](\Sigma) & z &\in \Omega^1[-1](I) \otimes \mathfrak{X}[1](\Sigma) \\ \lambda &\in C^\infty[1](I \times \Sigma) & \mu &\in \Omega^1[-1](I) \otimes C^\infty[1](\Sigma) \\ k^\dagger &\in C^\infty(I) \otimes \Omega^{N-1}[-1](\Sigma, \wedge^{N-2} \mathcal{V}) & c^\dagger &\in \Omega^1[-1](I) \otimes \Omega^{N-1}[-1](\Sigma, \wedge^{N-2} \mathcal{V}) \\ e^\dagger &\in C^\infty(I) \otimes \Omega^{N-1}[-1](\Sigma, \wedge^{N-1} \mathcal{V}) & y^\dagger &\in \Omega^1[-1](I) \otimes \Omega^{N-1}[-1](\Sigma, \wedge^{N-1} \mathcal{V}) \end{aligned} \qquad (66)$$

such that, for some $\sigma \in C^\infty(I) \otimes \Omega^1(\Sigma, \mathcal{V})$ and $B \in \Omega^1[-1](I) \otimes \Omega^1(\Sigma, \mathcal{V})$, they satisfy the *structural AKSZ constraints*:

$$\epsilon_n \{ (N-4)f^\dagger e^{N-5} d_\omega e + e^{N-4} d_\omega f^\dagger + e^{N-4} [u^\dagger, e] \} \qquad (67a)$$

$$\begin{aligned} + (\iota_z d_\omega \epsilon_n - [w - \iota_\xi u^\dagger, \epsilon_n])^{(a)} k_a^\dagger + X^{(a)} c_a^\dagger + (X^{(b)} f_b^\dagger)^{(a)} k_a^\dagger &= f^\dagger e^{N-4} \sigma + e^{N-3} B; \\ \epsilon_n e^{N-4} d_\omega e + X^{(a)} k_a^\dagger &= e^{N-3} \sigma; \end{aligned} \qquad (67b)$$

where $X = (L_\xi^\omega \epsilon_n - [c, \epsilon_n]) \in \Gamma(M, \mathcal{V})$, while $\epsilon_n \in \Gamma(M, \mathcal{V})$ is a fixed section, and the indices $\{(a), (n)\}$ denote components with respect to a basis $\{e_a, \epsilon_n\}$.

Remark 45. Observe that our target for the AKSZ construction for Palatini–Cartan theory is the BFV theory defined in Definition 26, whose space of fields $\mathcal{F}_{PC}^\partial$ is defined by the structural constraint (19). As a consequence, the BFV constraint must be imposed on the AKSZ fields at every point of I . As the AKSZ fields consists of a 0- and and 1-form component, along the interval, the structural constraints now has a 0-form and a 1- form component corresponding to (67b) and (67a), respectively. Despite the apparent complexity of these two equations, it is worth noting that they fix certain components of the AKSZ fields ω and u^\dagger . See Section 4.2 for an interpretation.

Remark 46. Recall that to define the BFV structure for PC theory we needed a fixed section $\epsilon_n \in \Gamma(\Sigma, \mathcal{V})$ (cf. Definition 25). Note that ϵ_n is not a field of the theory but is part of the structure that defines it (more like a coupling constant). For this reason, in the AKSZ construction ϵ_n does not depend on the coordinate x^n of the interval I . In the following, we will regard ϵ_n as a given section of $\Gamma(M, \mathcal{V})$ satisfying $d_I(\epsilon_n) = 0$.

Theorem 47. *The AKSZ data $\mathfrak{F}_{PC}^{\text{AKSZ}}(I; \mathfrak{F}_{PC}^\partial)$ are given by the quadruple*

$$\mathfrak{F}_{PC}^{\text{AKSZ}}(I; \mathfrak{F}_{PC}^\partial) = (\mathcal{F}_{PC}^{\text{AKSZ}}, S_{PC}^{\text{AKSZ}}, \varpi_{PC}^{\text{AKSZ}}, Q_{PC}^{\text{AKSZ}})$$

where:

$$\begin{aligned} \mathcal{F}_{PC}^{\text{AKSZ}} &\simeq T^*[-1](\text{Map}(I, \mathcal{F}_{PC}^\partial)) \\ \varpi_{PC}^{\text{AKSZ}} &= \int_{I \times \Sigma} \delta(e^{N-3} f^\dagger) \delta \omega + e^{N-3} \delta e \delta u^\dagger + \delta w \delta k^\dagger + \delta c \delta c^\dagger + \delta u^\dagger \delta(\iota_\xi k^\dagger) \\ &\quad + \delta \omega \delta(\iota_z k^\dagger) + \delta \omega \delta(\iota_\xi c^\dagger) - \delta \mu \epsilon_n \delta e^\dagger - \delta \lambda \epsilon_n \delta y^\dagger \\ &\quad + \iota_{\delta z} \delta(e e^\dagger) + \iota_{\delta \xi} \delta(f^\dagger e^\dagger) + \iota_{\delta \xi} \delta(e y^\dagger); \\ S_{PC}^{\text{AKSZ}} &= \int_{I \times \Sigma} w e^{N-3} d_\omega e + (N-3) c e^{N-4} f^\dagger d_\omega e + c e^{N-3} [u^\dagger, e] + c e^{N-3} d_\omega f^\dagger \\ &\quad + \iota_z e e^{N-3} F_\omega + \iota_\xi (e^{N-3} f^\dagger) F_\omega + \iota_\xi e e^{N-3} d_\omega u^\dagger + \epsilon_n \mu e^{N-3} F_\omega \\ &\quad + (N-3) \epsilon_n \lambda e^{N-4} f^\dagger F_\omega + \epsilon_n \lambda e^{N-3} d_\omega u^\dagger + [w, c] k^\dagger + \frac{1}{2} [c, c] c^\dagger \\ &\quad - \iota_z d_\omega c k^\dagger - [\iota_\xi u^\dagger, c] k^\dagger - \iota_\xi d_\omega w k^\dagger - \iota_\xi d_\omega c c^\dagger + \iota_z \iota_\xi F_\omega k^\dagger \\ &\quad + \frac{1}{2} \iota_\xi \iota_\xi d_\omega u^\dagger k^\dagger + \frac{1}{2} \iota_\xi \iota_\xi F_\omega c^\dagger - [w, \epsilon_n \lambda] e^\dagger - [c, \epsilon_n \mu] e^\dagger - [c, \epsilon_n \lambda] y^\dagger \\ &\quad + \iota_z d_\omega (\epsilon_n \lambda) e^\dagger + [\iota_\xi u^\dagger, \epsilon_n \lambda] e^\dagger + \iota_\xi d_\omega (\epsilon_n \mu) e^\dagger + \iota_\xi d_\omega (\epsilon_n \lambda) y^\dagger \\ &\quad + \iota_{[z, \xi]} e e^\dagger + \frac{1}{2} \iota_{[\xi, \xi]} f^\dagger e^\dagger + \frac{1}{2} \iota_{[\xi, \xi]} e y^\dagger \\ &\quad + \frac{1}{N-2} e^{N-2} d_I \omega + c d_I k^\dagger + d_I \omega \iota_\xi k^\dagger - \iota_{d_I \xi} e e^\dagger + d_I \lambda \epsilon_n e^\dagger. \end{aligned}$$

and Q_{PC}^{AKSZ} is defined as $Q_{PC}^{\text{AKSZ}} = Q_{PC}^{dR} + Q_{PC}^{\text{lift}}$ where Q_{PC}^{lift} is the tangent lift of Q_{PC}^∂ to $\text{Map}(T[1]I, \mathcal{F}_{PC}^\partial(\Sigma))$ and Q_{PC}^{dR} is the lift of the de Rham differential d_I .

Proof. This is a straightforward application of the AKSZ prescription outlined in Section 1.3. Using the transgression map we can build a symplectic form $\mathcal{F}_{PC}^{\text{AKSZ}}$

$$\varpi_{PC}^{\text{AKSZ}} = \int_{I \times \Sigma} \mathbf{e}^{N-3} \delta \mathbf{e} \delta \mathbf{w} + \delta \mathbf{c} \delta \mathbf{c}^\dagger + \delta \mathbf{w} \delta(\iota_{\mathbf{r}} \mathbf{c}^\dagger) - \delta \mathbf{l} \epsilon_n \delta \mathbf{y}^\dagger + \iota_{\delta \mathbf{r}} \delta(\mathbf{e} \mathbf{y}^\dagger) \quad (68)$$

from which we obtain the claimed expression using (65). Analogously the AKSZ action can be constructed using the transgression map from the boundary one-form α^∂ and from the boundary action S^∂ . Namely we have

$$\begin{aligned} S_{PC}^{\text{AKSZ}} &= \int_{I \times \Sigma} \frac{1}{N-2} \mathbf{e}^{N-2} d_I \mathbf{w} + c d_I \mathbf{c}^\dagger + d_I \mathbf{w} \iota_{\mathbf{r}} \mathbf{c}^\dagger - \iota_{d_I \mathbf{r}} \mathbf{e} \mathbf{y}^\dagger + d_I \mathbf{l} \epsilon_n \mathbf{y}^\dagger \quad (69) \\ &\quad + c d_\omega \mathbf{e} + \iota_{\mathbf{r}} \mathbf{c} \mathbf{e} F_\omega + \epsilon_n \mathbf{l} \mathbf{e} F_\omega + \frac{1}{2} [\mathbf{c}, \mathbf{c}] \mathbf{c}^\dagger - L_{\mathbf{r}}^{\mathbf{w}} \mathbf{c} \mathbf{c}^\dagger + \frac{1}{2} \iota_{\mathbf{r}} \iota_{\mathbf{r}} F_\omega \mathbf{c}^\dagger \\ &\quad - [\mathbf{c}, \epsilon_n \mathbf{l}] \mathbf{y}^\dagger + L_{\mathbf{r}}^{\mathbf{w}} (\epsilon_n \mathbf{l}) \mathbf{y}^\dagger + \frac{1}{2} \iota_{[\mathbf{r}, \mathbf{r}]} \mathbf{e} \mathbf{y}^\dagger. \end{aligned}$$

Again the claimed expression can be obtained straightforwardly from (65). \square

Remark 48. The invariance of the constraints (67b) and (67a) with respect to Q_{PC}^{AKSZ} is guaranteed by the invariance of the structural constraint on the boundary (19) with respect to Q_{PC}^{BFV} , and by the properties of the AKSZ construction.

From Theorem 14 we know that $\mathfrak{F}_{PC}^{\text{AKSZ}}(I; \mathfrak{F}_{PC}^\partial)$ yields a BV theory on the manifold $I \times \Sigma$. Furthermore, by Proposition 17 these data satisfy also the BV-BFV axioms of Equation (5).

Definition 49. We call *nondegenerate* AKSZ PC theory the data $\mathfrak{F}_{PC_*}^{\text{AKSZ}}$ obtained by restricting the space of fields of $\mathfrak{F}_{PC}^{\text{AKSZ}}(I; \mathfrak{F}_{PC}^\partial(\Sigma))$ to those maps whose μ component (as defined by Equation (65)) is nonvanishing.

In [CS19b] two of the authors proved that, using the natural symmetries of PC theory, the resulting BV theory \mathfrak{F}_{PC} does not satisfy the BV-BFV axioms (it is not a 1-extended BV theory) unless additional requirements on the fields are enforced. Next section will be devoted to the comparison between $\mathfrak{F}_{PC}(\Sigma \times I)$ and $\mathfrak{F}_{PC^*}^{\text{AKSZ}}(I; \mathfrak{F}_{PC}^{\partial}(\Sigma))$.

4.1. Comparison of BV data for PC theory. We want to compare the AKSZ-BV theory of Theorem 47 with the one proposed for PC-gravity by two of the authors [CS19b], which we briefly recall here. Let M be an N -dimensional manifold with $N > 2$.

Definition 50. We call *standard* BV theory for PC gravity the BV data

$$\mathfrak{F}_{PC}(M) = (\mathcal{F}_{PC}(M), S_{PC}(M), \varpi_{PC}(M), Q_{PC}(M))$$

where

$$\mathcal{F}_{PC}(M) := T^*[-1] (\Omega_{nd}^1(M, \mathcal{V}) \oplus \mathcal{A}(M) \oplus \mathfrak{X}[1](M) \oplus \Omega^0[1](M, \text{ad}P))$$

and the fields in the base are denoted by $(\mathbf{e}, \boldsymbol{\omega}, \boldsymbol{\xi}, \mathbf{c})$, while the corresponding variables in the cotangent fibre are denoted by $(\mathbf{e}^\dagger, \boldsymbol{\omega}^\dagger, \boldsymbol{\xi}^\dagger, \mathbf{c}^\dagger)$;

$$\begin{aligned} \varpi_{PC}(M) &= \int_M \delta \mathbf{e} \delta \mathbf{e}^\dagger + \delta \boldsymbol{\omega} \delta \boldsymbol{\omega}^\dagger + \delta \mathbf{c} \delta \mathbf{c}^\dagger + \iota_{\delta \boldsymbol{\xi}} \delta \boldsymbol{\xi}^\dagger; \\ S_{PC}(M) &= \int_M \frac{1}{N-2} \mathbf{e}^{N-2} F_\omega + (\iota_{\boldsymbol{\xi}} F_\omega - d_\omega \mathbf{c}) \boldsymbol{\omega}^\dagger - (L_\xi^\omega \mathbf{e} - [\mathbf{c}, \mathbf{e}]) \mathbf{e}^\dagger \\ &\quad + \int_M \frac{1}{2} (\iota_{\boldsymbol{\xi}} \iota_{\boldsymbol{\xi}} F_\omega - [\mathbf{c}, \mathbf{c}]) \mathbf{c}^\dagger + \frac{1}{2} \iota_{[\boldsymbol{\xi}, \boldsymbol{\xi}]} \boldsymbol{\xi}^\dagger. \end{aligned}$$

The explicit expression of the cohomological vector field Q_{PC} , defined by the equation $\iota_{Q_{PC}} \varpi_{PC} = \delta S_{PC}$, will be useful in the following:

$$\begin{aligned} Q_{PC} \mathbf{e} &= L_\xi^\omega \mathbf{e} - [\mathbf{c}, \mathbf{e}] \\ Q_{PC} \boldsymbol{\omega} &= \iota_{\boldsymbol{\xi}} F_\omega - d_\omega \mathbf{c} \\ Q_{PC} \mathbf{c} &= \frac{1}{2} \iota_{\boldsymbol{\xi}} \iota_{\boldsymbol{\xi}} F_\omega - \frac{1}{2} [\mathbf{c}, \mathbf{c}] \\ Q_{PC} \boldsymbol{\xi} &= \frac{1}{2} [\boldsymbol{\xi}, \boldsymbol{\xi}] \\ Q_{PC} \mathbf{e}^\dagger &= \mathbf{e}^{N-3} F_\omega + L_\xi^\omega \mathbf{e}^\dagger - [\mathbf{c}, \mathbf{e}^\dagger] \\ Q_{PC} \boldsymbol{\omega}^\dagger &= \mathbf{e}^{N-3} d_\omega \mathbf{e} - d_\omega \iota_{\boldsymbol{\xi}} \boldsymbol{\omega}^\dagger - [\mathbf{c}, \boldsymbol{\omega}^\dagger] + \iota_{\boldsymbol{\xi}} [\mathbf{e}, \mathbf{e}^\dagger] - \frac{1}{2} d_\omega \iota_{\boldsymbol{\xi}} \iota_{\boldsymbol{\xi}} \mathbf{c}^\dagger \\ Q_{PC} \mathbf{c}^\dagger &= -d_\omega \boldsymbol{\omega}^\dagger - [\mathbf{e}, \mathbf{e}^\dagger] - [\mathbf{c}, \mathbf{c}^\dagger] \\ Q_{PC} \boldsymbol{\xi}^\dagger &= F_{\omega \bullet} \boldsymbol{\omega}^\dagger - (d_{\omega \bullet} \mathbf{e}) \mathbf{e}^\dagger + \iota_{\boldsymbol{\xi}} F_{\omega \bullet} \mathbf{c}^\dagger + L_\xi^\omega \boldsymbol{\xi}^\dagger_\bullet + (d_\omega \iota_{\boldsymbol{\xi}} \boldsymbol{\xi}^\dagger)_\bullet. \end{aligned}$$

Here we used the symbol \bullet to remind the reader that $\boldsymbol{\xi}^\dagger$ is a one-form with values in densities, and on the right hand side we highlight the one-form part of the expression.

Remark 51. Throughout the analysis we should always keep in mind that, while Definition 50 is valid for any manifold M (possibly with boundary), the AKSZ theory obtained in Theorem 47 is by construction defined on a manifold diffeomorphic to a cylinder: $M = \Sigma \times I$. Furthermore, as we will see in this section, the fields in $\mathcal{F}_{PC^*}^{\text{AKSZ}}$ correspond to those in the standard BV theory for PC but with an additional constraint.

The product structure of M induces a splitting of fields:

$$\begin{aligned} \mathbf{e} &= \tilde{e} + \tilde{e}_n dx^n & \mathbf{e}^\dagger &= \tilde{e}^\dagger + \tilde{e}_n^\dagger dx^n \\ \boldsymbol{\omega} &= \tilde{\omega} + \tilde{\omega}_n dx^n & \boldsymbol{\omega}^\dagger &= \tilde{\omega}^\dagger + \tilde{\omega}_n^\dagger dx^n \\ \boldsymbol{\xi} &= \tilde{\xi} + \tilde{\xi}^n \partial_n & \boldsymbol{\xi}^\dagger &= \tilde{\xi}^\dagger + \tilde{\xi}_n^\dagger dx^n \end{aligned} \quad (70)$$

More compactly we can write field components in the x^n direction as $\tilde{\underline{e}}_n = \tilde{e}_n dx^n$, $\tilde{\underline{\xi}}^n = \tilde{\xi}^n \partial_n$ and so on. Observe that $\boldsymbol{\xi}^\dagger$ is a one-form with values in densities on M , so we can identify two dx^n contributions: we denote by $\tilde{\underline{\xi}}_n^\dagger$ the dx^n -component (of the one-form part) of $\boldsymbol{\xi}^\dagger$ and by $\tilde{\underline{\xi}}$ the rest, stressing that the image of $\tilde{\underline{\xi}}$ is nontrivial along dx^n . This decomposition allows us to define the maps

$$W_{\tilde{e}^{N-3}}^{i,j} : \Omega^i \left(M, \wedge^j \mathcal{V} \right) \rightarrow \Omega^{i+N-3} \left(M, \wedge^{j+N-3} \mathcal{V} \right); \quad W_{\tilde{e}^{N-3}}^{i,j}(v) = \tilde{e}^{N-3} \wedge v. \quad (71)$$

Let us now fix a nonzero section $\epsilon_n \in \Gamma(M, \mathcal{V})$ such that $d_I \epsilon_n = 0$. We will then restrict the field \tilde{e} not to have components parallel to ϵ_n . This is a restriction on the space of fields (it actually defines an open subspace). The nondegeneracy of e implies that (i) \tilde{e} and ϵ_n form a basis of V at every point, and (ii) \tilde{e}_n becomes a linear combination of \tilde{e} and ϵ_n , with nonzero ϵ_n -component. Denote by $X^{\{\mu\}}$ the components of a field X with respect to the basis given by \tilde{e} and ϵ_n (i.e. $X = X^{\{b\}} \tilde{e}_b + X^{\{n\}} \epsilon_n$).

Additionally, we consider the quantity

$$\mathfrak{W}^\dagger := \tilde{\omega}_n^\dagger - \tilde{\omega}_a^\dagger \tilde{e}_n^{\{a\}} dx^n - \iota_{\tilde{\xi}} \tilde{\omega}_n^\dagger + \tilde{c}_{an} \tilde{\xi}^n \tilde{e}_n^{\{a\}} dx^n. \quad (72)$$

Its meaning will become manifest with the following:

Definition 52. We denote by $\iota_R : \mathcal{F}_{PC}^{\text{res}} \rightarrow \mathcal{F}_{PC}$ the subspace of BV Palatini–Cartan fields defined by the following equations, which we call *PC structural constraints*:

$$\begin{aligned} \epsilon_n \tilde{e}^{(N-4)} d_{\tilde{\omega}} \tilde{e} - \epsilon_n \tilde{e}^{(N-4)} W_{\tilde{e}^{N-3}}^{-1}(\mathfrak{W}^\dagger) d\tilde{\xi}^n \\ + ([\tilde{c}, \epsilon_n] + L_{\tilde{\xi}}^{\tilde{\omega}}(\epsilon_n) - d_{\tilde{\omega}_n} \epsilon_n) \tilde{\xi}^n \}^{\{a\}} (\tilde{\omega}_a^\dagger - \tilde{c}_{an} \tilde{\xi}^n) \in \text{Im}(W_{\tilde{e}^{N-3}}^{1,1}) \end{aligned} \quad (73a)$$

$$\mathfrak{W}^\dagger \in \text{Im}(W_{\tilde{e}^{N-3}}^{1,1}) \quad (73b)$$

and by the condition that the metric $g_{\text{hor}} := \tilde{e}^* \eta$ is nowhere degenerate²⁵.

Remark 53. The PC structural constraints (73b) and (73a) are invariant under the action of Q_{PC} . Thus they define a BV theory

$$\mathfrak{F}_{PC}^{\text{res}} := (\mathcal{F}_{PC}^{\text{res}}, \varpi_{PC}^{\text{res}} = \iota_R^* \varpi_{PC}, S_{PC}^{\text{res}} = \iota_R^* S_{PC}, Q_{PC}^{\text{res}}) \quad (74)$$

where Q_{PC}^{res} is the restriction of Q_{PC} to $\mathcal{F}_{PC}^{\text{res}}$. We will call this theory the *restricted BV Palatini–Cartan theory*. The direct proof of the invariance of the constraints is lengthy and involved, yet we get this result for free as a corollary of the following theorem, which also specifies the relations between the three BV theories $\mathfrak{F}_{PC}^{\text{res}}$, \mathfrak{F}_{PC} and $\mathfrak{F}_{PC^*}^{\text{AKSZ}}$.

Theorem 54. *Upon choosing the same section $\epsilon_n \in \Gamma(M, \mathcal{V})$ and the same signature for g_{hor} in the three theories, the following diagram commutes*

$$\begin{array}{ccc} & & \mathcal{F}_{PC} \\ & \nearrow \varphi & \uparrow \iota_R \\ \mathcal{F}_{PC^*}^{\text{AKSZ}} & \xrightarrow{\varphi} & \mathcal{F}_{PC}^{\text{res}} \end{array} \quad (75)$$

²⁵Notice that the condition on g_{hor} will restrict the moduli space of solutions of the theory to an open subset.

Moreover, $\underline{\varphi}$ is a symplectomorphism and we have

$$\varpi^{\text{AKSZ}} = (\underline{\varphi} \circ \iota_R)^* \varpi_{PC}; \quad S_{PC}^{\text{AKSZ}} = (\underline{\varphi} \circ \iota_R)^* S_{PC}, \quad (76)$$

so that $\underline{\varphi}$ and ι induce a strong BV equivalence and a BV inclusion, respectively:

$$\mathfrak{F}_{PC^*}^{\text{AKSZ}} \xrightarrow{\underline{\varphi}} \mathfrak{F}_{PC}^{\text{res}} \quad \mathfrak{F}_{PC^*}^{\text{AKSZ}} \xrightarrow{\varphi} \mathfrak{F}_{PC}.$$

Remark 55 (Proof Strategy). In order to prove this, we will first show that there is an injective map $\varphi : \mathcal{F}_{PC^*}^{\text{AKSZ}} \rightarrow \mathcal{F}_{PC}$ such that $\varphi^* \varpi_{PC} = \varpi_{PC^*}^{\text{AKSZ}}$ and $\varphi^* S_{PC} = S_{PC^*}^{\text{AKSZ}}$. Note that, as a symplectomorphism, φ is then an immersion. Then we will show that $\mathcal{F}_{PC}^{\text{res}}$ is the image of this map, so that the PC structural constraints (73b) and (73a) are satisfied if and only if the AKSZ structural constraints (67b) and (67a) are. The fact that $\underline{\varphi}$ is a symplectomorphism preserving the action also proves indirectly that $\mathfrak{F}_{PC}^{\text{res}}$ is a BV theory.

Proof. Denoting by $\{\mathbf{e}, \boldsymbol{\omega}, \mathbf{c}, \boldsymbol{\xi}\}$ the fields in \mathcal{F}_{PC} (their antifields with a dagger), and following the notation of Equation (65) for the variables in $\mathcal{F}_{PC^*}^{\text{AKSZ}}$, we define the map $\varphi : \mathcal{F}_{PC^*}^{\text{AKSZ}} \rightarrow \mathcal{F}_{PC}$ in terms of the splitting (70) (with φ^* implicit on the right hand sides):

$$\varphi^* \mathbf{e} = \tilde{e} + \tilde{\underline{e}}_n \quad \varphi^* \boldsymbol{\omega} = \tilde{\omega} + \tilde{\underline{\omega}}_n \quad \varphi^* \mathbf{e}^\dagger = \tilde{e}^\dagger + \tilde{\underline{e}}_n^\dagger \quad (77a)$$

$$\varphi^* \boldsymbol{\omega}^\dagger = \tilde{\omega}^\dagger + \tilde{\underline{\omega}}_n^\dagger \quad \varphi^* \mathbf{c} = \tilde{c} \quad \varphi^* \mathbf{c}^\dagger = \tilde{c}^\dagger \quad (77b)$$

$$\varphi^* \boldsymbol{\xi} = \tilde{\xi} + \tilde{\underline{\xi}}^n \quad \varphi^* \boldsymbol{\xi}^\dagger = \tilde{\xi}^\dagger + \tilde{\underline{\xi}}_n^\dagger \quad (77c)$$

where, using again the underlined notation to signify that the quantity contains dx^n or ∂_n , and $a \in \{1, 2, \dots, N-1\}$:

$$\tilde{e} = e + \lambda\mu^{-1} f^\dagger \quad \tilde{\underline{e}}_n = \epsilon_n \underline{\mu} + \iota_{\underline{z}} e + \lambda\mu^{-1} z^a \underline{f}_a^\dagger \quad (78a)$$

$$\tilde{\omega} = \omega - \lambda\mu^{-1} u^\dagger \quad \tilde{\underline{\omega}}_n = \underline{\omega} - \iota_{\underline{\xi}} u^\dagger - \lambda\mu^{-1} z^a \underline{u}_a^\dagger \quad (78b)$$

$$\tilde{e}^\dagger = e^\dagger - \lambda\mu^{-1} y_n^\dagger \quad \tilde{\underline{e}}_n^\dagger = e^{N-3} \underline{u}^\dagger + \iota_{\underline{z}} e^\dagger - \lambda\mu^{-1} z^a \underline{y}_a^\dagger + (N-3)e^{N-4} \lambda\mu^{-1} f^\dagger \underline{u}^\dagger \quad (78c)$$

$$\tilde{\omega}^\dagger = k^\dagger \quad \tilde{\underline{\omega}}_n^\dagger = e^{N-3} \underline{f}^\dagger + \iota_{\underline{z}} k^\dagger + \iota_{\underline{\xi}} \underline{c}^\dagger \quad (78d)$$

$$\tilde{c} = c - \lambda\mu^{-1} \iota_{\underline{\xi}} u^\dagger \quad \tilde{\underline{c}}_n^\dagger = \underline{c}^\dagger \quad (78e)$$

$$\tilde{\xi}^a = \xi^a + \lambda\mu^{-1} z^a \quad \tilde{\underline{\xi}}^\dagger = e y^\dagger + f^\dagger e^\dagger - \underline{u}^\dagger k^\dagger + \underline{c}^\dagger \lambda\mu^{-1} u^\dagger \quad (78f)$$

$$\tilde{\underline{\xi}}^n = \tilde{\xi}^n \partial_n = \lambda\mu^{-1} \partial_n \quad \tilde{\underline{\xi}}_n^\dagger = e_n \underline{y}^\dagger + e^{N-3} f^\dagger \underline{u}^\dagger + f^\dagger \iota_{\underline{z}} e^\dagger + u^\dagger \iota_{\underline{z}} k^\dagger + c^\dagger \lambda\mu^{-1} z^a \underline{u}_a^\dagger \quad (78g)$$

The explicit, long but straightforward calculation needed to prove that φ is an inclusion of symplectic manifolds preserving the action functionals, i.e. such that $\varpi_{PC}^{\text{AKSZ}} = \varphi^* \varpi_{PC}$ and $S_{PC}^{\text{AKSZ}} = \varphi^* S_{PC}$, is given in Appendix A.

We then need to prove that $\text{Im}(\varphi) = \mathcal{F}_{PC}^{\text{res}}$. In other words we want to prove that the map defined in (78) will map a solution of the constraints (67b) and (67a) into a solution of (73b) and (73a). Applying (78) to the definition of \mathfrak{W}^\dagger as given in Equation (72) we get:

$$\begin{aligned} \mathfrak{W}^\dagger &= \tilde{\underline{\omega}}_n^\dagger - \tilde{\omega}_a^\dagger \tilde{e}_n^{\{a\}} dx^n - \iota_{\tilde{\underline{\xi}}} \tilde{\underline{\omega}}_n^\dagger + \tilde{c}_{an}^\dagger \tilde{\underline{\xi}}^n \tilde{e}_n^{\{a\}} dx^n \\ &= e^{N-3} \underline{f}^\dagger + \iota_{\underline{z}} k^\dagger + \iota_{\underline{\xi}} \underline{c}^\dagger - k_a^\dagger \underline{z}_a - \iota_{\underline{\xi}} \underline{c}^\dagger - \underline{c}_a^\dagger \lambda\mu^{-1} z^a + c_a^\dagger \lambda\mu^{-1} \underline{z}_a \\ &= e^{N-3} \underline{f}^\dagger, \end{aligned}$$

which is (73b). On the other hand, constraint (73a) is satisfied if (67b) and (67a) are, as it can be seen by direct inspection: using (78) we get²⁶

$$\begin{aligned}
& \epsilon_n \tilde{e}^{(N-4)} d_{\tilde{\omega}} \tilde{e} - \epsilon_n \tilde{e}^{(N-4)} W_{\tilde{e}^{N-3}}^{-1}(\mathfrak{W}^\dagger) d\tilde{\xi}^n \\
& + ([\tilde{c}, \epsilon_n] + L_{\tilde{\xi}}^\omega(\epsilon_n) - d_{\tilde{\omega}_n} \epsilon_n) \tilde{\xi}^n \{^a\} (\tilde{\omega}_a^\dagger - \tilde{c}_{an}^\dagger \tilde{\xi}^n) \\
& = \epsilon_n e^{N-4} d_\omega e + (N-4) \epsilon_n e^{N-5} \lambda \mu^{-1} f^\dagger d_\omega e + \epsilon_n e^{N-4} [\lambda \mu^{-1} u^\dagger, e] \\
& + \epsilon_n e^{N-4} d_\omega (\lambda \mu^{-1} f^\dagger) + (N-4) \epsilon_n e^{N-5} \lambda \mu^{-1} f^\dagger d_\omega (\lambda \mu^{-1} f^\dagger) \\
& + \epsilon_n f^\dagger e^{N-4} d_\omega (\lambda \mu^{-1}) + (N-4) \epsilon_n f^\dagger e^{N-5} \lambda \mu^{-1} f^\dagger d_\omega (\lambda \mu^{-1}) \\
& - \left([c, \epsilon_n] - [c, \epsilon_n]^{(b)} \lambda \mu^{-1} f_b^\dagger + [\lambda \mu^{-1} \iota_\xi u^\dagger, \epsilon_n] \right)^{(a)} (k_a^\dagger + c_a^\dagger \lambda \mu^{-1}) \\
& + \left(L_\xi^\omega(\epsilon_n) - L_\xi^\omega(\epsilon_n)^{(b)} \lambda \mu^{-1} f_b^\dagger + [\lambda \mu^{-1} \iota_\xi u^\dagger, \epsilon_n] \right)^{(a)} (k_a^\dagger + c_a^\dagger \lambda \mu^{-1}) \\
& - [w - \iota_\xi u^\dagger, \epsilon_n]^{(a)} k_a^\dagger \lambda \mu^{-1} \\
& = \epsilon_n e^{N-4} d_\omega e + (L_\xi^\omega(\epsilon_n) - [c, \epsilon_n])^{(a)} k_a^\dagger \\
& + \lambda \mu^{-1} \left(\epsilon_n \{ (N-4) f^\dagger e^{N-5} d_\omega e + e^{N-4} d_\omega f^\dagger + e^{N-4} [u^\dagger, e] \} \right. \\
& \left. + (\iota_z d_\omega \epsilon_n - [w - \iota_\xi u^\dagger, \epsilon_n])^{(a)} k_a^\dagger \right. \\
& \left. + (L_\xi^\omega(\epsilon_n) - [c, \epsilon_n])^{(a)} c_a^\dagger + \left((L_\xi^\omega(\epsilon_n) - [c, \epsilon_n])^{(b)} f_b^\dagger \right)^{(a)} k_a^\dagger \right) = (\spadesuit).
\end{aligned}$$

Using now the AKSZ constraints (67b) and (67a) we obtain

$$\begin{aligned}
(\spadesuit) & = e^{N-3} \sigma + \lambda \mu^{-1} (f^\dagger e^{N-4} \sigma + e^{N-3} B) \\
& = \tilde{e}^{N-3} (\sigma + \lambda \mu^{-1} B).
\end{aligned}$$

Comparing the first and the last line of this computation we get the desired constraint (73a). Hence φ defines a diffeomorphism $\underline{\varphi}: \mathcal{F}_{PC^*}^{\text{AKSZ}} \rightarrow \mathcal{F}_{PC}^{\text{res}}$. Indeed, the inverse of this map is readily found, as follows.

It is easy to find $k^\dagger = \tilde{\omega}^\dagger$, $\underline{c}^\dagger = \tilde{c}_n^\dagger$, and $\tilde{\xi}^n = \lambda \mu^{-1}$. Then we can write $e = \tilde{e} - \tilde{\xi}^n f^\dagger$, so that $\tilde{e}_n = \epsilon_n \mu + \iota_z \tilde{e}$, and taking $\{\tilde{e}_a, \epsilon_n\}$ as a basis, we have $z^a = \tilde{e}_n^a$ and $\mu = \tilde{e}_n^n$, which also implies $\lambda = \tilde{e}_n^n \tilde{\xi}^n$ and $\xi^a = \tilde{\xi}^a - \tilde{e}_n^a \tilde{\xi}^n$.

We now turn to equation (78d) which can be rewritten as

$$e^{N-3} \underline{f}^\dagger = \tilde{\omega}_n^\dagger - \iota_z k^\dagger - \iota_\xi \underline{c}^\dagger$$

Let us denote the known piece by $\tilde{\underline{A}} := -\iota_z k^\dagger - \iota_\xi \underline{c}^\dagger$, so that we have

$$\begin{cases} e = \tilde{e} - \tilde{\xi}^n f^\dagger \\ e^{N-3} \underline{f}^\dagger = \tilde{\omega}_n^\dagger + \tilde{\underline{A}} \end{cases} \implies \tilde{e}^{N-3} \underline{f}^\dagger = \tilde{\omega}_n^\dagger + \tilde{\underline{A}}$$

where we used that $f^\dagger f^\dagger = 0$. We see here that this equation can be solved only when

$$\tilde{\omega}_n^\dagger - \iota_z \tilde{\omega}^\dagger - \iota_\xi \tilde{c}_n^\dagger \in \text{Im}(W_{\tilde{e}^{N-3}}^{1,1}).$$

From the equations

$$\tilde{e}^\dagger = e^\dagger - \tilde{\xi}^n y^\dagger, \quad \tilde{e}_n^\dagger = e^{N-3} \underline{y}^\dagger + \iota_z e^\dagger - \lambda \mu^{-1} z^a \underline{y}_a^\dagger + (N-3) e^{N-4} \lambda \mu^{-1} f^\dagger \underline{y}^\dagger,$$

using again $e = \tilde{e} - \tilde{\xi}^n f^\dagger$, we get

$$\tilde{e}^{N-3} \underline{y}^\dagger = \tilde{e}_n^\dagger - \iota_z \tilde{e}^\dagger.$$

²⁶Note that $X^{\{a\}} = X^{(a)} - (X^{(b)} \lambda \mu^{-1} f_b^\dagger)^{(a)}$.

Since $\tilde{e}_n^\dagger - \iota_{\underline{z}}\tilde{e}^\dagger \in \Omega^{(N-2, N-1)}$, on which the map $W_{\tilde{e}^{N-3}}$ is surjective, we conclude that, up to components $p(\underline{u}^\dagger)$ in the kernel of $W_{\tilde{e}^{N-3}}$, we can find

$$u^\dagger = W_{\tilde{e}^{N-3}}^{-1}(\tilde{e}_n^\dagger - \iota_{\underline{z}}\tilde{e}^\dagger) + p\underline{u}^\dagger$$

However, we know that u^\dagger must satisfy the constraint (67a), which (implicitly but uniquely) fixes $p\underline{u}^\dagger$ as a function of $\tilde{e}, \tilde{\omega}, f^\dagger$. We can use this directly to solve

$$\omega = \tilde{\omega} + \tilde{\xi}^n u^\dagger = \tilde{\omega} + \tilde{\xi}^n (W_{\tilde{e}^{N-3}}^{-1}(\tilde{e}_n^\dagger - \iota_{\underline{z}}\tilde{e}^\dagger) + p\underline{u}^\dagger)$$

Analogously we can find \underline{w} and c as follows

$$\underline{w} = \tilde{\omega}_n + \iota_{\tilde{\xi}} (W_{\tilde{e}^{N-3}}^{-1}(\tilde{e}_n^\dagger - \iota_{\underline{z}}\tilde{e}^\dagger) + p\underline{u}^\dagger),$$

$$c = \tilde{c} + \tilde{\xi}^n \iota_{\tilde{\xi}} (W_{\tilde{e}^{N-3}}^{-1}(\tilde{e}_n^\dagger - \iota_{\underline{z}}\tilde{e}^\dagger) + p\underline{u}^\dagger).$$

Finally, we can conclude the calculation with y^\dagger and e^\dagger by inverting (78c), (78f) and (78g): it is useful to notice that it is possible to invert an equation of the form $e^\dagger(1 + \lambda X) = Y$ for some X, Y as $e^\dagger = Y(1 - \lambda X)$. However, we will not write down in full these last equations as we will not need them in what follows.

The BV theory $\mathfrak{F}_{PC_*}^{\text{AKSZ}}$ is obviously strongly equivalent to its image under the symplectomorphism φ , which is $\mathfrak{F}_{PC}^{\text{res}}$. Furthermore, since up to boundary terms Q_{PC} is the Hamiltonian vector field of S_{PC} , and the same holds for Q_{PC}^{AKSZ} and S_{PC}^{AKSZ} , we have that in the interior $M^\circ = M \setminus \partial M$ the compatibility $\varphi^* Q_{PC} = Q_{PC}^{\text{AKSZ}} \varphi^*$ is a consequence of $\varphi^* \varpi_{PC} = \varpi_{PC}^{\text{AKSZ}}$ and $\varphi^* S_{PC} = S_{PC}^{\text{AKSZ}}$. However, this is a local condition that then extends to the whole of $(M, \partial M)$ and φ is a BV inclusion. \square

Remark 56. The defining condition $\mu \neq 0$ and $g_{\text{hor}} = \tilde{e}^* \eta$ nondegenerate given in Definition 49, used in Theorem 54, are necessary in order to make \mathbf{e} non degenerate in the bulk, to build the symplectomorphism (78) (since $\epsilon_n^{[n]} = \mu^{-1}$).

Remark 57. The number of free components of $\iota_R^* \tilde{\omega}$ is $\frac{3N(N-1)}{2}$, since ω and w have respectively $N(N-1)$ and $\frac{N(N-1)}{2}$ free components. The $\frac{N(N-1)(N-3)}{2}$ missing components are those fixed by the condition in Equation (73a). Correspondingly, also $\iota_R^* \tilde{\omega}^\dagger$ has $\frac{3N(N-1)}{2}$ independent components: $\frac{N(N-1)}{2}$ coming from k^\dagger and $N(N-1)$ from f^\dagger .

4.2. An interpretation of the restricted theory. We now want to shed some light on the interpretation of the restricted theory $\mathfrak{F}_{PC}^{\text{res}}$, defined in Theorem 54.

Recall that among the Euler–Lagrange equations of the classical PC theory we have²⁷ $\mathbf{e}^{N-3} d_\omega \mathbf{e} = 0$, which, thanks to the nondegeneracy of \mathbf{e} , is equivalent to $d_\omega \mathbf{e} = 0$, i.e., the torsion-free condition for ω . Imposing this condition forces ω to correspond to the Levi-Civita connection for the metric $g_{\mu\nu} = \eta(\mathbf{e}_\mu, \mathbf{e}_\nu)$, which is used to recover the Einstein–Hilbert formulation of the theory. Note that this yields only a classical equivalence of the two theories, as the fluctuations might violate the condition $d_\omega \mathbf{e} = 0$ at the quantum level. Only by forcing this condition on the space of fields (i.e., by freezing the fluctuations that might violate it) may one recover the quantum Einstein–Hilbert theory.

However, one can consider a whole family of theories between PC and EH where only some part of the condition $d_\omega \mathbf{e} = 0$ is imposed on the fields, looking for a compromise.²⁸ that retains the good feature of PC of dealing with differential forms but yields a compatible boundary BFV theory as in EH [CS16].

²⁷We use boldface letters to denote fields in PC theory.

²⁸Imposing too few conditions out of $d_\omega \mathbf{e} = 0$ would not solve the compatibility problem with the boundary. Imposing too many generates other problems (see, e.g., [CS19c, Section 4.3], where the whole of $d_\omega \mathbf{e} = 0$ is imposed manually).

In particular, working on a cylinder $I \times \Sigma$, we may use the decomposition $\mathbf{e} = \tilde{\underline{e}}_n + \tilde{e}$, $\boldsymbol{\omega} = \tilde{\underline{\omega}}_n + \tilde{\omega}$. By choosing once and for all a nonzero section $\epsilon_n \in \Gamma(M, \mathcal{V})$ and requiring the components of \tilde{e} to span a transversal hyperplane in \mathcal{V} at each point, we may expand $\tilde{\underline{e}}_n$ in the basis (ϵ_n, \tilde{e}) ; moreover, we require \tilde{e} to define a nondegenerate metric $\eta(\tilde{e}, \tilde{e})$ at each point.²⁹ Observe that the splitting of fields $\mathbf{e}, \boldsymbol{\omega}$ induced by the cylinder structure also allows the definition of the maps $W_{\tilde{e}^{N-3}}^{i,j}$ given in Equation (71).

With these notations we may impose the constraint

$$\epsilon_n \tilde{e}^{N-3} d_{\tilde{\omega}} \tilde{e} \in \text{Im}(W_{\tilde{e}^{N-3}}^{1,1}), \quad (79)$$

which implements only some of the conditions in $d_{\boldsymbol{\omega}} \mathbf{e} = 0$.

Another interpretation of the constraints goes through considering a reduction of the fields instead of a restriction. Indeed we can also think of Equation (79) as a classical constraint that freezes certain components of the connection. We need the following

Definition 58. We define the space of reduced connections on a cylinder to be the quotient

$$\mathcal{A}^{\text{red}}(\Sigma \times I) := \mathcal{A}(\Sigma \times I) / \ker(W_{\tilde{e}^{N-3}}^{1,2}), \quad (80)$$

and denote by F_{PC}^{res} the fiber bundle

$$F_{PC}^{\text{res}} \longrightarrow \Omega_{nd}^1(\Sigma \times I, \mathcal{V}) \quad (81)$$

with typical fiber $\mathcal{A}^{\text{red}}(\Sigma \times I)$ obtained by reducing the fibers of the trivial bundle

$$\mathcal{A}(\Sigma \times I) \times \Omega_{nd}^1(\Sigma \times I, \mathcal{V}) \longrightarrow \Omega_{nd}^1(\Sigma \times I, \mathcal{V})$$

by $\ker(W_{\tilde{e}^{N-3}}^{1,2})$.

Proposition 59. Consider the splitting $\mathbf{e} = \tilde{e} + \tilde{\underline{e}}_n$, with $g_{hor} := \tilde{e}^* \eta$ nondegenerate. Then for every $([\omega], \tilde{e}, \tilde{\underline{e}}_n)$ there exists a unique $\omega \in \mathcal{A}(\Sigma \times I)$ such that

$$(N-3)\epsilon_n \wedge \tilde{e}^{N-4} \wedge d_{\omega} \tilde{e} \in \text{Im}(W_{\tilde{e}^{N-3}}^{1,1}), \quad (82)$$

which induces a section of the fibration:

$$\mathcal{A}(\Sigma \times I) \times \Omega_{nd}^1(\Sigma \times I, \mathcal{V}) \longrightarrow F_{PC}^{\text{res}}, \quad (83)$$

Proof. This is a straightforward adaptation of [CCS20, Theorem 17], which holds at every point in I . \square

Hence, imposing only some part of the equation $d_{\boldsymbol{\omega}} \mathbf{e} = 0$ produces an intermediate theory, that in view of Proposition 59 can be alternatively thought of as Palatini–Cartan theory for a tetrad and a *reduced* connection. However, in both interpretations, fixing a condition only on the classical fields does not produce a symplectic submanifold of the space of BV fields.

If we want to consistently restrict the BV theory of Definition 50 we first have to impose some condition on the antifields as well, in order to ensure that we have a nondegenerate BV form. One can show that (79) actually fixes the components of $\tilde{\omega}$ in the kernel of $W_{\tilde{e}^{N-3}}^{1,2}$. As a consequence, we can get a symplectic submanifold if, in addition to (79), we impose³⁰

$$\tilde{\omega}_n^\dagger \in \text{Im}(W_{\tilde{e}^{N-3}}^{1,1}). \quad (84)$$

²⁹A more physical requirement, as one would like the two boundary components of $I \times \Sigma$ to be space-like Cauchy surfaces, consists in choosing ϵ_n to be a time-like section and \tilde{e} to define a positive definite metric.

³⁰This condition requires that the antifield of $\tilde{\omega}$ be in the dual of the complement of the kernel of $W_{\tilde{e}^{N-3}}^{1,2}$, which is the image of $W_{\tilde{e}^{N-3}}^{1,1}$ because $\tilde{\omega}$ is tangent to the slice $\Sigma \times \{t\}$, and its antifield is of the form $\tilde{\omega}_n^\dagger dx^n$, with $\tilde{\omega}_n^\dagger \in \Omega^{N-2}(M, \wedge^{N-2}\mathcal{V})$.

The problem, though, is that (79) and (84) do not define a Q -submanifold, which is needed to have a BV theory. However, one can easily check that condition (79) is compatible with gauge transformations and diffeomorphisms upon using the Euler–Lagrange equations. This implies that it should be possible to correct (79)—and concurrently (84) because we want to preserve the condition that we get a symplectic submanifold—so as to obtain a Q -submanifold. The explicit solution to this problem is actually given by (73b) and (73a).

Remark 60. Observe that this solution might not be unique, as the choice of a structural constraint we made in Definition 25 was made to render the invariance of (19) more manifest. However, Theorem 54 tells us that a different choice of BFV structural constraint will provide a different extension of the constraint (82) in Palatini–Cartan theory.

4.3. Three dimensional case. When $N = 3$ some simplifications occur. Indeed, in this case the inclusion is actually an identity since there are no additional constraints on the field. Furthermore we know that the theory is strongly BV-equivalent, both in the bulk and on the boundary, to the topological BF theory, denoted here by $\mathfrak{F}_{BF'}^{\text{AKSZ}}$. Hence we can summarize the results in the following theorem.

Corollary 61. *The theories $\mathfrak{F}_{PC^*}^{\text{AKSZ}}$ and $\mathfrak{F}_{BF'}^{\text{AKSZ}}$ are strongly BV equivalent.*

Proof. The claim follows directly from Theorem 16 given the results of Theorem 54 and of [CS19a], which proves the strong equivalence (at all codimensions) of non-degenerate BF theory and PC gravity in three dimensions. \square

Pictorially we can describe the content of Corollary 61 as follows

$$\begin{array}{ccc}
 \mathcal{F}_{PC} & \xleftarrow{\psi} & \mathcal{F}_{BF'} \\
 \uparrow \phi & & \parallel \\
 \mathcal{F}_{PC^*}^{\text{AKSZ}} & & \mathcal{F}_{BF'}^{\text{AKSZ}} \\
 \downarrow B & & \downarrow B \\
 \mathcal{F}_{PC}^\partial & \xleftarrow{\psi^\partial} & \mathcal{F}_{BF'}^\partial
 \end{array}
 \quad (85)$$

where the arrows A represent the AKSZ constructions, the arrows B represent the BV-BFV reductions, while ψ , ψ^∂ and ϕ are the symplectomorphisms mentioned above.

APPENDIX A. LENGTHY CALCULATIONS

We prove here that the transformation (78) is a symplectomorphism between $\mathcal{F}^{\text{AKSZ}}$ and \mathcal{F}_R that preserves the action. In the computation we will use multiple times the following useful relation:

$$\epsilon_n^{[a]} = -z^a \epsilon_n^{[n]}, \quad \epsilon_n^{[n]} = \mu^{-1}.$$

We now prove that $\varphi^* \varpi_{PC} = \varpi_{PC}^{\text{AKSZ}}$.

$$\begin{aligned}
\varphi^* \varpi_{PC} &= \varphi^* \int_M \delta e \delta e^\dagger + \delta \omega \delta \omega^\dagger + \delta c \delta c^\dagger + \iota_{\delta \xi} \delta \xi^\dagger \\
&= \int_M \delta \tilde{e} \delta \tilde{e}^\dagger + \delta \tilde{e}_n \delta \tilde{e}^\dagger + \delta \tilde{\omega} \delta \tilde{\omega}^\dagger + \delta \tilde{\omega}_n \delta \tilde{\omega}^\dagger + \delta \tilde{c} \delta \tilde{c}^\dagger + \delta \tilde{\xi}^a \tilde{\xi}^\dagger + \delta \tilde{\xi}^n \tilde{\xi}^\dagger \quad (86) \\
&= \int_M \delta e \delta (e^{N-3} \underline{u}^\dagger) + \delta e \delta (\iota_{\underline{z}} e^\dagger) - \delta e \delta (\lambda \epsilon_n^{[a]} \underline{y}_a^\dagger) + \delta e \delta ((N-3) e^{N-4} \lambda \epsilon_n^{[n]} f^\dagger \underline{u}^\dagger) \\
&\quad + \delta (\lambda \epsilon_n^{[n]} f^\dagger) \delta (e \underline{u}^\dagger) + \delta (\lambda \epsilon_n^{[n]} f^\dagger) \delta (\iota_{\underline{z}} e^\dagger) - \delta (\lambda \epsilon_n^{[n]} f^\dagger) \delta (\lambda \epsilon_n^{[a]} \underline{y}_a^\dagger) \\
&\quad + \delta (\lambda \epsilon_n^{[n]} f^\dagger) \delta (\lambda \epsilon_n^{[n]} f^\dagger \underline{u}^\dagger) + \delta (\iota_{\underline{z}} e) \delta e^\dagger - \delta (\iota_{\underline{z}} e) \delta (\lambda \epsilon_n^{[n]} \underline{y}^\dagger) + \delta (\epsilon_n \underline{\mu}) \delta e^\dagger \\
&\quad - \delta (\epsilon_n \underline{\mu}) \delta (\lambda \epsilon_n^{[n]} \underline{y}^\dagger) + \delta (\lambda \epsilon_n^{[a]} \underline{f}_a^\dagger) \delta e^\dagger - \delta (\lambda \epsilon_n^{[a]} \underline{f}_a^\dagger) \delta (\lambda \epsilon_n^{[n]} \underline{y}^\dagger) \\
&\quad + \delta \omega \delta (e^{N-3} \underline{f}^\dagger) + \delta \omega \delta (\iota_{\underline{z}} k^\dagger) + \delta \omega \delta (\iota_{\underline{\xi}} \underline{c}^\dagger) - \delta (\lambda \epsilon_n^{[n]} u^\dagger) \delta (e^{N-3} \underline{f}^\dagger) \\
&\quad - \delta (\lambda \epsilon_n^{[n]} u^\dagger) \delta (\iota_{\underline{z}} k^\dagger) - \delta (\lambda \epsilon_n^{[n]} u^\dagger) \delta (\iota_{\underline{\xi}} \underline{c}^\dagger) + \delta \underline{w} \delta k^\dagger - \delta (\iota_{\underline{\xi}} \underline{u}^\dagger) \delta k^\dagger \\
&\quad - \delta (\lambda \epsilon_n^{[a]} \underline{u}_a^\dagger) \delta k^\dagger + \delta c \delta \underline{c}^\dagger - \delta (\iota_{\underline{\xi}} \lambda \epsilon_n^{[n]} u^\dagger) \delta \underline{c}^\dagger + \iota_{\delta \xi} \delta (e \underline{y}^\dagger) \\
&\quad + \iota_{\delta \xi} \delta (f^\dagger e^\dagger) - \iota_{\delta \xi} \delta (\underline{u}^\dagger k^\dagger) + \iota_{\delta \xi} \delta (\underline{c}^\dagger \lambda \epsilon_n^{[n]} u^\dagger) + \delta (\lambda \epsilon_n^{[a]}) \delta (e_a \underline{y}^\dagger) \\
&\quad + \delta (\lambda \epsilon_n^{[a]}) \delta (\underline{f}_a^\dagger e^\dagger) + \delta (\lambda \epsilon_n^{[a]}) \delta (\underline{u}_a^\dagger k^\dagger) + \delta (\lambda \epsilon_n^{[a]}) \delta (\underline{c}_a^\dagger \lambda \epsilon_n^{[n]} u^\dagger) \\
&\quad + \delta (\lambda \epsilon_n^{[n]}) \delta (e_n \underline{y}^\dagger) + \delta (\lambda \epsilon_n^{[n]}) \delta (e^{N-3} f^\dagger \underline{u}^\dagger) + \delta (\lambda \epsilon_n^{[n]}) \delta (f^\dagger \iota_{\underline{z}} e^\dagger) \\
&\quad + \delta (\lambda \epsilon_n^{[n]}) \delta (u^\dagger \iota_{\underline{z}} k^\dagger) + \delta (\lambda \epsilon_n^{[n]}) \delta (c^\dagger \lambda \epsilon_n^{[a]} \underline{u}_a^\dagger)
\end{aligned}$$

This expression should be compared with the symplectic form coming from the AKSZ construction:

$$\begin{aligned}
\varpi_{PC}^{\text{AKSZ}} &= \int_{I \times \partial M} \delta (e^{N-3} f^\dagger) \delta \omega + e^{N-3} \delta e \delta u^\dagger + \delta w \delta k^\dagger + \delta c \delta c^\dagger + \delta u^\dagger \delta (\iota_{\underline{\xi}} k^\dagger) \\
&\quad + \delta \omega \delta (\iota_{\underline{z}} k^\dagger) + \delta \omega \delta (\iota_{\underline{\xi}} \underline{c}^\dagger) - \delta \mu \epsilon_n \delta e^\dagger - \delta \lambda \epsilon_n \delta \underline{y}^\dagger \\
&\quad + \iota_{\delta z} \delta (e e^\dagger) + \iota_{\delta \xi} \delta (f^\dagger e^\dagger) + \iota_{\delta \xi} \delta (e \underline{y}^\dagger). \quad (87)
\end{aligned}$$

Almost all the terms in (87) can be directly found in $\varphi^* \varpi_{PC}$. The remaining terms can be identified using the following relations:

$$\begin{aligned}
\delta \underline{u}^\dagger \delta (\iota_{\underline{\xi}} k^\dagger) &= -\delta (\iota_{\underline{\xi}} \underline{u}^\dagger) \delta k^\dagger - \iota_{\delta \xi} \delta (\underline{u}^\dagger k^\dagger); \\
-\delta (\lambda \epsilon_n) \delta \underline{y}^\dagger &= \delta (e_a \lambda \epsilon_n^{[a]}) \delta \underline{y}^\dagger + \delta (\tilde{e}_n \lambda \epsilon_n^{[n]}) \delta \underline{y}^\dagger \\
&= \delta (\lambda \epsilon_n^{[a]}) \delta (e_a \underline{y}^\dagger) - \delta e \delta (\lambda \epsilon_n^{[a]} \underline{y}_a^\dagger) \\
&\quad + \delta (\lambda \epsilon_n^{[n]}) \delta (\tilde{e}_n \underline{y}^\dagger) - \delta \tilde{e}_n \delta (\lambda \epsilon_n^{[n]} \underline{y}^\dagger); \\
\iota_{\delta \underline{z}} \delta (e e^\dagger) &= \delta e \delta (\iota_{\underline{z}} e^\dagger) + \delta (\iota_{\underline{z}} e) \delta e^\dagger
\end{aligned}$$

All the other terms in (86) sum to zero because of the following identities:

$$\begin{aligned}
& \delta(\lambda\epsilon_n^{[n]}f^\dagger)\delta(e^{N-3}\underline{u}^\dagger) + (N-3)\delta e\delta(e^{N-4}\lambda\epsilon_n^{[n]}f^\dagger\underline{u}^\dagger) \\
& - \delta(\lambda\epsilon_n^{[n]}u^\dagger)\delta(e^{N-3}\underline{f}^\dagger) + \delta(\lambda\epsilon_n^{[n]})\delta(e^{N-3}f^\dagger\underline{u}^\dagger) = 0; \\
& \delta(\lambda\epsilon_n^{[n]}f^\dagger)\delta(\underline{u}_z e^\dagger) + \delta(\lambda\epsilon_n^{[a]}\underline{f}_a^\dagger)\delta e^\dagger + \delta(\lambda\epsilon_n^{[a]})\delta(\underline{f}_a^\dagger e^\dagger) + \delta(\lambda\epsilon_n^{[n]})\delta(f^\dagger\underline{u}_z e^\dagger) = 0; \\
& - \delta(\lambda\epsilon_n^{[n]}u^\dagger)\delta(\underline{u}_z k^\dagger) - \delta(\lambda\epsilon_n^{[a]}\underline{u}_a^\dagger)\delta k^\dagger + \delta(\lambda\epsilon_n^{[a]})\delta(\underline{u}_a^\dagger k^\dagger) + \delta(\lambda\epsilon_n^{[n]})\delta(u^\dagger\underline{u}_z k^\dagger) = 0; \\
& - \delta(\lambda\epsilon_n^{[n]}u^\dagger)\delta(\underline{u}_\xi \underline{c}^\dagger) - \delta(\underline{u}_\xi \lambda\epsilon_n^{[n]}u^\dagger)\delta \underline{c}^\dagger + \underline{u}_\xi \delta(\underline{c}^\dagger \lambda\epsilon_n^{[n]}u^\dagger) = 0; \\
& \delta(\lambda\epsilon_n^{[a]})\delta(\underline{c}_a^\dagger \lambda\epsilon_n^{[n]}u^\dagger) + \delta(\lambda\epsilon_n^{[n]})\delta(c^\dagger \lambda\epsilon_n^{[a]}\underline{u}_a^\dagger) = 0; \\
& - \delta(\lambda\epsilon_n^{[n]}f^\dagger)\delta(\lambda\epsilon_n^{[a]}\underline{y}_a^\dagger) - \delta(\lambda\epsilon_n^{[a]}f^\dagger)\delta(\lambda\epsilon_n^{[n]}y^\dagger) = 0; \\
& (N-3)\delta(\lambda\epsilon_n^{[n]}f^\dagger)\delta(e^{N-4}\lambda\epsilon_n^{[n]}f^\dagger\underline{u}^\dagger) = 0.
\end{aligned}$$

We go on to show that the symplectomorphism φ preserves the action i.e. $\varphi^*S_{PC} = S_{PC}^{\text{AKSZ}}$. We do it by direct inspection³¹:

$$\begin{aligned}
\varphi^*S_{PC} &= \varphi^* \int_M \frac{1}{N-2} e^{N-2} F_\omega + (d_\omega \mathbf{c} - \underline{u}_\xi F_\omega) \omega^\dagger + (L_\xi \mathbf{e} - [\mathbf{c}, \mathbf{e}]) \mathbf{e}^\dagger \quad (88) \\
& + \frac{1}{2} ([\mathbf{c}, \mathbf{c}] - \underline{u}_\xi \underline{u}_\xi F_\omega) \mathbf{c}^\dagger + \frac{1}{2} \iota_{[\xi, \xi]} \xi^\dagger \\
& = \int_M \tilde{e}^{N-3} \tilde{\underline{e}}_n F_{\tilde{\omega}} + \frac{1}{N-2} \tilde{e}^{N-2} \tilde{F}_{\tilde{\omega}_n} - \left(\iota_{\tilde{\xi}} F_{\tilde{\omega}} + F_{\tilde{\omega}_n} \tilde{\xi}_n - d_{\tilde{\omega}} \tilde{c} \right) \tilde{\omega}_n^\dagger \\
& - \left(\iota_{\tilde{\xi}} \tilde{F}_{\tilde{\omega}_n} - d_{\tilde{\omega}_n} \tilde{c} \right) \tilde{\omega}^\dagger + \left(L_{\tilde{\xi}} \tilde{e} + d_{\tilde{\omega}_n} \tilde{e} \tilde{\xi}_n + \tilde{e}_n d \tilde{\xi}_n - [\tilde{c}, \tilde{e}] \right) \tilde{\underline{e}}_n^\dagger \\
& + \left(\iota_{\tilde{\xi}} d_{\tilde{\omega}} \tilde{e}_n + \iota_{\tilde{\omega}_n} \tilde{\xi} \tilde{e} - d_{\tilde{\omega}_n} (\tilde{e}_n \tilde{\xi}_n) - [\tilde{c}, \tilde{e}_n] \right) \tilde{e}^\dagger \\
& - \left(\frac{1}{2} \iota_{\tilde{\xi}} \iota_{\tilde{\xi}} F_{\tilde{\omega}} + \iota_{\tilde{\xi}} F_{\tilde{\omega}_n} \tilde{\xi}_n - \frac{1}{2} [\tilde{c}, \tilde{c}] \right) \tilde{\underline{c}}^\dagger + \frac{1}{2} \iota_{[\tilde{\xi}, \tilde{\xi}]} \tilde{\xi}_n^\dagger + \frac{1}{2} \iota_{[\tilde{\xi}, \tilde{\xi}]} \tilde{\xi}_n^\dagger \\
& = \int_M e^{N-3} \epsilon_n \underline{\mu} F_\omega + e^{N-3} \underline{u}_z e F_\omega + e^{N-3} \lambda \epsilon_n^{[a]} \underline{f}_a^\dagger F_\omega + (N-3) e^{N-4} \lambda \epsilon_n^{[n]} f^\dagger \underline{e}_n F_\omega \\
& - e^{N-3} \underline{e}_n d_\omega (\lambda \epsilon_n^{[n]} u^\dagger) - e^{N-3} \lambda \epsilon_n^{[a]} \underline{f}_a^\dagger d_\omega (\lambda \epsilon_n^{[n]} u^\dagger) \\
& - (N-3) e^{N-4} \lambda \epsilon_n^{[n]} f^\dagger \underline{e}_n d_\omega (\lambda \epsilon_n^{[n]} u^\dagger) + \frac{1}{N-2} e^{N-2} \left(\partial_n \omega - \partial_n \lambda \epsilon_n^{[n]} u^\dagger + d_\omega \omega \right) \\
& - \frac{1}{N-2} e^{N-2} \left(d_\omega (\underline{u}_\xi \underline{u}^\dagger) + d_\omega (\lambda \epsilon_n^{[a]} \underline{u}_a^\dagger) + [\lambda \epsilon_n^{[n]} u^\dagger, \omega - \underline{u}_\xi \underline{u}^\dagger] \right) + e^{N-3} \lambda \epsilon_n^{[n]} f^\dagger F_{\omega_n} \\
& - \underline{u}_\xi F_\omega (e^{N-3} \underline{f}^\dagger + \underline{u}_z k^\dagger + \underline{u}_\xi \underline{c}^\dagger) + \underline{u}_\xi d_\omega (\lambda \epsilon_n^{[n]} u^\dagger) (e^{N-3} \underline{f}^\dagger + \underline{u}_z k^\dagger + \underline{u}_\xi \underline{c}^\dagger) \\
& - F_{\omega_a} \lambda \epsilon_n^{[a]} (e^{N-3} \underline{f}^\dagger + \underline{u}_z k^\dagger + \underline{u}_\xi \underline{c}^\dagger) + d_\omega (\lambda \epsilon_n^{[n]} u^\dagger)_a \lambda \epsilon_n^{[a]} (e^{N-3} \underline{f}^\dagger + \underline{u}_z k^\dagger + \underline{u}_\xi \underline{c}^\dagger) \\
& - F_{\tilde{\omega}_n} \lambda \epsilon_n^{[n]} e^{N-3} \underline{f}^\dagger - F_{\tilde{\omega}_n} \lambda \epsilon_n^{[n]} \underline{u}_z k^\dagger - F_{\tilde{\omega}_n} \lambda \epsilon_n^{[n]} \underline{u}_\xi \underline{c}^\dagger \\
& + d_\omega c (e^{N-3} \underline{f}^\dagger + \underline{u}_z k^\dagger + \underline{u}_\xi \underline{c}^\dagger) - [\lambda \epsilon_n^{[n]} u^\dagger, c] (e^{N-3} \underline{f}^\dagger + \underline{u}_z k^\dagger + \underline{u}_\xi \underline{c}^\dagger) \\
& - d_\omega (\underline{u}_\xi \lambda \epsilon_n^{[n]} u^\dagger) (e^{N-3} \underline{f}^\dagger + \underline{u}_z k^\dagger + \underline{u}_\xi \underline{c}^\dagger)
\end{aligned}$$

³¹We denote with ∂_n the de Rham differential on I (previously denoted with d_I) in order to be uniform with the notation of the fields (77).

$$\begin{aligned}
& -\iota_\xi \underline{\partial}_n \omega k^\dagger + \iota_\xi \underline{\partial}_n (\lambda \epsilon_n^{[n]} u^\dagger) k^\dagger - \iota_\xi d_\omega \underline{w} k^\dagger + \iota_\xi d_\omega (\iota_\xi \underline{u}^\dagger) k^\dagger \\
& + \iota_\xi d_\omega (\lambda \epsilon_n^{[a]} \underline{u}_a^\dagger) k^\dagger + \iota_\xi [\lambda \epsilon_n^{[n]} u^\dagger, \underline{w} - \iota_\xi \underline{u}^\dagger] k^\dagger - \underline{F}_{\tilde{\omega}_a n} \lambda \epsilon_n^{[a]} k^\dagger + \underline{\partial}_n c k^\dagger \\
& - \underline{\partial}_n (\iota_\xi \lambda \epsilon_n^{[n]} u^\dagger) k^\dagger + [\underline{w}, c] k^\dagger - [\iota_\xi \underline{u}^\dagger, c] k^\dagger - [\lambda \epsilon_n^{[a]} \underline{u}_a^\dagger, c] k^\dagger \\
& + [\underline{w} - \iota_\xi \underline{u}^\dagger, \iota_\xi \lambda \epsilon_n^{[n]} u^\dagger] k^\dagger \\
& + L_\xi^\omega e e^{N-3} \underline{u}^\dagger + L_\xi^\omega e \iota_{\underline{z}} e^\dagger - L_\xi^\omega e \lambda \epsilon_n^{[a]} \underline{y}_a^\dagger + (N-3) L_\xi^\omega e e^{N-4} \lambda \epsilon_n^{[n]} f^\dagger \underline{u}^\dagger \\
& + ((d_\omega e)_a \lambda \epsilon_n^{[a]} - d_\omega (e_a \lambda \epsilon_n^{[a]})) e^{N-3} \underline{u}^\dagger + ((d_\omega e)_a \lambda \epsilon_n^{[a]} - d_\omega (e_a \lambda \epsilon_n^{[a]})) \iota_{\underline{z}} e^\dagger \\
& + d_\omega (e_a \lambda \epsilon_n^{[a]}) \lambda \epsilon_n^{[a]} \underline{y}_a^\dagger - (N-3) d_\omega (e_a \lambda \epsilon_n^{[a]}) e^{N-4} \lambda \epsilon_n^{[n]} f^\dagger \underline{u}^\dagger + L_\xi^\omega (\lambda \epsilon_n^{[n]} f^\dagger) e^{N-3} \underline{u}^\dagger \\
& + L_\xi^\omega (\lambda \epsilon_n^{[n]} f^\dagger) \iota_{\underline{z}} e^\dagger - L_\xi^\omega (\lambda \epsilon_n^{[n]} f^\dagger) \lambda \epsilon_n^{[a]} \underline{y}_a^\dagger + (d_\omega (\lambda \epsilon_n^{[n]} f^\dagger))_a \lambda \epsilon_n^{[a]} e^{N-3} \underline{u}^\dagger \\
& + (d_\omega (\lambda \epsilon_n^{[n]} f^\dagger))_a \lambda \epsilon_n^{[a]} \iota_{\underline{z}} e^\dagger - [\iota_\xi \lambda \epsilon_n^{[n]} u^\dagger, e] (e^{N-3} \underline{u}^\dagger + \iota_{\underline{z}} e^\dagger) + \partial_n e \lambda \epsilon_n^{[n]} e^{N-3} \underline{u}^\dagger \\
& + \partial_n e \lambda \epsilon_n^{[n]} \iota_{\underline{z}} e^\dagger + [w - \iota_\xi u^\dagger, e] \lambda \epsilon_n^{[n]} e^{N-3} \underline{u}^\dagger + [w - \iota_\xi u^\dagger, e] \lambda \epsilon_n^{[n]} \iota_{\underline{z}} e^\dagger \\
& + \partial_n (\lambda \epsilon_n^{[n]} f^\dagger) \lambda \epsilon_n^{[n]} (e^{N-3} \underline{u}^\dagger + \iota_{\underline{z}} e^\dagger) + \underline{e}_n d (\lambda \epsilon_n^{[n]}) e^{N-3} \underline{u}^\dagger + \underline{e}_n d (\lambda \epsilon_n^{[n]}) \iota_{\underline{z}} e^\dagger \\
& + \lambda \epsilon_n^{[a]} f_a^\dagger d (\lambda \epsilon_n^{[n]}) e^{N-3} \underline{u}^\dagger + \lambda \epsilon_n^{[a]} f_a^\dagger d (\lambda \epsilon_n^{[n]}) \iota_{\underline{z}} e^\dagger - \underline{e}_n d (\lambda \epsilon_n^{[n]}) \lambda \epsilon_n^{[a]} \underline{y}_a^\dagger \\
& + (N-3) \underline{e}_n d (\lambda \epsilon_n^{[n]}) e^{N-4} \lambda \epsilon_n^{[n]} f^\dagger \underline{u}^\dagger - [c, e] e^{N-3} \underline{u}^\dagger - [c, e] \iota_{\underline{z}} e^\dagger + [c, e] \lambda \epsilon_n^{[a]} \underline{y}_a^\dagger \\
& - (N-3) [c, e] e^{N-4} \lambda \epsilon_n^{[n]} f^\dagger \underline{u}^\dagger - [c, \lambda \epsilon_n^{[n]} f^\dagger] e^{N-3} \underline{u}^\dagger - [c, \lambda \epsilon_n^{[n]} f^\dagger] \iota_{\underline{z}} e^\dagger \\
& + [\iota_\xi \lambda \epsilon_n^{[n]} u^\dagger, e] (e^{N-3} \underline{u}^\dagger + \iota_{\underline{z}} e^\dagger) \\
& + \iota_\xi d_\omega (\epsilon_n \underline{\mu}) e^\dagger + \iota_\xi d_\omega (\iota_{\underline{z}} e) e^\dagger + \iota_\xi d_\omega (\lambda \epsilon_n^{[a]} \underline{f}_a^\dagger) e^\dagger + d_{\omega_a} (\underline{e}_n) \lambda \epsilon_n^{[a]} e^\dagger \\
& + d_{\omega_a} (\lambda \epsilon_n^{[a]} \underline{f}_a^\dagger) \lambda \epsilon_n^{[a]} e^\dagger - \iota_\xi [\lambda \epsilon_n^{[n]} u^\dagger, \underline{e}_n] e^\dagger - \iota_\xi d_\omega (\underline{e}_n) \lambda \epsilon_n^{[n]} \underline{y}_n^\dagger \\
& - \iota_\xi d_\omega (\lambda \epsilon_n^{[a]} \underline{f}_a^\dagger) \lambda \epsilon_n^{[n]} \underline{y}_n^\dagger - \iota_{\underline{\partial}_n} \xi e e^\dagger + e_a \underline{\partial}_n (\lambda \epsilon_n^{[a]}) e^\dagger - \iota_{\underline{\partial}_n} \xi \lambda \epsilon_n^{[n]} f^\dagger e^\dagger \\
& + \iota_{\underline{\partial}_n} \xi e \lambda \epsilon_n^{[n]} \underline{y}_n^\dagger - \underline{\partial}_n (\tilde{e}_n \lambda \epsilon_n^{[n]}) e^\dagger - [\underline{w} - \iota_\xi \underline{u}^\dagger, \tilde{e}_n \lambda \epsilon_n^{[n]}] e^\dagger \\
& + \underline{\partial}_n (\tilde{e}_n \lambda \epsilon_n^{[n]}) \lambda \epsilon_n^{[n]} \underline{y}_n^\dagger - [c, \epsilon_n \underline{\mu}] e^\dagger - [c, \iota_{\underline{z}} e] e^\dagger - [c, \lambda \epsilon_n^{[a]} \underline{f}_a^\dagger] e^\dagger \\
& + [\iota_\xi \lambda \epsilon_n^{[n]} u^\dagger, \underline{e}_n] e^\dagger + [c, \underline{e}_n] \lambda \epsilon_n^{[n]} \underline{y}_n^\dagger \\
& - \frac{1}{2} \iota_\xi \iota_\xi F_\omega \underline{c}^\dagger + \frac{1}{2} \iota_\xi \iota_\xi d_\omega (\lambda \epsilon_n^{[n]} u^\dagger) \underline{c}^\dagger - \iota_\xi F_{\omega_a} \lambda \epsilon_n^{[a]} \underline{c}^\dagger \\
& + \iota_\xi d_\omega (\lambda \epsilon_n^{[n]} u^\dagger)_a \lambda \epsilon_n^{[a]} \underline{c}^\dagger - \iota_\xi F_{\tilde{\omega}_n} \lambda \epsilon_n^{[n]} \underline{c}^\dagger + \frac{1}{2} [c, c] \underline{c}^\dagger - [\iota_\xi \lambda \epsilon_n^{[n]} u^\dagger, c] \underline{c}^\dagger
\end{aligned}$$

$$\begin{aligned}
& -\xi^b \partial_b \xi^a e_a \underline{y}^\dagger - \xi^b \partial_b (\lambda \epsilon_n^{[a]}) e_a \underline{y}^\dagger - \lambda \epsilon_n^{[b]} \partial_b \xi^a e_a \underline{y}^\dagger - \lambda \epsilon_n^{[b]} \partial_b (\lambda \epsilon_n^{[a]}) e_a \underline{y}^\dagger \\
& - \lambda \epsilon_n^{[n]} \partial_n \xi^a e_a \underline{y}^\dagger - \xi^b \partial_b \xi^a \underline{f}_a^\dagger e^\dagger - \xi^b \partial_b (\lambda \epsilon_n^{[a]}) \underline{f}_a^\dagger e^\dagger - \lambda \epsilon_n^{[b]} \partial_b \xi^a \underline{f}_a^\dagger e^\dagger \\
& - \lambda \epsilon_n^{[b]} \partial_b (\lambda \epsilon_n^{[a]}) \underline{f}_a^\dagger e^\dagger - \lambda \epsilon_n^{[n]} \partial_n \xi^a \underline{f}_a^\dagger e^\dagger - \xi^b \partial_b \xi^a \underline{u}_a^\dagger k^\dagger - \xi^b \partial_b (\lambda \epsilon_n^{[a]}) \underline{u}_a^\dagger k^\dagger \\
& - \lambda \epsilon_n^{[b]} \partial_b \xi^a \underline{u}_a^\dagger k^\dagger + \lambda \epsilon_n^{[b]} \partial_b (\lambda \epsilon_n^{[a]}) \underline{u}_a^\dagger k^\dagger - \lambda \epsilon_n^{[n]} \partial_n \xi^a \underline{u}_a^\dagger k^\dagger \\
& - \lambda \epsilon_n^{[n]} \partial_n (\lambda \epsilon_n^{[a]}) \underline{u}_a^\dagger k^\dagger - \xi^b \partial_b \xi^a \underline{c}_a^\dagger \lambda \epsilon_n^{[n]} u^\dagger - \xi^b \partial_b (\lambda \epsilon_n^{[a]}) \underline{c}_a^\dagger \lambda \epsilon_n^{[n]} u^\dagger \\
& - \xi^a \partial_a (\lambda \epsilon_n^{[n]}) \tilde{e}_n \underline{y}^\dagger - \lambda \epsilon_n^{[a]} \partial_a (\lambda \epsilon_n^{[n]}) \tilde{e}_n \underline{y}^\dagger - \lambda \epsilon_n^{[n]} \partial_n (\lambda \epsilon_n^{[n]}) \tilde{e}_n \underline{y}^\dagger \\
& - \xi^a \partial_a (\lambda \epsilon_n^{[n]}) f^\dagger e^{N-3} \underline{u}^\dagger - \lambda \epsilon_n^{[a]} \partial_a (\lambda \epsilon_n^{[n]}) f^\dagger e^{N-3} \underline{u}^\dagger - \lambda \epsilon_n^{[n]} \partial_n (\lambda \epsilon_n^{[n]}) f^\dagger e^{N-3} \underline{u}^\dagger \\
& - \xi^a \partial_a (\lambda \epsilon_n^{[n]}) f^\dagger \underline{v}_z e^\dagger - \lambda \epsilon_n^{[a]} \partial_a (\lambda \epsilon_n^{[n]}) f^\dagger \underline{v}_z e^\dagger - \lambda \epsilon_n^{[n]} \partial_n (\lambda \epsilon_n^{[n]}) f^\dagger \underline{v}_z e^\dagger \\
& + \xi^a \partial_a (\lambda \epsilon_n^{[n]}) u^\dagger \underline{v}_z k^\dagger + \lambda \epsilon_n^{[a]} \partial_a (\lambda \epsilon_n^{[n]}) u^\dagger \underline{v}_z k^\dagger + \lambda \epsilon_n^{[n]} \partial_n (\lambda \epsilon_n^{[n]}) u^\dagger \underline{v}_z k^\dagger \\
& - \xi^a \partial_a (\lambda \epsilon_n^{[n]}) c^\dagger \lambda \epsilon_n^{[a]} \underline{u}_a^\dagger
\end{aligned}$$

We want to compare this with the AKSZ action:

$$\begin{aligned}
S_{PC}^{\text{AKSZ}} &= \int_{I \times \partial M} \underline{w} e^{N-3} d_\omega e + (N-3) c e^{N-4} \underline{f}^\dagger d_\omega e + c e^{N-3} [\underline{u}^\dagger, e] + c e^{N-3} d_\omega \underline{f}^\dagger \\
&+ \underline{v}_z e e^{N-3} F_\omega + \underline{v}_z (e^{N-3} \underline{f}^\dagger) F_\omega + \underline{v}_z e e^{N-3} d_\omega \underline{u}^\dagger + \epsilon_n \underline{\mu} e^{N-3} F_\omega \\
&+ (N-3) \epsilon_n \lambda e^{N-4} \underline{f}^\dagger F_\omega + \epsilon_n \lambda e^{N-3} d_\omega \underline{u}^\dagger + [\underline{w}, c] k^\dagger + \frac{1}{2} [c, c] \underline{c}^\dagger \\
&- \underline{v}_z d_\omega c k^\dagger - [\underline{v}_z \underline{u}^\dagger, c] k^\dagger - \underline{v}_z d_\omega \underline{w} k^\dagger - \underline{v}_z d_\omega c \underline{c}^\dagger + \underline{v}_z \underline{v}_z F_\omega k^\dagger \\
&+ \frac{1}{2} \underline{v}_z \underline{v}_z d_\omega \underline{u}^\dagger k^\dagger + \frac{1}{2} \underline{v}_z \underline{v}_z F_\omega \underline{c}^\dagger - [\underline{w}, \epsilon_n \lambda] e^\dagger - [c, \epsilon_n \underline{\mu}] e^\dagger - [c, \epsilon_n \lambda] \underline{y}^\dagger \\
&+ \underline{v}_z d_\omega (\epsilon_n \lambda) e^\dagger + [\underline{v}_z \underline{u}^\dagger, \epsilon_n \lambda] e^\dagger + \underline{v}_z d_\omega (\epsilon_n \underline{\mu}) e^\dagger + \underline{v}_z d_\omega (\epsilon_n \lambda) \underline{y}^\dagger + \underline{v}_z \underline{v}_z e e^\dagger \\
&+ \frac{1}{2} \underline{v}_z [\underline{v}_z, \underline{v}_z] \underline{f}^\dagger e^\dagger + \frac{1}{2} \underline{v}_z [\underline{v}_z, \underline{v}_z] e \underline{y}^\dagger + \frac{1}{N-2} e^{N-2} \partial_n \omega + c \partial_n k^\dagger + \partial_n \omega \underline{v}_z k^\dagger \\
&- \underline{v}_z \underline{v}_z e e^\dagger + \partial_n \lambda \epsilon_n e^\dagger. \tag{89}
\end{aligned}$$

We proceed as follows: we first check that all terms in (89) can be found in (88), then we show that all other terms in (88) sum to zero.

We can easily recognize many terms identically repeated in both expressions. Some other terms in (89) can be recovered in (88) using Leibniz rule and Cartan calculus.

$$\begin{aligned}
(N-3) c e^{N-4} \underline{f}^\dagger d_\omega e + c^{N-3} d_\omega \underline{f}^\dagger &= d_\omega c (e^{N-3} \underline{f}^\dagger); \\
-\frac{1}{N-2} e^{N-2} d_\omega (\underline{v}_z \underline{u}^\dagger) + L_\xi^\omega e e^{N-3} \underline{u}^\dagger &= + \underline{v}_z e e^{N-3} d_\omega \underline{u}^\dagger; \\
-\frac{1}{2} \underline{v}_z [\underline{v}_z, \underline{v}_z] \underline{u}^\dagger k^\dagger &= - \underline{v}_z d_\omega \underline{v}_z \underline{u}^\dagger k^\dagger + \frac{1}{2} \underline{v}_z \underline{v}_z d_\omega \underline{u}^\dagger k^\dagger;
\end{aligned}$$

$$\underline{v}_z [\underline{v}_z, \underline{v}_z] e e^\dagger = \underline{v}_z d_\omega \underline{v}_z e e^\dagger + L_\xi^\omega \underline{v}_z e e^\dagger$$

All the other relations involving terms of (88) are based on the expansion

$$\epsilon_n = e_a \epsilon_n^{(a)} + e_n \epsilon_n^{(n)}.$$

It is a long but rather easy computation to show that the remaining terms in (89) sum to zero. This is done by making repeated use of Cartan calculus and Leibniz rule. Notice also that some terms containing expressions of the form $\epsilon_n^{[a]} \epsilon_n^{[b]} X_{ab}$ vanish by antisymmetry.

REFERENCES

- [Ale+97] M. Aleksandrov, M. Kontsevich, A. Schwarz, and O. Zaboronsky. “The geometry of the master equation and topological quantum field theory”. *Int. J. Mod. Phys. A* (12 1997), pp. 1405–1430. DOI: 10.1142/S0217751X97001031.
- [AG14] K. Alkalaev and M. Grigoriev. “Frame-like Lagrangians and presymplectic AKSZ-type sigma models”. *International Journal of Modern Physics A* 29.18 (2014), p. 1450103. DOI: 10.1142/S0217751X14501036.
- [BBH95] G. Barnich, F. Brandt, and M. Henneaux. “Local BRST cohomology in the antifield formalism. I. General theorems”. *Comm. Math. Phys.* 174.1 (1995), pp. 57–91.
- [BG05] G. Barnich and M. Grigoriev. “Hamiltonian BRST and Batalin-Vilkovisky Formalisms for Second Quantization of Gauge Theories”. *Communications in Mathematical Physics* 254.3 (2005), pp. 581–601. DOI: 10.1007/s00220-004-1275-4.
- [BG11] G. Barnich and M. Grigoriev. “First order parent formulation for generic gauge field theories”. *Journal of High Energy Physics* 2011.1 (2011), p. 122. DOI: 10.1007/JHEP01(2011)122.
- [BF83] I. A. Batalin and E. S. Fradkin. “A generalized canonical formalism and quantization of reducible gauge theories”. *Physics Letters B* 122.2 (Mar. 1983), pp. 157–164. DOI: 10.1016/0370-2693(83)90784-0.
- [BV77] I. A. Batalin and G. A. Vilkovisky. “Relativistic S-matrix of dynamical systems with boson and fermion constraints”. *Physics Letters B* 69.3 (Aug. 1977), pp. 309–312. DOI: 10.1016/0370-2693(77)90553-6.
- [BV81] I. A. Batalin and G. A. Vilkovisky. “Gauge algebra and quantization”. *Physics Letters B* 102.1 (June 1981), pp. 27–31. DOI: 10.1016/0370-2693(81)90205-7.
- [BFW13] C. Blohmann, M. C. B. Fernandes, and A. Weinstein. “Groupoid symmetry and constraints in general relativity”. *Communications in Contemporary Mathematics* 15.01 (Jan. 2013), p. 1250061. ISSN: 1793-6683. DOI: 10.1142/s0219199712500617.
- [BW18] C. Blohmann and A. Weinstein. “Hamiltonian Lie algebroids” (2018). arXiv: 1811.11109 [math.SG].
- [BFR16] R. Brunetti, K. Fredenhagen, and K. Rejzner. “Quantum Gravity from the Point of View of Locally Covariant Quantum Field Theory”. *Communications in Mathematical Physics* 345.3 (2016), pp. 741–779. DOI: 10.1007/s00220-016-2676-x.
- [CCS20] G. Canepa, A. S. Cattaneo, and M. Schiavina. “Boundary structure of General Relativity in tetrad variables” (2020). To appear in *Advances in Theoretical and Mathematical Physics*. arXiv: 2001.11004 [math-ph].
- [CS19a] G. Canepa and M. Schiavina. “Fully extended BV-BFV description of General Relativity in three dimensions” (2019). arXiv: 1905.09333 [math-ph].
- [CSS18] A. S. Cattaneo, M. Schiavina, and I. Seliash. “BV equivalence between triadic gravity and BF theory in three dimensions”. *Letters in Mathematical Physics* 108 (Aug. 2018), pp. 1873–1884. DOI: 10.1007/s11005-018-1060-5.
- [CMR11] A. S. Cattaneo, P. Mnev, and N. Reshetikhin. “Classical and quantum Lagrangian field theories with boundary”. *PoS CORFU2011* (2011), p. 044. DOI: 10.22323/1.155.0044. arXiv: 1207.0239 [math-ph].
- [CMR14] A. S. Cattaneo, P. Mnev, and N. Reshetikhin. “Classical BV Theories on Manifolds with Boundary”. *Communications in Mathematical Physics* 332.2 (2014), pp. 535–603. ISSN: 1432-0916. DOI: 10.1007/s00220-014-2145-3.
- [CMR18] A. S. Cattaneo, P. Mnev, and N. Reshetikhin. “Perturbative Quantum Gauge Theories on Manifolds with Boundary”. *Communications in Mathematical Physics* 357.2 (Jan. 2018), pp. 631–730. DOI: 10.1007/s00220-017-3031-6.

- [CMR20] A. S. Cattaneo, P. Mnev, and N. Reshetikhin. “A Cellular Topological Field Theory”. *Communications in Mathematical Physics* 374.2 (2020), pp. 1229–1320. URL: <https://doi.org/10.1007/s00220-020-03687-3>.
- [CMW17] A. S. Cattaneo, P. Mnev, and K. Wernli. “Split Chern–Simons Theory in the BV-BFV Formalism”. *9th Summer School on Geometric, Algebraic and Topological Methods for Quantum Field Theory*. Mathematical Physics Studies. 2017, pp. 293–324. DOI: 10.1007/978-3-319-65427-0_9. arXiv: 1512.00588 [math.GT].
- [CMW19a] A. S. Cattaneo, P. Mnev, and K. Wernli. “Theta Invariants of lens spaces via the BV-BFV formalism” (2019). arXiv: 1810.06663 [math.AT].
- [CMW19b] A. S. Cattaneo, N. Moshayedi, and K. Wernli. “Globalization for Perturbative Quantization of Nonlinear Split AKSZ Sigma Models on Manifolds with Boundary”. *Communications in Mathematical Physics* 372.1 (2019), pp. 213–260. URL: <https://doi.org/10.1007/s00220-019-03591-5>.
- [CS16] A. S. Cattaneo and M. Schiavina. “BV-BFV approach to General Relativity: Einstein–Hilbert action”. *Journal of Mathematical Physics* 57.2 (2016), p. 023515. DOI: 10.1063/1.4941410.
- [CS17] A. S. Cattaneo and M. Schiavina. “On Time”. *Letters in Mathematical Physics* 107 (Feb. 2017), pp. 375–408. DOI: 10.1007/s11005-016-0907-x.
- [CS19b] A. S. Cattaneo and M. Schiavina. “BV-BFV approach to General Relativity: Palatini–Cartan–Holst action”. *Advances in Theoretical and Mathematical Physics* 23 (8 Dec. 2019), pp. 2025–2059. DOI: 10.4310/ATMP.2019.v23.n8.a3.
- [CS19c] A. S. Cattaneo and M. Schiavina. “The Reduced Phase Space of Palatini–Cartan–Holst Theory”. *Annales Henri Poincaré* 20.2 (2019), pp. 445–480. DOI: 10.1007/s00023-018-0733-z.
- [DeW67] B. S. DeWitt. “Quantum Theory of Gravity. I. The Canonical Theory”. *Phys. Rev.* 160 (5 Aug. 1967), pp. 1113–1148. DOI: 10.1103/PhysRev.160.1113. URL: <https://link.aps.org/doi/10.1103/PhysRev.160.1113>.
- [DGH90] A. Dresse, P. Gregoire, and M. Henneaux. “Path-integral equivalence between the extended and non-extended hamiltonian formalisms”. *Physics Letters B* 245.2 (1990), pp. 192–196.
- [GD00] M. Grigoriev and P. Damgaard. “Superfield BRST charge and the master action”. *Physics Letters B* 474.3 (2000), pp. 323–330.
- [Gri11] M. Grigoriev. “Parent formulation at the Lagrangian level”. *Journal of High Energy Physics* 2011.7 (2011), p. 61. DOI: 10.1007/JHEP07(2011)061.
- [Gri16] M. Grigoriev. *Presymplectic structures and intrinsic Lagrangians*. 2016. arXiv: 1606.07532 [hep-th].
- [HT92] M. Henneaux and C. Teitelboim. *Quantization of gauge systems*. 1992. ISBN: 978-0-691-03769-1.
- [Hen90] M. Henneaux. “Elimination of the auxiliary fields in the antifield formalism”. *Physics Letters B* 238.2 (1990), pp. 299–304.
- [IM19] R. Iraso and P. Mnev. “Two-Dimensional Yang–Mills Theory on Surfaces with Corners in Batalin–Vilkovisky Formalism”. *Communications in Mathematical Physics* 370.2 (2019), pp. 637–702. URL: <https://doi.org/10.1007/s00220-019-03687-3>.
- [Mne17] P. Mnev. “Lectures on Batalin–Vilkovisky formalism and its applications in topological quantum field theory” (July 2017). arXiv: 1707.08096 [math-ph].
- [MSW19] P. Mnev, M. Schiavina, and K. Wernli. “Towards holography in the BV-BFV setting”. *Annales Henri Poincaré* 21.3 (2019), pp. 993–1044. DOI: 10.1007/s00023-019-00862-8.
- [Sch08] F. Schätz. “BFV-Complex and Higher Homotopy Structures”. *Communications in Mathematical Physics* 286.2 (Dec. 2008), p. 399. ISSN: 1432-0916.

- [Sch09] F. Schätz. “Coisotropic Submanifolds and the BFV-Complex”. PhD thesis. Zurich, 2009. URL: <http://user.math.uzh.ch/cattaneo/schaetz.pdf>.
- [Sch15] M. Schiavina. “BV-BFV Approach to general relativity”. PhD thesis. Zurich, 2015. URL: <http://user.math.uzh.ch/cattaneo/schiavina.pdf>.
- [Sta97] J. Stasheff. “Homological reduction of constrained Poisson algebras”. *J. Differential Geom.* 45.1 (1997), pp. 221–240. DOI: 10.4310/jdg/1214459757.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, 8057 ZÜRICH,
SWITZERLAND
Email address: giovanni.canepa@math.uzh.ch

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, 8057 ZÜRICH,
SWITZERLAND
Email address: cattaneo@math.uzh.ch

INSTITUTE FOR THEORETICAL PHYSICS, ETH ZÜRICH, WOLFGANG PAULI STRASSE 27, 8092,
ZÜRICH, SWITZERLAND

DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, RÄMISTRASSE 101, 8092, ZÜRICH, SWITZERLAND
Email address: micschia@phys.ethz.ch