



*JGR Biogeosciences*

Supporting Information for:

## Impacts of Drying and Rewetting on the Radiocarbon Signature of Respired CO<sub>2</sub> and Implications for Incubating Archived Soils

Jeffrey Beem-Miller<sup>1</sup>    Marion Schruppf<sup>1</sup>    Alison Hoyt<sup>1,2</sup>    Georg Guggenberger<sup>3</sup>  
Susan Trumbore<sup>1,4</sup>

<sup>1</sup>Max Planck Institute for Biogeochemistry, Jena, Germany

<sup>2</sup>Lawrence Berkeley National Laboratory, Berkeley, CA, USA

<sup>3</sup>Institute of Soil Science, Leibniz Universität Hannover, Hannover, Germany

<sup>4</sup>Department of Earth System Sciences, University of California, Irvine, CA, USA

*June 15, 2021*

## Contents

Supplemental methods . . . . .	2
Text S1: Linear mixed models . . . . .	2
Linear Mixed Model (LMM) results . . . . .	3
Table S1: LMM marginal means for enclosure period $\Delta^{14}\text{C-CO}_2$ . . . . .	3
Table S2: LMM marginal means for control and treatment $\Delta^{14}\text{C-CO}_2$ (2 <sup>nd</sup> enclosure period only) . . . . .	3
Table S3: Summary of storage duration LMM with Oak Ridge samples . . . . .	4
Table S4: Summary of storage duration LMM without Oak Ridge samples . . . . .	4
Text S2: Comparing differences in $\Delta^{14}\text{C-CO}_2$ across control and treatment samples for the Hainich-Dün time series . . . . .	5
Figure S1: 95% confidence intervals for LMM contrasts of Hainich-Dün forest time series data . . . . .	6
Figure S2: 95% confidence intervals for LMM contrasts of Hainich-Dün forest time series . . . . .	7
Supplemental respiration rate results . . . . .	8
Figure S3: Respiration rates for Experiment 1 and Experiment 2 (rewetting pulse respiration rates shown as a cumulative average for all samples) . . . . .	8
Figure S4: Respiration rates for Experiment 3 . . . . .	9
Treatment effects on $\delta^{13}\text{C-CO}_2$ . . . . .	10
Fig. S5 $\delta^{13}\text{C-CO}_2$ of rewetting pulse and 2 <sup>nd</sup> enclosure period . . . . .	10
Fig. S6 Time series of control and treatment $\delta^{13}\text{C-CO}_2$ (Experiments 1 and 2) . . . . .	11
Additional factors influencing treatment effects on $\Delta^{14}\text{C-CO}_2$ . . . . .	12

Fig. S7 Change in $\Delta^{14}\text{C-CO}_2$ in relation to cumulative soil carbon respired . . . . .	12
Text S3: Change in second enclosure period $\Delta^{14}\text{C-CO}_2$ as a function of field-moisture content	13
Fig. S8 Change in $\Delta^{14}\text{C-CO}_2$ relative to the change in moisture content (control - treatment)	14
Site data, soil properties, and supporting references for all samples (Experiments 1, 2, and 3) . . .	15
Table S5 . . . . .	15
References . . . . .	16

## Supplemental methods

### Text S1: Linear mixed models

We compared the results of the paired mean difference analysis approach discussed in the main text with a linear mixed modeling framework. For the first set of models we set  $\Delta^{14}\text{C-CO}_2$  observed in the second enclosure period as the response variable, and used sample ID as a random intercept term to account for the imbalance in the number of laboratory replicates analyzed for control versus treatment incubations. For fixed effects, we assessed the interaction of ecosystem type with treatment, as well as the three-way interaction of treatment, ecosystem type, and experiment for the combine dataset of Experiment 1 and Experiment 2 samples (Eq. S1). We evaluated the significance of the treatment effect by looking at the contrasts between control and treatment samples across experiments but within ecosystem types. These models were also run for  $\delta^{13}\text{C-CO}_2$ .

We also used the linear mixed modeling framework to assess changes in  $\Delta^{14}\text{C-CO}_2$  and  $\delta^{13}\text{C-CO}_2$  between enclosure periods. For these models we extended our initial model by adding enclosure period as an additional dependent variable. These models were restricted to the experiments and treatments where we measured the response variable in both enclosure periods (Experiment 1 treatment samples, and all Experiment 2 samples). We looked at the overall significance of the parameter estimates as well as the contrasts from this model by each experiment, treatment, and ecosystem type.

We tested the effect of storage duration on observed  $\Delta^{14}\text{C-CO}_2$  using a combined dataset of Experiments 1 and Experiment 3 samples. We used  $\Delta^{14}\text{C-CO}_2$  observed in the second enclosure period for all samples except the Experiment 3 treatment samples for which only a single enclosure period was observed. We constructed a linear mixed model with storage duration, treatment, and the interaction of these two variables as fixed effects. As with the previous models we allowed for a random intercept term for each sample. We did not include ecosystem type in this model as all of the grassland samples were collected at the same point in time. We also excluded the effect of experiment, since this could lead to a spurious relationship due to the change in  $\Delta^{14}\text{C}$  of atmospheric  $\text{CO}_2$  over time and the fact that samples were collected and analyzed at different times. This model was run first with and then without the Oak Ridge samples, as we considered these samples to be a separate population as they contain  $^{14}\text{C}$  from a labelling experiment in addition to atmospheric  $^{14}\text{C}$ .

All statistical analyses were performed in R (R Core Team 2019). We used the package lme4 (Bates et al. 2015) to perform the mixed modeling, and for contrast analysis we used the package emmeans (Lenth 2021). When performing statistical tests we employed Tukey’s honestly significant difference test to account for multiple comparisons and the Kenward-Roger method for estimating degrees of freedom, which has shown to perform well for small sample sizes (Kenward and Roger 1997).

## Linear Mixed Model (LMM) results

**Table S1: LMM marginal means for enclosure period  $\Delta^{14}\text{C-CO}_2$**

Enclosure period $^{14}\text{C}$								
Mixed model means and 95% CIs								
Period	Treatment	Experiment	Type	mean	SE	df	lower.CL	upper.CL
2nd	control	1	forest	93.8	7.8	23.0	77.7	109.9
2nd	treatment	1	forest	82.2	6.9	14.7	67.5	97.0
1st	treatment	1	forest	91.4	8.1	25.3	74.7	108.1
2nd	control	2	forest	44.0	8.2	25.7	27.2	60.9
1st	control	2	forest	20.2	8.2	25.7	3.4	37.1
2nd	treatment	2	forest	56.7	8.2	25.7	39.9	73.6
1st	treatment	2	forest	55.3	8.2	25.7	38.5	72.1
2nd	control	1	grassland	54.5	7.8	23.0	38.4	70.6
2nd	treatment	1	grassland	77.8	6.9	14.7	63.0	92.5
1st	treatment	1	grassland	75.0	7.0	15.5	60.2	89.9
2nd	control	2	grassland	20.8	8.1	25.4	4.1	37.5
1st	control	2	grassland	10.4	8.1	25.4	-6.3	27.0
2nd	treatment	2	grassland	40.3	8.1	25.4	23.7	57.0
1st	treatment	2	grassland	39.6	8.1	25.4	22.9	56.3

**Table S2: LMM marginal means for control and treatment  $\Delta^{14}\text{C-CO}_2$  (2<sup>nd</sup> enclosure period only)**

14C of control and treatment samples (Experiments 1 & 2)							
Mixed model means and 95% CIs							
Treatment	Experiment	Type	mean	SE	df	lower.CL	upper.CL
control	1	forest	93.8	6.0	18.4	81.3	106.3
treatment	1	forest	82.2	5.4	12.6	70.5	93.9
control	2	forest	43.0	6.3	20.8	29.9	56.1
treatment	2	forest	55.7	6.3	20.8	42.6	68.8
control	1	grassland	54.5	6.0	18.4	42.0	67.0
treatment	1	grassland	77.8	5.4	12.6	66.1	89.5
control	2	grassland	21.8	6.3	20.8	8.7	34.9
treatment	2	grassland	41.4	6.3	20.8	28.3	54.5

Table S3: Summary of storage duration LMM with Oak Ridge samples

```
## $emtrends
## treat.bi dur.trend SE df lower.CL upper.CL
## control      12.18 4.46 44.7 3.205 21.2
## treatment     8.46 4.48 45.8 -0.569 17.5
##
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95
##
## $contrasts
## contrast          estimate SE df t.ratio p.value
## control - treatment    3.73 1.31 61.4 2.855 0.0059
##
## Degrees-of-freedom method: kenward-roger
```

Table S4: Summary of storage duration LMM without Oak Ridge samples

```
## $emtrends
## treat.bi dur.trend SE df lower.CL upper.CL
## control      8.59 4.63 36.9 -0.786 18.0
## treatment     5.61 4.54 34.4 -3.605 14.8
##
## Degrees-of-freedom method: kenward-roger
## Confidence level used: 0.95
##
## $contrasts
## contrast          estimate SE df t.ratio p.value
## control - treatment    2.98 3.9 44.5 0.765 0.4485
##
## Degrees-of-freedom method: kenward-roger
```


## Text S2: Comparing differences in $\Delta^{14}\text{C-CO}_2$ across control and treatment samples for the Hainich-Dün time series

Contrast analysis of the LMM output shows that control-treatment differences are not significant for the forest samples, but are significant for the grassland samples in both Experiment 1 and Experiment 2 (first two columns of **Fig. S1**). While the results are similar to the paired difference approach used in the main text, the paired difference approach found both forest and grassland differences to be significant.

Comparing the differences observed over time when samples were treated the same at both timepoints (columns “ /t Ctl” and “ /t Trt”, **Fig. S1**), we see significant differences for both the samples that were never air-dried (“ /t Ctl”) and the samples that were air-dried and rewet (“ /t Trt”). Both differences are positive, i.e.  $\Delta^{14}\text{C-CO}_2$  declined for both control and treatment samples over the period 2011 to 2019. However, we see that the difference over time appear smaller for the forest samples when comparing the difference between the treatment samples (26‰) to the difference between the control samples (49‰), although the confidence intervals overlap substantially. We believe this provides support for reliability of the archived technique when looking at changes in  $\Delta^{14}\text{C-CO}_2$  over time across samples that have been air-dried and rewetted.


Finally, when comparing treatment samples that have never been air-dried (final two columns of figure), the estimated differences are skewed higher or are no longer significant. Specifically, we fail to detect a significant change in the grassland  $\Delta^{14}\text{C-CO}_2$  over time when comparing control samples from 2011 to air-dried and rewet samples from 2019 (penultimate column, “ /t Ex1 Ctl - Ex2 Trt”). Looked at the other way, i.e. treatment  $\Delta^{14}\text{C-CO}_2$  from the 2011 grassland samples compared to control grassland samples in 2019, the difference is substantially exaggerated (last column): 58‰ vs. 27‰ (ctl-ctl) or 38‰ (trt-trt). The difference is also greater for forest samples for both of these cross-treatment comparisons. These differences imply it is important to treat the soils from all time points the same in regards to air-drying and rewetting when constructing a time series using  $\Delta^{14}\text{C-CO}_2$  measured on archived soils in order to minimize bias.

Figure S1: 95% confidence intervals for LMM contrasts of Hainich-Dün forest time series data



*Caption:* Contrasts shown compare treatment and control samples within Experiments 1 and 2 (first two columns, respectively), control samples between 2011 and 2019 (third column), treatment samples between 2011 and 2019 (fourth column), control samples from Experiment 1 (2011) to treatment samples from Experiment 2 (2019) (fifth column), and treatment samples from Experiment 1 (2011) to control samples from Experiment 2 (2019) (sixth column).


Figure S2: 95% confidence intervals for LMM contrasts of Hainich-Dün forest time series



*Caption:* Alternate version of Fig. 5 in main text. Instead of pooled standard deviations, the error bars here show the 95% confidence intervals estimated from the linear mixed model.

## Supplemental respiration rate results


Figure S3: Respiration rates for Experiment 1 and Experiment 2 (rewetting pulse respiration rates shown as a cumulative average for all samples)



*Caption:* CO<sub>2</sub> concentrations for Experiment 1 control samples were only measured once during the pre-incubation period, in contrast to daily measurements for all other samples. Pre-incubation respiration rates are shown here calculated as cumulative averages for the whole pre-incubation period for ease of comparison across all treatments in both Experiment 1 and Experiment 2.




Figure S4: Respiration rates for Experiment 3



*Caption:* Experiment 3 storage duration treatment samples were only incubated for a single enclosure period, as the results of Experiment 1 and Experiment 2 showed no significant difference in  $\Delta^{14}\text{C-CO}_2$  between the rewetting pulse  $\text{CO}_2$  released during the pre-incubation period and the  $\text{CO}_2$  respired during the second enclosure period. The grassland storage duration treatment samples (blue dotted line) respired an equivalent amount of  $\text{CO}_2$  in just 3 d as the corresponding control-3 samples respired during the rewetting pulse period and the second enclosure period combined. Consequently those incubations were stopped after the first  $\text{CO}_2$  measurement point.


## Treatment effects on $\delta^{13}\text{C-CO}_2$

Fig. S5  $\delta^{13}\text{C-CO}_2$  of rewetting pulse and 2<sup>nd</sup> enclosure period



*Caption:* Points are means; error bars show the minimum and maximum of laboratory duplicates.


Fig. S6 Time series of control and treatment  $\delta^{13}\text{C}\text{-CO}_2$  (Experiments 1 and 2)



*Caption:* Points are means; error bars show pooled standard deviations.

## Additional factors influencing treatment effects on $\Delta^{14}\text{C-CO}_2$

Fig. S7 Change in  $\Delta^{14}\text{C-CO}_2$  in relation to cumulative soil carbon respired



*Caption:* Error bars show minimum and maximum values measured for laboratory duplicates, while points show the mean. Lines connect mean pre-incubation and second enclosure period observations for a single sample. Lines parallel to the x-axis indicate a lack of trend in  $\Delta^{14}\text{C-CO}_2$  with the amount of carbon respired, while differences between open and filled symbols show the impact of treatments on both the amount of carbon respired and  $\Delta^{14}\text{C-CO}_2$ . Note that pre-incubation  $\Delta^{14}\text{C-CO}_2$  was not measured for the control-1 samples in 2011. Plot limits exclude outlier point (HEW22 control-2, pre-incubation) for improved legibility.


### Text S3: Change in second enclosure period $\Delta^{14}\text{C-CO}_2$ as a function of field-moisture content

Differences in field moisture content of samples could be related to the magnitude of the shift in  $\Delta^{14}\text{C-CO}_2$  observed between control and treatment sample, as control sample field moisture content varied. All treatment samples were air-dried in the laboratory prior to rewetting: a change in moisture content of zero percent water holding capacity (%WHC) to 60 %WHC. In contrast, moisture adjustment of control samples was made from field moisture, thus, for example, control samples with lower field moisture contents received a correspondingly greater water addition than wetter control samples.

In order to control for the variance in field moisture content of control samples, we looked at the relationship of the difference in the second enclosure period  $\Delta^{14}\text{C-CO}_2$  observed between control and treatment samples and the change in moisture content of the control samples. If the shift in  $\Delta^{14}\text{C-CO}_2$  observed in response to the air-drying and rewetting treatment were a linear function of the change in moisture content, the differences between control and treatment  $\Delta^{14}\text{C-CO}_2$  should be smaller for samples with lower field moisture. However, we did not observe any consistent relationship between the difference in  $\Delta^{14}\text{C-CO}_2$  and field moisture (**Fig. S8**).

We observed the strongest trend in the Experiment 2 grassland samples, but the trend was opposite to what we expected: differences in  $\Delta^{14}\text{C-CO}_2$  between treatment samples and control samples were greater for drier samples than wetter samples (**Fig. S8**). Experiment 2 forest samples showed the expected trend, but it did not appear to be linear (**Fig. S8**). Given the relatively low sample number when considered within treatment and ecosystem groups (Experiment 1  $n = 6$ , Experiment 2  $n = 3$ ), we do not consider these trends to be significant, but the data from Experiment 2 suggest that the relationship between the change in  $\Delta^{14}\text{C-CO}_2$  and the magnitude of rewetting warrents further study.

Fig. S8 Change in  $\Delta^{14}\text{C-CO}_2$  relative to the change in moisture content (control - treatment)



*Caption:* Differences in  $\Delta^{14}\text{C-CO}_2$  are shown as means; error bars show pooled standard deviations. All samples were rewetted to 60% of water holding capacity (WHC) prior to incubation, but control samples were rewetted from field moisture whereas treatment samples were rewetted after air-drying. Data from Experiment 3 are not shown as field moisture content was unknown for the majority of samples (Table S5).

Site data, soil properties, and supporting references for all samples (Experiments 1, 2, and 3)

Table S5

Experiment	Collection date	Treatment incubation date	Control incubation laboratory	Treatment incubation laboratory	Control AMS facility	Treatment AMS facility	Latitude	Longitude	Region	Site	Ecosystem	ID	Incubation replicates	Soil order	Depth						article size distributio						
															Sieved	Top	Bottom	Field moisture		Incubation moisture		Organic C	Total N	Sand	Silt	Clay	Reference
															< 2mm	cm	gravimetric	gravimetric	% water holding capacity	g kg <sup>-1</sup>	g kg <sup>-1</sup>						
															WRB name												
1	2011	2018	MPI-BGC	MPI-BGC	P	P	53.09	13.63	Central Germany	Schorfheide-Chorin	forest	SEW11	2	Cambisol	Yes	0	10	0.26	0.26	60	31.3	1.3	884	85	31	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	52.90	13.85	Central Germany	Schorfheide-Chorin	forest	SEW34	2	Albeluvisol	Yes	0	10	0.24	0.24	60	16.4	0.7	889	69	42	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	52.90	13.93	Central Germany	Schorfheide-Chorin	forest	SEW43	2	Cambisol	Yes	0	10	0.30	0.30	60	18.4	1.1	810	121	69	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	53.12	13.68	Central Germany	Schorfheide-Chorin	grassland	SEG38	2	Cambisol	Yes	0	10	0.25	0.27	60	22.8	2.2	838	72	89	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	53.12	13.84	Central Germany	Schorfheide-Chorin	grassland	SEG40	2	Luvisol	Yes	0	10	0.26	0.27	60	21.3	2.0	710	192	98	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	52.98	13.83	Central Germany	Schorfheide-Chorin	grassland	SEG46	2	Cambisol	Yes	0	10	0.31	0.34	60	24.3	2.3	644	210	146	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	51.34	10.36	Central Germany	Hainich-Dün	forest	HEW22	2	Luvisol	Yes	0	10	0.38	0.37	60	23.3	1.7	68	747	184	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	51.11	10.45	Central Germany	Hainich-Dün	forest	HEW41	2	Luvisol	Yes	0	10	0.40	0.42	60	23.4	1.9	34	754	210	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	51.10	10.46	Central Germany	Hainich-Dün	forest	HEW42	2	Stagnosol	Yes	0	10	0.34	0.36	60	24.3	1.7	60	760	184	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	51.28	10.45	Central Germany	Hainich-Dün	grassland	HEG10	2	Vertisol	Yes	0	10	0.47	0.61	60	43.7	4.0	30	532	436	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	51.08	10.57	Central Germany	Hainich-Dün	grassland	HEG32	2	Cambisol	Yes	0	10	0.52	0.54	60	40.0	3.8	17	640	340	Solly et al. 20	
1	2011	2018	MPI-BGC	MPI-BGC	P	P	51.29	10.38	Central Germany	Hainich-Dün	grassland	HEG48	2	Cambisol	Yes	0	10	0.55	0.56	60	41.6	4.0	50	488	465	Solly et al. 20	
2	2019	2019	MPI-BGC	MPI-BGC	P	P	51.34	10.36	Central Germany	Hainich-Dün	forest	HEW22	2	Luvisol	Yes	0	10	0.38	0.37	60	23.3	1.7	68	747	184	Solly et al. 20	
2	2019	2019	MPI-BGC	MPI-BGC	P	P	51.11	10.45	Central Germany	Hainich-Dün	forest	HEW41	2	Luvisol	Yes	0	10	0.40	0.42	60	23.4	1.9	34	754	210	Solly et al. 20	
2	2019	2019	MPI-BGC	MPI-BGC	P	P	51.10	10.46	Central Germany	Hainich-Dün	forest	HEW42	2	Stagnosol	Yes	0	10	0.34	0.36	60	24.3	1.7	60	760	184	Solly et al. 20	
2	2019	2019	MPI-BGC	MPI-BGC	P	P	51.28	10.45	Central Germany	Hainich-Dün	grassland	HEG10	2	Vertisol	Yes	0	10	0.23	0.22	60	43.7	4.0	30	532	436	Solly et al. 20	
2	2019	2019	MPI-BGC	MPI-BGC	P	P	51.08	10.57	Central Germany	Hainich-Dün	grassland	HEG32	2	Cambisol	Yes	0	10	0.17	0.23	60	40.0	3.8	17	640	340	Solly et al. 20	
2	2019	2019	MPI-BGC	MPI-BGC	P	P	51.29	10.38	Central Germany	Hainich-Dün	grassland	HEG48	2	Cambisol	Yes	0	10	0.19	0.22	60	41.6	4.0	50	488	465	Solly et al. 20	
3	2011	2018	MPI-BGC	MPI-BGC	P	P	51.34	10.51	Central Germany	Hainich-Dün	forest	HEW26	2	Luvisol	Yes	0	10	0.34	0.36	60	24.4	1.6	54	796	150	Solly et al. 20	
3	2011	2018	MPI-BGC	MPI-BGC	P	P	51.18	10.38	Central Germany	Hainich-Dün	forest	HEW47	2	Stagnosol	Yes	0	10	0.43	0.45	60	32.5	2.4	46	632	323	Solly et al. 20	
3	2011	2018	MPI-BGC	MPI-BGC	P	P	51.22	10.37	Central Germany	Hainich-Dün	grassland	HEG20	3	Stagnosol	Yes	0	10	0.47	0.45	60	27.2	2.3	102	661	239	Solly et al. 20	
3	2011	2018	MPI-BGC	MPI-BGC	P	P	51.11	10.43	Central Germany	Hainich-Dün	grassland	HEG33	3	Cambisol	Yes	0	10	0.47	0.47	60	40.1	3.8	29	618	353	Solly et al. 20	
3	2011	2018	MPI-BGC	MPI-BGC	P	P	51.21	10.39	Central Germany	Hainich-Dün	grassland	HEG6	3	Stagnosol	Yes	0	10	0.41	0.45	60	20.8	2.0	45	698	257	Solly et al. 20	
3	2008	2018	UCI	UCI	UCI	UCI	35.98	-79.09	Duke FACE	Duke FACE control	forest	120	1	Ultic Alfisol	Yes	5	15	0.95	0.95		16.6	0.8				Hopkins et al	
3	1999	2009	UCI	UCI	UCI	UCI	42.54	-72.18	Harvard Forest	Harvard Forest	forest	NWN-1 Ap (bag)	1	Inceptisol	Yes	0	16				60.0					Gaudinski et.	
3	1999	2009	UCI	UCI	UCI	UCI	42.54	-72.18	Harvard Forest	Harvard Forest	forest	NWN-1 Ap #27	1	Inceptisol	Yes	0	16				60.0					Gaudinski et.	
3	1999	2009	UCI	UCI	UCI	UCI	42.54	-72.18	Harvard Forest	Harvard Forest	forest	NWN-2 Ap #34	1	Inceptisol	Yes	0	16				60.0					Gaudinski et.	
3	1999	2009	UCI	UCI	UCI	UCI	42.54	-72.18	Harvard Forest	Harvard Forest	forest	NWN-1 Ap #44	1	Inceptisol	Yes	0	16				60.0					Gaudinski et.	
3	2004	2018	UCI	UCI	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA 6E C	1	Inceptisol	No	0	5	0.28	0.28		24.9	1.2				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA 2B C	1	Inceptisol	No	0	5	0.30	0.30		24.9	1.2				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA 3B C	1	Inceptisol	No	0	5	0.49	0.49		24.9	1.2				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA 5B C	1	Inceptisol	No	0	5	0.26	0.26		24.9	1.2				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB 4B C	1	Ultisol	No	0	5	0.34	0.34		24.9	1.0				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB 5B C	1	Ultisol	No	0	5	0.25	0.25		24.9	1.0				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB 8B C	1	Ultisol	No	0	5	0.34	0.34		24.9	1.0				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB 3E C	1	Ultisol	No	0	5	0.36	0.36		24.9	1.0				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB 7E C	1	Ultisol	No	0	5	0.18	0.18		24.9	1.0				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA 4E	1	Inceptisol	No	0	5	0.26			24.9	1.2				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA 6E	1	Inceptisol	No	0	5	0.30			24.9	1.2				Cisneros-Doz	
3	2004	2018	UCI	UCI	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA 8E	1	Inceptisol	No	0	5	0.22			24.9	1.2				Cisneros-Doz	
3	2004	2009	UCI	USGS Menlo Park	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA2B-C_IT2	1	Inceptisol	No	0	5	0.30			24.9	1.2				Cisneros-Doz	
3	2004	2009	UCI	USGS Menlo Park	UCI	UCI	35.94	-84.33	Oak Ridge	TVA	forest	TVA3-C_IT1	1	Inceptisol	No	0	5	0.49			24.9	1.2				Cisneros-Doz	
3	2004	2009	UCI	USGS Menlo Park	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB4B-C_IT2	1	Ultisol	No	0	5	0.34			24.9	1.0				Cisneros-Doz	
3	2004	2009	UCI	USGS Menlo Park	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB5-C_IT2	1	Ultisol	No	0	5	0.25			24.9	1.0				Cisneros-Doz	
3	2004	2009	UCI	USGS Menlo Park	UCI	UCI	35.97	-84.27	Oak Ridge	Walker Branch	forest	WB8-C_IT2	1	Ultisol	No	0	5	0.34			24.9	1.0				Cisneros-Doz	
3	2009	2018	UCI	UCI	UCI	UCI	37.03	-119.27	Sierra Nevada	Musick	forest	MA	3	Ultic Haploxeralf	Yes	5	20	0.07	0.33	50	27.4	1.0	600	270	150	Koarashi et a	
3	2009	2018	UCI	UCI	UCI	UCI	37.03	-119.19	Sierra Nevada	Shaver	forest	SA	3	Pachic Xerumbrept	Yes	5	20	0.07	0.31	50	29.4	1.2	800	150	50	Koarashi et a	



## References

Bates, Douglas, Martin Mächler, Ben Bolker, and Steve Walker. 2015. “Fitting Linear Mixed-Effects Models Using lme4.” *Journal of Statistical Software* 67 (1): 1–48. <https://doi.org/10.18637/jss.v067.i01>.

Kenward, Michael G., and James H. Roger. 1997. “Small Sample Inference for Fixed Effects from Restricted Maximum Likelihood.” *Biometrics* 53 (3). [Wiley, International Biometric Society]: 983–97. <http://www.jstor.org/stable/2533558>.

Lenth, Russell V. 2021. *Emmeans: Estimated Marginal Means, Aka Least-Squares Means*. <https://CRAN.R-project.org/package=emmeans>.