

Farm use of calcium hydroxide as an effective barrier against pathogens

著者	MATSUZAKI Shinji, AZUMA Kento, LIN Xuguang, KURAGANO Masahiro, UWAI Koji, YAMANAKA Shinya, TOKURAKU Kiyotaka
journal or publication title	SCIENTIFIC REPORTS
volume	11
number	1
year	2021
URL	http://hdl.handle.net/10258/00010407

doi: [info:doi/10.1038/s41598-021-86796-w](https://doi.org/10.1038/s41598-021-86796-w)



OPEN

Farm use of calcium hydroxide as an effective barrier against pathogens

Shinji Matsuzaki, Kento Azuma, Xuguang Lin, Masahiro Kuragano, Koji Uwai, Shinya Yamanaka & Kiyotaka Tokuraku

Livestock farming is affected by the occurrence of infectious diseases, but outbreaks can be prevented by proper sanitary control measures. Calcium hydroxide ($\text{Ca}(\text{OH})_2$), commonly called slaked lime, powder is traditionally used as a disinfectant to prevent infectious diseases in livestock. Since $\text{Ca}(\text{OH})_2$ can inactivate a wide variety of pathogens, has a small environmental impact, does not require a disinfection tank (i.e., can be spread directly on the ground) and is produced inexpensively worldwide, it is used for the prevention of epidemics on farms worldwide. Water is essential for the strong alkalinity that underlies its disinfecting effect, but it is unknown how much water is required under field conditions. In addition, $\text{Ca}(\text{OH})_2$ reacts with carbon dioxide in the environment, reducing its pH, but it is unclear how long its degradation takes under actual field use. Thus, we measured the water adsorption ability of $\text{Ca}(\text{OH})_2$ -based disinfectants and its relation to disinfectant activity, as assessed by colony counts and live/dead staining and observation. We found that 15–20% (w/w) water in $\text{Ca}(\text{OH})_2$ was necessary for disinfection to occur in practice. Moreover, we found that the pH of $\text{Ca}(\text{OH})_2$ decreased within about two weeks to one month under actual use in practical conditions and lost its ability to disinfect. We further showed that granules prepared from $\text{Ca}(\text{OH})_2$ and zeolite maintained high alkalinity more than twice as long as calcium powder. These findings will help to establish a suitable method of applying $\text{Ca}(\text{OH})_2$ to protect farms from infectious diseases.

Livestock infectious diseases are prevalent worldwide (Fig. 1). For example, recently, the global impact of African swine fever (ASF; Fig. 1, red circles) in 2018 was severe, as 6.8 million pigs were killed, and since 900 million livestock pigs were raised that year, 0.75% were killed by ASF alone^{1,2}. The destruction of the environment caused by an increase in livestock is regarded as a critical management issue^{3,4}, so farms need to be protected against livestock infectious diseases to prevent the unnecessary death of livestock. Since many farms exist in regions with unfavorable economic conditions, disease prevention using inexpensive and universal disinfectant systems is needed to increase livestock food supply production.

It is well known that strong alkalinity inactivates various pathogens such as bacteria and viruses. After a foot-and-mouth (FMD) disease outbreak occurred in 2010, with 290,000 livestock killed in Miyazaki, Japan⁵, the Japanese Ministry of Agriculture, Forestry, and Fisheries strongly recommended the use of $\text{Ca}(\text{OH})_2$ to prevent livestock infectious diseases. In this so-called “stand-by disinfection” system, $\text{Ca}(\text{OH})_2$ powder is spread at the entrances to a farm and around livestock houses, or related facilities as a livestock hygiene service center at a density of approximately 0.5–1.0 kg/m² (Supplementary Fig. 1). The strong alkalinity (pH 12.4, 25 °C) of $\text{Ca}(\text{OH})_2$ prevents the invasion of pathogens brought to the premises by cars, people, wild animals, etc., and acts as a barrier against various infectious diseases by inactivating bacteria and viruses. *Escherichia coli*, *Salmonella*, and highly pathogenic avian influenza are inactivated in 24 h, 2 h, and 5–10 min, respectively^{6–8}.

Disinfection by $\text{Ca}(\text{OH})_2$ is related to the release of hydroxyl ions in an aqueous environment^{6,8,9}. However, no studies have examined the disinfectant properties of solid $\text{Ca}(\text{OH})_2$ under stand-by disinfection conditions, that is, when a particulate-bound pathogen is placed into contact with $\text{Ca}(\text{OH})_2$ powder, which occurs during field application. Rather, the standard protocols for evaluating disinfectants, as stipulated by the European Norm (EN), the American Society for Testing and Materials (ASTM), and the Association of Official Analytical Chemists (AOAC), consist of dropping disinfectant solutions on agar or mixing a solution with liquid medium.

$\text{Ca}(\text{OH})_2$ costs 109 US dollars/ton, which is approximately one twentieth of the cost of using a surfactant as a disinfectant (2066 dollars/ton)¹⁰. In addition, $\text{Ca}(\text{OH})_2$ can be spread directly on the ground, i.e. no water tank is

Department of Applied Science, Muroran Institute of Technology, Muroran, Hokkaido, Japan. email: syama@mmm.muroran-it.ac.jp; tokuraku@mmm.muroran-it.ac.jp




Figure 1. Locations of lime production and outbreaks of infectious diseases in domestic animals. Countries with lime production over 1 million tons per year, and locations of outbreaks of foot-and-mouth disease, avian influenza, classical swine fever and African swine fever in 2018–2019 (lime production was quoted from Mineral Commodity Summaries 2019¹¹, and the location of livestock infectious diseases was quoted from the World Animal Health Information System Interface², and we drew the world map summarizing them).

required. Since $\text{Ca}(\text{OH})_2$ does not significantly affect soil microflora¹¹ because it reacts with CO_2 in the environment to form nontoxic calcium carbonate (CaCO_3), it is an environmentally friendly disinfectant. Furthermore, since limestone, which is the raw material for $\text{Ca}(\text{OH})_2$ production, is produced in large quantities worldwide¹² (Fig. 1, open triangles), $\text{Ca}(\text{OH})_2$ can be widely used on farms.

In spite of its many strengths, there are several disadvantages of using $\text{Ca}(\text{OH})_2$ as a disinfectant: (1) $\text{Ca}(\text{OH})_2$ powder is scattered by wind, (2) the amount of water required for stand-by disinfection conditions is unknown, and (3) $\text{Ca}(\text{OH})_2$ and its environmental transformation product CaCO_3 , which lacks disinfection activity, are indistinguishable using common methods. By overcoming these disadvantages, realistic and effective barriers can be built to protect farms around the world against the threat of infectious diseases.

To the best of our knowledge, this report is the first to examine the effect of water content the disinfection ability of $\text{Ca}(\text{OH})_2$. We used a stand-by disinfection strategy that considers those conditions under which $\text{Ca}(\text{OH})_2$ has the highest disinfection activity. The disinfection effects of $\text{Ca}(\text{OH})_2$ (slaked lime) powder and newly developed slaked lime-based granules (SLBGs)¹³ were evaluated. SLBGs were developed to prevent scattering by wind and zeolite was added as an expected deodorant effect and for the long-term persistence of pH. These evaluations were made under conditions mimicking actual stand-by disinfection conditions, while appropriate water content was also examined.

Results and discussion

Preparation of SLBGs. SLBGs, which scatter in the wind less readily than powdered $\text{Ca}(\text{OH})_2$, were made from $\text{Ca}(\text{OH})_2$ powder and natural zeolite powder with a high shear wet granulator without any binder, according to our recent report¹³. In this study, sieved granules of 1.0 to 2.0 mm were used for the tests.

pH changes persist for up to 1 month following $\text{Ca}(\text{OH})_2$ application by stand-by disinfection. To date, there have been no reports examining how long the high pH underlying disinfection effects is maintained under stand-by disinfection conditions. First, we evaluated the persistence of pH changes occurring in $\text{Ca}(\text{OH})_2$ powder applied by stand-by disinfection (Fig. 2a) to soil, grass, a road, and in front of a barn (Supplementary Fig. 2). Farm vehicles and people pass frequently on a road and in front of a barn, but not on soil and grass. In front of a barn, livestock pass, in addition to vehicles and people. In this test (Fig. 2b), the high alkalinity of $\text{Ca}(\text{OH})_2$ -treated surfaces began to decrease after approximately two weeks. After 44 days, their pH had decreased to approximately 9, which is comparable to the pH of CaCO_3 . In front of the barn, the pH dropped to 8.99 after 22 days (Fig. 2b, closed triangles). These results suggest that the disinfection effect resulting from the high alkalinity of $\text{Ca}(\text{OH})_2$ lasted only approximately 2 weeks to 1 month under stand-by disinfection conditions. In another study, the pH and disinfection effect of $\text{Ca}(\text{OH})_2$ powder decreased after alternating wet and dry conditions¹⁴, suggesting that the decrease in pH during the one month was due to environmental conditions. The fastest drop in pH in front of the barn may have been due to $\text{Ca}(\text{OH})_2$ contact with livestock manure, as the CO_2 generated by heterotrophic microorganisms in manure can neutralize hydroxyl ions¹⁵.

In the outdoor test, the results varied depending on the conditions at the test location, for example, rainfall fluctuation, the frequency of passing vehicles, and contact with manure. To determine the persistence of pH changes under constant conditions, we carried out an indoor test with conditions similar to those in stand-by disinfection (Fig. 2c). In the indoor test, the pH of the $\text{Ca}(\text{OH})_2$ powder remained high through day 21, but decreased to 9.63 on day 35 (Fig. 2d, black squares). Since this result was similar to that observed in the outdoor test (Fig. 2b), the indoor test was considered to represent the conditions of the actual outdoor stand-by disinfection. The pH of SLBGs after 76 days was 11.58 (Fig. 2d, white squares), suggesting that the disinfecting effect of SLBGs was maintained for more than twice longer than $\text{Ca}(\text{OH})_2$ powder. These results suggest that the $\text{Ca}(\text{OH})_2$




Figure 2. Evaluation of pH persistence under stand-by disinfection conditions. (a) Typical outdoor test. $\text{Ca}(\text{OH})_2$ powder was spread at 1 kg/m^2 . (b) In the outdoor test, the change in pH after treatment with $\text{Ca}(\text{OH})_2$ powder was evaluated on soil, grass, a road, and in front of a barn. (c) Schematic of the indoor test to measure the pH following disinfectant treatment. To mimic the outdoor conditions, this system consisted of a tray, artificial grass, and a paper box. For the treatments, disinfectants were placed in a paper box. Water ($200 \text{ cm}^3/\text{day}$) was applied, equal the annual precipitation in Hokkaido prefecture, Japan. (d) In the indoor test, the pH after treatment with $\text{Ca}(\text{OH})_2$ powder or SLBGs was evaluated. Data represent the mean from three separate experiments. Error bars show the standard deviation.

powder could maintain high alkalinity for only up to about one month under the stand-by disinfection conditions tested in this study while the SLBGs we developed maintained high alkalinity for at least twice as long. In the process of preparing SLBGs, we used water and dried it at 60°C overnight (see methods section). During this process, $\text{Ca}(\text{OH})_2$ in SLBGs is expected to be partially carbonated. Nevertheless, SLBGs maintained high alkalinity for longer periods of time than $\text{Ca}(\text{OH})_2$ powder (Fig. 2d). This result suggests that granulation with zeolite provides advantages, by preventing carbonation caused during the preparation process. By optimizing the preparation process, it will be possible to SLBGs with higher $\text{Ca}(\text{OH})_2$ content and longer-lasting high alkalinity.

Water is required for the disinfection activity of $\text{Ca}(\text{OH})_2$. To directly evaluate the disinfectant activity associated with the pH changes induced by the $\text{Ca}(\text{OH})_2$ treatments, we investigated in vitro bactericidal effects. In this study, *E. coli*, which is highly resistant to $\text{Ca}(\text{OH})_2$ ^{6–8}, was used as a model pathogen to evaluate the disinfectant activity of the treatment. In previous studies, *E. coli* and *Salmonella* underwent a $6 \log_{10}$ and $4 \log_{10}$ decrease, respectively, after 10 min of exposure to a $\text{Ca}(\text{OH})_2$ solution in an in vitro experiment⁷, suggesting that $\text{Ca}(\text{OH})_2$ is an effective disinfectant. However, in that report⁷, it was unclear whether $\text{Ca}(\text{OH})_2$ exerted its disinfectant effect under stand-by disinfection conditions (Supplementary Fig. 1, Fig. 2) because the evaluation was performed using $\text{Ca}(\text{OH})_2$ solution⁷. Moreover, a separate study reported that after $\text{Ca}(\text{OH})_2$ powder was added to livestock feces, that the powder became inactive against *E. coli* and *Salmonella* within 6 h¹⁶. However, water was also abundant in that system, while the inactivation was due to some other factors such as CO_2 emission by microbes in the feces. Therefore, we developed a new method (Supplementary Fig. 3) to evaluate the disinfection effects of dry $\text{Ca}(\text{OH})_2$ powder, SLBGs, and CaCO_3 powder on *E. coli* under stand-by disinfection conditions. Details of the new method and its verification are explained in the Supplementary methods and Supplementary Figs. 4–8, respectively. At the initiation of contact between the disinfectants and *E. coli*, bacterial abundance was $4\text{--}5 \log_{10}$ colony-forming units (CFU)/mL in all samples (Fig. 3a, 0 h). After 3 h of contact, abundance was similar in all samples, including the control (i.e., without any disinfectants) sample (Fig. 3a, 3 h). These results (Fig. 3a) clearly demonstrate that dry $\text{Ca}(\text{OH})_2$ could not inactivate *E. coli*, indicating that $\text{Ca}(\text{OH})_2$ disinfectant activity requires water.

Next, we evaluated the disinfection effect when water was added under the same conditions (Fig. 3b). The abundance of *E. coli* was approximately $4\text{--}5 \log_{10}$ CFU/mL before incubation (Fig. 3b, 0 s). After exposure to wet CaCO_3 powder for 3 h in the presence of water ($5 \mu\text{L}$ of water per 1.5 mg of CaCO_3), the abundance of *E. coli* was $4.7 \log_{10}$ CFU/mL (Fig. 3b, 10,800 s), suggesting that CaCO_3 did not exert a disinfection effect, even in the presence of water (Fig. 3b, dotted bars). This result revealed that a weakly alkaline ($\sim \text{pH } 9$) CaCO_3 solution had no disinfecting effect on *E. coli*. We also confirmed that $\text{Ca}(\text{OH})_2$, whose pH dropped to 9.28 in the door test after 63 days, had no disinfecting effect (Fig. 3c, diagonal bars). Upon exposure to $\text{Ca}(\text{OH})_2$ powder in the presence of water, the abundance of *E. coli* was reduced to $2.5 \log_{10}$ CFU/mL in 640 s and decreased below the detection limit after 1,200 s of contact (Fig. 3b, diagonal bars). After the addition of SLBGs to *E. coli* in the presence of water, the bacterial abundance decreased to $2.2 \log_{10}$ CFU/mL in 3600 s, to $1.5 \log_{10}$ CFU/mL in 5400 s, and below the detection limit after 7,200 s (Fig. 3b, gray bars). These results (Fig. 3a,b) confirm that disinfection by $\text{Ca}(\text{OH})_2$ requires the presence of water. Furthermore, monitoring the potential environmental degradation




Figure 3. The effect of water on disinfection. **(a)** *E. coli* abundance (\log_{10} colony-forming units, CFU) after contact without (control) or with disinfectant (CaCO_3 , Ca(OH)_2 , or SLBGs) with no water added (dry condition) for 3 h. The dashed line indicates the detection limit estimated from the dilution ratio. **(b)** *E. coli* abundance after contact with disinfectant (CaCO_3 , Ca(OH)_2 , or SLBGs) with water added (wet condition) for up to 3 h. **(c)** *E. coli* abundance after contact without (control) with low pH Ca(OH)_2 (in this Ca(OH)_2 , pH was reduced to about 9.28 in the door test for 63 days in Fig. 2d) with water added (wet condition) for up to 3 h. **(d)** Live/dead assay with disinfectant (control, CaCO_3 , Ca(OH)_2 , or SLBGs) in dry and wet conditions after 3 h. Images were captured using a fluorescence microscope equipped with a 20 \times objective. Bar, 100 μm . Data in **a**, **b**, and **c** represent the mean and SD (error bars) from three separate experiments. Alive and Dead in **a**, **b**, and **c** indicate the range of CFU in which the number of colonies was observed and the range of CFU below the detection limit ($2 \log_{10}$ CFU/mL) in which no colonies were observed, respectively.

of $\text{Ca}(\text{OH})_2$ to CaCO_3 is also important to ensure adequate stand-by disinfection, because CaCO_3 (Fig. 3b) and $\text{Ca}(\text{OH})_2$ with reduced pH (Fig. 3c) did not show disinfection activity under this degraded condition.

Next, we performed a live/dead assay to confirm whether disinfection occurred concomitantly with the changes in pH under the same conditions. In this experiment, live and dead *E. coli* were stained green and red using DMAO and EthD-III solutions in Bacteria Live/Dead Staining Kit, respectively. DMAO is a green fluorescent dye that stains both live and non-membrane-damaged bacteria, and EthD-III is a red fluorescent dye that stains only dead bacteria with membrane damage¹⁷. *E. coli* were stained green in the presence of each disinfectant in the dry condition (Fig. 3d, dry condition), suggesting that *E. coli* survived dry conditions (Fig. 3a). In the wet condition, *E. coli* also stained green in the control and CaCO_3 samples (Fig. 3d, wet condition). In contrast, *E. coli* treated with $\text{Ca}(\text{OH})_2$ powder or SLBGs were stained red (Fig. 3d, wet condition), suggesting that these disinfectants were effective in wet conditions. We then confirmed the presence of *E. coli* in dry conditions by microscopic analysis of all samples (Supplementary Fig. 9), and also confirmed that the abundance of *E. coli* decreased after exposure to $\text{Ca}(\text{OH})_2$ powder or SLBGs in wet conditions (Supplementary Fig. 9). One possible explanation for these results is that the cell wall/membrane was disrupted by high alkalinity in wet conditions in the presence of $\text{Ca}(\text{OH})_2$ powder or SLBGs. $\text{Ca}(\text{OH})_2$ solution can inactivate bacteria by effectively destroying the cell wall/membranes¹⁸, supporting this result.

High levels of humidity mediate $\text{Ca}(\text{OH})_2$ disinfectant activity. Stand-by disinfection is performed under variable environmental conditions. Many water molecules are adsorbed to the surface of $\text{Ca}(\text{OH})_2$ particles under high humidity, which may contribute to the disinfection effect. Two studies reported that $\text{Ca}(\text{OH})_2$ powder exhibited disinfection effects under moist conditions, although the exact moisture content was not reported^{19,20}. To investigate the effect of humidity on disinfection activity under stand-by disinfection conditions, we evaluated the viability of *E. coli* at 20%, 40%, 60%, 80%, and 90% humidity using thermo-hygrostat chamber (Fig. 4a). Soil surface temperature is approximately 5 °C higher than air temperature²¹, so we performed this evaluation at 35 °C, to represent the summer ground temperature in Japan. When *E. coli* was exposed to the disinfectants for 3 h, colonies were detected in all samples at 20%, 40%, 60%, and 80% humidity (Fig. 4a). However, at 90% humidity, colonies were not detected in the $\text{Ca}(\text{OH})_2$ powder (Fig. 4a, diagonal bars) and SLBG (Fig. 4a, gray bars) samples but were detected in the control (Fig. 4a, white bars) and CaCO_3 (Fig. 4a, dotted bars) samples, suggesting that these two disinfectants were effective under high humidity (~90%), without water being present in a liquid phase.

Figure 4b shows the results of a confirmatory live/dead assay. *E. coli* stained green under all conditions in the control, CaCO_3 powder, $\text{Ca}(\text{OH})_2$ powder, and SLBG samples at 20%, 40%, 60% and 80% humidity. However, at 90% humidity, *E. coli* stained green in the control and CaCO_3 powder samples but red in the $\text{Ca}(\text{OH})_2$ powder and SLBG samples, again suggesting that the latter two disinfectants are effective at 90% humidity. Microscopy observation of *E. coli* (Supplementary Fig. 10) suggest that *E. coli* was inactivated without any change in cell morphology. This result implies that contact with the solid $\text{Ca}(\text{OH})_2$ surface holding water partially destroyed the cell wall/membrane of *E. coli*, resulting in cytoplasmic leakage and cell death.

To determine how much water adsorption is required for the observed effect, we also measured the amount of water adsorbed by the samples at 90% humidity (Fig. 4c). After 0.5 h of incubation, adsorbed water accounted for 3.0% of the CaCO_3 mass (Fig. 4c, dotted bars), 9.0% of the $\text{Ca}(\text{OH})_2$ mass (Fig. 4c, diagonal bars) and 15% of the SLBG mass (Fig. 4c, gray bars). After 3 h of incubation, when the disinfection effect was confirmed (Fig. 4c), 6.6%, 15%, and 20% water had adsorbed to the CaCO_3 , $\text{Ca}(\text{OH})_2$, and SLBGs, respectively. Zeolite, one of the raw materials of SLBGs, is an aluminosilicate crystal composed of SiO_4 and AlO_4 in a tertiary structure that is also used for the removal of ammonia and CO_2 ^{22–24}. The adsorption of water to SLBGs may have been higher than to $\text{Ca}(\text{OH})_2$ since zeolite adsorbs water due to its hygroscopicity. Overall, these results (Fig. 4) reveal that 15–20%wt of water content in $\text{Ca}(\text{OH})_2$ or SLBG disinfectant is necessary for disinfection. These results indicate potential best practices regarding the use of $\text{Ca}(\text{OH})_2$ in the field.

On-site detection of the disinfection effect by a pH indicator. Since the persistence of pH as a basis for the disinfection effect (Fig. 3c) varies depending on the situation of stand-by disinfection and on the type of $\text{Ca}(\text{OH})_2$ disinfectant (Fig. 2), it is important to periodically evaluate the disinfection activity using, in this case, the persistence of, or changes to, pH as a proxy. However, periodic pH measurement using a pH meter is time consuming and expensive, so we developed a simple evaluation method using a pH indicator. The detailed preparation method of the pH indicator will be reported elsewhere (Uwai et al., manuscript in preparation), but the results obtained using the pH indicator are shown in Fig. 5 and Supplementary Movie 1. Spray application of the pH indicator stained the SLBGs prepared from $\text{Ca}(\text{OH})_2$ and $\text{Ca}(\text{OH})_2$ powder blue, but the SLBGs prepared from CaCO_3 and CaCO_3 powder stained red (Fig. 5). With this spray, farmers will be able to quickly determine the degradation of $\text{Ca}(\text{OH})_2$ and make necessary adjustments. In addition, if farmers have overspread $\text{Ca}(\text{OH})_2$, this adjustment may save using it.

Conclusion

In this study, we found that the pH of $\text{Ca}(\text{OH})_2$ decreases within about two weeks to one month under stand-by disinfection conditions and loses its disinfecting activity while 15–20% (w/w) water is required for the disinfecting effect. We further confirmed that the pH persistence of SLBGs was more than twice as long as $\text{Ca}(\text{OH})_2$ powder. These results will help to establish an appropriate application of $\text{Ca}(\text{OH})_2$ to protect farms from infectious diseases.




Figure 4. The effect of humidity on disinfection. **(a)** *E. coli* abundance (log₁₀ CFU) after contact with disinfectant (control, CaCO₃, Ca(OH)₂, or SLBGs) under various humidity conditions (20%, 40%, 60%, 80%, or 90%) for 3 h. The detection limit was 2 log₁₀ CFU/mL (dashed line). **(b)** Live/dead assay with the disinfectant (control, CaCO₃, Ca(OH)₂, or SLBGs) under various humidity conditions (20%, 40%, 60%, 80%, or 90%) after 3 h. Images were captured using a fluorescence microscope equipped with a 20× objective. Bar, 100 μm. **(c)** Water content of disinfectant (CaCO₃, Ca(OH)₂, or SLBGs) over time at 35 °C and 90% humidity. Data in **a** and **c** represent the mean and SD (error bars) from three separate experiments.

Methods

Materials. Ca(OH)₂ powder was purchased from Hokkaido Lime Co., Ltd. (Tomakomai, Japan), CaCO₃ powder was purchased from Kanto Chemical Co., Inc. (Tokyo, Japan), and zeolite was purchased from Hokkaido Zeolite Co., Inc. (Otaru, Japan). Ca(OH)₂ powder and SLBGs were subdivided into screw cap tubes or plastic bags and stored in a sealed state.




Figure 5. Detection of carbonation of $\text{Ca}(\text{OH})_2$ using a pH indicator. SLBGs containing $\text{Ca}(\text{OH})_2$ or CaCO_3 (left) or powder consisting of $\text{Ca}(\text{OH})_2$ or CaCO_3 (left) (right) were sprayed with the pH indicator we developed (Uwai et al., manuscript in preparation). Granules and powder containing $\text{Ca}(\text{OH})_2$ instantly turned blue from red due to their strong alkalinity, but granules and powder containing CaCO_3 remained red (Supplementary Movie 1).

Preparation of slaked lime-based granules (SLBGs). SLBGs were prepared using a stirring granulator (WB-5, Pacific Machinery & Engineering, Narashino, Japan). First, 750 g of $\text{Ca}(\text{OH})_2$ powder and 750 g of zeolite were premixed for 60 s in the granulator. After 500 mL of deionized water was added to the mixed powder, granulation was carried out for 300 s, using spindle and chopper rotation speeds of 167 and 3000 rpm, respectively. The granules were dried in an oven at 60 °C overnight. In this study, granules of 1.0 to 2.0 mm were used for the biological tests. The effects of granulation conditions on the physical properties of the granules (namely, granule size distribution and hardness) are described in detail elsewhere¹³.

Evaluation of the persistence of pH changes induced by disinfectants. $\text{Ca}(\text{OH})_2$ powder was used for stand-by disinfection at four outdoor locations (soil, grass, road, and in front of a barn, Supplementary Fig. 1). For stand-by disinfection, 0.5–1 kg/m² $\text{Ca}(\text{OH})_2$ powder was spread, such that 1 kg of $\text{Ca}(\text{OH})_2$ powder for 1 m² was spread at the four outdoor locations (Fig. 2a). To measure pH, 1 g of disinfectant was collected in a 15 mL plastic tube, 2.5 mL of distilled water was added, and the mixture was stirred. The suspension was left for 1 h, and the pH of the supernatant was measured by a pH meter (Docu-pH meter, Sartorius, New York, USA).

For the indoor experiment, a paper box was placed on artificial plastic grass (440–0011, Mizushima, Osaka, Japan), and 62.5 g of disinfectant was spread on the paper box (Fig. 2c). Water was sprayed daily (200 cm³/day) to match the annual precipitation in Hokkaido, Japan²⁵. This amount was determined as follows. Since average rainfall in Hokkaido from 1986 to 2015 was 114.8 cm/year (0.315 cm/day) and the size of the paper box was 25 × 25 cm, the daily watering rate was set to 25 × 25 × 0.32 cm = 200 cm³. Water (200 cm³) was applied once a day using a watering can. pH was measured in the same way as in the outdoor experiment.

Evaluation of the disinfection effect under stand-by disinfection conditions. Our method to evaluate the effectiveness of disinfection (Supplementary Fig. 3) was established according to the conditions described in Supplementary Figs. 6–10. One milliliter of bacterial suspension was dispensed into a 1.5 mL plastic tube and centrifuged (9,100 g, 2 min, 4 °C). Then, the supernatant was removed. The pellet was resuspended in 1 mL of phosphate-buffered saline (PBS) and centrifuged (2,300 g, 2 min, 4 °C), and the supernatant was removed again. The obtained pellets were dried in an incubator (37 °C, 3 h). To increase the contact area with the disinfectant, the dried cells were homogenized for 5 s with a Powermasher II (Nippi, Tokyo, Japan). Disinfectant (1.5 mg) was added to the homogenized dried cells, vortexed for 3 s and statically incubated for various periods. After incubation, the homogenized samples were instantly neutralized by adding 1 mL of 0.1 M phosphate buffer (pH 6.8). The neutralized solution was then diluted tenfold and 100-fold. Then, 0.1 mL of the diluted samples were applied to plastic Petri dish (90 × 20 mm), and the number of colonies was counted after overnight incubation at 37 °C. Finally, *E. coli* log₁₀ number of CFUs was determined.

Escherichia coli strains. Since *E. coli* is the most commonly used gram-negative bacterial model for pathogenic bacteria²⁶, *E. coli* XL-1 blue was used as the model bacterium in this study. *E. coli* XL-1 blue was purchased from GMBiolab Co., Ltd. (Taichung, Taiwan). First, 100 μL of *E. coli* XL-1 blue that had been stored at – 80 °C was thawed and precultured (37 °C, 80 rpm, overnight) in LB medium (10 g/L polypeptone, 5 g/L yeast extract, 5.0 g/L NaCl, pH 7.0). Next, a primary culture (LB medium, 37 °C, 80 rpm, 15 h) was inoculated with 100 μL of the preculture in LB medium. The primary culture after the logarithmic growth phase, in which the number of viable bacteria was maximized, was used for the experiment.

Evaluation of disinfection in the presence of water. For the dry condition, 1.5 mg of disinfectant was added to the homogenized dry cells in a 1.5 mL tube, which was then closed. For the wet condition, 1.5 mg of disinfectant and 5 μL of water were added to the homogenized dry cells, and the tube was closed and statically incubated for various periods. For the controlled-humidity conditions, 1.5 mg of disinfectant was added to the homogenized dry cells, and the tube was placed in a thermo-hygrostat chamber (SH-222, Espec, Osaka, Japan) for 3 h. In this incubator, the 1.5 mL tube was opened. After incubation, the sample was neutralized instantly by adding 1 mL of 0.1 M phosphate buffer. The resulting neutralized solution was diluted tenfold and 100-fold with 0.1 M phosphate buffer, 0.1 mL of the diluted sample was applied to a plate containing LB agar medium, and the number of colonies was counted after incubation. The *E. coli* log₁₀ number of CFUs was then determined.

Fluorescence microscopy. The disinfection effect was confirmed using a Bacteria Live/Dead Staining Kit (Takara Bio Inc., Kusatsu, Japan). After the samples were statically incubated with disinfectant, the disinfectant was neutralized with 1 mL of 0.1 M phosphate buffer. The desired sample dilution was obtained by adding 60 μ L of 0.1 M phosphate buffer to 40 μ L of the neutralized suspension. Next, 1 μ L of staining solution (DMAO:EthD-III:0.85% NaCl = 1:2:8) was added to the diluted solution, which was then incubated in the dark (room temperature, 15 min). Images were captured using a conventional fluorescence microscope (ECLIPSE Ni; Nikon, Tokyo, Japan) equipped with a CMOS camera (DS-Qi2; Nikon) and an objective lens (Plan Fluor ELWD 20 \times /0.45, Nikon) together with NIS-Elements ver. 4.5 software (Nikon). All procedures were performed at room temperature (22 \pm 2 $^{\circ}$ C). Immunofluorescence images were analyzed using NIS-Elements software (Nikon) and Image J ver. 1.52a (NIH, Bethesda, USA).

Water adsorption by the disinfectant. To determine the water content adsorbed by the disinfectant at 35 $^{\circ}$ C and 90% humidity, 1.5 mg of disinfectant was placed in a 1.5 mL tube and incubated in a thermo-hygrostat chamber (SH-222, Espec). Then, the weight of the tube was measured. The water content was determined as the ratio of the weight of water adsorbed on the disinfectant to the weight of the disinfectant that adsorbed the water.

Preparation of the pH indicator. The preparation of the pH indicator for the on-site assessment of Ca(OH)₂ pH indicator is described elsewhere (Uwai et al., manuscript in preparation).

Statistics. Evaluations of the persistence of pH changes, disinfection effects, and water absorption were performed in three independent experiments (triplicate of each experiment). The data were analyzed using Excel software (Microsoft, Redmond, USA). In this paper, we defined CFU below the detection limit (2 log₁₀ CFU/mL) as a significant disinfectant effect.

Received: 19 June 2020; Accepted: 19 March 2021

Published online: 12 April 2021

References

1. Food and Agriculture Organization of the United Nations. Live Animals Statistics. <http://www.fao.org/faostat/en/#data/QA> (2017).
2. World Animal Health Information System Interface. WAHIS Interface country information exceptional epidemiological events. https://www.oie.int/wahis_2/public/wahid.php/Countryinformation/Countryreports (2019).
3. Willett, W. et al. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *Lancet* **393**, 447–492 (2019).
4. Springmann, M. et al. Options for keeping the food system within environmental limits. *Nature* **562**, 519–525 (2018).
5. Muroga, N. et al. The 2010 foot-and-mouth disease epidemic in Japan. *J. Vet. Med. Sci.* **74**, 399–404 (2012).
6. Bennett, D. D. et al. Effects of lime on *Salmonella enteritidis* survival in vitro. *J. Appl. Poult. Res.* **12**, 65–68 (2003).
7. Bean, C. L. et al. Class B alkaline stabilization to achieve pathogen inactivation. *Int. J. Environ. Res. Public Health* **4**, 53–60 (2007).
8. Takeda, S. Prevention and control of highly pathogenic avian influenza by the use of lime. *J. Soc. Inorg. Mater.* **16**, 49–54 (2009).
9. Siqueira, J. F. & Lopes, H. P. Mechanisms of antimicrobial activity of calcium hydroxide: a critical review. *Int. Endod. J.* **32**, 361–369 (1999).
10. Research and Statistics Department Minister's Secretariat Ministry of Economy Trade and Industry. Yearbook of Current Production Statistics Chemical Industry (2018).
11. Mori, M., Sakagami, Y., Hamazaki, Y. & Jojima, T. Evaluation of the influence of sprinkling powdered slaked lime on microorganisms for the prevention of domestic animal infectious diseases. *Environ. Technol.* **40**, 3094–3104 (2018).
12. United States Geological Survey. Mineral Commodity Summaries 2019. https://prd-wret.s3-us-west-2.amazonaws.com/assets/palladium/production/atoms/files/mcs2019_all.pdf. <https://doi.org/10.1007/978-3-540-47108-0-4> (2019).
13. Yamanaka, S., Hirano, S., Uwai, K. & Tokuraku, K. Design of calcium hydroxide-based granules for livestock sanitation. *Case Studies Chem. Environ. Eng.* **2**, 100005 (2020).
14. Ruenphet, S., Paditporn, K., Punyadarsaniya, D., Jantafong, T. & Takehara, K. Bactericidal and virucidal efficacies of food additive grade calcium hydroxide under various concentrations, organic material conditions, exposure duration, and its stability. *Vet. World* **12**, 1383–1389 (2019).
15. Nyberg, K. A., Vinnerås, B., Lewerin, S. S., Kjellberg, E. & Albin, A. Treatment with Ca(OH)₂ for inactivation of *Salmonella Typhimurium* and *Enterococcus faecalis* in soil contaminated with infected horse manure. *J. Appl. Microbiol.* **110**, 1515–1523 (2011).
16. Hakim, H. et al. Accuracy of the evaluation method for alkaline agents' bactericidal efficacies in solid, and the required time of bacterial inactivation. *J. Vet. Med. Sci.* **79**, 244–247 (2017).
17. Adamski, M. & Pietr, S. J. Biodiversity of bacteria associated with eight *Pleurotus ostreatus* (Fr.) P. Kumm. strains from Poland, Japan and the USA. *Polish J. Microbiol.* **68**, 71–81 (2019).
18. National Lime Association. Biosolids and Sludge <https://www.lime.org/lime-basics/uses-of-lime/enviromental/biosolids-and-sludge/>. (2019).
19. Ookubo, Y. & Tojo, H. Evaluation of slaked lime for “trapping” disinfection effect on bacteria and avian influenza virus. *J. Jpn. Soc. Poult. Dis.* **45**, 84–90 (2009).
20. Thammakarn, C. et al. Efficacy of scallop shell powders and slaked lime for inactivating avian influenza virus under harsh conditions. *Arch. Virol.* **160**, 2577–2581 (2015).
21. Kondo, M. & Suzuki, S. A quantitative study on an existence effect of the urban vegetation by measuring the surface temperature. *J. Jpn. Inst. Landsc. Archit.* **47**, 1–15 (1983).
22. Bae, T. H. et al. Evaluation of cation-exchanged zeolite adsorbents for post-combustion carbon dioxide capture. *Energy Environ. Sci.* **6**, 128–138 (2013).
23. Burgess, R. M. et al. Use of zeolite for removing ammonia and ammonia-caused toxicity in marine toxicity identification evaluations. *Arch. Environ. Contam. Toxicol.* **47**, 440–447 (2004).
24. Hiraki, T., Nosaka, A., Okinaka, N. & Akiyama, T. Synthesis of zeolite-X from waste metals. *ISIJ Int.* **49**, 1644–1648 (2009).
25. Japan Ministry of Land, Infrastructure, Transport and Tourism. <http://www.mlit.go.jp/common/001319365.pdf>.

26. Zhou, S., Huang, S., Li, X., Angelidaki, I. & Zhang, Y. Microbial electrolytic disinfection process for highly efficient *Escherichia coli* inactivation. *Chem. Eng. J.* **342**, 220–227 (2018).

Acknowledgments

This research was supported by grants from the Project of the NARO Bio-oriented Technology Research Advancement Institution (research program on the development of innovative technology) (29012C), the Northern Advancement Center for Science and Technology, and the Muroran Advancement Center of Industrial Technology and Management.

Author contributions

S.M. performed the biological experiments and wrote the paper. K.A. prepared the SLBGs and evaluated the persistence of pH. X.L. and M.K. provided advice on the biological experiments and wrote the paper. K.U. was involved in creating the pH indicator and wrote the paper. S.Y. and K.T. designed the project, performed the experimental work, analyzed the data, and wrote the paper.

Competing interests

K.U., S.Y., and K.T. established a startup company (Core Lab, Muroran, Japan) that manufactures and sells the deliverables of this project and have been appointed directors of that company. S.M., K.A., X.L., and M.K. have no competing interests as defined by Nature Research or other interests that might be perceived to influence the results and/or discussion reported in this paper.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-86796-w>.

Correspondence and requests for materials should be addressed to S.Y. or K.T.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021