

1-1-2016

Mitochondrial genome of an Atlantic white shark (*Carcharodon carcharias*)

Nicholas J. Marra
Cornell University

Minghui Wang
Cornell University

Qi Sun
Cornell University

Paulina D. Pavinski Bitar
Cornell University

Michael J. Stanhope
Cornell University

See next page for additional authors

Follow this and additional works at: https://nsuworks.nova.edu/cnso_bio_facarticles



Part of the [Biology Commons](#)


NSUWorks Citation

Marra, Nicholas J.; Minghui Wang; Qi Sun; Paulina D. Pavinski Bitar; Michael J. Stanhope; and Mahmood S. Shivji. 2016. "Mitochondrial genome of an Atlantic white shark (*Carcharodon carcharias*)."
Mitochondrial DNA Part B: Resources 1, (1): 717-719. doi:10.1080/23802359.2016.1222248.

This Article is brought to you for free and open access by the Department of Biological Sciences at NSUWorks. It has been accepted for inclusion in Biology Faculty Articles by an authorized administrator of NSUWorks. For more information, please contact nsuworks@nova.edu.

Authors

Nicholas J. Marra, Minghui Wang, Qi Sun, Paulina D. Pavinski Bitar, Michael J. Stanhope, and Mahmood S. Shivji



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

Mitochondrial genome of an Atlantic white shark (*Carcharodon carcharias*)

Nicholas J. Marra, Minghui Wang, Qi Sun, Paulina D. Pavinski Bitar, Michael J. Stanhope & Mahmood S. Shivji

To cite this article: Nicholas J. Marra, Minghui Wang, Qi Sun, Paulina D. Pavinski Bitar, Michael J. Stanhope & Mahmood S. Shivji (2016) Mitochondrial genome of an Atlantic white shark (*Carcharodon carcharias*), *Mitochondrial DNA Part B*, 1:1, 717-719, DOI: [10.1080/23802359.2016.1222248](https://doi.org/10.1080/23802359.2016.1222248)

To link to this article: <https://doi.org/10.1080/23802359.2016.1222248>



© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 22 Nov 2016.



Submit your article to this journal [↗](#)



Article views: 933



View related articles [↗](#)




View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

Mitochondrial genome of an Atlantic white shark (*Carcharodon carcharias*)

Nicholas J. Marra^{a,b} , Minghui Wang^c, Qi Sun^c, Paulina D. Pavinski Bitar^a, Michael J. Stanhope^{a*} and Mahmood S. Shivji^{b*}

^aDepartment of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY, USA; ^bSave Our Seas Shark Research Center and Guy Harvey Research Institute, Nova Southeastern University, Dania Beach, FL, USA; ^cComputational Biology Service Unit, Biotechnology Resource Center, Cornell University, Ithaca, NY, USA

ABSTRACT

Here we report the first full mitochondrial genome sequence for a white shark caught in the Atlantic Ocean. The mitochondrial genome is 16,745 bp in length and contains 13 protein-coding genes, 22 tRNA genes, 2 rRNA genes, and a non-coding control region. The base composition of this mtDNA lineage is A: 30.7%, C: 26.9%, G: 13.8%, and T: 28.6%. In concordance with previous population genetic studies, the Atlantic caught individual forms a separate lineage from individuals caught on either side of the Pacific Ocean.

ARTICLE HISTORY

Received 23 June 2016
Accepted 5 August 2016

KEYWORDS

Mitochondrial genome;
MITObim; *Carcharodon carcharias*

Mitochondrial genes exhibit a lower rate of evolution in elasmobranchs (Martin et al. 1992), thus complete mitochondrial genomes may be preferable to using a subset of the mitochondrial genome for identifying intraspecific variation (Feutry et al. 2014). The white shark, *Carcharodon carcharias*, is CITES listed and has been the subject of population genetic studies using the mitochondrial DNA (mtDNA) control region and microsatellite loci (Pardini et al. 2001; Jorgensen et al. 2010; Blower et al. 2012; O'Leary et al. 2015). These studies identified multiple mtDNA lineages but the only full mtDNA genome sequences for white shark come from the Pacific Ocean (off the coast of Taiwan, Chang et al. 2014; off the coast of Mexico, Diaz-Jaimes et al. 2014). We characterize the first full mitochondrial genome sequence for a white shark from another ocean basin, the Atlantic Ocean, in an effort to identify additional areas of the mitochondrial genome that could be exploited for identification of genetic variation in this species.

The individual sequenced in this study (accession number OC130-Cc_H stored at -80°C at Nova Southeastern University, College of Natural Sciences and Oceanography), was illegally fished off the coast of Delaware, USA in 2007 and confiscated by the National Oceanic and Atmospheric Administration Office for Law Enforcement but made available for this work (see Richards et al. 2013 for additional details). As part of a white shark nuclear genome sequencing project underway in our laboratory, Illumina HiSeq 2500

single end reads, 450 bp in length (produced by overlapping 250 bp reads) were used to assemble the mitochondrial genome. We identified reads from mtDNA by mapping reads to the RefSeq white shark mtDNA genome (NC_022415.1) using Bowtie2 (Langmead & Salzberg 2012). The mtDNA reads were used to produce a mitochondrial genome assembly with the program MITObim v. 1.8 (Hahn et al. 2013). This assembly was annotated using DOGMA (Wyman et al. 2004) as in Diaz-Jaimes et al. (2014) and annotations were confirmed by comparison to the gene annotation for the RefSeq sequence. Gene order followed that of Diaz-Jaimes et al. (2014) and that seen in most vertebrates. Using MUSCLE (Edgar 2004), we conducted pairwise alignments of the Atlantic and two Pacific shark sequences. Additionally, we reconstructed a phylogenetic tree with all RefSeq mitochondrial genomes from members of the Lamnidae (Figure 1) using PHYML (Guindon & Gascuel 2003).

The Atlantic shark sequence (gb: KX389266) is 16,745 bp in length and contains 13 protein-coding, 22 tRNA, and 2 rRNA genes, and a non-coding control region. The Atlantic sequence had a similar level of sequence identity with the Mexican (99.0%) and Taiwanese (98.9%) sequences while the two Pacific sequences are almost identical (99.8%). All subsequent comparisons herein discussed are to the RefSeq (Taiwanese) sequence. There are 174 nucleotide differences between the two genomes. As is the case with all of the lamnid sequences, *Cox1* has a GTG start codon in all three white shark sequences.

CONTACT Mahmood Shivji  mahmood@nova.edu  Save Our Seas Shark Research Center and Guy Harvey Research Institute, Nova Southeastern University, 8000 North Ocean Drive, Dania Beach, FL, 33004, USA; Michael Stanhope  mjs297@cornell.edu  Department of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY, 14853, USA

*These authors are co-senior authors of this paper.

© 2016 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. We used MUSCLE to align the mtDNA genome sequences from all members of the Lamniformes present in RefSeq along with the three white shark sequences. Sequences from *Cetorhinus maximus* and *Carcharias taurus* were included as an outgroup. This alignment was then used to produce a Maximum Likelihood Tree with the PHYML plugin for Geneious using the GTR + G model of evolution and 1000 bootstraps. Support values are displayed on a scale of 1 to 100 for the 1000 bootstraps and the root of the tree was placed at the base of the split between the Lamnidae and the two outgroup species. Each label includes the species name and identifying GI and RefSeq numbers for the sequence used. The two previously published white shark sequences from Mexico and Taiwan clustered together and were sister to the Atlantic caught individual used in this study.

The phylogeny depicts the Atlantic sequence as a separate lineage from the Mexican and Taiwanese lineages.

ORCID

Nicholas J. Marra  <http://orcid.org/0000-0002-1953-0787>

Acknowledgements

We thank L. Natanson and the NOAA Office for Law Enforcement for the shark sample.

Disclosure statement

We have no conflicts of interest to report and the authors are solely responsible for the content and writing of this paper.

Funding

This work was supported by the Save Our Seas Foundation and Guy Harvey Ocean Foundation.

References

- Blower DC, Pandolfi JM, Bruce BD, Gomez-Cabrera MDC, Ovenden JR. 2012. Population genetics of Australian white sharks reveals fine-scale spatial structure, transoceanic dispersal events and low effective population sizes. *Mar Ecol Prog Ser.* 455:229–244.
- Chang C-H, Shao K-T, Lin Y-S, Fang Y-C, Ho H-C. 2014. The complete mitochondrial genome of the great white shark, *Carcharodon carcharias* (Chondrichthyes, Lamnidae). *Mitochondrial DNA.* 23:357–358.
- Diaz-Jaimes P, Hinojosa-Alvarez S, Sánchez-Hernández X, Hoyos-Padilla M, García-De-León FJ. 2014. The complete mitochondrial DNA of white shark (*Carcharodon carcharias*) from Isla Guadalupe, Mexico. *Mitochondrial DNA.* 1736:1–2.

- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32:1792–1797.
- Feutry P, Kyne PM, Pillans RD, Chen X, Naylor GJP, Grewe PM. 2014. Mitogenomics of the Speartooth Shark challenges ten years of control region sequencing. *BMC Evol Biol.* 14:232.
- Guindon S, Gascuel O. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol.* 52:696–704.
- Hahn C, Bachmann L, Chevreur B. 2013. Reconstructing mitochondrial genomes directly from genomic next-generation sequencing reads: a baiting and iterative mapping approach. *Nucleic Acids Res.* 41:e129.
- Jorgensen SJ, Reeb CA, Chapple TK, Anderson S, Perle C, Van Sommeran SR, Fritz-Cope C, Brown AC, Klimley AP, Block BA. 2010. Philopatry and migration of Pacific white sharks. *Proc Biol Sci.* 277:679–688.
- Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 9:357–359.
- Martin AP, Naylor GJ, Palumbi SR. 1992. Rates of mitochondrial DNA evolution in sharks are slow compared with mammals. *Nature.* 357:153–155.
- O’Leary SJ, Feldheim KA, Fields AT, Natanson LJ, Wintner S, Hussey N, Shivji MS, Chapman DD. 2015. Genetic diversity of white sharks, *Carcharodon carcharias*, in the Northwest Atlantic and Southern Africa. *J Hered.* 106:258–265.
- Pardini AT, Jones CS, Noble LR, Kreiser B, Malcolm H, Bruce BD, Stevens JD, Cliff G, Scholl MC, Francis M, et al. 2001. Sex-biased dispersal of great white sharks. *Nature.* 412:139–140.
- Richards VP, Suzuki H, Stanhope MJ, Shivji MS. 2013. Characterization of the heart transcriptome of the white shark (*Carcharodon carcharias*). *BMC Genomics.* 14:697.
- Wyman SK, Jansen RK, Boore JL. 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics.* 20:3252–3255.