


## **$^{87}\text{Sr}/^{86}\text{Sr}$ dating of the Alcácer do Sal Formation (upper Miocene, mainland Portugal)**



### **Datação $^{87}\text{Sr}/^{86}\text{Sr}$ da Formação de Alcácer do Sal (Miocénico superior, Portugal continental)**

Ressurreição, R.<sup>1</sup>, Legoinha, P.<sup>2</sup>, Dias, R.<sup>1</sup>, Santos, J.<sup>3</sup>, Ribeiro, S.<sup>3</sup>, Patinha, C.<sup>3</sup>,  
Miranda, R.<sup>4</sup>

**DOI:**

Received em 31/12/2020 / Aceite em 17/05/2021

Published online em maio de 2021

© 2021 LNEG – Laboratório Nacional de Energia e Geologia IP

Artigo original  
Original article

**Abstract:** The Lower Tagus Basin was subjected to several marine transgressions along the Neogene, related to positive eustatic oscillations and also controlled by the regional tectonic activity. In the Alcácer do Sal-Sines region, previous studies identified an episode of marine sedimentation interpreted as having occurred during the late Serravalian to early Tortonian interval, based on biostratigraphic data. This episode is represented in the Alcácer do Sal Formation. The study of the SMS-12-01B borehole, drilled for mineral prospection purposes, located about 10 km NNE of Melides, allowed the identification of a 3 m thick layer of sediments containing marine fossils.  $^{87}\text{Sr}/^{86}\text{Sr}$  determinations of oyster shells point to deposition about 11.5 Ma ago, validating previous proposals for the age of the formation. These data correspond to the first numerical ages obtained for the Alcácer do Sal Formation, allowing a solid correlation with other known marine units in the Lower Tagus and Algarve basins related to the same transgression episode.

**Keywords:** Lower Tagus Basin, upper Miocene, Alcácer do Sal Formation,  $^{87}\text{Sr}/^{86}\text{Sr}$  ages.

**Resumo:** Ao longo do Neogénico, a Bacia do Baixo Tejo foi afectada por várias transgressões marinhas, relacionadas com oscilações eustáticas positivas articuladas com a tectónica regional. Na região de Alcácer do Sal-Sines, estudos anteriores identificaram um episódio de sedimentação marinha interpretado como tendo ocorrido durante o Serravaliano final a Tortoniano inicial, com base em dados bioestratigráficos. Este episódio está representado pela Formação de Alcácer do Sal. No estudo da sondagem SMS-12-01B, efectuada para prospecção mineral e localizada a cerca de 10 km a NNE de Melides, foi identificado um nível de sedimentos com 3 m de espessura contendo fósseis marinhos. A determinação de razões isotópicas  $^{87}\text{Sr}/^{86}\text{Sr}$  em conchas de ostras indica idade próxima de 11,5 Ma, validando interpretações anteriores sobre a provável idade da formação. Corresponde à primeira idade numérica obtida para esta unidade sedimentar, permitindo uma correlação sólida com outras unidades conhecidas nas bacias do Baixo Tejo e do Algarve associadas ao mesmo episódio transgressivo.

**Palavras-chave:** Bacia do Baixo Tejo, Miocénico superior, Formação de Alcácer do Sal, idades  $^{87}\text{Sr}/^{86}\text{Sr}$ .

<sup>3</sup> GeoBioTec, Departamento de Geociências, Universidade de Aveiro, 3810-193 Aveiro, Portugal.

<sup>4</sup> IDL – Instituto Dom Luiz, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal.

\*Corresponding author/Autor correspondente: ricardo.ressurreicao@lneg.pt

### **1. Introduction**

The Lower Tagus Basin is an elongated tectonic depression with general trending NNE-SSW, being the largest Cenozoic sedimentary basin in Portugal (*e.g* Pais *et al.*, 2012). It was formed during the Middle Eocene as a consequence of the Iberia-Eurasia collision, in the context of the Alpine orogeny event responsible for the genesis of the Cantabrian-Pyrenean Cordillera (*e.g* Cunha, 1992, 2019; Barbosa, 1995; de Vicente *et al.*, 2011, 2018).

In the southwestern/distal sector of the Lower Tagus Basin, the Cenozoic sedimentary record, is strongly related to sea level oscillations. Since the Aquitanian until the middle Tortonian, the basin was partially invaded by Atlantic waters for several times, as shown by the interbedding of marine and alluvial facies (*e.g* Cunha *et al.*, 2009; Pais *et al.*, 2012). An extensive stratigraphic analysis based on faunal assemblage and other methods, such as K-Ar dating and comparison of  $^{87}\text{Sr}/^{86}\text{Sr}$  ratios with the curve of Sr isotopic composition of seawater through time, among other data, has been crucial to the definition and dating of several depositional sequences (*e.g* Antunes *et al.*, 2000; Pais *et al.*, 2012, Legoinha and Flores, 2014). In the Lisbon-Setúbal region, the geological record shows the existence of ten sequences that can be correlated with the third-order cycles of Haq *et al.* (1987) (*op. cit.*).

To the south of the Arrábida Chain the record is less diversified and evidence of only one transgression exists. Between Alcácer do Sal and Sines, this event is represented by predominant coastal to marine siliciclastic sediments detected both in outcrops and boreholes, corresponding to the Alcácer do Sal Formation. This unit has been considered of late Serravalian to early Tortonian age based solely on biostratigraphic data (Antunes, 1983; Antunes and Mein, 1983).

<sup>1</sup> Laboratório Nacional de Energia e Geologia (LNEG), Apartado 7586 - Alfragide, 2610-999 Amadora, Portugal.

<sup>2</sup> GeoBioTec, Departamento de Ciências da Terra, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829 - 516 Caparica, Portugal.

In the Algarve basin (south of mainland Portugal), the upper Serravallian is overlain by lower Tortonian marine deposits, through an erosive discontinuity and a basal conglomerate with phosphatic nodules and glauconite, yielding *Globigerina concina* and cf. *Neogloboquadrina acostaensis*, and an oyster shell Sr isotopic age of 10.7 (+0.8 / -1.2) Ma (Armenteros *et al.*, 2019).

In this paper we present new numerical ages for this unit given by  $^{87}\text{Sr}/^{86}\text{Sr}$  measured in oyster shells sampled from the SMS-12-01B borehole (Alcácer do Sal-Melides region), being the first time this sedimentary unit is dated with a technique that provides numerical values. The importance of these results for a better understanding of the Alcácer do Sal Formation will be discussed, as well as its meaning in the context of the Lower Tagus Basin geological evolution. Figure 1 shows the Alcácer do Sal Formation geographic distribution, its cartographic relationship with other Cenozoic units and the location of the SMS-12-01B borehole.

## 2. The Alcácer do Sal Formation: general aspects and problematics

This lithostratigraphic unit was first proposed by Antunes (1983). In general, these sediments consist of micaceous sandstones, often clayey, with frequent carbonate-rich layers and fossiliferous sandstones, deposited in a shallow marine environment, as indicated by the faunas of ostracods, foraminifers, lamellibranchs, gastropods, fishes and mammals. The identification of small mammals at Cerrado da Pedra (Alcácer do Sal), cf. *Metaxytherium medium* (DESMAREST), *Galerix socialis* (von MEYER), *Prolagus oenningensis* (KONIG), *Heteroxerus aff. rubricatus* CRUSAFONT et VILLALTA, *Armantomys tricristatus* LOPEZ-MARTINEZ, *Miodyromys hamadryas* (MAJOR), *Megacricetodon aff. minor* (SCHAUB), *Megacricetodon cf. ibericus* (SCHAUB), *Hispanomys aguirrei* (SESE-BENITO)




Figure 1. Simplified geological map of the study area showing the SMS-12-01B borehole location. 1, Paleozoic; 2, Mesozoic; 3, Paleogene (Vale do Guizo Formation); 4, late Serravallian-early Tortonian (Alcácer do Sal Formation); 5, Messinian (Esbarrendadoiro and Melides formations); 6, Zanclean-Gelasian (Ulme, Alvalade, Galé and Vale Figueira formations); 7, Holocene (aeolian and beach sands); 8, Holocene (alluvium). Cenozoic stratigraphy according to Antunes and Pais (1983), Antunes *et al.* (1991), Pimentel (1997), Pais *et al.* (2012), Ressurreição (2018), Cunha (2019). Adapted from published geological maps at 1:50 000 scale (Antunes and Pais, 1983; Antunes *et al.*, 1991; Inverno *et al.*, 1986; Dias *et al.*, 2011).

Figura 1. Mapa geológico simplificado da área de estudo com a localização da sondagem SMS-12-01B. 1, Paleozóico; 2, Mesozóico; 3, Paleogénico (Formação de Vale do Guizo); 4, Serravaliano final-Tortoniano inicial (Formação do Alcácer do Sal); 5, Messiniano (formações de Melides e Esbarrendadoiro); 6, Zancleano-Gelasiano (formações de Ulme, Alvalade, Galé e Vale Figueira); 7, Holocénico (areias eólicas e de praia); 8, Holocénico (aluvões). Estratigrafia do Cenozóico com base em Antunes e Pais (1983), Antunes *et al.* (1991), Pimentel (1997), Pais *et al.* (2012), Ressurreição (2018), Cunha (2019). Adaptado da cartografia geológica publicada na escala 1:50 000 (Antunes e Pais, 1983; Antunes *et al.*, 1991; Inverno *et al.*, 1986; Dias *et al.*, 2011).

allowed a relatively precise stratigraphic positioning, compatible with the mammal unit MN8 (Antunes and Mein, 1983), corresponding to the late Serravalian to early Tortonian, between 11.9 and 11.1 Ma (Agustí *et al.*, 2001; Alba *et al.*, 2011; Casanovas-Vilar *et al.*, 2011).

It correlates with the upper part of the allostratigraphic unit UBS10 (Unconformity Bounded Sequence), proposed by Cunha (1992) and updated in subsequent studies (Cunha *et al.*, 2009; Pais *et al.*, 2010; Pais *et al.*, 2012; Cunha, 2019). The Alcácer do Sal Formation is limited at the top and bottom by erosional surfaces. Usually, it overlies, by disconformity, the Paleogene (Vale do Guizo Formation – UBS7 to 8) and is covered by the Upper Miocene (Melides/Esbarrendadoiro formations – UBS12) or Plio-Quaternary (Alvalade/Ulme/Galé formations – UBS13) (Antunes and Pais, 1983; Pimentel, 1997; Dias *et al.*, 2011, 2016; Ressurreição 2018). Figure 2 presents the stratigraphic correlation between the several Cenozoic units in the study sector and adjacent sedimentation areas.

The exact geographical extension of the Alcácer do Sal Formation is still unknown. There is an apparent geographic continuity of Miocene marine/paralic sediments along the eastern margin of the Lower Tagus Basin, between the Lavre and Alcácer do Sal regions, extending south to the Sines area as shown by the published geologic maps at 1:50 000 scale (Zbyszewski and Ferreira, 1972; Zbyszewski *et al.*, 1976; Antunes and Pais, 1983; Inverno *et al.*, 1986, 1993; Carvalhosa and Zbyszewski, 1994; Dias *et al.*, 2011). Despite this continuity, the resolution of the stratigraphic knowledge is different from place to place.

The coastal and marine sediments ascribed to the Alcácer do Sal Formation (deposition during the late Serravalian to early Tortonian) were identified in the Alcácer do Sal, Melides and Sines regions (Antunes, 1983; Inverno *et al.*, 1993; Dias *et al.*, 2011, 2016). However, the age of the Melides sediments has been matter of debate. Besides the cited studies, Dollfus *et al.* (1904) considered deposition during the Tortonian; Cachão (1995) defined the “Melides Calcarenites” of Serravalian age, but older than the Alcácer do Sal Formation.

To the north of the Alcácer do Sal area, in the right margin of the Sado River, the mapping of the Miocene marine sediments is older and a “Vindobonian” age was proposed, based on its fossil content (Zbyszewski *et al.*, 1972; Zbyszewski and Ferreira, 1976; Carvalhosa and Zbyszewski, 1994). These are the “Complexo argilo-gresoso de Palma e do Vale de Marateca”, “Complexo flúvio-marinho de Lavre” and “Complexo argiloso marinho de Lavre” (*op cit.*). The “Vindobonian” age, which corresponds to the Langhian to Tortonian time interval (Glaessner, 1953), covers a time span larger than that presently attributed to the Alcácer do Sal Formation. Whilst we can reasonably assume that the sediments occurring near Alcácer do Sal are part of this unit (following Antunes, 1983), for those positioned in the Lavre region, about 50 km to NNE, this correlation might not be accurate, and a better stratigraphic resolution should be pursued with the aid of techniques that can provide numerical ages.

Despite its small surficial (outcrop) representation, borehole data suggests a significant extension and continuity for this unit below younger sediments, covering a large area located to the west

Erat./Era	Syst./Per.	Series/Epoch	Stage/Age	Ma	Depositional sequences	Mammal Zone	Lower Tagus Basin		Alvalade Basin	UBS (Cunha, 1992)
							Lisbon-Almada	Setúbal Peninsula western littoral	Lower Tagus Basin meridional sector and Coastal Alentejo	
Cenozoic	Quat.	Holocene					Alluvium	Alluvium	Alluvium	
		U. Upper	0,0117					Eolian sands	Beach sands	Eolian sands
		Chibanian	0,129						Eolianites	
		M. Calabrian	0,774					Terraces	Terraces	
		L. Gelasian	1,80		MN 17				Marco Furado Fm. Belverde Conglomerate	
	Neogene	Pliocene		2,58	MN 16				Vale Figueira Fm.	
		U. Piacenzian	3,60		MN 15				Santa Marta Fm.	
		L. Zanclean	5,333		MN 14				Galé Fm. / Ulme Fm.	
		Messinian	7,246		MN 13				Rio da Prata sands	
		U. Tortonian	11,63		MN 12				Melides Fm. / Esbarrendadoiro Fm.	
Paleogene	Miocene	Serravalian		T2	MN 10	Cabo Ruivo Sands Bráo de Prata Sands Marvila Limestones		Ribeira das Lages deposits Penedo Norte glauconitic dep.		
		M. Langhian	13,82	T1	MN 9					
		L. Burdigalian	15,97		MN 7 + 8				Alcácer do Sal Fm.	
		Aquitanian	20,44		S2	Grilos Sandstones				
		Oligocene	23,03		S1	Xabregas Blue Clays Quinta das Conchas limestones		Penedo Norte deposits		
	Eocene	U. Chattian	27,82		L1	MN 6				
		L. Rupelian	33,9		B2	MN 5	Vale de Chelas Sands Musgueira Limestones			
		U. Priabonian	37,71		B1	MN 4	Placuna miocénica Casal Vistoso Limestones			
		M. Bartonian	41,3			MN 3	Quinta do Bacalhau Sands Forno do Tijoelo Blue Clays			
		L. Lutetian	47,8				Banco Real			
	Paleocene	U. Ypresian	56,0		B0		Estefânia Sands			
		M. Thetian	59,2		A2	MN 2	Prazeres Clay Fm.			
		L. Seländig	61,6		A1	MN 1				
		L. Danian	66,0				Fm. Benfica	Fm. Benfica	Vale do Guizo Fm.	

Figure 2. Stratigraphic correlation between the Cenozoic units in the sector under study and adjacent sedimentation areas. Modified from Cunha *et al.* (2009), Pais *et al.* (2010), Pais *et al.* (2012), Ressurreição (2018), Cunha (2019).

Figura 2. Correlação estratigráfica entre as unidades cenozóicas existentes no sector em estudo e áreas de sedimentação adjacentes. Modificado de Cunha *et al.* (2009), Pais *et al.* (2010), Pais *et al.* (2012), Ressurreição (2018), Cunha (2019).

and south of Alcácer do Sal and to the west of the Grândola range, until the present-day coastline.

### 3. Description of the SMS-12-01B borehole

The SMS-12-01B borehole (Fig. 3) was drilled by the Colt Resources mining company for massive sulphides prospection purposes. It is located at the Brejinho de Água area, near the Porqueira geodesic vertex, about 10 km NNE of Melides and 22 km SW of Alcácer do Sal. The borehole is 441 m deep, starting at an elevation of 70 m. At elevation -151 m, Paleozoic metasediments of the South Portuguese Zone were intercepted. Above these, from -151 to -81 m Paleogene reddish and whitish conglomerates, sandstones and mudstones with carbonate cement occur. Between -81 and 11 m there is a succession of yellowish, whitish and brownish clayey sands and silts, sometimes with carbonate cement, interpreted as deposited during the Miocene. From 29 to 70 m, yellowish, whitish and brownish Plio-Pleistocene sandstones exist (UBS13 allostratigraphic unit); no record was obtained between 11 and 29 m. The surficial horizon corresponds to Holocene aeolian sands, however, their thickness at the borehole location is negligible. This stratigraphic interpretation is based on knowledge obtained from regional surface surveys and previously published studies (*e.g.* Antunes, 1983; Pimentel, 1997; Dias *et al.*, 2011, 2016; Ressurreição *et al.*, 2014; Ressurreição, 2018).

It is not clear if at the SMS-12-01B location, the Miocene succession (92 m thick) corresponds entirely to the Alcácer do Sal Formation.




Figure 3. SMS-12-01B borehole log description.

Figura 3. Descrição do log da sondagem SMS-12-01B.

However, since neither ruptures nor major lithological changes were identified in the well, we assume a thickness of approximately 90 m for this unit. According to Antunes (1983), this unit has a thickness of at least 69 m at Alcácer do Sal and of 146 m at Asseiceira-Albergaria (respectively 22 km and 15 km NE of the SMS-12-01B location). Around elevation -40 m, there is a 3 m thick horizon rich in bioclasts, being the only fossiliferous level identified in the Miocene succession. In this horizon fragments of oysters, benthonic foraminifers (*Elphidium crispum*, *Cibicides rufulgens*, *Ammonia beccarii*), ostracods and fish teeth were identified, which are compatible with shallow marine environment, with moderately warm waters, generally of low energy levels, in a sheltered gulf with depth not exceeding the photic zone (~20 m), sandy bottoms and salinity sometimes lower




Figure 4. Siliciclastic sandstone cemented by carbonate sampled from the SMS-12-01B borehole, containing oyster shells (*Crassostrea*) A, B and C.

Figura 4. Amostra de arenito siliciclástico com cimento carbonatado da sondagem SMS-12-01B, contendo as conchas de ostreídeos (*Crassostrea*) A, B e C.

than normal (as indicated by *Terebralia* and oysters), as stated by Antunes (1983). The absence of planktonic foraminifera also excludes an open ocean environment.

### 3. Methods and analytical results: Sr isotopic analysis and conversion of $^{87}\text{Sr}/^{86}\text{Sr}$ ratios to numerical ages

Three shells (A, B, and C) of *Crassostrea* from the fossiliferous sediment SMS-12-01B were used for Sr isotopic analysis at the Laboratory of Isotope Geology of the University of Aveiro (Portugal).

The shells were collected from a 12 cm thick portion of the borehole (Fig. 4) and occupy the same stratigraphic position. Shell A was the largest one and it was sampled into four different aliquots (1, 2, 3, and 4) to be analysed separately. From shells B and C only one aliquot was obtained in each case.

To avoid hypothetical alteration or contamination in the surface of the shell, a 1 cm hole was drilled in each specimen and the inner material was sampled. All the samples were coarse crushed into 1 - 3 mm sized fragments in agate mortar.

About 0.5 g from the fragmented samples were previously cleaned with Milli-Q water and then submitted to an acid-leaching process to remove the soluble labile Sr component from diagenetic or alteration provenance. Four ml of 5 M hydrochloric acid solution was added to each sample in a centrifuge tube at room temperature for some minutes. After that, the liquid was discarded, and the sample was rinsed three times with Milli-Q water to remove all the acid. Following leaching, and to ensure that only carbonate fraction was sampled, a weak acid was used for the dissolution. Each sample was dissolved with 10 ml acetic acid (1M – environmental grade), in a PTFE vessel during 1 - 2 h in a hotplate. For the  $^{87}\text{Sr}/^{86}\text{Sr}$  analysis, the solution was transferred to a new vessel, dried and the precipitate was redissolved in 3 ml of nitric acid (conc.) and dried. Sr-Resin (50 - 100  $\mu\text{m}$ , Eichrom) ion chromatography columns were used to separate Sr from the other elements. Strontium obtained this way was loaded onto Ta filaments with phosphoric acid, and the isotopic ratio was measured with a VG Sector 54 thermal ionization mass spectrometer in dynamic mode. Typical runs consisted in the acquisition of 60 isotopic ratios, with peak measurements at 1 - 2 V for  $^{88}\text{Sr}$ , and the analytical data were corrected for mass fractionation relative to  $^{88}\text{Sr}/^{86}\text{Sr} = 0.1194$ . All reagents used in the global procedure were sub-boiling distilled, and the water was produced by a Milli-Q Element (Millipore) apparatus. The procedure average blanks levels were less than 2 ng, which are

negligible compared to the total amount of Sr processed in the analytical protocol. During this study, the SRM-987 standard gave an average value of <sup>87</sup>Sr/<sup>86</sup>Sr = 0.710245 ± 0.000015 (conf. lim. = 95%; N = 12).

The obtained <sup>87</sup>Sr/<sup>86</sup>Sr ratios were converted to numerical ages (Tab. 1) based on the works by Howarth and McArthur (1997), McArthur *et al.* (2001), and McArthur *et al.* (2012), and using the Look-Up Table Version 3:10 / 99.

<sup>87</sup>Sr/<sup>86</sup>Sr ratios obtained in four portions of shell A go from 0.708869 to 0.708822. By comparing these values with the curve of variation of the Sr isotopic composition of seawater, as proposed in the quoted works, the obtained age values vary between 10.73 and 12.76 Ma. Measurements in the other two smaller shells gave <sup>87</sup>Sr/<sup>86</sup>Sr ratios of 0.708889 (shell B) and 0.708921 (shell C), that would correspond to ages of ca. 10 Ma and 9 Ma, respectively.

4. Discussion and conclusions  
The <sup>87</sup>Sr/<sup>86</sup>Sr ages indicate that the sampled Miocene sediment, considering its sedimentological properties and geographical location, corresponds to the Alcácer do Sal Formation. They also strengthen the plausibility of a correlation between the Melides sediments and this unit, as done by other authors (Antunes, 1983; Dias *et al.*, 2011, 2016).

The obtained ages, 8.64 - 12.76 Ma, imply a larger time interval than that proposed by the biostratigraphic data collected from the Alcácer do Sal Formation (Antunes, 1983; Antunes and Mein 1983), meaning that the unit could extend to the upper Tortonian. At first sight, this would have implied the maintenance of the same depositional environment for approximately 4 Ma and a very low sedimentation rate, which seems unlikely, in the place where the sample was collected.

A reasonable hypothesis is that the larger shell has better preserved in its inner parts the original strontium isotopic composition, whilst the smaller shells could have been affected by interaction with post-depositional fluids, which are usually expected to contribute to increase the <sup>87</sup>Sr/<sup>86</sup>Sr ratio (*e.g* Macdougall, 1991, and references therein), therefore leading, in the case of the shells studied in this work, to apparent younger "ages". The "ages" obtained in shells B and C could also be explained by a process of differential enrichment in the precipitation of Sr<sup>2+</sup> transported by diffusion and/or advection from younger stratigraphic levels (Baker *et al.*, 1982). Regardless

of the origin and the precise mechanism of Sr mobility, the larger shell is most likely giving results more closely representative of the isotopic composition of the seawater contemporary of the oysters, and consequently, its <sup>87</sup>Sr/<sup>86</sup>Sr values may be viewed as the most reliable in the determination of the true age of the sedimentation.

Despite the mentioned uncertainties, Sr mean ages of the larger shell point to a deposition at 11.53 (+0.28 / -0.29) Ma ago (Fig. 5), in the early Tortonian, not totally excluding the upper Serravalian.

The Alcácer do Sal Formation is associated with the transgressive event also represented in the Lisbon - Setúbal area by the "Marvila sandstones", "Braço de Prata Sands", "Penedo Norte Glauconitic deposits" and "Ribeira das Lages deposits", which largely correspond to the T1 depositional sequence and with the third-order cycle 3.1 of Haq *et al.* (1987) (Legoinha, 2001). The paleogeographic reconstruction during the early Tortonian was proposed by Pais *et al.* (2012) and Cunha (2019) (Fig. 6).

The data here presented have high relevance for the characterization of the Alcácer do Sal Formation since it validates previous interpretations based on biostratigraphy with new numerical ages. However, for a more complete understanding of this unit, namely its spatial and temporal evolution, more extensive sampling and numerical dating should be done. The verification if the marine Miocene sediments that crop out along the left margin of the Lower Tagus Basin correspond to this transgressive event has a significant implication on the knowledge of this sedimentary basin evolution. Since the mentioned sediments reach the Lavre region, located about 80 km far from the present coastline, they are related to one of the broadest transgressions that affected the basin.

## Acknowledgments

The authors are grateful for the financial support of GeoBioTec (UIDB/04035/2020). We would like to thank the late João Pais for the several discussions about the Neogene problematics in the Lower Tagus and Alvalade basins. We also thank Pedro Proença e Cunha, Nuno Pimentel and Ildefonso Armenteros for the constructive comments and suggestions that greatly improved the manuscript.

Table 1. Sr isotope ratios and corresponding ages using the curve of variation of the Sr isotopic composition of seawater.

Tabela 1. Razões isotópicas Sr e idades correspondentes utilizando a curva de variação da composição isotópica de Sr da água do mar.

Sample	Specimen	<sup>87</sup> Sr/ <sup>86</sup> Sr	Error (2σ)	Age (Ma)	Age error a)	Age error b)
A-1	A (large shell)	<b>0.708869</b>	0.000021	<b>10.73</b>	+0.23 / -0.26	+0.98 / -1.14
A-2	A (large shell)	<b>0.708822</b>	0.000020	<b>12.76</b>	+0.33 / -0.41	+1.70 / -1.40
A-3	A (large shell)	<b>0.708851</b>	0.000024	<b>11.30</b>	+0.28 / -0.25	+1.54 / -1.06
A-4	A (large shell)	<b>0.708850</b>	0.000013	<b>11.34</b>	+0.28 / -0.26	+0.92 / -0.66
B	B (small shell)	<b>0.708921</b>	0.000017	<b>8.64</b>	+0.35 / -0.46	+1.07 / -1.50
C	C (small shell)	<b>0.708889</b>	0.000017	<b>10.00</b>	+0.28 / -0.32	+0.87 / -1.02

Notes: Sr isotope ratios were converted to numerical ages based on the works by Howarth and McArthur (1997), McArthur *et al.* (2001) and McArthur *et al.* (2012) and using the Look-Up Table Version 3:10/99 of R. J. Howarth and J. M. McArthur. Age error a) assumes only the error of the curve of variation of the Sr isotopic composition of seawater; age error b) assumes both the error of the curve and the analytical error of Sr isotopic analysis in this work.

Notas: As razões isotópicas de Sr foram convertidas em idades numéricas com base nos trabalhos de Howarth e McArthur (1997), McArthur *et al.* (2001) e McArthur *et al.* (2012) e na utilização do documento "Look-Up Table Version 3:10/99", de R. J. Howarth e J. M. McArthur. O erro da idade a) assume unicamente o erro da curva de variação da composição isotópica de Sr da água do mar; o erro da idade b) assume conjuntamente os erros dessa curva e o erro da análise isotópica de Sr neste trabalho.




Figure 6. Paleogeographic reconstruction of the Lower Tagus Basin for the early Tortonian (adapted from Pais et al., 2012 and Cunha, 2019).

Figura 6. Reconstrução paleogeográfica da Bacia do Baixo Tejo para o Tortoniano inicial (adaptado de Pais et al., 2012 e Cunha, 2019).




Figure 5. Graph projection of Sr ages and average for shell A, using age error a) from table 1 (assumes only the error of the curve of variation of the Sr isotopic composition of seawater). MN zonation based on Alba et al. (2011) and Casanovas-Vilar et al. (2011).

Figura 5. Projeção gráfica das idades de Sr e da média aritmética para a concha A, utilizando o erro da idade a) constante na tabela 1 (assume unicamente o erro da curva de variação da composição isotópica de Sr da água do mar). Zonação MN baseada em Alba et al. (2011) e Casanovas-Vilar et al. (2011).

## References

- Agustí, J., Cabrera, L., Garcés, M., Krijgsman, W., Oms, O., Parés, J. M., 2001. A calibrated mammal scale for the Neogene of Western Europe. *State of the art Earth-Science Reviews*, **52**: 247-260. [https://doi.org/10.1016/S0012-8252\(00\)00025-8](https://doi.org/10.1016/S0012-8252(00)00025-8).
- Alba, D. M., Casanovas-Vilar, I., Robles, J. M., Moyà-Solà, S., 2011. Parada 3. El Aragoniense superior y la transición con el Vallesiense: Can Mata y la exposición paleontológica de els Hostalets de Pierola. *Paleontología i Evolució*, Memòria especial, **6**: 95-109.
- Antunes, M. T., 1983. *Notícia explicativa da Folha 39-C Alcácer do Sal, Carta Geológica de Portugal na escala 1:50 000*. Serviços Geológicos de Portugal, Lisboa, 21.
- Antunes, M. T., Pais, J., 1983. *Folha 39-B Alcácer do Sal, Carta Geológica de Portugal na escala 1:50 000*. Serviços Geológicos de Portugal, Lisboa.
- Antunes, M. T., Mein, P., 1983. Petits mammifères de la fin du Miocène moyen dans les dépôts marins de Cerrado da Pedra, Alcácer do Sal. Conséquences stratigraphiques. *Comunicações dos Serviços Geológicos de Portugal*, **69**(2): 373-379.
- Antunes, M. T., Pais, J., Gonçalves, F., Oliveira, J. T., 1991. *Folha 39-D Torrão, Carta Geológica de Portugal na escala 1:50 000*. Serviços Geológicos de Portugal, Lisboa.
- Armenteros, I., Dabrio, C., Legoinha, P., González-Delgado, J. A., Martínez-Graña, A. M., Civis, J., Pais, J., 2019. Facies and sequence analysis of Miocene open-shelf warm-temperate carbonates in Portimão (Lagos-Portimão Formation, Portugal). *Facies*, **65**: 33. <https://doi.org/10.1007/s10347-019-0575-2>.
- Baker, P. A., Gieskes, J. M., Elderfield, H., 1982. Diagenesis of carbonate in deep-sea sediments—Evidence from Sr/Ca ratio and interstitial dissolved Sr<sup>2+</sup> data. *Journal of Sedimentary Petrology*, **52**: 71-82.
- Barbosa, B., 1995. *Alostratigrafia e litostratigrafia das unidades continentais da Bacia terciária do baixo Tejo. Relações com o eustatismo e a tectónica*. Dissertação de Doutoramento, Univ. de Lisboa, 253.
- Carvalho, A., Zbyszewski, G., 1994. *Folha 35-D Montemor-o-Novo, Carta Geológica de Portugal na escala 1:50 000*. Serviços Geológicos de Portugal, Lisboa.
- Casanovas-Vilar, I., Alba, D. M., Garcés, M., Robles, J. M., Moyà-Solà, S., 2011. Updated chronology for the Miocene hominoid radiation in Western Eurasia. *Proceedings of the National Academy of Sciences of the U.S.A.*, **108**: 5554-5559. <https://doi.org/10.1073/pnas.1018562108>.
- Cunha, P. P., 1992. *Estratigrafia e sedimentologia dos depósitos do Cretáceo Superior e Terciário de Portugal Central, a leste de Coimbra*. Dissertação de Doutoramento, Univ. Coimbra, 262.
- Cunha, P. P., 2019. Cenozoic Basins of Western Iberia: Mondego, Lower Tejo and Alvalade basins. In: Quesada, C., Oliveira, J. T. (Eds.). *The Geology of Iberia: A Geodynamic Approach*. Regional Geology Reviews, Cenozoic Basins, Springer International Publishing, **4**(4):105-130. <https://doi.org/10.1007/978-3-030-11190-8>.
- Cunha, P. P., Pais, J., Legoinha, P., 2009. Evolução geológica de Portugal continental durante o Cenozoico-sedimentação aluvial e marinha numa margem continental passiva (Iberia ocidental). *Simpósio sobre el Margen Ibérico Atlántico*, MIA09, 11-20.
- de Vicente, G., Cloetingh, S., Van Wees, J. D., Cunha, P. P., 2011. Tectonic classification of Cenozoic Iberian foreland basins. *Tectonophysics*, **502**(1-2): 38-61. <https://doi.org/10.1016/j.tecto.2011.02.007>.
- de Vicente, G., Cunha, P. P., Muñoz-Martín, A., Cloetingh, S. A. P. L., Olaiz, A., Vegas, R., 2018. The Spanish–Portuguese Central System: an example of intense intraplate deformation and strain partitioning. *Tectonics*, **37**: 4444-4469. <https://doi.org/10.1029/2018TC005204>.
- Dias, R. P. (Coord.), Oliveira, J. T., Leal, J., Romão J., Inverno, C., Manuppella, G., Machado, S., Rodrigues, A., Bartolomeu, A., Pais, J., Ressurreição, R., 2011. *Folha 42-A Grândola, Carta Geológica de Portugal na escala 1:50 000*. Laboratório Nacional de Energia e Geologia, Lisboa.
- Dias, R. P., Oliveira, J. T., Matos, J. X., Ressurreição, R., Pereira, Z., Machado, S., Pais, J., Manuppella, G., 2016. *Notícia Explicativa da Folha 42-A Grândola, Carta Geológica de Portugal na escala 1:50 000*. Laboratório Nacional de Energia e Geologia, 106.
- Dollfus, G., Cotter, I. C. B., Gomes, I. P., 1903-1904. Mollusques tertiaires du Portugal: Planches de Céphalopodes, Gastéropodes et Pelecypodes laissées par F.A. Pereira da Costa, accompagnées d'une explication sommaire et d'une esquisse géologique. *Memórias/Comissão do Serviço Geológico de Portugal*, 48.
- Glaessner, M. F., 1953. Time Stratigraphy and the Miocene Epoch. *GSA Bulletin*, **64**(6): 647-658. [https://doi.org/10.1130/0016-7606\(1953\)64\[647:TATME\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1953)64[647:TATME]2.0.CO;2).
- Haq, B. U., Hardenbol, J., Vail, P. R., 1987. Chronology of fluctuating sea levels since the Triassic (250 million years ago to present). *Science*, **235**: 1156-1167.
- Howarth, R. J., McArthur, J. M., 1997. Statistics for strontium isotope stratigraphy. A robust LOWESS fit to the marine Sr-isotope curve for 0-

- 206 Ma, with look up table for te derivation of numerical age. *Journal of Geology*, **105**: 441-456. <https://doi.org/10.1086/515938>.
- Inverno, M. C., Manuppella, G., Zbyszewski, 1986. *Folha 42-C Santiago do Cacém, Carta Geológica de Portugal na escala 1:50 000*. Lisboa, Serviços Geológicos de Portugal.
- Inverno, M. C., Manuppella, G., Zbyszewski, G., Pais, J., Ribeiro, M. L., 1993. *Notícia Explicativa da Folha 42-C Santiago do Cacém, Carta Geológica de Portugal na escala 1:50 000*. Lisboa, Serviços Geológicos de Portugal.
- Legoinha, P., 2001. *Biostratigrafia de Foraminíferos do Miocénico de Portugal (Baixo Tejo e Algarve)*. Dissertação de Doutoramento, Universidade Nova de Lisboa, Faculdade de ciências e Tecnologia, 239.
- Legoinha, P., Flores, A.-J., 2014. Refinement of the biostratigraphy and biochronology of the Belverde borehole (Setúbal Peninsula, Portugal) using calcareous nannofossil data. In: Rocha, R., Pais, J., Kullberg, J. C., Finney, S. (Eds.), *STRATI 2013*, Springer Geology, 1119-1122. [https://doi.org/10.1007/978-3-319-04364-7\\_214](https://doi.org/10.1007/978-3-319-04364-7_214).
- Macdougall, J. D., 1991. Radiogenic isotopes in seawater and sedimentary systems. *Mineralogical Association of Canada Short Course Handbook*, **19**: 337-364.
- McArthur, J. M., Howarth, R. J., Bailey, T. R., 2001. Strontium isotope stratigraphy: LOWESS version 3: best fit to the marine Sr-isotope curve for 0-509 Ma and accompanying look-up table for deriving numerical age. *The Journal of Geology*, **109**: 155-170. <https://doi.org/10.1086/319243>.
- McArthur, J. M., Howarth, R. J., Shields, G. A., 2012. Strontium isotope stratigraphy. In: Gradstein, F. M., Ogg, J. G., Schmitz, M., Ogg, G. (Eds.), *The Geologic Time Scale 2012*, Elsevier, Amsterdam, 127-143. <https://doi.org/10.1016/b978-0-444-59425-9.00007-x>.
- Pais, J., Cunha, P. P., Legoinha, P., 2010. Litostratigrafia do Cenozoico de Portugal. In: Neiva, J. M. C., Ribeiro, A., Victor, L. M., Noronha, F., Ramalho, M. (Eds.), *Ciências Geológicas: Ensino e Investigação*, **I**: 365-376.
- Pais, J., Cunha, P. P., Pereira, D., Legoinha, P., Dias, R. P., Moura, D., Brum da Silveira, A., Kullberg, J. C., González-Delgado, J. A., 2012. *The Paleogene and Neogene of Western Iberia (Portugal): A Cenozoic Record in the European Atlantic Domain*. Springer Briefs in Earth Sciences, Springer, 158. <https://doi.org/10.1007/978-3-642-22401-0>.
- Pimentel, N. L., 1997. *O Terciário da Bacia do Sado – Sedimentologia e Análise Tectono-Sedimentar*. Tese de Doutoramento, Univ. Lisboa, 381.
- Ressurreição, R., 2018. *Evolução tectono-estratigráfica cenozóica do litoral alentejano (sector Melides-Odemira) e enquadramento no regime geodinâmico actual*. Tese de Doutoramento, FCUL, 296.
- Ressurreição, R., Dias, R., Cabral, J., Pais, J., 2014. Estratigrafia do Cenozóico no Alentejo litoral (sector Melides-Vila Nova de Milfontes). IX CNG/2º CoGePLIP, Porto 2014, *Comunicações Geológicas*, **101(I)**: 547-550.
- Zbyszewski, G., Ferreira, O. V., 1972. *Folha 39-A Águas de Moura, Carta Geológica de Portugal na escala 1:50 000*. Serviços Geológicos de Portugal, Lisboa.
- Zbyszewski, G., Ferreira, O. V., Carvalhosa, A., 1976. *Folha 35-B Mora, Carta Geológica de Portugal na escala 1:50 000*. Serviços Geológicos de Portugal, Lisboa.