

PALEOGENOMICS

Ten millennia of hepatitis B virus evolution

Arthur Kocher^{1,2,3*}, Luka Papac^{2,3}, Rodrigo Barquera^{2,3}, Felix M. Key^{2,4}, Maria A. Spyrou^{2,3,5}, Ron Hübner², Adam B. Rohrlach^{2,3,6}, Franziska Aron², Raphaela Stahl², Antje Wissgott², Florian van Bömmel⁷, Maria Pfefferkorn⁷, Alissa Mittnik^{2,8,9}, Vanessa Villalba-Mouco^{2,10}, Gunnar U. Neumann^{2,3}, Maité Rivollat^{2,11}, Marieke S. van de Loosdrecht², Kerttu Majander^{2,12}, Rezeda I. Tukhbatova^{2,13}, Lyazzat Musralina^{2,3,14,15}, Ayshin Ghalichi^{2,3}, Sandra Penske^{2,3}, Susanna Sabin², Megan Michel^{2,3,9}, Joscha Gretzinger^{2,3}, Elizabeth A. Nelson², Tiago Ferraz^{2,16}, Kathrin Nägele^{2,3}, Cody Parker^{2,17}, Marcel Keller^{2,18}, Evelyn K. Guevara^{2,19}, Michal Feldman^{2,5}, Stefanie Eisenmann^{2,3}, Eirini Skourtanioti^{2,3}, Karen Giffin^{2,3}, Guido Alberto Gneccchi-Ruscione^{2,3}, Susanne Friederich²⁰, Vittoria Schimmenti²¹, Valery Khartanovich²², Marina K. Karapetian²³, Mikhail S. Chaplygin²⁴, Vladimir V. Kufteer²⁵, Aleksandr A. Khokhlov²⁶, Andrey A. Chizhevsky²⁷, Dmitry A. Stashenkov²⁸, Anna F. Kochkina²⁸, Cristina Tejedor-Rodríguez²⁹, Íñigo García-Martínez de Lagrán³⁰, Héctor Arcusa-Magallón³¹, Rafael Garrido-Pena³², José Ignacio Royo-Guillén³³, Jan Nováček^{34,35}, Stéphane Rottier³¹, Sacha Kacki^{31,36}, Sylvie Saintot³⁷, Elena Kaverzheva³⁸, Andrej B. Belinskiy³⁹, Petr Velemínský⁴⁰, Petr Limburský⁴¹, Michal Kostka⁴², Louise Loe⁴³, Elizabeth Popescu⁴⁴, Rachel Clarke⁴⁴, Alice Lyons⁴⁴, Richard Mortimer⁴⁴, Antti Sajantila^{19,45}, Yadira Chinique de Armas⁴⁶, Silvia Teresita Hernandez Godoy^{47,48}, Diana I. Hernández-Zaragoza^{49,50}, Jessica Pearson⁵¹, Didier Binder⁵², Philippe Lefranc⁵³, Anatoly R. Kantorovich⁵⁴, Vladimir E. Maslov⁵⁵, Luca Lai^{56,57}, Magdalena Zoledziewska⁵⁸, Jessica F. Beckett⁵⁹, Michaela Langová⁴¹, Alžběta Danielisová⁴¹, Tara Ingman⁶⁰, Gabriel García Atiénzar⁶¹, María Paz de Miguel Ibáñez⁶¹, Alejandro Romero^{61,62}, Alessandra Sperduti^{63,64}, Sophie Beckett^{65,66,67}, Susannah J. Salter^{65,68}, Emma D. Zilivinskaya²⁵, Dmitry V. Vasil'ev⁶⁹, Kristin von Heyking⁷⁰, Richard L. Burger⁷¹, Lucy C. Salazar⁷¹, Luc Amkreutz⁷², Masnav Navruzbekov⁷³, Eva Rosenstock⁷⁴, Carmen Alonso-Fernández⁷⁵, Vladimir Slavchev⁷⁶, Alexey A. Kalmykov³⁹, Biaslan Ch. Atabiev⁷⁷, Elena Batieva⁷⁸, Micaela Alvarez Calmet⁷⁹, Bastien Llamas^{80,81,82}, Michael Schultz^{83,84}, Raiko Krauß⁸⁵, Javier Jiménez-Echevarría⁷⁵, Michael Francken⁸⁶, Svetlana Shnaider⁸⁷, Peter de Knijff⁸⁸, Eveline Altena⁸⁸, Katrien Van de Vijver^{89,90,91}, Lars Fehren-Schmitz^{92,93}, Tiffany A. Tung⁹⁴, Sandra Lösch⁹⁵, Maria Dobrovolskaya⁵⁵, Nikolaj Makarov⁵⁵, Chris Read⁹⁶, Melanie Van Twest⁶⁵, Claudia Sagona⁹⁷, Peter C. Ramsi⁹⁸, Murat Akar⁹⁹, K. Aslihan Yener¹⁰⁰, Eduardo Carmona Ballesterio^{101,102}, Francesco Cucca^{58,103}, Vittorio Mazzarello¹⁰³, Pilar Utrilla¹⁰⁴, Kurt Rademaker¹⁰⁵, Eva Fernández-Domínguez³⁶, Douglas Baird⁵¹, Patrick Semal⁸⁹, Lourdes Márquez-Morfin¹⁰⁶, Mirjana Rokсандić^{46,107,108}, Hubert Steiner¹⁰⁹, Domingo Carlos Salazar-García^{110,111,112}, Natalia Shishlina^{22,38}, Yilmaz Selim Erdal¹¹³, Fredrik Hallgren¹¹⁴, Yavor Boyadzhiev¹¹⁵, Kamen Boyadzhiev¹¹⁵, Mario Küßner³⁴, Duncan Sayer¹¹⁶, Päivi Onkamo^{117,118}, Robin Skeates³⁶, Manuel Rojo-Guerra²⁹, Alexandra Buzhilova²³, Elmira Khussainova¹⁵, Leyla B. Djansugurova¹⁵, Arman Z. Beisenov¹¹⁹, Zainolla Samashev^{120,121}, Ken Massy¹²², Marcello Mannino^{123,124}, Vyacheslav Moiseyev²², Kristiina Mannermaa¹²⁵, Oleg Balanovsky^{126,127,128}, Marie-France Deguilloux¹¹, Sabine Reinhold¹²⁹, Svend Hansen¹²⁹, Egor P. Kitov^{25,119}, Miroslav Dobeš⁴¹, Michal Ernée⁴¹, Harald Meller²⁰, Kurt W. Alt^{130,131,132}, Kay Prüfer^{2,3}, Christina Warinner^{2,3,133}, Stephan Schiffels^{2,3}, Philipp W. Stockhammer^{2,3,122}, Kirsten Bos^{2,3}, Cosimo Posth^{2,5}, Alexander Herbig^{2,3}, Wolfgang Haak^{2,3,134}, Johannes Krause^{2,3*}, Denise Kühnert^{1,2,3,135*}

Hepatitis B virus (HBV) has been infecting humans for millennia and remains a global health problem, but its past diversity and dispersal routes are largely unknown. We generated HBV genomic data from 137 Eurasians and Native Americans dated between ~10,500 and ~400 years ago. We date the most recent common ancestor of all HBV lineages to between ~20,000 and 12,000 years ago, with the virus present in European and South American hunter-gatherers during the early Holocene. After the European Neolithic transition, Mesolithic HBV strains were replaced by a lineage likely disseminated by early farmers that prevailed throughout western Eurasia for ~4000 years, declining around the end of the 2nd millennium BCE. The only remnant of this prehistoric HBV diversity is the rare genotype G, which appears to have reemerged during the HIV pandemic.

The World Health Organization (WHO) estimates that in 2015, 257 million people were living with chronic hepatitis B virus (HBV) infection, which causes close to 1 million deaths each year (1). HBV is transmitted through contact with bodily fluids,

mainly in sexual and perinatal contexts (2), and has no known environmental or animal reservoir. Its spread is therefore tightly linked to the dispersal of humans, whose past population dynamics and migrations have likely shaped the genetic diversity of this partially

double-stranded DNA virus, which is currently classified into nine genotypes associated with characteristic ethno-geographic ranges (Fig. 1) (3, 4). However, the temporal and geographic context of HBV origins in humans, as well as its major routes of dissemination in the past, remain widely debated (5–10). Recent studies have retrieved HBV DNA from archaeological human remains (11–16), providing new avenues to address questions about HBV evolution and phylogeographic history. In particular, these studies revealed the presence of HBV in Europe as early as the Neolithic and ancient HBV lineages that are now seemingly extinct. Ancient DNA data permits molecular clock calibration, and the time to the most recent common ancestor (tMRCA) of all known HBV lineages has been dated to between ~21 thousand years ago (ka) and ~9 ka (14). However, the extent of the past diversity of this virus remains generally unknown because only 19 ancient HBV genomes with a limited temporal and geographic distribution have been reconstructed to date.

The MRCA of all known HBV lineages

Here, we report genomic evidence of HBV in the skeletal remains of 137 individuals from Eurasia and the Americas dated to between ~10,500 and ~400 years ago (Fig. 1, fig. S1 and data S1). Despite advances in molecular virology and numerous sequences from present-day HBV genomes, assessing the phylogenetic relationships among HBV genotypes has proven challenging (7, 17–20), and doubts have been cast about its evolutionary rate and molecular clock-like behavior (9, 16, 21). Nevertheless, most HBV phylogenetic reconstructions have recovered a topology in which HBV genotypes typically found in Native Americans (genotypes F and H) represent a sister clade to the rest of worldwide HBV diversity (which we refer to as the Eurasian branch) (18). This topology was supported by a study incorporating 12 ancient HBV genomes (14) and was also reconstructed in this work (Fig. 2 and figs. S2 and S3). In particular, the monophyly of the American HBV branch, comprising all ancient genomes from the Americas dating back to as early as ~9 ka from the Cuncacha rock shelter in the Andean highlands (CUN002), was highly supported. However, deep nodes within the Eurasian branch were not well resolved, pointing to plausible alternative topologies in which some of the earliest Eurasian lineages would have diverged before the American branch (figs. S4 and S5) (22). Our results confirm that HBV genomic data do exhibit a clear temporal structure when incorporating samples spanning several thousand years (fig. S3). Using the best-fitting uncorrelated relaxed clock model, we estimate the tMRCA of HBV, corresponding to the divergence of American and Eurasian HBV branches, to be between ~16 and ~12 ka [95%

highest posterior density (HPD)] (table S1), which is within the range of previous findings (14). This suggests that contacts between ancestral Eurasians and First Americans occurred until at least shortly before the Bølling-Allerød interstadial (~15 to 13 ka), a period of warming corresponding to widespread human expansion in North America (23, 24). However, studies of ancient human genomes indicate that the ancestors of the First Americans likely began diverging from their closest Eurasian relatives between ~25 and 18 ka, possibly re-

flecting an extended isolation in a Beringian refugium during the Last Glacial Maximum, before dispersing into and across the Americas (25–27). The use of a time-dependent rate (TDR) model yielded an estimate of ~20 to 17 ka for the HBV tMRCA (95% HPD), which was more consistent in this regard. This suggests that not accounting for the time dependency of the evolutionary rate may have led to an underestimation of deep divergence times. However, model selection favored the use of a relaxed clock over a TDR model (log BF: 405)

(22). Taken together, these results point to a scenario in which the MRCA of all HBV strains examined to date existed around the end of the Pleistocene and gave rise to one or several lineages that spread across Eurasia and eventually reached Africa and Oceania, and to another lineage that spread into the Americas with early settlers of this continent.

Our findings challenge the view that current HBV diversity reflects early human dispersals out of Africa. This model is supported in particular by the exclusive association of HBV

¹Transmission, Infection, Diversification and Evolution Group, Max Planck Institute for the Science of Human History, 07745 Jena, Germany. ²Department of Archaeogenetics, Max Planck Institute for the Science of Human History, 07745 Jena, Germany. ³Department of Archaeogenetics, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany. ⁴Max Planck Institute for Infection Biology, 10117 Berlin, Germany. ⁵Archaeo- and Palaeogenetics group, Institute for Archaeological Sciences, Eberhard Karls University Tübingen, 72070 Tübingen, Germany. ⁶ARC Centre of Excellence for Mathematical and Statistical Frontiers, School of Mathematical Sciences, University of Adelaide, Adelaide, SA 5005, Australia. ⁷Division of Hepatology, Department of Medicine II, Leipzig University Medical Center, Leipzig, Germany. ⁸Department of Genetics, Harvard Medical School, Boston, MA, USA. ⁹Department of Human Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA. ¹⁰Institute of Evolutionary Biology, CSIC-Universitat Pompeu Fabra, Barcelona, Spain. ¹¹Université de Bordeaux, CNRS, PACEA UMR 5199, Pessac, France. ¹²Institute of Evolutionary Medicine (IEM), University of Zürich, 8057 Zürich, Switzerland. ¹³Laboratory of Structural Biology, Kazan Federal University, Kazan, Russia. ¹⁴Al-Farabi Kazakh National University, Almaty, Kazakhstan. ¹⁵Institute of Genetics and Physiology, 050060 Almaty, Kazakhstan. ¹⁶Departamento de Genética e Biologia Evolutiva, Instituto de Biociências, Universidade de São Paulo, São Paulo, SP, Brazil. ¹⁷Arizona State University School of Human Evolution and Social Change, Tempe Arizona, USA. ¹⁸Estonian Biocentre, Institute of Genomics, University of Tartu, Tartu, Estonia. ¹⁹Department of Forensic Medicine, University of Helsinki, Helsinki, Finland. ²⁰State Office for Heritage Management and Archaeology Saxony-Anhalt and State Museum of Prehistory, D-06114 Halle, Germany. ²¹Museo Archeologico Regionale “Antonino Salinas”, 90133 Palermo, Italy. ²²Peter the Great Museum of Anthropology and Ethnography (Kunstkamera) RAS, 199034 St. Petersburg, Russia. ²³Anuchin Research Institute and Museum of Anthropology, Lomonosov Moscow State University, Moscow, Russia. ²⁴Sterlitamak Museum of Local History, Sterlitamak, Russia. ²⁵Institute of Ethnology and Anthropology, Russian Academy of Sciences, Moscow, Russia. ²⁶Samara State University of Social Sciences and Education, Samara, Russia. ²⁷Institute of Archaeology named after A. Kh. Khalikov, Tatarstan Academy of Sciences, Kazan, Russia. ²⁸Samara Museum for Historical and Regional Studies named after P. V. Alabin, Samara, Russia. ²⁹Department of Prehistory and Archaeology, Faculty of Philosophy and Letters, University of Valladolid, Spain. ³⁰Department of Prehistory and Archaeology, National Distance Education University (UNED), Madrid, Spain. ³¹Private Technical Archaeologist, 50196 La Muela, Spain. ³²Department of Prehistory and Archaeology, Faculty of Philosophy and Letters, Autonomous University of Madrid, Spain. ³³Technical Archaeologist, Government of Aragón, Spain. ³⁴Thuringian State Office for Heritage Management and Archaeology, 99423 Weimar, Germany. ³⁵University Medical School Göttingen, Institute of Anatomy and Cell Biology, 37075 Göttingen, Germany. ³⁶Department of Archaeology, Durham University, South Road, Durham, DH1 3LE, UK. ³⁷INRAP, ARAR UMR 5138, Maison de l’Orient et de la Méditerranée, Lyon, France. ³⁸State Historical Museum, Moscow, Russia. ³⁹Nasledie Cultural Heritage Unit, 355006 Stavropol, Russia. ⁴⁰Department of Anthropology, The National Museum, Prague, Czech Republic. ⁴¹Institute of Archaeology of the Czech Academy of Sciences, Prague, Czech Republic. ⁴²The City of Prague Museum, Prague, Czech Republic. ⁴³Oxford Archaeology South, Janus House, Osney Mead, Oxford, OX2 0ES, UK. ⁴⁴Oxford Archaeology East, Bar Hill, Cambridge, CB23 8SQ, UK. ⁴⁵Forensic Medicine Unit, Finnish Institute of Health and Welfare, Helsinki, Finland. ⁴⁶Department of Anthropology, University of Winnipeg, Winnipeg, MB, Canada. ⁴⁷Grupo de Investigación y Desarrollo, Dirección Provincial de Cultura, Matanzas, Cuba. ⁴⁸Universidad de Matanzas, Matanzas, Cuba. ⁴⁹Molecular Genetics Laboratory, Escuela Nacional de Antropología e Historia (ENAH), Mexico City, Mexico. ⁵⁰Immunogenetics Unit, Técnicas Genéticas Aplicadas a la Clínica (TGAC), Mexico City, Mexico. ⁵¹Department of Archaeology, Classics and Egyptology, University of Liverpool, Liverpool L69 7WZ, UK. ⁵²Université Côte d’Azur, CNRS, CEPAM UMR 7264, Nice, France. ⁵³University of Strasbourg, CNRS, Archimède UMR 7044, Strasbourg, France. ⁵⁴Department of Archaeology, Faculty of History, Lomonosov Moscow State University, 119192 Moscow, Russia. ⁵⁵Institute of Archaeology, Russian Academy of Sciences, Moscow 117292, Russia. ⁵⁶Department of Anthropology, University of South Florida, Tampa, FL, USA. ⁵⁷Department of Anthropology, University of North Carolina at Charlotte, Charlotte, NC, USA. ⁵⁸Istituto di Ricerca Genetica e Biomedica–CNR, Monserrato, Italy. ⁵⁹Private contractor, Cagliari, Sardinia, Italy. ⁶⁰Koç University, Research Center for Anatolian Civilizations, Istanbul 34433, Turkey. ⁶¹Institute for Research in Archaeology and Historical Heritage (INAPH), University of Alicante, 03690, Alicante, Spain. ⁶²Departamento de Biotecnología, Facultad de Ciencias, Universidad de Alicante, 03690, Alicante, Spain. ⁶³Bioarchaeology Service, Museum of Civilizations, Rome, Italy. ⁶⁴Dipartimento Asia Africa e Mediterraneo, Università di Napoli L’Orientale, Napoli, Italy. ⁶⁵Sedgeford Historical and Archaeological Research Project, Old Village Hall, Sedgeford, Hunstant PE36 5LS, UK. ⁶⁶Melbourne Dental School, University of Melbourne, Victoria 3010 Australia. ⁶⁷Cranfield Forensic Institute, Cranfield Defence and Security, Cranfield University, College Road, Cranfield, MK43 0AL, UK. ⁶⁸Department of Veterinary Medicine, University of Cambridge, Cambridge CB3 0ES, UK. ⁶⁹Astrakhan State University, Astrakhan, Russia. ⁷⁰SNSB, State Collection for Anthropology and Palaeoanatomy, 80333 Munich, Germany. ⁷¹Department of Anthropology, Yale University, New Haven, CT 06511, USA. ⁷²National Museum of Antiquities, 2301 EC Leiden, Netherlands. ⁷³Institute of History, Archaeology and Ethnography, Dushanbe, Tajikistan. ⁷⁴Freie Universität Berlin, Einstein Center Chronoi, 14195 Berlin, Germany. ⁷⁵Cronos S.C. Arqueología y Patrimonio, Burgos, Spain. ⁷⁶Regional Historical Museum Varna, Varna, BG-9000, Bulgaria. ⁷⁷Institute for Caucasus Archaeology, 361401 Nalchik, Republic Kabardino-Balkaria, Russia. ⁷⁸Azov History, Archaeology and Palaeontology Museum-Reserve, Azov 346780, Russia. ⁷⁹Museo de Sitio Pucllana, s/n Miraflores, 18 Lima, Peru. ⁸⁰Australian Centre for Ancient DNA, School of Biological Sciences and The Environment Institute, Adelaide University, Adelaide, SA 5005, Australia. ⁸¹Centre of Excellence for Australian Biodiversity and Heritage (CABA), University of Adelaide, Adelaide, SA 5005, Australia. ⁸²National Centre for Indigenous Genomics, Australian National University, Canberra, ACT 0200, Australia. ⁸³University Medical School Göttingen, Institute of Anatomy and Embryology, 37075 Göttingen, Germany. ⁸⁴Institute of Biology, University of Hildesheim, Germany. ⁸⁵Institute for Prehistory, Early History and Medieval Archaeology, University of Tübingen, 72070 Tübingen, Germany. ⁸⁶State Office for Cultural Heritage Baden-Württemberg, 78467 Konstanz, Germany. ⁸⁷ArchaeoZoology in Siberia and Central Asia–ZooSCAN, CNRS–IAET SB RAS International Research Laboratory, IRL 2013, Novosibirsk, Russia. ⁸⁸Department of Human Genetics, Leiden University Medical Center, Leiden, 2333 ZC, Netherlands. ⁸⁹Royal Belgian Institute of Natural Sciences, Brussels, Belgium. ⁹⁰Center for Archaeological Sciences, University of Leuven, Belgium. ⁹¹Dienst Archeologie–Stad Mechelen, Belgium. ⁹²UCSC Paleogenomics Laboratory, Department of Anthropology, University of California at Santa Cruz, Santa Cruz, CA 95064, USA. ⁹³UCSC Genomics Institute, University of California at Santa Cruz, Santa Cruz, CA 95064, USA. ⁹⁴Department of Anthropology, Vanderbilt University, Nashville, TN 37235, USA. ⁹⁵Department of Physical Anthropology, Institute of Forensic Medicine, University of Bern, Bern, Switzerland. ⁹⁶Applied Archaeology School of Science, Institute of Technology Sligo, Ireland. ⁹⁷School of Historical and Philosophical Studies, University of Melbourne, Victoria 3010, Australia. ⁹⁸Institute of Prehistoric and Historical Archaeology, University of Vienna, Austria. ⁹⁹Department of Archaeology, Hatay Mustafa Kemal University, Alahan-Antakya, Hatay 31060, Turkey. ¹⁰⁰Institute for the Study of the Ancient World (ISAW), New York University, New York, NY 10028, USA. ¹⁰¹Territorial Service of Culture and Tourism from Valladolid, Castilla y León Regional Government, C/ San Lorenzo, 5, 47001, Valladolid, Spain. ¹⁰²Department of History, Geography and Communication, University of Burgos, Paseo de Comendadores, s/n 09001 Burgos (Burgos), Spain. ¹⁰³Dipartimento di Scienze Biomediche, Università di Sassari, Sassari, Italy. ¹⁰⁴Área de Prehistoria, P3A DGA Research Group, IPH, University of Zaragoza, C/ Pedro Cerbuna 12, 50009, Zaragoza, Spain. ¹⁰⁵Department of Anthropology, Michigan State University, East Lansing, MI 48824, USA. ¹⁰⁶Osteology Laboratory, Post Graduate Studies Division, Escuela Nacional de Antropología e Historia (ENAH), Mexico City, Mexico. ¹⁰⁷Caribbean Research Institute, University of Winnipeg, Winnipeg, MB, Canada. ¹⁰⁸DFG Center for Advanced Studies “Words, Bones, Genes, Tools,” University of Tübingen, Tübingen, Germany. ¹⁰⁹South Tyrol Provincial Heritage Service, South Tyrol, Italy. ¹¹⁰Grupo de Investigación en Prehistoria IT-1223-19 (UPV-EHU)/KERBASQUE-Basque Foundation for Science, Vitoria, Spain. ¹¹¹Departament de Prehistòria, Arqueologia i Història Antiga, Universitat de València, València, Spain. ¹¹²Department of Geological Sciences, University of Cape Town, Cape Town, South Africa. ¹¹³Human_G Laboratory, Department of Anthropology, Hacettepe University, Ankara 06800, Turkey. ¹¹⁴The Cultural Heritage Foundation, 722 12 Västerås, Sweden. ¹¹⁵National Archaeological Institute with Museum at the Bulgarian Academy of Sciences, Sofia 1000, Bulgaria. ¹¹⁶School of Natural Sciences, University of Central Lancashire, Preston, UK. ¹¹⁷Department of Biosciences, University of Helsinki, 00014 Helsinki, Finland. ¹¹⁸Department of Biology, University of Turku, 20500 Turku, Finland. ¹¹⁹Institute of archaeology named after A. Kh. Margulan, 44 Almaty, Kazakhstan. ¹²⁰Branch of Institute of Archaeology named after A.Kh. Margulan, 24 of 511 Nur-Sultan, Kazakhstan. ¹²¹State Historical and Cultural Museum-Reserve “Berel”, Katon-Karagay district, East Kazakhstan region, Kazakhstan. ¹²²Institut für Vor- und Frühgeschichtliche Archäologie und Provinzialrömische Archäologie, Ludwig-Maximilians-Universität München, 80539 Munich, Germany. ¹²³Department of Archeology and Heritage Studies, Aarhus University, 8270 Højbjerg, Denmark. ¹²⁴Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig, Germany. ¹²⁵Department of Cultures, University of Helsinki, 00014 Helsinki, Finland. ¹²⁶Research Centre for Medical Genetics, Moscow, Russia. ¹²⁷Biobank of North Eurasia, Moscow, Russia. ¹²⁸Vavilov Institute of General Genetics, Moscow, Russia. ¹²⁹Eurasia Department, German Archaeological Institute, Berlin, Germany. ¹³⁰Danube Private University, Center of Natural and Cultural Human History, A - 3500 Krems-Stein, Austria. ¹³¹Integrative Prehistory and Archaeological Science, Spalenring 145, CH-4055 Basel, Switzerland. ¹³²Department of Biomedical Engineering (DBE), Universitätsspital Basel (HFZ), CH-4123 Allschwil, Switzerland. ¹³³Department of Anthropology, Harvard University, Cambridge, MA 02138, USA. ¹³⁴School of Biological Sciences, University of Adelaide, Adelaide, SA, 5005, Australia. ¹³⁵European Virus Bioinformatics Center (EVBC), Jena, Germany.

*Corresponding author. Email: kocher@shh.mpg.de (A.K.); krause@shh.mpg.de (J.K.); kuehnert@shh.mpg.de (D.K.)




Fig. 1. Geographic location, time period, and lineage of ancient HBV genomes. (A and B) Lineages from (A) Eurasia and (B) the Americas. (C) Main distribution of present-day HBV genotypes [adapted from (4, 14)].

subgenotype C4 with the Aboriginal people of Australia, suggesting that this subgenotype may have been carried by the first settlers of Australia at least ~50 ka (5, 20). Instead, in accordance with previous findings (14), our results indicate that all known modern and ancient HBV strains descend from a lineage that began to diversify at a more recent stage of human history and that subgenotype C4 was introduced in the Australian continent after ~4.5 ka (Fig. 2). Nevertheless, the age of the observed MRCA only represents a lower limit for the earliest presence of HBV in humans. Whether the latter has been preceded by long coevolution, a recent spillover from another animal species, or any intermediate scenario remains an open question. Other viruses from the Hepadnaviridae family have been recovered from a wide range of vertebrates, but none of them appear to represent an ancestral zoonotic source for the human HBV (8).

HBV circulated widely in western Eurasia as early as 10 ka

The retrieval of HBV genomes from around 10 ka in different parts of Europe and Anatolia indicates that the virus was widespread in western Eurasia at that time (Fig. 1 and fig. S1). The oldest HBV strains recovered in Europe form two distinct clades (Fig. 2, fig. S2, and table S2): one that was found in three hunter-gatherers (HGs) from northwestern Russia, Belgium, and Doggerland (Mesolithic 1) and another that was found in an HG from western Russia (Mesolithic 2). These two lineages are placed within the Eurasian branch as sister groups to the modern strains found in nonhuman primates (NHPs) from Southeast Asia and Africa, respectively. The position of NHP HBV lineages within human HBV diversity has been observed in most previous phylogenetic reconstructions and is thought to reflect spillover events from

humans to NHPs (7, 22, 28). The HBV genome reconstructed from an early Anatolian farmer forms a separate lineage recovered at a phylogenetic position intermediate to the two European Mesolithic clades. Between ~9 and 7.5 ka, HBV strains found in HGs from Karelia (northwestern Russia), Sweden, Luxembourg, and Sicily all belonged to the Mesolithic 2 clade. Thus, although our data do not allow detailed phylogeographic inference, they suggest that during the early Holocene, HBV strains could spread over large parts of western Eurasia within a few thousand years. This is consistent with evidence of genetic connections between Europe and the Near East that predate the Neolithic transition (29, 30) and with the observed genetic cline from Western to Eastern HGs (31). Our results further highlight that Mesolithic populations likely formed a network through which pathogens could spread.




Fig. 2. HBV phylogeny and genetic profile of infected individuals. (A) Time-calibrated phylogenetic tree of HBV obtained by using a skyline coalescent tree prior and a lognormal relaxed clock. Main clades were collapsed and annotated with their typical present-day geographic location and the number of ancient genomes they contain. Posterior node supports and date estimates (gray bars, indicating 95% HPD) are reported. Additional

time intervals written on deep nodes are 95% HPD estimates obtained with a time-dependent rate model. **(B)** Principal components analysis plot of modern and ancient western Eurasians summarizing the genetic variation of a subset of individuals for which human genetic data was available. Individuals are colored according to the lineage of the HBV strain they carried, as in the tree.

It has been suggested that most human-adapted pathogens emerged after the Neolithic transition in association with sedentary lifestyles, increased contact with domesticated animals, and higher population densities, a phenomenon sometimes referred to as the “first epidemiological transition” (32–34). Our finding of widespread HBV in HG populations indicates that HBV was present before the advent of agriculture and animal husbandry in different parts of the world. Today, HBV rarely causes lethal fulminant hepatitis but rather asymptomatic infections that may evolve into chronic forms, sometimes developing into liver complications and possible liver failure after decades of infection (1, 2). Although it is difficult to extrapolate from present-day medical studies what the clinical impact of a pathogen would have been in the past—given different diets, disease burdens, and life expectancies—the virus has likely exhibited similar pathophysiological features. Consequently, our findings are consistent with

the view that although small HG communities could not sustain highly epidemic “crowd” diseases, they could maintain chronic infectious agents (35, 36).

A replacement of HBV diversity occurred with the Neolithic transition in Europe

Our data show that HBV remained widespread in Europe after the Neolithic transition (8 to 7 ka), with numerous strains recovered from early European farmers (EEF) across the continent (Fig. 3, fig. S1, and data S1). All of these strains belong to a single HBV lineage that does not descend from previously observed Mesolithic strains (Figs. 2 and 3 and fig. S2). We refer to this HBV lineage as the Western-Eurasian Neolithic-to-Bronze Age (WENBA) lineage. This transition is also observable at a microscale in Grotta dell’Uzzo (Sicily), where HBV strains recovered from Neolithic individuals are unrelated to a Late Mesolithic strain identified at the same site (figs. S1 and S2). This suggests that the HBV

strains observed in EEFs were not acquired from local HGs in different areas but were rather disseminated by EEFs themselves. Although EEFs ultimately derived from early agricultural populations in the Near East (37, 38), the strain we retrieved from an Anatolian farmer dated to ~10 ka was not ancestral to the WENBA lineage (Fig. 2). Therefore, even if EEFs were indeed key in disseminating WENBA strains, whether this lineage originated in Near Eastern centers of early agriculture or in another location along EEF’s expansion routes remains to be determined. Furthermore, given the current sample availability for this period, a scenario in which the WENBA lineage would have originated and disseminated among European HGs shortly before the Neolithic transition cannot be completely excluded.

We also found WENBA HBV strains in two HGs from transitional Neolithic contexts in western Russia dated to ~7.2 and ~6.4 ka (JAZ001 and MUR007), as well as on both




Fig. 3. Spatiotemporal distribution of ancient western Eurasian HBV strains. (A) Time-calibrated phylogenetic tree (Eurasian branch). Lineages containing ancient HBV genomes are colored. (B) Histogram showing the number of recovered ancient HBV genomes belonging to each lineage through time. (C) Geographic distribution of ancient HBV genomes within different time-periods, colored by lineage.

sides of the Greater Caucasus Mountain range and in Anatolia as early as ~ 5.6 ka (fig. S1). In general, phylogenetic relationships among HBV sublineages within the WENBA clade do not exhibit a strong geographical structure (fig. S2), nor do they seem to reflect the material culture or genetic profile of the individuals in which they were found (fig. S6). Furthermore, our phylodynamic reconstruction indicates that after an initial growth phase, the transmission of WENBA HBV reached an equilibrium from ~ 7.5 to ~ 3.5 ka (fig. S7). Overall, this suggests that HBV strains disseminated by EEFs quickly spread throughout much of western Eurasia beyond the limits of the

European agricultural expansion, where they became endemic and continued to circulate widely across different populations, for several thousand years. In particular, we do not observe substantial changes in the HBV genetic landscape associated with the expansion of steppe-related ancestry that dramatically altered the genetic profile of Europeans from ~ 5 ka onward (Fig. 2, fig. S2, and data S1) (37). Sexual and perinatal transmission have likely always been the major mechanisms of HBV infection in humans, but cultural practices involving contact with blood [such as tattooing (39)] or nonsexual violent interactions (40) could also have played a role in the spread of

the virus in the past. In general, our findings attest to a degree of interconnectivity among prehistoric populations of different origins, subsistence modes, and cultures that allowed for the dissemination of directly transmitted pathogens.

The collapse of WENBA HBV during the 2nd millennium BCE

After the Early Neolithic (8 to 7 ka), the WENBA HBV lineage prevailed in most parts of western Eurasia for more than 4000 years (Fig. 3). However, the latest occurrence of a WENBA strain in our dataset is dated to ~ 3.3 ka, after which this lineage is no longer observed (figs.

S1 and S2). By contrast, genotype A, which we first observed at the eastern edge of Europe and in the Near East between ~5 and ~3.5 ka, still appears after ~2.5 ka, by which time it had reached the Carpathian Basin in central Europe. Around the same date, we first observed genotype D in two individuals from the Italian Alps, as well as in various locations in the western steppe, before prevailing in large parts of Europe during the Medieval period. Thus, it seems that as most WENBA HBV lineages disappeared by the end of the 2nd millennium BCE, genotypes A and D subsequently spread from eastern reservoirs to eventually reach western regions that had previously only harbored WENBA strains (22).

The second half of the 2nd millennium BCE bears witness to major cultural shifts in the archaeological record in western Eurasia, including the sudden disappearance of tell settlements in the Carpathian Basin (41), the expansion of the Urnfield culture and the increase of military conflicts in large parts of Europe (42–45), the breakdown of the Terramare culture in northern Italy (46), and the so-called Late Bronze Age collapse of most state societies in the eastern Mediterranean region and Near East (47, 48). Some of these societal transformations could have been triggered by underlying phenomena such as climatic events (49) or the spread of epidemic diseases (50) and were likely associated with substantial shifts in population densities, trans-regional networks, and modes and scales of human mobility. The observed decline of WENBA HBV diversity, as well as our phylodynamic reconstruction (fig. S7), further point to major changes in epidemiological dynamics over large parts of western Eurasia during this period. However, although our data suggests that new lineages disseminated across Europe only later on, the lack of observations around 3 ka (Fig. 3) could reflect sampling biases related to the widespread adoption of cremation practices around that time (42–44) rather than a decrease of HBV prevalence. Searching for the virus in a large number of systematically dated samples across this period could help to better characterize the process that ultimately led to the renewal of western Eurasian HBV diversity after the end of the 2nd millennium BCE.

Recent reemergence of the WENBA HBV lineage

The majority of HBV strains circulating in western Eurasia today belong to genotypes A and D (3, 4), thus only reflecting a relatively recent part of the phylogeographic history of this virus. However, our results show that despite the seemingly complete disappearance of WENBA HBV strains around the end of the 2nd millennium BCE, one lineage descending from this clade has persisted to the present.

The latter gave rise to a group of modern strains classified as genotype G (Fig. 2 and fig. S2), a rare, recently described genotype for which the biology is poorly understood (51). First discovered in patients from France and the United States, genotype G was later found in other parts of Europe, the Americas, and in Asia, making its geographic origin unclear (52). Despite its wide distribution, genotype G exhibits remarkably low genetic diversity (53), suggesting a recent reemergence after thousands of years of low-level persistence. Furthermore, genotype G has mostly been found in HIV-positive patients, and phylodynamic patterns have pointed to a sharp increase of its dissemination co-occurring with the HIV pandemic, possibly associated with highly sexually active groups and injection-drug users (52).

Genotype G has sometimes been referred to as “aberrant” because of its distinctive genomic features: a 36-nucleotide insertion near the 5′ end of the core gene and two nonsense mutations in the precore region (51, 54). These changes inhibit production of the immunotolerant e antigen (HBeAg), which appears to be essential for the establishment of a persistent HBV infection, and alter the structure of the HBV core protein, which may impair packaging and replication of the viral genetic material (54, 55). This likely explains why in the vast majority of cases genotype G occurs in co-infections with other HBV genotypes, which can provide the HBeAg and core protein production functions lacking in genotype G (54–56). We identified similar insertions and stop codons in 14 ancient HBV genomes ranging in age between ~7 and 3.5 ka, which form the WENBA subclade from which genotype G descends (fig. S8). Additionally, most of these ancient genomes were found in individuals showing signs of infections with several HBV variants (fig. S8 and data S2) (22). Cases of mixed infection were exclusively found in individuals carrying WENBA HBV strains, among which they were very frequent (22 of 83 individuals, likely underestimating the true frequency). In all cases, both major and minor strains appeared to belong to the WENBA lineage, and sequencing data were partially supporting a ~40–base pair insertion at the 5′ end of the core gene (table S3 and data S1).

Therefore, although genotype G is considered rare today, it seems that the cotransmission of its ancestral form together with another HBeAg⁺ WENBA strain was a common epidemiological feature of HBV between ~7.5 and 3.5 ka. Notably, this functionally limited variant persisted until today, whereas the rest of the WENBA HBV diversity seemingly went extinct. Virologic studies indicate that genotype G tends to outcompete HBeAg-producing strains during late HBV in-

fection stages after anti-HBeAg seroconversion (56–58). Although these short-term selection patterns parallel the survival of this lineage over thousands of years, the latter may rather be related to less deterministic factors. One of the closest Bronze Age ancestors of genotype G was recovered at the archaeological site of Shagara in the eastern European forest zone (SGR003) (figs. S1 and S2), a location where the present-day widespread genotype A was already circulating (SGR004). Genotype A is the most common genotype found in mixed infections with genotype G today (55, 57). The discovery of ancestral forms of both genotypes at the same archaeological site, albeit from different individuals and time periods, may indicate that this viral association had already formed during prehistory in eastern Europe.

Conclusions

This study demonstrates the value of large-scale paleogenomic analyses for studying the phylogeographic history of HBV. DNA enrichment allowed us to reconstruct large proportions of more than 100 ancient HBV genomes from a variety of skeletal tissues, opening possibilities for future paleovirologic studies. We show that HBV was already widely present in humans during the early Holocene and that its phylogeographic history reflects several well-known human migrations and demographic events, including the expansion of First American populations in the Americas and the Neolithic transition in Europe, but not others, such as later Bronze Age steppe ancestry expansions. Furthermore, our results reveal patterns that were not expected on the basis of human genetic and archaeological data alone, such as the near complete renewal of western Eurasian HBV diversity around the end of the 2nd millennium BCE. These findings highlight that the reconstruction of ancient viral diversity has great potential to contribute to our understanding of human history.

REFERENCES AND NOTES

1. World Health Organization (WHO), *Global Hepatitis Report* (WHO, 2017).
2. D. Lavanchy, M. Kane, in *Hepatitis B Virus in Human Diseases*, Y.-F. Liaw, F. Zoulim, Eds. (Springer International Publishing, Cham, 2016), *Molecular and Translational Medicine*, pp. 187–203.
3. A. Kramvis, *Intervirolgy* **57**, 141–150 (2014).
4. S. Velkov, J. J. Ott, U. Protzer, T. Michler, *Genes (Basel)* **9**, 495 (2018).
5. M. Littlejohn, S. Locarnini, L. Yuen, *Cold Spring Harb. Perspect. Med.* **6**, a021360 (2016).
6. S. Locarnini, M. Littlejohn, M. N. Aziz, L. Yuen, *Semin. Cancer Biol.* **23** (6 Pt B), 561–575 (2013).
7. D. Paraskevis et al., *Hepatology* **57**, 908–916 (2013).
8. A. Rasche, A.-L. Sander, V. M. Corman, J. F. Drexler, *J. Hepatol.* **70**, 501–520 (2019).
9. P. Simmonds, *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **356**, 1013–1026 (2001).
10. G. Zehender et al., *World J. Gastroenterol.* **20**, 7622–7634 (2014).

11. G. Kahila Bar-Gal *et al.*, *Hepatology* **56**, 1671–1680 (2012).
12. R. Barquera *et al.*, *Curr. Biol.* **30**, 2078–2091.e11 (2020).
13. B. Krause-Kyora *et al.*, *eLife* **7**, e36666 (2018).
14. B. Mühlemann *et al.*, *Nature* **557**, 418–423 (2018).
15. J. Neukamm *et al.*, *BMC Biol.* **18**, 108 (2020).
16. Z. Patterson Ross *et al.*, *PLOS Pathog.* **14**, e1006750 (2018).
17. P. L. Bollyky, E. C. Holmes, *J. Mol. Evol.* **49**, 130–141 (1999).
18. B. A. Godoy, J. R. R. Pinho, N. J. R. Fagundes, *Virus Res.* **276**, 197776 (2020).
19. D. Paraskevis *et al.*, *Mol. Phylogenet. Evol.* **93**, 44–54 (2015).
20. L. K. W. Yuen *et al.*, *Mol. Biol. Evol.* **36**, 942–954 (2019).
21. R. Bouckaert, M. V. Alvarado-Mora, J. R. Pinho, *Antivir. Ther.* **18** (3 Pt B), 497–503 (2013).
22. Materials and methods are available as supplementary materials.
23. D. Palacios *et al.*, *Earth Sci. Rev.* **203**, 103113 (2020).
24. M. R. Waters, *Science* **365**, eaat5447 (2019).
25. B. Llamas *et al.*, *Sci. Adv.* **2**, e1501385 (2016).
26. J. V. Moreno-Mayar *et al.*, *Nature* **553**, 203–207 (2018).
27. M. Raghavan *et al.*, *Science* **349**, aab3884 (2015).
28. B. F. de Carvalho Dominguez Souza *et al.*, *J. Hepatol.* **68**, 1114–1122 (2018).
29. M. Feldman *et al.*, *Nat. Commun.* **10**, 1218 (2019).
30. Q. Fu *et al.*, *Nature* **534**, 200–205 (2016).
31. I. Mathieson *et al.*, *Nature* **555**, 197–203 (2018).
32. R. Barrett, C. W. Kuzawa, T. McDade, G. J. Armelagos, *Annu. Rev. Anthropol.* **27**, 247–271 (1998).
33. J. Diamond, *Nature* **418**, 700–707 (2002).
34. F. M. Key *et al.*, *Nat. Ecol. Evol.* **4**, 324–333 (2020).
35. M. C. Inhorn, P. J. Brown, *Annu. Rev. Anthropol.* **19**, 89–117 (1990).
36. N. D. Wolfe, C. P. Dunavan, J. Diamond, *Nature* **447**, 279–283 (2007).
37. W. Haak *et al.*, *Nature* **522**, 207–211 (2015).
38. I. Lazaridis *et al.*, *Nature* **513**, 409–413 (2014).
39. A. Deter-Wolf, B. Robitaille, L. Krutak, S. Galliot, *J. Archaeol. Sci. Rep.* **5**, 19–24 (2016).
40. K. W. Alt *et al.*, *Sci. Rep.* **10**, 2131 (2020).
41. F. Gogáltn, *Studia Hercynia* **23**, 198–214 (2019).
42. G. Capuzzo, J. A. Barceló, *World Archaeol.* **47**, 622–641 (2015).
43. F. Falkenstein, in *Ancestral Landscape. Burial mounds in the Copper and Bronze Ages (Central and Eastern Europe–Balkans–Adriatic–Aegean, 4th–2nd millennium B.C.)* Proceedings of the International Conference held in Udine, 15 to 18 May 2008 (Maison de l’Orient et de la Méditerranée Jean Pouilloux, 2012), Travaux de la Maison de l’Orient et de la Méditerranée. Série recherches archéologiques, pp. 329–340.
44. H. Fokkens, *Antiquity* **71**, 360–373 (1997).
45. S. Hansen, in *Hillforts and Weaponry in the Early and Middle Bronze Age*, S. Hansen, R. Krause, Eds. (Habelt, 2019).
46. A. Cardarelli, in *Scienze dell’Antichità* (Edizioni Quasar, 2009), vol. 15, pp. 449–520.
47. E. H. Cline, *1177 B.C.: The Year Civilization Collapsed* (Princeton Univ. Press, 2015).
48. J. Driessen *et al.*, *An Archaeology of Forced Migration. Crisis-Induced Mobility and the Collapse of the 13th c. BCE Eastern Mediterranean* (PLU, 2018).
49. D. Kaniewski, E. V. Campo, in *3.2 ka BP Megadrought and the Late Bronze Age Collapse*, H. Weiss, Ed. (Oxford Univ. Press, 2017).
50. P. Norrie, in *A History of Disease in Ancient Times: More Lethal than War*, P. Norrie, Ed. (Springer International Publishing, 2016), pp. 61–101.
51. L. Stuyver *et al.*, *J. Gen. Virol.* **81**, 67–74 (2000).
52. J. M. Wolf, S. De Carli, V. R. Z. B. Pereira, D. Simon, V. R. Lunge, *J. Viral Hepat.* **28**, 393–399 (2021).
53. M. Cornelissen *et al.*, *BMC Infect. Dis.* **16**, 268 (2016).
54. K. Li *et al.*, *J. Virol.* **81**, 9202–9215 (2007).
55. T. Sakamoto *et al.*, *J. Viral Hepat.* **20**, e27–e36 (2013).
56. Y. Tanaka *et al.*, *Virology* **376**, 408–415 (2008).
57. H. Kato *et al.*, *Hepatology* **35**, 922–929 (2002).
58. M. Sugiyama *et al.*, *Hepatology* **45**, 929–937 (2007).

ACKNOWLEDGMENTS

We thank M. Harbeck, L. Bondioli, R. Risch, and H. Duda for their contribution to the archaeological interpretation; R. Heyne and H. Heyne for comments concerning clinical manifestation of hepatitis B infections; B. Mühlemann and T. C. Jones for helpful discussion on genome assemblies and mixed infections; and M. O’Reilly for her help on the design of the figures. **Funding:** The research was funded by the Max Planck Society, the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (771234–PALEORIDER, to W.H.; 805268–CoDisEASe to K. Bos; 834616–ARCHCAUCASUS to S.H.), the Slovak Academy of Sciences and the European Union’s Seventh Framework Programme and Marie Curie Actions under the Programme SASPRO (1340/03/03 to P.C.R.), the ERA.NET RUS Plus–S&T programm of the European Union’s Seventh Framework Programme (277–BIOARCAUCASUS to S.Re. and S.H.), the Werner Siemens Stiftung (“Paleochemistry”, to CW), the Award Praemium Academiae of the Czech Academy of Sciences (to M.E.), the Institute of Archaeology of the Czech Academy of Sciences (RVO 67985912, to M.Dobe.), the Russian Foundation for Basic Research (19-09-00354a, to M.K.K. and V.V.K.; 19-78-10053 to SSh), the German Research Foundation (DFG-HA-5407/4-1–INTERACT to W.H. and RE2688/2 to S.Re.), the French National Research Agency (ANR-17-FRAL-0010–INTERACT, to M.F.D., M.Ri., S.Ro., S.Sai., D.Bi., and P.Le.), the Wenner-Gren Dissertation Fieldwork Grant (9558 to S.Sab.), and the Ministry of Education and Science

of the Republic of Kazakhstan (AP08856654 to L.B.D., L.M., and E.Kh. and AP08857177 to A.Z.B.). **Author contributions:** D.K. and J.K. initiated and supervised the study. S.F., V.Sc., V.K., M.K.K., M.S.C., V.V.K., A.A.Kh., A.A.C., D.A.S., A.F.K., C.T.R., Í.G.-M.d.L., H.A.M., R.G.P., J.I.R.G., J.N., S.Ro., S.K., S.Sai., E.Ka., A.B.B., P.V., P.Li., M.Ko., L.Lo., E.P., R.C., A.L., R.M., Y.C.d.A., S.T.H.G., D.I.H.Z., J.P., D.Bi., P.Le., A.R.K., V.E.M., L.La., M.Z., J.F.B., M.L., A.D., T.I., G.G.A., M.P.d.M.I., A.R., A.Sp., S.B., S.Sai., E.D.Z., D.V.V., K.v.H., R.L.B., L.C.S., L.A., M.N., E.R., C.A.F., V.Si., A.A.Ka., B.C.A., E.B., M.A.C., M.S., R.K., J.J.E., M.Fr., S.Sh., P.d.K., E.A., K.V.d.V., L.F.S., T.A.T., S.L., M.Dobr., N.M., C.R., M.V.T., C.S., P.C.R., M.A., K.A.Y., E.C.B., F.C., V.Ma., P.U., K.R., D.Ba., P.S., L.M.M., M.Ro., H.S., D.C.S.G., N.S., Y.S.E., F.H., Y.B., K.Boy., M.Kü., D.S., P.O., R.Sk., M.R.G., A.B., L.B.D., A.Z.B., Z.S., K.Mas., M.Ma., V.Mo., K.Man., S.Re., S.H., E.P.K., M.Dobe., M.E., H.M., and K.W.A. conducted the archaeological/anthropological work or the curation of the archeological material. J.K., W.H., C.Po., K.Bos., P.W.S., S.Sc., C.W., M.F.D., O.B., E.Kh., E.F.D., B.L., A.Sa., E.B., E.K.G., L.M., and M.Ri. organized the sample collection for genetic analyses. R.B., L.P., M.A.S., F.A., R.St., A.W., A.M., V.V.M., G.U.N., M.Ri., M.S.v.d.L., K.Maj., R.I.T., L.M., A.G., S.P., S.Sab., M.Mi., J.G., E.A.N., T.F., K.N., C.Pa., M.Ke., E.K.G., M.Fe., S.E., E.S., K.G., G.A.G.R., D.I.H.Z., B.L., E.F.D., O.B., M.F.D., and W.H. participated in the laboratory work, sample management, and gathering of contextual information. A.K., F.M.K., M.A.S., and R.H. performed HBV screening of the sequencing data, which was supervised by A.H.; A.K. and K.P. performed the bioinformatic analyses. A.K. and D.K. performed the phylogenetic analyses. L.P. performed the human population genetics analyses. A.K. and A.B.R. performed the statistical analyses. All authors contributed to the interpretation of the data. A.K. wrote the first version of the manuscript, which was edited by D.K., J.K., W.H., A.H., C.Po., K.Bos., P.W.S., S.Sc., C.W., F.v.B., M.P., L.P., R.B., F.M.K., and M.A.S., and all authors contributed to its improvement. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** The sequencing data generated in this study has been deposited at the European Nucleotide Archive (www.ebi.ac.uk/ena/data/view/PRJEB45699). Complete HBV sequence alignments are available in the supplementary materials.

SUPPLEMENTARY MATERIALS

[science.org/doi/10.1126/science.abi5658](https://doi.org/10.1126/science.abi5658)
Materials and Methods
Supplementary Text
Figs. S1 to S10
Tables S1 to S5
References (59–254)
Data S1 to S4
MDAR Reproducibility Checklist

[View/request a protocol for this paper from Bio-protocol.](https://www.bio-protocol.org)

18 March 2021; accepted 17 August 2021
10.1126/science.abi5658

Ten millennia of hepatitis B virus evolution

Arthur KocherLuka PapacRodrigo BarqueraFelix M. KeyMaria A. SpyrouRon HüblerAdam B. RohrlachFranziska AronRaphaela StahlAntje WissgottFlorian van BömmelMaria PfefferkornAlissa MitnikVanessa Villalba-MoucoGunnar U. NeumannMaité RivollatMarieke S. van de LoosdrechtKerttu MajanderRezeda I. TukhbatovaLyazzat MusralinaAyshin GhalichiSandra PenskeSusanna SabinMegan MichelJoscha GretzingerElizabeth A. NelsonTiago FerrazKathrin NägeleCody ParkerMarcel KellerEvelyn K. GuevaraMichal FeldmanStefanie EisenmannEirini SkourtaniotiKaren GiffinGuido Alberto Gneccchi-RuscioneSusanne FriederichVittoria SchimmentiValery KhartanovichMarina K. KarapetianMikhail S. ChaplyginVladimir V. KufferinAleksandr A. KhokhlovAndrey A. ChizhevskyDmitry A. StashenkovAnna F. KochkinaCristina Tejedor-RodríguezÍñigo García-Martínez de LagránHéctor Arcusa-MagallónRafael Garrido-PenaJosé Ignacio Royo-GuillénJan Nová#ekStéphane RottierSacha KackiSylvie SaintotElena KaverznevaAndrej B. BelinskiyPetr VelemínskýPetr LimburskýMichal KostkaLouise LoeElizabeth PopescuRachel ClarkeAlice LyonsRichard MortimerAntti SajantilaYadira Chinique de ArmasSilvia Teresita Hernandez GodoyDiana I. Hernández-ZaragozaJessica PearsonDidier BinderPhilippe LefrancAnatoly R. KantorovichVladimir E. MaslovLuca LaiMagdalena ZoledziowskaJessica F. BeckettMichaela LangováAlžběta DanielisováTara IngmanGabriel García AtiénzarMaria Paz de Miguel IbáñezAlejandro RomeroAlessandra SperdutiSophie BeckettSusannah J. SalterEmma D. ZilivinskayaDmitry V. Vasil'evKristin von HeykingRichard L. BurgerLucy C. SalazarLuc AmkreutzMasnav NavruzbekovEva RosenstockCarmen Alonso-FernándezVladimir SlavchevAlexey A. KalmykovBiaslan Ch. AtabievElena BatievaMicaela Alvarez CalmetBastien LlamasMichael SchultzRaiko KraußJavier Jiménez-EchevarríaMichael FranckenSvetlana ShnaiderPeter de KnijffEveline AltenaKatrien Van de VijverLars Fehren-SchmitzTiffany A. TungSandra LöschMaria DobrovolskayaNikolaj MakarovChris ReadMelanie Van TwestClaudia SagonaPeter C. RamsMurat AkarK. Aslihan YenerEduardo Carmona BallesterFrancesco CuccaVittorio MazzarelloPilar UtrillaKurt RademakerEva Fernández-DomínguezDouglas BairdPatrick SemalLourdes Márquez-MorfinMirjana RoksandicHubert SteinerDomingo Carlos Salazar-GarcíaNatalia ShishlinaYilmaz Selim ErdalFredrik HallgrenYavor BoyadzhievKamen BoyadzhievMario KüßnerDuncan SayerPäivi OnkamoRobin SkeatesManuel Rojo-GuerraAlexandra BuzhilovaElmira KhussainovaLeyla B. DjansugurovaArman Z. BeisenovZainolla SamashevKen MassyMarcello ManninoVyacheslav MoiseyevKristiina MannermaaOleg BalanovskyMarie-France DeguillouxSabine ReinholdSvend HansenEgor P. KitovMiroslav DobešMichal ErnéeHarald MellerKurt W. AltKay PrüferChristina WarinnerStephan SchiffelsPhilipp W. StockhammerKirsten BosCosimo PosthAlexander HerbigWolfgang HaakJohannes KrauseDenise Kühnert

Science, 374 (6564),

Ancient DNA traces the history of hepatitis B

Hepatitis B virus (HBV) infections represent a worldwide human health concern. To study the history of this pathogen, Kocher *et al.* identified 137 human remains with detectable levels of virus dating between 400 and 10,000 years ago. Sequencing and analyses of these ancient viruses suggested a common ancestor between 12,000 and 20,000 years ago. There is no evidence indicating that HBV was present in the earliest humans as they spread out of Africa; however, HBV was likely present in human populations before farming. Furthermore, the virus was present in the Americas by about 9000 years ago, representing a lineage sister to the viral strains found in Eurasia that diverged about 20,000 years ago. —LMZ

View the article online

<https://www.science.org/doi/10.1126/science.abi5658>

Permissions

<https://www.science.org/help/reprints-and-permissions>

Use of think article is subject to the [Terms of service](#)

Science (ISSN) is published by the American Association for the Advancement of Science. 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2021 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works