

Detection of Red-spotted grouper nervous necrosis virus (RGNNV) in shrimp and squid of the Mediterranean Sea

Carolina Johnstone^{1*}, Montse Pérez², Marta Arizcun³, Cristina García-Ruiz¹, Cristina Ciércoles¹, Elena Chaves-Pozo^{3*}.

¹Oceanographic Centre of Malaga, Instituto Español de Oceanografía – Consejo Superior de Investigaciones Científicas (IEO-CSIC), Fuengirola, Málaga, Spain



and Shellfish

²Oceanographic Centre of Vigo, IEO-CSIC, Vigo, Spain.

³Oceanographic Centre of Murcia, IEO-CSIC, Mazarrón, Murcia, Spain. *elena.chaves@ieo.es

Nervous necrosis virus (NNV) is a betanodavirus that causes viral encephalopathy and retinopathy (VER) or viral nervous necrosis (VNN) in fish. Genotypes are discriminated according to genomic RNA2 encoding the coat protein. The RGNNV and its reassortant with SJNNV genotype currently threat Mediterranean aquaculture. The virus can be transmitted through water, between fish, or through contaminated food including trash shellfish.



HOST SPECIES: in the Alboran Sea during the ∕ Fish Mediterranean International Trawl A Shellfish Survey (MEDITS) in 2015. Total COLLECTION AREA: RNA was isolated from frozen ocular CADIZ tissue. NNV was detected through ALBORAN sequencing using primers that SICILY discriminate RNA2 of RGNNV and ADRIATIC SJNNV (1.2). RGNNV was detected AEGEAN LEVANTINE in all three species analyzed and in eight of nine locations studied.

Invertebrate animals were collected

NNV is detected for the first time in the Alboran Sea.

In the Mediterranean Sea. **RGNNV** is widely distributed between areas and to date has been detected in four Classes and seven Families of marine invertebrates. But this study detected, for the first time, the RGNNV in squid and shrimp.

Clustering of RGNNV RNA2 sequences isolated from Mediterranean fish and shellfish suggest that RGNNV is being widely transmitted between shellfish, fish, and ecosystems.

ACKNOWLEDGEMENTS: This study was financed by the NODAMED project (IEO), 🕯 COMPANY PARTIES

European Commission

and co-funded by the European Maritime and Fisheries Fund (EMFF)



Detection of RGNNV in shellfish collected in the Mediterranean Sea.

	Classification ^a	Species	Collection Area	Tissue ^b	Sequence ^c	Reference
	Mollusca Bivalvia					
	Mytiloida Mytilidae	Mytilus galloprovincialis	Greece, Aegean Sea.	n.i.	MK766986	(3)
		Xenostrobus securis	Sicily, Italy.	Hepatopancreas	MG195159	(4)
			Italy, Adriatic Sea.	Hepatopancreas	MT759757-	(5)
					MT759760	
0	Veneroida, Veneridae	Ruditapes philipinarum	Italy, Adriatic Sea.	Hepatopancreas	MG195167-	(4)
					MG195186	
0	Mollusca Gastropoda					
	Neogastropoda Muricidae	Stramonita haemastoma	Greece, Aegean Sea.	n.i.	MK766973	(3)
	Mollusca Cephalopoda					
	Myopsida Loliginidae	Alloteuthis media	Spain, Alboran Sea.	Ocular tissue	MZ546624-	This study
					MZ546628	
N	Oegopsida	Abralia veranyi	Spain, Alboran Sea.	Ocular tissue	MZ546629	This study
	Enoploteuthidae					
	Arthropoda Malacostraca					
3	Decapoda Pandalidae	Plesionika heterocarpus	Spain, Alboran Sea.	Ocular tissue	MZ546630-	This study
					$MZ546642^d$	
	Decapoda Portunidae	Portunus pelagicus	Israel, Levantine	Nerve tissue	MH663494,	(6)
-A	5		Sea.		MH663495,	
`					MH970545,	
					MH970544	

REFERENCES: (1) López-Jimena, et al., 2010. J. Appl. Microbiol. 109: 1361. (2) Moreno, et al., 2014. Appl. Environ. Microbiol. 80:6560. (3) Bitchava et al., 2019. J. Fish Dis. 42:1773. (4) Volpe et al., 2018. J. Fish Dis. 41:603. (5) Errani et al., 2020. J. Appl. Microbiol. 130:994. (6) Berzak et al., 2019. Dis. Aquat. Organ. 133:7.