

Ammonium regulates the development of pine roots through hormonal crosstalk and differential expression of transcription factors in the apex

Francisco Ortigosa¹  | César Lobato-Fernández¹  | Hitomi Shikano²  |
 Concepción Ávila¹  | Shu Taira²  | Francisco M. Cánovas¹  | Rafael A. Cañas^{1,3} 

¹Grupo de Biología Molecular y Biotecnología, Departamento de Biología Molecular y Bioquímica, Universidad de Málaga, Campus Universitario de Teatinos, Málaga, Spain

²Faculty of Food and Agricultural Sciences, Fukushima University, Kanayagawa, Fukushima, Japan

³Integrative Molecular Biology Lab, Departamento de Biología Molecular y Bioquímica, Universidad de Málaga, Campus Universitario de Teatinos, Málaga, Spain

Correspondence

Rafael A. Cañas, Departamento de Biología Molecular y Bioquímica, Facultad de Ciencias, Universidad de Málaga, Campus Universitario de Teatinos s/n, E-29071 Málaga, Spain.
 Email: rcanas@uma.es

Funding information

CONSORCIO DE BIBLIOTECAS UNIVERSITARIAS DE ANDALUCÍA (CUBA) Funding for open access charge; Ministerio de Ciencia e Innovación, Grant/Award Numbers: BIO2015-73512-JIN MINECO/AEI/FEDER, RTI2018-094041-B-I00, UE, EQC2018-004346-P; Junta de Andalucía, Grant/Award Number: BIO-114; Universidad de Málaga Funding for open access charge, Grant/Award Number: UMAJ11, FEDER, FSE, Junta de Andalucía

Abstract

Ammonium is a prominent source of inorganic nitrogen for plant nutrition, but excessive amounts can be toxic for many species. However, most conifers are tolerant to ammonium, a relevant physiological feature of this ancient evolutionary lineage. For a better understanding of the molecular basis of this trait, ammonium-induced changes in the transcriptome of maritime pine (*Pinus pinaster* Ait.) root apex have been determined by laser capture microdissection and RNA sequencing. Ammonium promoted changes in the transcriptional profiles of multiple transcription factors, such as *SHORT-ROOT*, and phytohormone-related transcripts, such as *ACO*, involved in the development of the root meristem. Nano-PALDI-MSI and transcriptomic analyses showed that the distributions of IAA and CKs were altered in the root apex in response to ammonium nutrition. Taken together, the data suggest that this early response is involved in the increased lateral root branching and principal root growth, which characterize the long-term response to ammonium supply in pine. All these results suggest that ammonium induces changes in the root system architecture through the IAA-CK-ET phytohormone crosstalk and transcriptional regulation.

KEYWORDS

conifer, laser capture microdissection, nitrogen nutrition, phytohormones, *Pinus pinaster*, root development, Transcriptomics

1 | INTRODUCTION

Nitrogen (N) is a vitally important nutrient for all living organisms because it is a constituent of essential biomolecules such as nucleic acids, proteins, amino acids, porphyrins and hormones, among others (Miller & Cramer, 2005). This nutrient is indispensable for the proper

growth and development of plants, and it can be assimilated from different kinds of sources, including organic (peptides, amino acids and urea) and inorganic (nitrate and ammonium) forms (Hachiya & Sakakibara, 2017; Näsholm et al., 1998). Together with nitrate (NO_3^-), ammonium (NH_4^+) is one of the main forms of inorganic N available for plants, and the relative proportions of these elements

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Plant, Cell & Environment* published by John Wiley & Sons Ltd.

in soil can vary depending on biological and climate conditions (Bijlsma et al., 2000).

Experimental evidence strongly suggest that NH_4^+ is perceived and recognized by plant cells as a signal that promotes physiological and morphological changes in plants (Liu & von Wirén, 2017). However, NH_4^+ at millimolar levels usually causes toxicity in most plants. The effects caused by an excessive NH_4^+ availability include decreased plant growth, leaf chlorosis, decreased root/shoot ratios, decreased root gravitropism and altered root system architecture (RSA) (Esteban et al., 2016). Regarding the changes in RSA, the inhibition of root elongation, enhancement of lateral root (LR) branching and impaired root gravitropism are commonly observed (Y. Liu & von Wirén, 2017).

Root elongation involves three interconnected biological processes: cell division, cell expansion and cell differentiation (Youssef et al., 2018). Both cell division and cell expansion are affected by NH_4^+ through two independent determinants: (i) a decreased capacity for protein N-glycosylation and (ii) an increased production of reactive oxygen species (ROS) (Jia & von Wirén, 2020). In *Arabidopsis*, root growth inhibition in response to NH_4^+ represses root cell production by decreasing the meristem size and the number of dividing cells without altering the cell division rate (Y. Liu et al., 2013). Furthermore, NH_4^+ also decreases the number of root cap cells (Y. Liu et al., 2013). In this sense, several transcription factors (TFs) have been described to play key roles in root cap development, such as BEARSKIN (BRN) and SOMBRERO (SMB) (Bennett et al., 2010, 2014; Kamiya et al., 2016). However, the transcriptional pathway regulating NH_4^+ root cap development has not yet been deciphered.

In addition, auxins (IAAs) play a prominent role in cell division and cell expansion (Wang & Ruan, 2013). IAAs can be synthesized in the root meristem and transported toward upstream adjacent regions through specific transporters, such as AUXIN RESPONSE 1 (AUX1) and PINFORMED 1-7 (PIN1-7) (Grunewald & Friml, 2010). IAAs induce the expression of PLETHORA 1-4 (PLT1-4) TFs in the root meristem, which are responsible for cell proliferation maintenance (Galinha et al., 2007). Other TFs have been described to play key roles in the stem cell niche and the quiescent center (QC) maintenance to produce all tissues required to form a mature root, including the TFs SHORT-ROOT (SHR) and SCARECROW (SCR) (Sablowski, 2011). SHR expression is localized in a zone of the stele that constitutes the central part of the root and stem (Kim et al., 2020; Miyashima et al., 2011). When SHR transcripts are translated, SHR proteins move into adjacent cells (pericycle cells, endodermis, QC and phloem pole) to activate the expression of SCR (Cui et al., 2007; Helariutta et al., 2000; Nakajima et al., 2001; Sena et al., 2004). SHR also plays key roles in phloem development, controlling the asymmetric cell division process for sieve element development by the regulation of NARS1 and SND2 NAC-type TFs (Kim et al., 2020). This fact is relevant because these cell types constitute the phloem (Greb, 2020), which is the main means of IAA transport (Chapman & Estelle, 2009). However, Y. Liu et al. (2013) showed that NH_4^+ does not affect the IAA content at QC, where the maximum level of this phytohormone in roots is localized (Zhou et al., 2010).

As was mentioned above, LR branching is enhanced by NH_4^+ supply (Araya et al., 2016). When the quadruple ammonium transporter (AMT) knockout lines, that exhibit a severe reduction in LR branching, were complemented with *AtAMT1.1* or *AtAMT1.3*, only *AtAMT1.3*-complemented plants were able to restore the LR branching phenotype induced by NH_4^+ (Lima et al., 2010), suggesting that the NH_4^+ -induced LR branching signaling events are dependent on *AtAMT1.3* (Lima et al., 2010). Recently, it has been demonstrated that under NH_4^+ supply, RSA changes mediated by IAAs transport are highly linked to root acidification (Jia et al., 2020; Meier et al., 2020), which seems to be the molecular basis for the AMT transporter effect on LR branching since NH_4^+ uptake is accompanied by pH imbalance. In addition, IAAs have been suggested to be involved in this signaling event due to the repression of PIN2 under a supply of NH_4^+ (Y. Liu et al., 2013; Zou et al., 2013), which is provoked by the PIN2 hyperphosphorylation (Ötvös et al., 2021). Regarding root agravitropic response, NH_4^+ promotes the downregulation of AUX1 and PIN2, which are two pivotal IAA transporters and are subject to the antagonistic action between PIN2 and ARG1. ARG1 is involved in the transduction of the root gravity signal and required for normal AUX1 expression and basipetal IAA transport in root apices, promoting an asymmetric auxin flow (Y. Liu et al., 2013; Zou et al., 2013). All these evidence highlight the importance of IAA in orchestrating the root system configuration in response to this nutritional stimulus. However, the transcriptional regulatory mechanisms that control this response remain unclear.

Together with IAAs, cytokinins (CKs) have been previously established to be important in root plant growth and vascular development (Kamada-Nobusada et al., 2013; Mao et al., 2020; Miyashima et al., 2019). CK biosynthesis and activity in plants are closely related to N availability (Kamada-Nobusada et al., 2013; Takei et al., 2004). In the roots of rice, NH_4^+ nutrition leads to the accumulation of different CKs and CK-derived compounds (Kamada-Nobusada et al., 2013). During active growth of *Arabidopsis*, the content of CKs first increased in the vascular system, thus reflecting CKs transport from roots to shoots (Shtratnikova et al., 2015). Recently, it has been reported that CK signaling in the early protophloem-sieve element cell files of *Arabidopsis* root procambial tissue promotes the expression of several DOF TFs (Miyashima et al., 2019). Together with the IAA-responsive dependent HD-ZIP III proteins, these TFs compose a transcriptional network that integrates spatial information of the hormonal domains and microRNA (miRNA) gradients, which are essential for root vascular development (Miyashima et al., 2019).

Previous works in *Arabidopsis*, based on root protoplast generation and cell-sorting using FACS coupled to expression studies, revealed cell-specific responses to different processes such as N nutrition (Gifford et al., 2008; Walker et al., 2017) or plant immunity (Rich-Griffin et al., 2020). Regarding N nutrition, co-expression network approaches revealed that within the root cortex N controlled a wide range of processes including phytohormone responses, while in pericycle regulatory networks were related to formation of new organs, organ structure development, and establishment of cell localization (Walker et al., 2017) showing that phytohormones are

candidates signaling for cell-specific responses to N (Gifford et al., 2008).

Conifers are an ancient group of gymnosperms that cover vast regions in the Northern Hemisphere, and they have exceptional ecologic and socioeconomic importance (Farjon, 2018). Since conifers represent a differentiated evolutionary lineage of plants, they exhibit substantial differences in N metabolism with respect to angiosperms such as *Arabidopsis*. For instance, conifers lack glutamine biosynthesis in the plastid of photosynthetic cells (Cánovas et al., 2007), and most species prefer NH_4^+ over NO_3^- as the main source of inorganic N (Hawkins & Robbins, 2010; Kronzucker et al., 1997). This is the case for maritime pine (*Pinus pinaster* Aiton) (Ortigosa et al., 2020; Warren & Adams, 2002), a southwestern Mediterranean conifer employed as a model in studies of N nutrition and metabolism and for which a large body of genomics resources are available (Cañas et al., 2017; Ortigosa et al., 2020; 2021). Previous transcriptomic studies of pine seedling tissues provided an overview of the gene expression distribution in different pine tissues but also it was possible to highlight the relationships between gene expression patterns and function in a tissue-dependent manner (Cañas et al., 2014; 2017).

Maritime pine seedlings under NH_4^+ supply accumulate more biomass than those fed with NO_3^- what is related with a higher uptake of NH_4^+ than NO_3^- (Ortigosa et al., 2020). In maritime pine roots, a long-term supply of NH_4^+ induced the expression of transcripts related to defense, such as *antimicrobial peptide 1* (*PpAMP1*) (Canales et al., 2010), which is not found in dicots and can regulate the NH_4^+ uptake (Canales et al., 2011). There also was a close relationship between NH_4^+ -responsive genes and genes involved in amino acid metabolism, particularly those involved in asparagine biosynthesis and utilization (Canales et al., 2010). Additionally, NH_4^+ promotes changes in the epitranscriptome that mainly regulate the translational response and growth, including the repression of 1-aminocyclopropane-1-carboxylic acid (ACC) oxidase (ACO), the last enzyme in the ethylene (ET) biosynthesis pathway (Ortigosa et al., 2021). Considering this background, the main goal of the present work was to decipher the molecular mechanisms underlying the early response to NH_4^+ in the root apex and its relationship with root development in maritime pine.

2 | MATERIALS AND METHODS

2.1 | Plant material

Maritime pine seeds (*P. pinaster* Aiton) from "Sierra Segura y Alcaraz" (Albacete, Spain) were provided by the *Área de Recursos Genéticos Forestales* of the Spanish *Ministerio de Agricultura, Pesca y Alimentación*. Seed germination was carried out following the protocol described elsewhere (Cañas et al., 2006). Seedlings were grown in vermiculite in plant growth chambers (Aralab Fitoclima 1200, Rio de Mouro, Portugal) under 16 h light photoperiod, a

luminal intensity of $125 \mu\text{mol m}^{-2} \text{s}^{-1}$, a constant temperature of 23°C , 50% relative humidity and watered twice a week with distilled water. One-month-old maritime pine seedlings were used for the experiments. Pine seedlings were randomly subdivided into three different groups, relocated into forestall seedbeds and irrigated with 80 ml of a solution that contains macronutrients and micronutrients without any N source (1.16 mM KCl; 0.63 mM KH_2PO_4 ; 0.35 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$; 0.17 mM $\text{CaCl}_2 \cdot \text{H}_2\text{O}$; 80 μM EDTA- FeSO_4 ; 25.9 μM H_3BO_3 ; 10.2 μM $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$; 1.3 μM $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$; 0.7 μM $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$; 0.1 μM $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$). After 3 days of acclimation, the control group was irrigated with 80 ml of water and the experimental group was supplied with 80 ml of 3 mM NH_4Cl . Root samples were collected 24 h after the treatment and immediately frozen in liquid N (Figure 1).

Ten seedlings were collected and pooled per each sample for RT-qPCR validation. The same experiment was carried out three independent times. The screening condition and adequate development of each experiment was verified through the gene expression analysis by RT-qPCR of control genes following previous results (Ortigosa et al., 2021) (Figure 1). To perform laser capture microdissection (LCM), seedling's root apices were cut, and tissues (5–6 mm) were imbibed in a specimen holder with Tissue-Tek optimal cutting temperature embedding medium (Sakura Finetek) and immediately frozen in liquid N for cryostat sectioning. Frozen samples were stored at -80°C until use.

2.2 | Root phenotyping in response to nitrogen supply

For the root phenotypes under different nitrogen sources (NH_4^+ and NO_3^-), the pine seedlings were cultivated in a growth chamber with a 16/8 h light/dark photoperiod, light intensity of $125 \mu\text{mol m}^{-2} \text{s}^{-1}$, constant temperature of 23°C , and 50% relative humidity (Aralab Fitoclima 1200). One-month-old seedlings ($n = 30$) were randomly transplanted into seedbeds for each experimental condition: distilled water, 3 mM KNO_3 and 3 mM NH_4Cl . Each experimental condition group was irrigated twice per week with 50 ml of the corresponding N solution or 50 ml of distilled water for 74 days. After treatment, seedling roots were dissected and whole roots, primary root (PR) and LRs were weighted. The PR length and number of LRs was manually assessed (Figure S1).

2.3 | Whole root RNA isolation

Total RNA from roots of maritime pine seedlings was isolated following the protocol described by Liao et al. (2004) and modified by Canales et al. (2012). RNA concentration and purity were determined spectrophotometrically using a Nanodrop ND-1000 (Thermo Scientific). Purity of the preparation was assessed by determining the 260/280 and 260/230 ratios. The integrity was assessed by electrophoresis.




FIGURE 1 Experimental design. Timeline from germination to harvest. RT-qPCR data for the validation of the experimental design are included. *Glutamine synthetase 1b* (*PpGS1b*); *NADH-dependent glutamate synthase* (*PpNADH-GOGAT*); *antimicrobial peptide 1* (*PpAMP1*); *S-adenosyl methionine synthase* (*PpSAMS*) Significant differences were determined with a *t*-test (* $p < 0.05$; ** $p < 0.01$). Error bars show SE with $n = 3$. Isolated regions from the root apex: cap (RC), meristem (RM), developing cortex (RDC) and developing vessels (RDV). RT-qPCR, quantitative reverse-transcription polymerase chain reaction [Color figure can be viewed at wileyonlinelibrary.com]

2.4 | Laser capture microdissection, RNA isolation and low-input RNA-seq

LCM procedure was carried through as previously described (Cañas et al., 2014). Full step protocol is described in Supporting Information Methods S1. Four different tissue areas were isolated by microdissection corresponding to the root cap (RC), meristem (RM), developing cortex (RDC) and developing vessels (RDV) areas (Supporting Information Video S1).

All RNA extractions from the microdissection procedure were carried out using manufacturer's instruction protocol (non-LCM) for the RNAqueous-Micro RNA Isolation Kit (Ambion). RNA quality, DNA contamination and first quantification were performed via RNA Pico Assay for the 2100 Bioanalyzer (Agilent). Quantification was verified via a Qubit RNA BR (Broad-Range) Assay Kit (Invitrogen). RNA samples with RNA integrity number (RIN) higher than 7 were used for subsequent RNA sequencing, messenger RNA amplification and complementary DNA (cDNA) synthesis 1 ng of RNA for each sample.

The low input RNA-seq was carried out by Novogen (Hong Kong). The cDNA synthesis and amplification, and the library

preparation was made with the SMART-Seq™ v4 Ultra™ Low Input RNA Kit for Sequencing (Takara) following the manufacturer's instructions. RNA sequencing was made in a NovaSeq. 6000 sequencer according to the manufacturer's instructions for paired-end reads (Illumina). The 24 samples were sequenced producing paired-end reads of 150 bp length. The sequencing output is shown in Table S1.

The raw reads were trimmed (quality and contamination) using SeqTrimBB software (<https://github.com/rafnunser/seqtrimbb>). Only the pairs in which both reads passed the quality test were further analyzed ($Q > 20$). Trimmed reads are shown in Table S1. These reads were assembled using Trinity 2.11 (Haas et al., 2013). Contigs lower than 400 pb were eliminated, for the rest of contigs the redundancy was reduced using CD-HIT-EST software (Fu et al., 2012). This transcriptome shotgun assembly project has been deposited at DDBJ/EMBL/GenBank under the accession GJFX00000000. The version described in this paper is the first version, GJFX01000000. The final transcriptome was used as the reference for the read mapping that was performed with BWA using the MEM option (Li & Durbin, 2009). The read count was obtained with the python script *sam2counts* (<https://github.com/vsbuffalo/sam2counts>). Differentially expressed (DE) transcripts were identified using the edgeR

package for R, the transcripts were normalized by cpm and filtered; 2 cpm in at least two samples (Robinson et al., 2010). Each sample was from a single seedling in a different experimental replicate. The samples were basically grouped by tissue and nutritional condition (Figure S2). For the tissue-treatment interaction only the transcripts with $FDR < 0.05$ and the three experimental replicates with the same expression sense than the final logFC (positive or negative) were considered as DE. For the tissue analysis same parameters have been considered. The DE transcripts were used to construct a gene co-expression network. An unsigned network has been carried out using the R package WGCNA soft-thresholding power value of 9 (Langfelder & Horvath, 2008). From the network, the 10% of the transcripts with more connections in each module were considered hub genes.

These RNA-Seq data have been deposited in the NCBI's Gene Expression Omnibus (Edgar et al., 2002) and are accessible through GEO Series with the accession number GSE175587 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE175587>). Additionally, RNA-seq and network results are accessible through a database in html format that can be installable with R packages and downloaded from GitHub (<https://github.com/ceslobfer/Rootapp>).

Total RNA (1 ng) was retrotranscribed and amplified to verify the expression analyses for several DE genes by RT-qPCR. The cDNA synthesis and amplification protocol was carried out using the Conifer RNA Amplification (CRA+) protocol previously described by Cañas et al. (2014). Full step protocol is described in Supporting Information Methods S1. The amplification process was monitored using the ERCC RNA Spike-in kit (Thermo Scientific) according to manufacturer's instructions. The primers used for cDNA synthesis and amplification are listed in Table S2.

2.5 | Functional annotation and enrichment analyses

The assembled transcriptome was functionally annotated with BLAST2GO (Götz et al., 2008) using DIAMOND software with *blastx* option (Buchfink et al., 2015) against the NCBI's plants-*nr* database (NCBI Resource Coordinators, 2016). Blast results were considered valid with $e < 1.0E-6$. Singular enrichment analysis (SEA) of the GO terms was made in the AGRIGO v2.0 web tool under standard parameters using as GO term reference the whole assembled transcriptome annotation (Tian et al., 2017). Representative enriched GO was determined using REVIGO with 0.5 as dispensability cutoff value (Supek et al., 2011).

2.6 | RT-qPCR

The cDNA synthesis was performed using 1 µg of total RNA and iScript™ cDNA Synthesis Kit (Bio-Rad) following manufacturer's instructions. The qPCR primers were designed following the MIQE guidelines (Bustin et al., 2009). The primers are listed in Table S2.

qPCRs were carried out using 10 ng of cDNA and 0.4 mM of primers and 2X SsoFast™ EvaGreen® Supermix (Bio-Rad) in a total volume of 10 µl. Relative quantification of gene expression was performed using thermocycler CFX 384™ Real-Time System (Bio-Rad). The qPCR program was as follows: 3 min at 95°C (1 cycle), 1 s at 95°C and 5 s at 60°C (50 cycles) and a melting curve from 60°C to 95°C, to generate the dissociation curve to confirm the specific amplification of each individual reaction. The analyses were carried out as described by Cañas et al. (2014) using the MAK3 model in the R package *qpcR* (Ritz & Spiess, 2008). Normalization for gene expression of experimental design viability was performed using geometric mean of two reference genes, a *Saposisin-like aspartyl protease* (pp_199988) and *Myosin heavy chain-related* (pp_58489) that were previously tested for maritime pine (Granados et al., 2016). Normalization for gene expression of LCM isolated tissues was performed using as reference gene a *SKP1/ASK1 family protein* (pp_18128) that was previously tested for maritime pine LCM samples (Granados et al., 2016). For the RT-qPCR analysis, three technical replicates of each sample and three biological replicates were made.

2.7 | Imaging of phytohormones in roots sections

The main phytohormones were localized in root apex cuts using nano-particle assisted laser desorption/ionization mass spectrometry imaging (Nano-PALDI-MSI). The elongation zones, which corresponded to the region from 2.5 to 5 mm of the root apex, were embedded in super cryoembedding medium (SCEM; Leica Biosystems) and frozen in liquid nitrogen. The specimen block was cut into 10 µm sections using a cryostat (NX-70; Thermo Scientific) set at -23°C in the chamber and at -25°C on the object holder. The sections were gently mounted on slides coated with indium tin oxide (ITO) (Bruker Daltonik GmbH). Optical images of the sections were obtained by a virtual slide scanner (Nanozimmer-SQ, Hamamatsu Photonics) before analysis by Nano-PALDI-MSI.

For Nano-PALDI-MSI, iron oxide-based nanoparticles (Fe-NPs) were prepared by stirring aqueous solutions of FeCl₂·4H₂O (5 ml, 100 mM; FUJIFILM Wako Pure Chemical), and 3-aminopropyltriethoxysilane (5 ml; γ-APTES; Shin-Etsu Chemical) at room temperature for 1 h. The resulting precipitate was washed several times with ultrapure water, resuspended in methanol (Moritake et al., 2007).

One milligram of Fe-NPs was resuspended in 1 ml methanol and sprayed on pine root tissue sections on ITO-coated glass slides with an airbrush (nozzle caliber, 0.2 mm). To obtain images, each data point on the section were irradiated with 200 laser shots in the positive ion detection mode of the mass spectrometer. Only signals between 80 and 800 *m/z* were analyzed to detect the correlated IAA (*m/z* 176.3), cytokinin (*m/z* 221.3), ACC (*m/z* 102.8), salicylic acid (SA) (*m/z* 139.4), JA (*m/z* 211.0) and ABA (*m/z* 265.0) as protonated ions, respectively. For each section, approximately 101,800 data points were obtained, 5 µm apart. The MS image was reconstructed from the obtained MS spectra with a mass bin width of $m/z \pm 0.1$ from the

exact mass using flexImaging 4.0 (Bruker Daltonik GmbH). The accurate mass of the ions was used for image generation, and mass accuracy and root-mean-square error (RMSE) were automatically calculated by the imaging software to avoid false-positive signals (Shiono & Taira, 2020). Comparisons of MS images were derived from the relative intensity for each signal normalized by the highest intensity spot on the slide. The peak intensity value of the spectra was normalized by dividing them with the total ion current (TIC) to achieve semi-quantitative analysis between control- and NH_4^+ -treated roots.

2.8 | Phylogenetic analyses

Evolutionary analyses were performed in MEGA7 (Kumar et al., 2016). The protein sequence alignment was made with Muscle (Edgar, 2004). The evolutionary history was inferred using the Neighbor-Joining method (Saitou & Nei, 1987). The bootstrap consensus tree inferred from 1000 replicates is taken to represent the evolutionary history of the taxa analyzed (Felsenstein, 1985).

3 | RESULTS

3.1 | Tissue-specific transcriptomic response to NH_4^+ supply

In this study, tissue-specific transcriptome changes triggered by NH_4^+ nutrition were analyzed in the maritime pine root apex by a combination of laser capture microdissection and high-throughput RNA sequencing. Prior proceeding with tissue isolation by LCM, each experimental replicate was validated through the expression analysis of transcripts that were expected to be upregulated by NH_4^+ supply at 24 h post-irrigation (Figure 1) (Canales et al., 2011; Ortigosa et al., 2021). As expected, NH_4^+ induced the accumulation of transcripts coding for glutamine synthetase 1b (*PpGS1b*), antimicrobial peptide 1 (*PpAMP1*) and S-adenosyl methionine synthase (*PpSAMS*). Although not significant, the NADH glutamate synthase (*PpNADH-GOGAT*) expression also increased with NH_4^+ supply.

As described in the Materials and Methods section, root apices from seedlings that were irrigated with 3 mM NH_4^+ and harvested at 24 h were used for the isolation of four different tissues by LCM, namely, RC, RM, RDC and RDV (Video S1). The differential expression results are shown in Dataset S1. A total of 295 DE transcripts were identified in the low-input RNA-seq analysis (Figure 2A,B), of which 182 DE transcripts were downregulated (Figure 2A) and 113 were upregulated (Figure 2B). Among the isolated tissues, RDC showed the highest number of DE transcripts (107 upregulated and 70 downregulated) while RDV had the lowest response when seedlings were treated with NH_4^+ (12 upregulated transcripts). Interestingly, only two genes were upregulated in all root tissues, namely, two splicing isoforms from a common gene with unknown function.

The analysis of the DE transcripts revealed 8 expression patterns (Figure 2c). Among the groups with the most significant differences, the cluster 2 had transcripts repressed in RC by NH_4^+ , and it contained several TFs, such as *PpSHR*, *PpDOF11* and *PpGATA20-like*, and transcripts related to phytohormones, such as *PpVAN3/FKD2* (Figure 2c; Dataset S2). In cluster 3, transcripts related to root development, such as *PpBAM1*, and phytohormones such as *flavanone 3-dioxygenase*, were found to be repressed by NH_4^+ in RC and RDC (Figure 2c; Dataset S2). In RDC and RM, NH_4^+ induced the transcripts framed into cluster 4, which are mainly related to defense, such as chitinases of class I and IV (Figure 2C; Dataset S2).

To validate our transcriptomic analysis, the expression value of several DE transcripts was corroborated by RT-qPCR (Figure 2d; Dataset S1). The downregulation of *PpSHR* (pp_346622) and *PpNPF3.1* (pp_58258) in RC; *PpDOF12* (pp_85381) and *PpACO* (pp_202153) in RM; and *PpBLISTER-like* (pp_170664) in RDC was confirmed. The upregulation of transcripts coding a *cycloeucalenol cycloisomerase* (*PpCPI1*; pp_189071) in RC and *PpAMP1* (pp_580007) in RDC was also corroborated.

3.2 | Functional enrichment analysis of tissue-specific transcriptomic response to NH_4^+ supply

The functional study of DE transcripts, including the SEA results, is summarized in Figure 3. In the most responsive tissues (RC, RM and RDC), NH_4^+ induced a general alteration in the amounts of transcripts coding TFs involved in development (most of them downregulated) and transcripts related to different phytohormones (Figure 3a–c). In the less responsive tissue (RDV) but also in RM and RDC, NH_4^+ promoted the accumulation of defense-related transcripts (Figure 3b–d). In the SEA results, no significant GO terms were found for DE transcripts upregulated in RC and RDV and downregulated in RM and RDV. The complete results of the SEA analyses are shown in Dataset S3. The largest number of SEA results were for transcripts downregulated in RC and upregulated in RDC. The downregulated transcripts in RC were significantly enriched in GO terms in the biological process (BP) category, such as “biological regulation” (GO:0065007), “gene expression” (GO:0010467), “response to hormone” (GO:0009725), “regulation of hormone levels” (GO:0010817) and “regulation of transcription, DNA-templated” (GO:0006355) (Figure 3e). The terms enriched in the molecular function (MF) category included “transcription factor activity” (GO:0003700), “protein binding” and “DNA binding” (GO:0005515 and GO:0003677) (Figure 3f). In the BP category for the downregulated transcripts in RDC only the “cytoskeleton organization” term (GO:0007010) was significantly enriched; however, for the upregulated transcripts in this tissue, numerous enriched functions were significant, such as those involved in defense responses such as “induction of programmed cell death” (GO:0012502), “cell killing” (GO:0001906), “defense response” (GO:0006952) and “cell wall macromolecule catabolic process” (GO:0016998) and “amino sugar catabolic process” (GO:0046348) (Figure 3e). Regarding the MF category for the




FIGURE 2 Low-input RNA-seq results in the different tissues of maritime pine root tip. (a) Venn diagram of downregulated differentially expressed (DE) transcripts. (b) Venn diagram of upregulated DE transcripts. (c) Heatmap and hierarchical clustering of DE transcripts. Expression values correspond to median CPM between biological/experimental replicates. The hierarchical clustering was made employing Ward's minimum variance method. (d) Experimental validation of low-input RNA-seq results for 8 DE transcripts through RT-qPCR. RC, the root cap; RDC, root developing cortex; RDV, root developing vessels; RM, root meristem; RT-qPCR, quantitative reverse-transcription polymerase chain reaction. *SHORT-ROOT* (*PpSHR*); *nitrate transporter 1/peptide transporter family 3.1* (*PpNPF3.1*); *cycloeucaenol cycloisomerase* (*PpCPI1*); *DOF transcription factor 12* (*PpDOF12*); *NAC transcription factor 38* (*PpNAC38*); *aminocyclopropane-1-carboxylic acid oxidase* (*PpACO*); *antimicrobial peptide 1* (*PpAMP1*); *BLISTER-like transcription factor* (*PpBLISTER-like*) [Color figure can be viewed at wileyonlinelibrary.com]

upregulated transcripts in RDC, there were significant GO terms related to the response to biotic stress, such as “chitinase activity” (GO:0004568), “chitin binding” (GO:0008061), “mannose binding” (GO:0005537) and “hydrolase activity, acting on glycosyl bonds” (GO:0016798) (Figure 3f). Interestingly, the SEA results of the upregulated transcripts in RM were similar but limited for defense responses, which was also found for the upregulated transcripts in RDC (Figure 3e,f). Finally, the only enriched GO terms for the cellular component category were “extracellular region” (GO:0005576) and “extracellular space” (GO:0005615) for the transcripts upregulated in RDC tissue (Dataset S3).

3.3 | RNA-seq analysis in function of the root tissue

Although it has been studied more in depth in a previously work (Cañas et al., 2017), the transcriptional changes underlying different

tissues that form the root apex of maritime pine (RC, RM, RDC and RDV) have been briefly analyzed (Dataset S1). The comparison of the DE transcripts showed a set of characteristic genes of each tissue. This gene core was composed of 1014, 2529 and 2692 DE transcripts for RC, RM, RDC and RDV, respectively (Figure S3A). The functional study of the core DE transcripts revealed the main functions for these tissues (Dataset S3). Some of the enriched GO terms for RC are “aminoglycan metabolic process” (GO:0006022), “cell wall organization or biogenesis” (GO:0071554), “dehiscence” (GO:0009900) or “polysaccharide catabolic process” (GO:0000272) (Figure S3B). For RDC gene core, GO terms such as “anatomical structure formation involved in morphogenesis” (GO:0048646), “catabolic process” (GO:0009056) or “response to stimulus” (GO:0050896) were observed (Figure S3B). On the other hand, RDV was the tissue that showed the highest number of DE transcripts when compared to the rest of the tissues, which presented enrichment GO terms such as “auxin influx” (GO:0060919), “carbohydrate metabolic process” (GO:0005975), “fluid transport” (GO:0042044) or




FIGURE 3 Resume of the main functional results obtained in the low-input RNA-Seq analysis from LCM isolated samples. Functions and DE transcripts in (a) the root cap (RC), (b) root meristem (RM), (c) root developing cortex (RDC) and (d) root developing vessels (RDV). Blue arrows indicate downregulated transcripts, red arrows upregulated ones. (e) Significant GO terms from Biological Processes category after a SEA analysis. (f) Significant GO terms from Molecular Function category after a SEA analysis. DE, differentially expressed; LCM, laser capture microdissection; SEA, singular enrichment analysis [Color figure can be viewed at wileyonlinelibrary.com]

“phloem or xylem histogenesis” (GO:0010087), among others (Figure S3B). Interestingly, in RM only 2 DE transcripts were identified as characteristic and both transcripts are splicing variants of the same gene encoding a bHLH TF, similar to the *Arabidopsis* FAMA TF (AT3G24140) (Figure S3C).

3.4 | Transcription factors affected by NH_4^+ in the apex of maritime pine roots

The DE transcripts were individually analyzed looking to identify different kinds of regulators, such as TFs and transcripts involved in the phytohormone response. From a total of 295 DE transcripts, 31 TFs were identified (Figure 4, Table 1, Dataset S1). All 10 TFs identified in RC were repressed in the presence of NH_4^+ . However, in RM and RDC, 6 and 4 TF transcripts were downregulated while 5 and 6 TF transcripts were upregulated, respectively. Interestingly, a high abundance of TFs related to root growth and development was observed (Table 1), such as the strong downregulation of *PpSHR* in RC, *INCURVATA-like* and homeobox-leucine zipper protein *HAT-like* both in RM and involved in meristem developmental regulation, and the TF

identified as *zinc finger C2H2 SHOOT GRAVITROPISM* in RC related to the gravitropism response. *PpSHR* was the most highly repressed TF ($-11 \log_{10} \text{FC}$). Furthermore, the induction of two transcriptional repressors of different plant developmental processes was observed, with *PpNAC31* identified as an *SMB-like* NAC TF in RM and an *OFP17-like* repressor in RDC. In addition, novel TFs in maritime pine were identified. Two of them belong to the DOF-family and were named as *PpDOF11* and *PpDOF12*, and they were both downregulated in RC and RM, respectively. Phylogenetic analysis of these two new DOF-type TFs revealed that *PpDOF11* is grouped with *AtDOF1.4* (AT1G28310) and *OsDOF7.2* (LOC_Os07g32510) into subfamily E and that *PpDOF12* is grouped with members of subfamily A of *P. pinaster*: *PpDOF4*, *PpDOF7*, *PpDOF8* and *PpDOF10* (Figure S4, Table S3). Additionally, a member of the NAC family (*PpNAC38*) that was upregulated in RM tissue was identified. This TF has homology to *BRN1* (AT1G33280) (e-value: $7e-96$) and *BRN2* (AT4G10350) (e-value: $3e-97$) of *Arabidopsis thaliana* and is framed within the NAC subfamily C (Figure S5, Table S4).

A gene co-expression network was constructed to identify hub TFs and putatively determine the processes controlled by them. Thirteen transcripts were identified as hubs corresponding




FIGURE 4 Expression heatmaps of differentially expressed transcriptional regulators and hormone-related transcripts. (a) Transcription factors and other putative regulators. (b) Transcripts involved in hormone metabolism, transport and signaling. CPM expression values were normalized by transcript. RC, the root cap; RDC, root developing cortex; RDV, root developing vessels; RM, root meristem [Color figure can be viewed at wileyonlinelibrary.com]

to eight different TFs: *PpDOF1*, *PpDOF2*, *PpDOF11*, *PpGATA20-like*, *PpINCURVATA-like*, *PpMyb149*, *PpSGR/IDD16* and *PpSHR*. All of them were DE for NH_4^+ treatment in different tissues except RDV but all of them were more expressed in RDV than in the rest of tissues except *PpGATA20-like* (Dataset S1). The functions and expression localization of the most correlated transcripts ($>|0.9|$) were analyzed (Figure S6; Dataset S4). Most of the transcripts were correlated with *DOF* TFs, *PpINCURVATA-like* and *PpSGR/IDD16* but also belonged to the RDV gene core (1879 from 2362) (Figure S6A). Thus, the enriched functions were mainly related with RDV functions such as “auxin influx” (GO:0060919), “carbohydrate metabolic process” (GO:0005975), “fluid transport” (GO:0042044) or “phloem or xylem histogenesis” (GO:0010087) (Figure S6B; Dataset S5). Interestingly, transcripts correlated with *PpGATA20-like* were enriched in functions such as “gene expression” (GO:0010467) and “ribosome biogenesis” (GO:0042254) (Figure S6B; Dataset S5).

3.5 | Phytohormone-related genes

Twenty-six DE transcripts related to phytohormone pathways were identified (Table 1), of which 19 were upregulated and 7 were downregulated (Figure 4, Dataset S1). Most of these DE transcripts were related to IAA (Table 1). Some interesting examples are the downregulation of transcripts encoding proteins required for PIN transporters localization, such as VAN3/FKD2 Auxin canal/PH2 (pp_60103) in RC and the upregulation of a *PpCPI1* (pp_189071) in RC, which is related to PIN transporters endocytosis. In addition, in RDC, the repression of transcripts coding for the IAA receptor TIR1/AFB (pp_238715) and for a sulfotransferase (pp_208269) with a high similarity degree to SULT202B1 (AT3G45070) (e-value: $5e-72$) was observed.

CK-related genes were the second most represented phytohormone-related transcripts, and strong repression of CK riboside 5'-monophosphate phosphoribohydrolase (pp_222714) (-6.9

TABLE 1 Most representative DE transcripts coding for TFs and hormone related proteins

TF in maritime pine	Gene ID	Tissue	Transcript in <i>A. thaliana</i>	Gene ID	Process involved	References
PpDOF11	pp_58878	RC	HCA2/DOF5.6	AT5G62940	Root tip vascular proliferation: radial growth of protophloem sieve elements	Guo et al. (2009) Miyashima et al. (2019)
PpSHR	pp_246622	RC	SHR	AT4G37650	- Quiescent center cells specification - Asymmetric cell division related to root radial pattern formation - Phloem development	Fukaki et al. (1998) Helariutta et al. (2000) Cui et al. (2007) Kim et al. (2020)
PpGATA20-like	pp_204819	RC	GATA20	AT2G18380	Root tip vascular development	Smit et al. (2020)
PpSGR/IDD16	pp_174625	RC	IDD14	AT1G68130	- Lateral organ morphogenesis and gravitropic responses - Establishment of auxin gradients (auxin biosynthesis and transport)	Cui et al. (2013)
PpPHR1-like	pp_64350	RDC	PHR1	AT4G28610	Phosphate starvation	Rubio et al. (2001)
PpOPF17	pp_80824	RDC	OPF17	AT2G30395	Plant growth and development	Wang et al. (2011)
PpBLISTER-like	pp_170664	RDC	BLISTER	AT3G23980	Plant vegetative growth	Schatlowski et al. (2010) Hong et al. (2019)
PpDOF2	pp_119525	RDC	PEAR1/DOF2.4	AT2G37590	Root tip vascular proliferation: radial growth of protophloem sieve elements	Miyashima et al. (2019)
PpDOF12	pp_85381	RM	CDF3	AT3G47500	Involved in abiotic stress tolerance and nitrogen response	Corrales et al. (2017) Dominguez-Figueroa et al. (2020)
PpDOF1	pp_233816	RM	OPB3	AT3G55370	Regulator of phyB and cry1 signaling pathways	Ward et al. (2005)
PpHAT-like	pp_141831	RM	HAT1	AT4G17460	- Inflorescence meristem regulation, floral meristem determination, gynoecium medial domain development - Repressor of BR responsive genes and ABA biosynthesis	Zúñiga-Mayo et al. (2012) Zhang et al. (2014) Tan et al. (2018)
PpINCURVATA-like	pp_85012	RM	ATBH-15	AT1G52150	Root meristem patterning and putatively in the regulation of procambial and vascular tissue formation or maintenance.	Ochando et al. (2006)
PpNAC31	pp_168080	RM	SMB	AT1G79580	Cellular maturation of root cap regulation	Willemssen et al. (2008) Bennett et al. (2010) Fendrych et al. (2014)
PpNAC38	pp_72282	RM	BRN1/BRN2	AT1G33280	Cellular maturation of root cap regulation	Bennett et al. (2010)
PpVAN3/FKD2	pp_60103	RC and RM	VAN3/FORKED2	AT5G13300	Plant vascular tissue formation by PIN transporters localization	Steynen and Schultz (2003) Naramoto et al. (2009) Hou et al. (2010)
Cytokinin riboside 5'-monophosphate phosphoribohydrolase	pp_222714	RC	LOG3	AT2G37210	Cytokinin activation	Kuroha et al. (2009)

TABLE 1 (Continued)

TF in maritime pine	Gene ID	Tissue	Transcript in <i>A. thaliana</i>	Gene ID	Process involved	References
<i>Cycloaucalenol cycloisomerase</i>	<i>pp_189071</i>	RC	CPII	AT5G50375	PIN2 endocytosis	Men et al. (2008)
<i>Sulfotransferase</i>	<i>pp_208269</i>	RDC	SULT20B1	AT3G45070	Flavonoids sulfation	Hashiguchi et al. (2013)
<i>PpABA2</i>	<i>pp_198945</i>	RDC and RM	ABA2	AT1G52340	Biosynthesis of abscisic acid	González-Guzmán et al. (2002)
<i>PpCYP90B1</i>	<i>pp_247190</i>	RDC	CYP90B1	AT3G50660	Biosynthesis of brassinosteroids	Fujita et al. (2006)
<i>PpTIR1</i>	<i>pp_238715</i>	RDC	TIR1	AT3G62980	Auxin signalling pathway	Dharmasiri et al. (2005) Salehin et al. (2015)
<i>PpCYP87A3</i>	<i>pp_81114</i>	RDC	CYP87A2	AT1G12740	Putatively involved in hormone biosynthesis and/or auxin signaling	Chaban et al. (2003), Bak et al. (2011)
<i>GASA gibberellin regulated cysteine rich protein</i>	<i>pp_111379</i>	RDC	GASA10	AT5G59845	Putatively involved in response to phytohormones	Zhang and Wang (2017)
<i>HMG-CoA reductase</i>	<i>pp_144669</i>	RDC	HMGR	AT2G17370	Regulatory enzyme of the cytokinin and brassinosteroid biosynthesis pathways	Antolin-Llovera et al. (2011)
<i>PpABA2</i>	<i>pp_209084</i>	RDC	ABA2	AT1G52340	Biosynthesis of abscisic acid	González-Guzmán et al. (2002)
<i>Cytokinin dehydrogenase</i>	<i>pp_215376</i>	RDC	CKX1	AT2G41510	Catalyzes the oxidation of cytokinins	Werner et al. (2003)
<i>PpACO</i>	<i>pp_202153</i>	RM	ACO	AT2G19590	Ethylene biosynthesis	Houben and Van de Poel (2019)
<i>Flavonoid 3'-monooxygenase</i>	<i>pp_203885</i>	RM	F3'H	AT5G07990	Involved in flavonoids biosynthesis pathway from naringenin to eriodictyol and dihydrokaempferol to dihydroquercetin	Schoenbohm et al. (2000)
<i>EXORDIUM</i>	<i>pp_129078</i>	RM	EXO	AT4G08950	Putatively involved in brassinosteroids response in roots	Coll-García et al. (2004)

Abbreviations: RC, the root cap; RDC, root developing cortex; RDV, root developing vessels; RM, root meristem.

logFC), a CK-activating enzyme, was observed in RC. However, up-regulation of a transcript coding for a CK dehydrogenase (pp_215376), a CK-inactivating enzyme, was also observed in RDC. The repression of transcripts related to brassinosteroids and ethylene (ET) biosynthesis, for example, cytochrome P450 90B1 (pp_247190) in RDC and *PpACO* (pp_202153) in RM was also observed. In addition, transcripts related to the gibberellin (GA) response and abscisic acid (ABA) biosynthesis were upregulated, such as a GASA gibberellin-regulated cysteine-rich protein (pp_111379) in RDC and two ABA2 coding transcripts that might be involved in ABA biosynthesis (pp_198945, and pp_209084) in RM and RDC.

3.6 | Phytohormone detection in maritime pine root apex

To corroborate whether NH_4^+ nutrition altered the spatial allocation of several phytohormones as the transcriptomic data suggest, the distribution of multiple phytohormones was determined by Nano-PALDI-MSI in the apex of maritime pine roots (Shiono & Taira, 2020). In comparison to control seedlings, changes in the patterns of IAAs, CK (tZ, *trans*-Zeatin), ACC, ABA, SA and jasmonic acid (JA) were observed in the presence of NH_4^+ (Figure 5 and S5). In the control plants, an IAA gradient was observed, the maximum level was detected in the most distal area of the principal root apex (Figure 5a); however, in the NH_4^+ -treated roots, the IAA maximum was detected in a more distant zone from the root tip (Figure 5a). CK (tZ) showed a wide distribution in the roots of control plants (Figure 5b). In the NH_4^+ -treated seedlings, CK was distributed in the outer tissues of the root mainly below the root tip (Figure 5b). Regarding ET, its precursor ACC was found in a very restricted area of the root tip in control and NH_4^+ -treated plants, showing no changes when plants were supplied with NH_4^+ (Figure 5c). ABA was detected mainly in the RDC tissue in both control and treated seedlings, and it was higher under NH_4^+ supply and distributed below the root apex (Figure 5d). SA and JA phytohormone distributions were also affected by the presence of NH_4^+ (Figure S7). SA was located at the end of the root tip, while JA was also detected in the elongation zone in control plants. When the seedlings were supplied with NH_4^+ , SA showed a vaguer spatial distribution with respect to the observed distribution in the control plants (Figure S7A) while JA tended to slightly increase in the outermost parts of the root apex ends and disappeared from the elongation zone (Figure S7B).

3.7 | Maritime pine root phenotype

Maritime pine root morphological studies showed no differences between treatments regarding whole root biomass accumulation (Figure 6a). When the weight of the principal roots (PRs) was measured, NH_4^+ -fed seedlings exhibited a significant reduction in PR weight compared to water- and NO_3^- -treated plants (Figure 6b) and a statistically higher root length when pine seedlings were supplied

with either inorganic nitrogen form (Figure 6c). Moreover, NH_4^+ promoted an increase in LR number and weight compared to the water and NO_3^- treatments (Figure 6d,e), although no statistically significant differences were observed regarding LR density (LRD) between treatments (Figure 6f).

4 | DISCUSSION

Despite the obvious environmental and economic importance of conifers, little is known about their response to diverse nutrients at the molecular and developmental levels. Since they represent an ancient lineage of gymnosperms, conifers are of particular interest from an evolutionary perspective because they have common but also differential responses with angiosperm models. The aim of the present work was to explore the early response of maritime pine to NH_4^+ nutrition in the different tissues of the root apex. For this purpose, tissue isolation was performed by LCM and combined with low-input RNA-seq to determine the local transcriptomic response in the growth area of the root. This strategy avoids the dilution effect of transcripts with low and much localized expression, revealing new genes not previously annotated (Cañas et al., 2017), such as *PpDOF11-12* and *PpNAC38*. Clearly a dilution effect can be observed when compared the expression of marker genes in the whole roots (Figure 1) and in the specific tissue transcriptomics (Figure S8A). Supporting this, from 47,118 transcripts used in the tissue-specific RNA-seq, 8,491 transcripts were not previously identified in whole root transcriptomic experiments (Dataset S1). A similar number of DE transcripts was obtained in the present work (295) when compared to the same NH_4^+ nutrition treatment over the whole root (350) (Ortigosa et al., 2021) (Figure S8B). However, only three transcripts with the same identifiers are shared, although there are different transcripts from the same gene that are DE in both analyses, for example, *PpAMP1* (RDC: pp_58007/whole root: pp_58005) (Figure S8C).

The response to NH_4^+ in RDC involved the expression of defense-related genes as previously described in maritime pine and other model plants (Figure 3c,e,f) (Canales et al., 2010; Ortigosa et al., 2021; Patterson et al., 2010; Ravazzolo et al., 2020). Obviously, this response might be linked to the peripheral localization of RDC in the roots since the defense proteins encoded by the expressed transcripts develop their functions in the extracellular media (Dataset S3). Some examples are the upregulation of different *ginkbilobin-like/embryo abundant protein* transcripts (e.g., pp_252647), *chitinases* (e.g., pp_117806) and the *PpAMP1* (pp58007), which are consistent with results previously reported in maritime pine roots (Canales et al., 2010; Ortigosa et al., 2021).

In angiosperm model plants, NH_4^+ alters RSA by inhibiting root elongation, stimulating LR branching and affecting root hair development (Y. Liu & von Wirén, 2017). RSA principally targets root system processes, such as root elongation, root gravitropism and LR branching, appearing to occur in the root tip (Li et al., 2016). Based on the presented transcriptomic analyses, early root exposure to NH_4^+




FIGURE 5 Phytohormone localization in root apex sections. The main phytohormones were localized in root apex cuts of control and 3 mM NH_4^+ treated seedlings. The phytohormone identification and imaging was made using nano-particle assisted laser desorption/ionization mass spectrometry imaging (Nano-PALDI-MSI). (a) Auxin/Indole-3-acetic acid (IAA); (b) cytokinin (CK); (c) aminocyclopropane-1-carboxylic acid (ACC) and (d) abscisic acid (ABA). Size bars correspond to 500 μm . Column graphs show relative hormone accumulations respect to control samples in the approximate LCM areas. Dashed lines mark hormone accumulation in the control roots. LCM, laser capture microdissection; RC, root cap; RDC, root developing cortex; RDV, root developing vessels; RM, root meristem [Color figure can be viewed at wileyonlinelibrary.com]

caused a wide impact on the expression of TFs related to root growth and development in pine (Figure 2 and Table 1). Thus, NH_4^+ caused a strong repression of *PpSHR* (pp_246622) in RC tissue, a GRAS-type TF involved in the regulation and coordination of root development, including phloem differentiation (Kim et al., 2020). Interestingly, in RDC tissue, it was observed that *PpBAM-like* (pp_19744) transcripts were also severely affected by NH_4^+ (Dataset S1), which is consistent with previous cell-specific transcriptomic profiles observed in *Arabidopsis* (Brady et al., 2007). BAM1/2 kinase receptors are required for SHR-dependent formative divisions in roots and for the proper

CYCD6;1 expression, which is required to promote division in cortex endodermal initial daughter cells (Crook et al., 2020).

Additionally, a progressive reduction in the abundance of IAA efflux carriers (PINs) in *Arabidopsis shr* mutants has been described (Lucas et al., 2011), and NH_4^+ negatively affects the expression of PIN2 and AUX1 coding transcripts (Y. Liu et al., 2013). The expression of genes coding for PIN transporters was not affected in pine roots. Recently, it has been linked IAA signaling pathway to acidification (Jia et al., 2020; Meier et al., 2020). However, based on our results, no significant changes were observed in the expression of pH-related




FIGURE 6 Root growth and architecture parameters. Pine seedlings were cultivated for 74 days and irrigated once a week with 50 ml of water, 3 mM KNO_3 and 3 mM NH_4Cl , respectively. Six different parameters were measured after harvesting: (a) weight of whole roots; (b) weight of primary root; (c) primary root length; (d) number of lateral roots; (e) weight of lateral roots; (f) lateral root density. Significant differences were determined with a one-way ANOVA. Letters above the conditions show significant differences based on a Turkey post-hoc test ($p < 0.05$) with $n = 30$. ANOVA, analysis of variance [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]

genes. However, it was observed that the expression of coding transcripts for proteins involved in IAA transporter localization and components of the IAA signaling pathway were affected by NH_4^+ . This is the case for *PpCPI1* (pp_189071), which is upregulated in RC and required for PIN2 endocytosis (Men et al., 2008), and the strong repression of *VAN3/PH2/FKD2* (pp_60103) in RC and RM tissues (Dataset S2), which is crucial in *Arabidopsis* for the vascular leaf pattern formation by the proper PIN transporter localization (Hou et al., 2010). These results suggest that NH_4^+ triggers an alteration in polar auxin transport (PAT) involving a highly coordinated transcriptomic response between tissues of the pine root. The nano-PALDI-MSI data confirmed that under a supply of NH_4^+ , PAT was impaired, thus promoting a putative local increase in IAA presumably in RC and/or RM tissues where cambial development takes place based on IAA-related transcripts and TF expression (Figures 5a and 7), which could suggest a different regulatory mechanism of

those pH-dependent described in rice and *Arabidopsis* (Jia et al., 2020; Meier et al., 2020). One possible alternative for this PAT alteration could be IAA conjugation with sugars/amino acids as previously described (Di et al., 2021; Tamura et al., 2010). However, our data revealed nonsignificant expression changes of IAA conjugating genes (Dataset S1). This suggests that the NH_4^+ concentration used in this study is not excessive for maritime pine compared with those applied before (>5 mM NH_4^+) in rice and *Arabidopsis* (Di et al., 2021; Tamura et al., 2010), which is supported by the non-inhibition of PR growth under NH_4^+ supply (Figure 6c). Therefore, other mechanisms could be acting during the observed PAT process in the root apex of maritime pine.

In addition, our results differ from those previously described in *Arabidopsis* where the presence of NH_4^+ does not seem to affect the distribution of IAAs in the root tip (Y. Liu et al., 2013). This finding could be related to root vascular development and root cell

differentiation since adjustments to IAA flux are involved in root cambium development (Marhava et al., 2018), and SCR, a partner of SHR, is involved in the expression and polar localization of PINs (Xu et al., 2006).

Following this line of argument, it is interesting to mention that four maritime pine DOFs were significantly downregulated: *PpDOF11* (pp_58878) and *PpDOF12* (pp_85381) in RC and RM tissues respectively, and *PpDOF1* (pp_233816) and *PpDOF2* (pp_119525) in RDC (Figure 2), which were preferentially expressed in tissues close to vascular vessels (Figures S9–S12). Therefore, it is tempting to suggest that these DOFs may be involved in the development of root vessels, as described for PHLOEM EARLY DOF (PEAR) proteins, which are regulated by CK levels (Miyashima et al., 2019). This is also supported by the network analysis and the functions of the most correlated transcripts to these DOFs (Figure S6B). Interestingly, the CK distribution under NH_4^+ supply (Figure 5B) correlates with the

repression of CK riboside 5'-monophosphate phosphoribohydrolase (pp_222714), which is a CK activator, in RC (Kuroha et al., 2009) and the upregulation of a CK dehydrogenase (pp_215376), which is involved in CK degradation (Werner et al., 2003), in RDC (Figure 3a,c). All these results suggest that NH_4^+ could promote an alteration of CK spatial distribution in maritime pine roots, which could be the reason for the wide repression of DOF TFs in RC and RM tissues and its possible role as regulators of cambium development (Figure 7). Phytohormone imaging showed that CK and IAA were distributed in different zones under a supply of NH_4^+ (Figure 5a,b), which was possibly due to the opposite roles of these phytohormones in the root meristem since CK negatively regulates QC specification and functions by modulating the IAA response (Zhang et al., 2013).

Additionally, homeobox-leucine zipper (HD-Zip) and NAC-domain TFs are described to take part during development processes (Bennett et al., 2010; Ochando et al., 2006). In *Arabidopsis*, several




FIGURE 7 Schematic representation of the transcriptional response induced by NH_4^+ supply in maritime pine seedling roots 24 h after irrigation. Transcripts in red letters are upregulated. Transcripts in blue letters are downregulated. The root tip tissues are the root cap (RC) root meristem (RM) and root developing cortex (RDC). Results for root developing vessels (RDV) are not shown since they were very limited [Color figure can be viewed at wileyonlinelibrary.com]

HATs play important roles in development including the regulation of meristem and procambial maintenance and/or formation (Prigge et al., 2005; Roodbarkelari & Groot, 2017). Accordingly, two HD-Zip coding transcripts identified as *HAT-like* (pp_141831) and *INCURVATA-like* (pp_85012) were downregulated in RM tissue (Table 1). These findings are consistent with the repression of TFs related to the root vascular development process and suggest that NH_4^+ could affect PR vascular development during early exposure (Figure 7). Nevertheless, long-term NH_4^+ nutrition promotes PR growth but causes a decrease in PR weight (Figure 6b,c), suggesting that root radial growth could be affected by this transcriptional network. In future studies, cell division and elongation should be analyzed in the root these zones. Unfortunately, the exact anatomy of root meristem with the quiescent center position has not yet been defined in conifers.

Interestingly, in RM, the upregulation of two NAC TFs, *PpNAC31* and *PpNAC38*, similar to the *Arabidopsis* *SMB* and *BRN* genes (Table 1, Figure S4), is consistent with the expression patterns of *SMB*, *BRN1* and *BRN2*, which largely overlap in *Arabidopsis* roots (Bennett et al., 2010). *SMB* represses stem cell-like divisions in the root cap daughter cells (Bennett et al., 2014; Kamiya et al., 2016), and together with *BRN1* and *BRN2*, it regulates cellular maturation of the root cap in *Arabidopsis* (Bennett et al., 2010). These findings are consistent with the reduction in the number of root cap cells under NH_4^+ , as described by Y. Liu et al. (2013), thus providing a possible explanation for this observation in *Arabidopsis*.

In rice, the endogenous ABA content alleviates NH_4^+ rice toxicity, promoting NH_4^+ assimilation by stimulating the GS/GOGAT cycle and enhancing antioxidant activities (superoxide dismutase, ascorbate peroxidase and catalase) (Sun et al., 2020). In the maritime pine root apex under NH_4^+ supply, ABA levels increased (Figure 5d), and transcripts encoding ABA2 enzymes (e.g., pp_198956) involved in ABA biosynthesis were upregulated in RM and RDC, and several putative ABA-induced TFs (e.g., pp_233345) were upregulated in RC and RM (Dataset S1). However, the GS/GOGAT cycle was apparently not induced, although it was observed in the whole root response to NH_4^+ supply (Ortigosa et al., 2021) (Figure 1). In general, there was no expression changes in genes involved in N uptake and assimilation at the root apex of pine in response to NH_4^+ supply. In angiosperms, the upregulation of GS/GOGAT cycle has been described in specific tissues such as root epidermis around root hair zone in rice (Ishiyama et al., 1998; Ishiyama, Inoue, Tabuchi, et al., 2004) and *Arabidopsis* (IshiyamaInoue, Inoue, Watanabe-Takahashi, et al., 2004; Konishi et al., 2017). This raises the question of whether conifers might have a different response to NH_4^+ availability at the root apex than angiosperms.

Regarding antioxidant metabolism, different transcripts involved in ascorbate production were downregulated (Dataset S1), including *PpBLISTER-like* TF (pp_170664) in RDC tissue (Purdy et al., 2011) and two *L-gulonolactone oxidases* (pp_183087 and pp_234650) in RC and RDC tissues (Aboobucker et al., 2017). This finding suggests that the glutathione-ascorbate cycle is not operative in maritime pine roots exposed to NH_4^+ , which is consistent with the levels of oxidized glutathione observed under long-term experiments (Ortigosa et al., 2020).

Finally, ET is involved in NH_4^+ -responsive pathways in rice (Sun et al., 2017) and in cambial development in poplar trees (Love et al., 2009). In *Arabidopsis*, the ET levels increased under a supply of NH_4^+ , which reduced lateral root formation because of the repression of auxin transporter 1 (*AUX1*) (Li et al., 2016). In contrast, the results in maritime pine suggest a reduction in ET biosynthesis following NH_4^+ application. A transcript encoding an ACO was repressed in RM, which is consistent with proteomic and epitranscriptomic results in maritime pine whole roots (Ortigosa et al., 2021). This finding is consistent with the downregulation of *BLISTER-like* TF and *L-gulonolactone oxidases* since ascorbate is one of the ACO enzyme substrates (Dilley et al., 2013). However, ACC was not affected by NH_4^+ treatment (Figure 5c). Accordingly, NH_4^+ stimulated the LR components (number and weight) and PR length (Figure 5) in maritime pine, which is consistent with the increased number of root apices under NH_4^+ nutrition compared to NO_3^- nutrition in *Pinus massoniana* Lamb. (Ren et al., 2020). Taken together, the results suggest that maritime pine reduces ET biosynthesis under NH_4^+ supply, which could be linked to the PAT and CK alteration observed (Figure 4a,b) as well as to the expression patterns observed for TFs related to cambium development (Table 1) due to the importance of IAA-ET-CK crosstalk during root development (J. Liu et al., 2017).

5 | CONCLUSIONS

NH_4^+ nutrition is a complex process in plants since NH_4^+ is a nutrient that can be toxic when supplied in excess. However, conifers in general are quite tolerant to NH_4^+ . Therefore, the identifying of the differential mechanisms underlying NH_4^+ tolerance in conifers would be enormously helpful for improving NH_4^+ tolerance not only in these trees but also in other crops. In contrast to other plant models, supplying NH_4^+ to pine represses ET and ascorbate production and alters IAA and CK patterns, thus regulating a transcriptional network in the short term. Consequently, the growth and development of PR and LRs are stimulated in pine.

The expression profiles of genes related to root cambium development and phytohormones suggest a molecular mechanism underlying changes in the RSA phenotype that includes IAA-CK-ET crosstalk and a transcriptional network at least during the early root response to NH_4^+ supply. The results reported in the present study tentatively link *SHR*, and other TFs to NH_4^+ nutrition and its phenotypic effect on RSA, which likely affect early vascular development (Figure 7).

This study provides new and valuable data to unravel the mechanisms involved in the response of maritime pine to NH_4^+ nutrition. However, further research efforts are required to reach a full understanding of the molecular basis of NH_4^+ tolerance.

ACKNOWLEDGEMENTS

This study was funded by Spanish *Ministerio de Ciencia e Innovación*, grant numbers BIO2015-73512-JIN MINECO/AEI/FEDER, UE; RTI2018-094041-B-I00 and EQC2018-004346-P. Funding for open

access charge: Universidad de Málaga/CBUA. FO was supported by grants from the Universidad de Málaga (Programa Operativo de Empleo Juvenil vía SNJG, UMAJ11, FEDER, FSE, Junta de Andalucía) and BIO-114, Junta de Andalucía.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from different databases, supporting information and from the corresponding author upon reasonable request. RNA-Seq data have been deposited in the NCBI's Gene Expression Omnibus and are accessible through GEO Series with the accession number GSE175587 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE175587>). The transcriptome shotgun assembly project has been deposited at DDBJ/EMBL/GenBank under the accession GJFX00000000. The version described in this paper is the first version, GJFX01000000.

ORCID

Francisco Ortigosa  <http://orcid.org/0000-0002-7517-7594>

César Lobato-Fernández  <http://orcid.org/0000-0001-9576-7236>

Hitomi Shikano  <http://orcid.org/0000-0001-8973-3355>

Concepción Ávila  <http://orcid.org/0000-0001-8817-7529>

Shu Taira  <http://orcid.org/0000-0002-9243-0395>

Francisco M. Cánovas  <http://orcid.org/0000-0002-4914-2558>

Rafael A. Cañas  <http://orcid.org/0000-0001-9727-5585>

REFERENCES

- Aboobucker, S.I., Suza, W.P. & Lorence, A. (2017) Characterization of two *Arabidopsis* L-gulonolactone oxidases, AtGULO3 and AtGULO5, involved in ascorbate biosynthesis. *Reactive Oxygen Species (Apex, NC)*, 4(12), 389–417. Available from: <https://doi.org/10.20455/ros.2017.861>
- Antolín-Llovera, M., Leivar, P., Arró, M., Ferrer, A., Boronat, A. & Campos, N. (2011) Modulation of plant HMG-CoA reductase by protein phosphatase 2A: positive and negative control at a key node of metabolism. *Plant Signaling & Behavior*, 6(8), 1127–1131. Available from: <https://doi.org/10.4161/psb.6.8.16363>
- Araya, T., Kubo, T., von Wirén, N. & Takahashi, H. (2016) Statistical modeling of nitrogen-dependent modulation of root system architecture in *Arabidopsis thaliana*. *Journal of Integrative Plant Biology*, 58(3), 254–265. Available from: <https://doi.org/10.1111/jipb.12433>
- Bak, S., Beisson, F., Bishop, G., Hamberger, B., Höfer, R. & Paquette, S. et al. (2011) Cytochromes P450. In: *The Arabidopsis Book*/American Society of Plant Biologists. Vol. 9. Rockville, MD: American Society of Plant Biologists. <https://doi.org/10.1199/tab.0144>
- Bennett, T., van den Toorn, A., Sanchez-Perez, G.F., Campilho, A., Willemsen, V., Snel, B. et al. (2010) SOMBRERO, BEARSKIN1, and BEARSKIN2 regulate root cap maturation in *Arabidopsis*. *The Plant Cell*, 22(3), 640–654. Available from: <https://doi.org/10.1105/tpc.109.072272>
- Bennett, T., van den Toorn, A., Willemsen, V. & Scheres, B. (2014) Precise control of plant stem cell activity through parallel regulatory inputs. *Development*, 141(21), 4055–4064. Available from: <https://doi.org/10.1242/dev.110148>
- Bijlsma, R.J., Lambers, H. & Kooijman, S.A.L.M. (2000) A dynamic whole-plant model of integrated metabolism of nitrogen and carbon. 1. Comparative ecological implications of ammonium-nitrate interactions. *Plant and Soil*, 220(1), 49–69.
- Brady, S.M., Orlando, D.A., Lee, J.Y., Wang, J.Y., Koch, J., Dinneny, J.R. et al. (2007) A high-resolution root spatiotemporal map reveals dominant expression patterns. *Science*, 318(5851), 801–806. Available from: <https://doi.org/10.1126/science.1146265>
- Buchfink, B., Xie, C. & Huson, D.H. (2015) Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, 12(1), 59–60. Available from: <https://doi.org/10.1038/nmeth.3176>
- Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M. et al. (2009) The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clinical Chemistry*, 55(4), 611–22. Available from: <https://doi.org/10.1373/clinchem.2008.112797>
- Cánovas, F.M., Ávila, C., Cantón, F.R., Cañas, R.A. & de la Torre, F. (2007) Ammonium assimilation and amino acid metabolism in conifers. *Journal of Experimental Botany*, 58(9), 2307–2318. Available from: <https://doi.org/10.1093/jxb/erm051>
- Canales, J., Ávila, C. & Cánovas, F.M. (2011) A maritime pine antimicrobial peptide involved in ammonium nutrition. *Plant, Cell & Environment*, 34(9), 1443–1453. Available from: <https://doi.org/10.1111/j.1365-3040.2011.02343.x>
- Canales, J., Flores-Monterroso, A., Rueda-López, M., Ávila, C. & Cánovas, F.M. (2010) Identification of genes regulated by ammonium availability in the roots of maritime pine trees. *Amino Acids*, 39(4), 991–1001. Available from: <https://doi.org/10.1007/s00726-010-0483-9>
- Canales, J., Rueda-López, M., Craven-Bartle, B., Ávila, C. & Cánovas, F.M. (2012) Novel insights into regulation of asparagine synthetase in conifers. *Frontiers in Plant Science*, 3, 100. Available from: <https://doi.org/10.1007/s00726-010-0483-9>
- Cañas, R.A., Canales, J., Gómez-Maldonado, J., Ávila, C. & Cánovas, F.M. (2014) Transcriptome analysis in maritime pine using laser capture microdissection and 454 pyrosequencing. *Tree Physiology*, 34(11), 1278–1288. Available from: <https://doi.org/10.1093/treephys/tpt113>
- Cañas, R.A., de la Torre, F., Cánovas, F.M. & Cantón, F.R. (2006) High levels of asparagine synthetase in hypocotyls of pine seedlings suggest a role of the enzyme in re-allocation of seed-stored nitrogen. *Planta*, 224(1), 83–95. Available from: <https://doi.org/10.1007/s00425-005-0196-6>
- Cañas, R.A., Li, Z., Pascual, M.B., Castro-Rodríguez, V., Ávila, C., Sterck, L. et al. (2017) The gene expression landscape of pine seedling tissues. *The Plant Journal*, 91(6), 1064–1087. Available from: <https://doi.org/10.1111/tbj.13617>
- Chaban, C., Waller, F., Furuya, M. & Nick, P. (2003) Auxin responsiveness of a novel cytochrome P450 in rice coleoptiles. *Plant Physiology*, 133(4), 2000–2009. Available from: <https://doi.org/10.1104/pp.103.022202>
- Chapman, E.J. & Estelle, M. (2009) Mechanism of auxin-regulated gene expression in plants. *Annual Review of Genetics*, 43, 265–285. Available from: <https://doi.org/10.1146/annurev-genet-102108-134148>
- Coll-García, D., Mazuch, J., Altmann, T. & Müssig, C. (2004) EXORDIUM regulates brassinosteroid-responsive genes. *FEBS Letters*, 563(1–3), 82–86. Available from: [https://doi.org/10.1016/S0014-5793\(04\)00255-8](https://doi.org/10.1016/S0014-5793(04)00255-8)
- Corrales, A.R., Carrillo, L., Lasiera, P., Nebauer, S.G., Dominguez-Figueroa, J., Renau-Morata, B. et al. (2017) Multifaceted role of cycling DOF factor 3 (CDF3) in the regulation of flowering time and abiotic stress responses in *Arabidopsis*. *Plant, Cell & Environment*, 40(5), 748–764. Available from: <https://doi.org/10.1111/pce.12894>
- Crook, A.D., Willoughby, A.C., Hazak, O., Okuda, S., VanDerMolen, K.R., Soyars, C.L. et al. (2020) BAM1/2 receptor kinase signaling drives CLE peptide-mediated formative cell divisions in *Arabidopsis* roots. *Proceedings of the National Academy of Sciences*, 117(51),

- 32750–32756. Available from: <https://doi.org/10.1073/pnas.2018565117>
- Cui, D., Zhao, J., Jing, Y., Fan, M., Liu, J., Wang, Z. et al. (2013) The *Arabidopsis* IDD14, IDD15, and IDD16 cooperatively regulate lateral organ morphogenesis and gravitropism by promoting auxin biosynthesis and transport. *PLoS Genetics*, 9(9), e1003759. Available from: <https://doi.org/10.1371/journal.pgen.1003759>
- Cui, H., Levesque, M.P., Vernoux, T., Jung, J.W., Paquette, A.J., Gallagher, K.L. et al. (2007) An evolutionarily conserved mechanism delimiting SHR movement defines a single layer of endodermis in plants. *Science*, 316(5823), 421–425. Available from: <https://doi.org/10.1126/science.1139531>
- Dharmasiri, N., Dharmasiri, S., Weijers, D., Lechner, E., Yamada, M., Hobbie, L., Ehrismann, J.S., Jürgens, G. & Estelle, M. (2005) Plant development is regulated by a family of auxin receptor F box proteins. *Developmental Cell*, 9(1), 109–119. Available from: <https://doi.org/10.1016/j.devcel.2005.05.014>
- Di, D.-W., Li, G., Sun, L., Wu, J., Wang, M., Kronzucker, H.J. et al. (2021) High ammonium inhibits root growth in *Arabidopsis thaliana* by promoting auxin conjugation rather than inhibiting auxin biosynthesis. *Journal of Plant Physiology*, 261, 153415. Available from: <https://doi.org/10.1016/j.jplph.2021.153415>
- Dilley, D.R., Wang, Z., Kadirjan-Kalbach, D.K., Ververidis, F., Beaudry, R. & Padmanabhan, K. (2013) 1-Aminocyclopropane-1-carboxylic acid oxidase reaction mechanism and putative post-translational activities of the ACCO protein. *AoB Plants*, 5, Available from: <https://doi.org/10.1093/aobpla/plt031>
- Domínguez-Figueroa, J., Carrillo, L., Renau-Morata, B., Yang, L., Molina, R.V., Marino, D. et al. (2020) The *Arabidopsis* transcription factor CDF3 is involved in nitrogen responses and improves nitrogen use efficiency in tomato. *Frontiers in Plant Science*, 11, 1825. Available from: <https://doi.org/10.3389/fpls.2020.601558>
- Edgar, R., Domrachev, M. & Lash, A.E. (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Research*, 30(1), 207–210. Available from: <https://doi.org/10.1093/nar/30.1.207>
- Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32(5), 1792–1797. Available from: <https://doi.org/10.1093/nar/gkh340>
- Esteban, R., Ariz, I., Cruz, C. & Moran, J.F. (2016) Mechanisms of ammonium toxicity and the quest for tolerance. *Plant Science*, 248, 92–101. Available from: <https://doi.org/10.1016/j.plantsci.2016.04.008>
- Farjon, A. (2018) The Kew review: conifers of the world. *Kew Bulletin*, 73, 8. Available from: <https://doi.org/10.1007/s12225-018-9738-5>
- Felsenstein, J. (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, 39, 783–791. Available from: <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>
- Fendrych, M., Van Hautegeem, T., Van Durme, M., Olvera-Carrillo, Y., Huysmans, M., Karimi, M. et al. (2014) Programmed cell death controlled by ANAC033/SOMBRERO determines root cap organ size in *Arabidopsis*. *Current Biology*, 24(9), 931–940. Available from: <https://doi.org/10.1016/j.cub.2014.03.025>
- Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. (2012) CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics*, 28(23), 3150–3152. Available from: <http://doi.org/10.1093/bioinformatics/bts565>
- Fujita, S., Ohnishi, T., Watanabe, B., Yokota, T., Takatsuto, S., Fujioka, S. et al. (2006) *Arabidopsis* CYP90B1 catalyses the early C-22 hydroxylation of C27, C28 and C29 sterols. *The Plant Journal*, 45(5), 765–774. Available from: <https://doi.org/10.1111/j.1365-313X.2005.02639.x>
- Fukaki, H., Wysocka-Diller, J., Kato, T., Fujisawa, H., Benfey, P.N. & Tasaka, M. (1998) Genetic evidence that the endodermis is essential for shoot gravitropism in *Arabidopsis thaliana*. *The Plant Journal*, 14(4), 425–430. Available from: <https://doi.org/10.1046/j.1365-313x.1998.00137.x>
- Galinha, C., Hofhuis, H., Luitjen, M., Willemsen, V., Blilou, I., Heidstra, R. et al. (2007) PLETHORA proteins as dose-dependent master regulators of *Arabidopsis* root development. *Nature*, 449(7165), 1053–1057. Available from: <https://doi.org/10.1038/nature06206>
- Gifford, M.L., Dean, A., Gutierrez, R.A., Coruzzi, G.M. & Birnbaum, K.D. (2008) Cell-specific nitrogen responses mediate developmental plasticity. *Proceedings of the National Academy of Sciences*, 105(2), 803–808. Available from: <https://doi.org/10.1073/pnas.0709559105>
- González-Guzmán, M., Apostolova, N., Bellés, J.M., Barrero, J.M., Piqueras, P., Ponce, M.R., Micol, J.L., Serrano, R. & Rodríguez, P.L. (2002) The short-chain alcohol dehydrogenase ABA2 catalyzes the conversion of xanthoxin to abscisic aldehyde[W]. *The Plant Cell*, 14(8), 1833–1846. Available from: <https://doi.org/10.1105/tpc.002477>
- Götz, S., García-Gómez, J.M., Terol, J., Williams, T.D., Nagaraj, S.H., Nueda, M.J. et al. (2008) High-throughput functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Research*, 36(10), 3420–3435. Available from: <https://doi.org/10.1093/nar/gkn176>
- Granados, J.M., Ávila, C., Cánovas, F.M. & Cañas, R.A. (2016) Selection and testing of reference genes for accurate RT-qPCR in adult needles and seedlings of maritime pine. *Tree Genetics & Genomes*, 12(3), 1–15. Available from: <https://doi.org/10.1007/s11295-016-1018-7>
- Greb, T. (2020) Plant development: how phloem patterning occurs. *Current Biology*, 30, 217–219. Available from: <https://doi.org/10.1016/j.cub.2020.01.021>
- Grunewald, W. & Friml, J. (2010) The march of the PINs: developmental plasticity by dynamic polar targeting in plant cells. *The EMBO Journal*, 29(16), 2700–2714. Available from: <https://doi.org/10.1038/emboj.2010.181>
- Guo, Y., Qin, G., Gu, H. & Qu, L.J. (2009) Dof5. 6/HCA2, a Dof transcription factor gene, regulates interfascicular cambium formation and vascular tissue development in *Arabidopsis*. *The Plant Cell*, 21(11), 3518–3534. Available from: <https://doi.org/10.1105/tpc.108.064139>
- Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J. et al. (2013) De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols*, 8(8), 1494–1512. Available from: <https://doi.org/10.1038/nprot.2013.084>
- Hachiya, T. & Sakakibara, H. (2017) Interactions between nitrate and ammonium in their uptake, allocation, assimilation, and signaling in plants. *Journal of Experimental Botany*, 68(10), 2501–2512. Available from: <https://doi.org/10.1093/jxb/erw449>
- Hashiguchi, T., Sakakibara, Y., Hara, Y., Shimohira, T., Kurogi, K., Akashi, R., Liu, M.C. & Suiko, M. (2013) Identification and characterization of a novel kaempferol sulfotransferase from *Arabidopsis thaliana*. *Biochemical and Biophysical Research Communications*, 434(4), 829–835. Available from: <https://doi.org/10.1016/j.bbrc.2013.04.022>
- Hawkins, B.J. & Robbins, S. (2010) pH affects ammonium, nitrate and proton fluxes in the apical region of conifer and soybean roots. *Physiologia Plantarum*, 138(2), 238–247. Available from: <https://doi.org/10.1111/j.1399-3054.2009.01317.x>
- Helariutta, Y., Fukaki, H., Wysocka-Diller, J., Nakajima, K., Jung, J., Sena, G. et al. (2000) The SHORT-ROOT gene controls radial patterning of the *Arabidopsis* root through radial signaling. *Cell*, 101(5), 555–567. Available from: [https://doi.org/10.1016/S0092-8674\(00\)80865-X](https://doi.org/10.1016/S0092-8674(00)80865-X)
- Hong, Z.H., Qing, T., Schubert, D., Kleinmanns, J.A. & Liu, J.X. (2019) BLISTER-regulated vegetative growth is dependent on the protein kinase domain of ER stress modulator IRE1A in *Arabidopsis thaliana*.

- PLOS Genetics*, 15(12), e1008563. Available from: <https://doi.org/10.1371/journal.pgen.1008563>
- Hou, H., Erickson, J., Meservy, J. & Schultz, E.A. (2010) FORKED1 encodes a PH domain protein that is required for PIN1 localization in developing leaf veins. *The Plant Journal*, 63(6), 960–973. Available from: <https://doi.org/10.1111/j.1365-313X.2010.04291.x>
- Houben, M. & Van de Poel, B. (2019) 1-Aminocyclopropane-1-carboxylic acid oxidase (ACO): the enzyme that makes the plant hormone ethylene. *Frontiers in Plant Science*, 10, 695. Available from: <https://doi.org/10.3389/fpls.2019.00695>
- Ishiyama, K., Hayakawa, T. & Yamaya, T. (1998) Expression of NADH-dependent glutamate synthase protein in the epidermis and exodermis of rice roots in response to the supply of ammonium ions. *Planta*, 204(3), 288–294. Available from: <https://doi.org/10.1007/s004250050258>
- Ishiyama, K., Inoue, E., Tabuchi, M., Yamaya, T. & Takahashi, H. (2004) Biochemical background and compartmentalized functions of cytosolic glutamine synthetase for active ammonium assimilation in rice roots. *Plant and Cell Physiology*, 45(11), 1640–1647. Available from: <https://doi.org/10.1093/pcp/pch190>
- Ishiyama, K., Inoue, E., Watanabe-Takahashi, A., Obara, M., Yamaya, T. & Takahashi, H. (2004) Kinetic properties and ammonium-dependent regulation of cytosolic isoenzymes of glutamine synthetase in *Arabidopsis*. *Journal of Biological Chemistry*, 279(16), 16598–16605. Available from: <https://doi.org/10.1074/jbc.M313710200>
- Jia, L.T., Xie, Y.M., Wang, Z., Luo, L., Zhang, C., Pelissier, P.M. et al. (2020) Rice plants respond to ammonium stress by adopting a helical root growth pattern. *The Plant Journal*, 104(4), 1023–1037. Available from: <https://doi.org/10.1111/tpj.14978>
- Jia, Z. & von Wirén, N. (2020) Signaling pathways underlying nitrogen-dependent changes in root system architecture: from model to crop species. *Journal of Experimental Botany*, 71(15), 4393–4404. Available from: <https://doi.org/10.1093/jxb/eraa033>
- Kamada-Nobusada, T., Makita, N., Kojima, M. & Sakakibara, H. (2013) Nitrogen-dependent regulation of de novo cytokinin biosynthesis in rice: the role of glutamine metabolism as an additional signal. *Plant and Cell Physiology*, 54(11), 1881–1893. Available from: <https://doi.org/10.1093/pcp/pct127>
- Kamiya, M., Higashio, S.Y., Isomoto, A., Kim, J.M., Seki, M., Miyashima, S. et al. (2016) Control of root cap maturation and cell detachment by BEARSKIN transcription factors in *Arabidopsis*. *Development*, 143(21), 4063–4072. Available from: <https://doi.org/10.1242/dev.142331>
- Kim, H., Zhou, J., Kumar, D., Jang, G., Ryu, K.H., Sebastian, J. et al. (2020) SHORTRoot-mediated intercellular signals coordinate phloem development in *Arabidopsis* roots. *The Plant Cell*, 32(5), 1519–1535. Available from: <https://doi.org/10.1105/tpc.19.00455>
- Konishi, N., Ishiyama, K., Beier, M.P., Inoue, E., Kanno, K., Yamaya, T. et al. (2017) Contribution of two glutamine synthetase isozymes to ammonium assimilation in *Arabidopsis* roots. *Journal of Experimental Botany*, 68(3), 613–625. Available from: <https://doi.org/10.1093/jxb/erw454>
- Kronzucker, H.J., Siddiqi, M.Y. & Glass, A.D. (1997) Conifer root discrimination against soil nitrate and the ecology of forest succession. *Nature*, 385(6611), 59–61. Available from: <https://doi.org/10.1038/385059a0>
- Kumar, S., Stecher, G. & Tamura, K. (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870–1874. Available from: <https://doi.org/10.1093/molbev/msw054>
- Kuroha, T., Tokunaga, H., Kojima, M., Ueda, N., Ishida, T., Nagawa, S. et al. (2009) Functional analyses of LONELY GUY cytokinin-activating enzymes reveal the importance of the direct activation pathway in *Arabidopsis*. *The Plant Cell*, 21(10), 3152–3169. Available from: <https://doi.org/10.1105/tpc.109.068676>
- Langfelder, P. & Horvath, S. (2008) WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics*, 9, 559. Available from: <https://doi.org/10.1186/1471-2105-9-559>
- Li, G., Kronzucker, H.J. & Shi, W. (2016) Root developmental adaptation to Fe toxicity: mechanisms and management. *Plant Signaling & Behavior*, 11(1), e1117722. Available from: <https://doi.org/10.1080/15592324.2015.1117722>
- Li, H. & Durbin, R. (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, 25(14), 1754–1760. Available from: <https://doi.org/10.1093/bioinformatics/btp324>
- Liao, Z., Chen, M., Guo, L., Gong, Y., Tang, F., Sun, X. et al. (2004) Rapid isolation of high-quality total RNA from *Taxus* and *Ginkgo*. *Preparative Biochemistry and Biotechnology*, 34(3), 209–214. Available from: <https://doi.org/10.1081/PB-200026790>
- Lima, J.E., Kojima, S., Takahashi, H. & von Wirén, N. (2010) Ammonium triggers lateral root branching in *Arabidopsis* in an AMMONIUM TRANSPORTER1; 3-dependent manner. *The Plant Cell*, 22(11), 3621–3633. Available from: <https://doi.org/10.1105/tpc.110.076216>
- Liu, J., Moore, S., Chen, C. & Lindsey, K. (2017) Crosstalk complexities between auxin, cytokinin, and ethylene in *Arabidopsis* root development: From experiments to systems modeling, and back again. *Molecular Plant*, 10(12), 1480–1496. Available from: <https://doi.org/10.1016/j.molp.2017.11.002>
- Liu, Y., Lai, N., Gao, K., Chen, F., Yuan, L. & Mi, G. (2013) Ammonium inhibits primary root growth by reducing the length of meristem and elongation zone and decreasing elemental expansion rate in the root apex in *Arabidopsis thaliana*. *PLOS One*, 8(4), e61031. Available from: <https://doi.org/10.1371/journal.pone.0061031>
- Liu, Y. & von Wirén, N. (2017) Ammonium as a signal for physiological and morphological responses in plants. *Journal of Experimental Botany*, 68(10), 2581–2592. Available from: <https://doi.org/10.1093/jxb/erx086>
- Love, J., Björklund, S., Vahala, J., Hertzberg, M., Kangasjärvi, J. & Sundberg, B. (2009) Ethylene is an endogenous stimulator of cell division in the cambial meristem of *Populus*. *Proceedings of the National Academy of Sciences*, 106(14), 5984–5989. Available from: <https://doi.org/10.1073/pnas.0811660106>
- Lucas, M., Swarup, R., Paponov, I.A., Swarup, K., Casimiro, I., Lake, D. et al. (2011) Short-root regulates primary, lateral, and adventitious root development in *Arabidopsis*. *Plant Physiology*, 155(1), 384–398. Available from: <https://doi.org/10.1104/pp.110.165126>
- Mao, C., He, J., Liu, L., Deng, Q., Yao, X., Liu, C. et al. (2020) OsNAC2 integrates auxin and cytokinin pathways to modulate rice root development. *Plant Biotechnology Journal*, 18(2), 429–442. Available from: <https://doi.org/10.1111/pbi.13209>
- Marhava, P., Bassukas, A.E.L., Zourelidou, M., Kolb, M., Moret, B., Fastner, A. et al. (2018) A molecular rheostat adjusts auxin flux to promote root protophloem differentiation. *Nature*, 558(7709), 297–300. Available from: <https://doi.org/10.1038/s41586-018-0186-z>
- Meier, M., Liu, Y., Lay-Pruitt, K.S., Takahashi, H. & von Wirén, N. (2020) Auxin-mediated root branching is determined by the form of available nitrogen. *Nature Plants*, 6(9), 1136–1145. Available from: <https://doi.org/10.1038/s41477-020-00756-2>
- Men, S., Boutté, Y., Ikeda, Y., Li, X., Palme, K., Stierhof, Y.D. et al. (2008) Sterol-dependent endocytosis mediates post-cytokinetic acquisition of PIN2 auxin efflux carrier polarity. *Nature Cell Biology*, 10(2), 237–244. Available from: <https://doi.org/10.1038/ncb1686>
- Miller, A.J. & Cramer, M.D. (2005) Root nitrogen acquisition and assimilation. In: *Root physiology: From gene to function*. Dordrecht: Springer. pp. 1–36. Available from: https://doi.org/10.1007/1-4020-4099-7_1
- Miyashima, S., Koi, S., Hashimoto, T. & Nakajima, K. (2011) Non-cell-autonomous microRNA165 acts in a dose-dependent manner to

- regulate multiple differentiation status in the *Arabidopsis* root. *Development*, 138(11), 2303–2313. Available from: <https://doi.org/10.1242/dev.060491>
- Miyashima, S., Roszak, P., Sevilem, I., Toyokura, K., Blob, B., Heo, J.O. et al. (2019) Mobile PEAR transcription factors integrate positional cues to prime cambial growth. *Nature*, 565(7740), 490–494. Available from: <https://doi.org/10.1038/s41586-018-0839-y>
- Moritake, S., Taira, S., Ichiyana, Y., Morone, N., Song, S.Y., Hatanaka, T. et al. (2007) Functionalized nano-magnetic particles for an in vivo delivery system. *Journal of Nanoscience and Nanotechnology*, 7(3), 937–944. Available from: <https://doi.org/10.1166/jnn.2007.216>
- Nakajima, K., Sena, G., Nawy, T. & Benfey, P.N. (2001) Intercellular movement of the putative transcription factor SHR in root patterning. *Nature*, 413(6853), 307–311. Available from: <https://doi.org/10.1038/35095061>
- Naramoto, S., Sawa, S., Koizumi, K., Uemura, T., Ueda, T., Friml, J., Nakano, A. & Fukuda, H. (2009) Phosphoinositide-dependent regulation of VAN3 ARF-GAP localization and activity essential for vascular tissue continuity in plants. *Development*, 136(9), 1529–1538. Available from: <https://doi.org/10.1242/dev.030098>
- Näsholm, T., Ekblad, A., Nordin, A., Giesler, R., Högborg, M. & Högborg, P. (1998) Boreal forest plants take up organic nitrogen. *Nature*, 392(6679), 914–916. Available from: <https://doi.org/10.1038/31921>
- NCBI Resource Coordinators. (2016) Database resources of the national center for biotechnology information. *Nucleic Acids Research*, 44(D1), D7–D19.
- Ochando, I., Jover-Gil, S., Ripoll, J.J., Candela, H., Vera, A., Ponce, M.R. et al. (2006) Mutations in the microRNA complementarity site of the INCURVATA4 gene perturb meristem function and adaxialize lateral organs in *Arabidopsis*. *Plant Physiology*, 141(2), 607–619. Available from: <https://doi.org/10.1104/pp.106.077149>
- Ortigosa, F., Lobato-Fernández, C., Pérez-Claros, J.A., Cantón, F.R., Ávila, C., Cánovas, F.M. et al. (2021) Epitranscriptome changes triggered by ammonium nutrition regulate the proteome response of maritime pine roots. *bioRxiv*, Available from: <https://doi.org/10.1101/2021.04.20.440618>
- Ortigosa, F., Valderrama-Martín, J.M., Urbano-Gámez, J.A., García-Martín, M.L., Ávila, C., Cánovas, F.M. et al. (2020) Inorganic nitrogen form determines nutrient allocation and metabolic responses in maritime pine seedlings. *Plants*, 9, 481. Available from: <https://doi.org/10.3390/plants9040481>
- Ötvös, K., Marconi, M., Vega, A., O'Brien, J., Johnson, A., Abualia, R. et al. (2021) Modulation of plant root growth by nitrogen source-defined regulation of polar auxin transport. *The EMBO Journal*, 40(3), e106862. Available from: <https://doi.org/10.15252/embj.2020106862>
- Patterson, K., Cakmak, T., Cooper, A., Lager, I.D.A., Rasmusson, A.G. & Escobar, M.A. (2010) Distinct signalling pathways and transcriptome response signatures differentiate ammonium- and nitrate-supplied plants. *Plant, Cell & Environment*, 33(9), 1486–1501. Available from: <https://doi.org/10.1111/j.1365-3040.2010.02158.x>
- Prigge, M.J., Otsuga, D., Alonso, J.M., Ecker, J.R., Drews, G.N. & Clark, S.E. (2005) Class III homeodomain-leucine zipper gene family members have overlapping, antagonistic, and distinct roles in *Arabidopsis* development. *The Plant Cell*, 17(1), 61–76. Available from: <https://doi.org/10.1105/tpc.104.026161>
- Purdy, S.J., Bussell, J.D., Nelson, D.C., Villadsen, D. & Smith, S.M. (2011) A nuclear-localized protein, KOLD SENSITIV-1, affects the expression of cold-responsive genes during prolonged chilling in *Arabidopsis*. *Journal of Plant Physiology*, 168(3), 263–269. Available from: <https://doi.org/10.1016/j.jplph.2010.07.001>
- Ravazzolo, L., Trevisan, S., Forestan, C., Varotto, S., Sut, S., Dall'Acqua, S. et al. (2020) Nitrate and ammonium affect the overall maize response to nitrogen availability by triggering specific and common transcriptional signatures in roots. *International Journal of Molecular Sciences*, 21(2), 686. Available from: <https://doi.org/10.3390/ijms21020686>
- Ren, Q., Zhou, Y. & Zhou, X. (2020) Combined transcriptome and proteome analysis of masson Pine (*Pinus massoniana* Lamb.) Seedling root in response to nitrate and ammonium supplementations. *International Journal of Molecular Sciences*, 21(20), 7548. Available from: <https://doi.org/10.3390/ijms21207548>
- Rich-Griffin, C., Eichmann, R., Reitz, M.U., Hermann, S., Woolley-Allen, K., Brown, P.E. et al. (2020) Regulation of cell type-specific immunity networks in *Arabidopsis* roots. *Plant Cell*, 32(9), 2742–2762. Available at <https://doi.org/10.1105/tpc.20.00154>
- Ritz, C. & Spiess, A.N. (2008) qpcR: an R package for sigmoidal model selection in quantitative real-time polymerase chain reaction analysis. *Bioinformatics*, 24(13), 1549–1551. Available from: <https://doi.org/10.1093/bioinformatics/btn227>
- Robinson, M.D., McCarthy, D.J. & Smyth, G.K. (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1), 139–140. Available from: <https://doi.org/10.1093/bioinformatics/btp616>
- Roodbarkelari, F. & Groot, E.P. (2017) Regulatory function of homeodomain-leucine zipper (HD-ZIP) family proteins during embryogenesis. *New Phytologist*, 213(1), 95–104. Available from: <https://doi.org/10.1111/nph.14132>
- Rubio, V., Linhares, F., Solano, R., Martín, A.C., Iglesias, J., Leyva, A. et al. (2001) A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. *Genes & Development*, 15(16), 2122–2133. Available from: <https://doi.org/10.1101/gad.204401>
- Sablowski, R. (2011) Plant stem cell niches: from signalling to execution. *Current Opinion in Plant Biology*, 14(1), 4–9. Available from: <https://doi.org/10.1016/j.pbi.2010.08.001>
- Saitou, N. & Nei, M. (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406–425. Available from: <https://doi.org/10.1093/oxfordjournals.molbev.a040454>
- Salehin, M., Bagchi, R. & Estelle, M. (2015) SCFTIR1/AFB-based auxin perception: mechanism and role in plant growth and development. *The Plant Cell*, 27(1), 9–19. Available from: <https://doi.org/10.1105/tpc.114.133744>
- Schatlowski, N., Stahl, Y., Hohenstatt, M.L., Goodrich, J. & Schubert, D. (2010) The CURLY LEAF interacting protein BLISTER controls expression of polycomb-group target genes and cellular differentiation of *Arabidopsis thaliana*. *The Plant Cell*, 22(7), 2291–2305. Available from: <https://doi.org/10.1105/tpc.109.073403>
- Schoenbohm, C., Martens, S., Eder, C., Forkmann, G. & Weisshaar, B. (2000) Identification of the *Arabidopsis thaliana* flavonoid 3'-hydroxylase gene and functional expression of the encoded P450 enzyme. *Biological Chemistry*, 381(8), 749–753. Available from: <https://doi.org/10.1515/BC.2000.095>
- Sena, G., Jung, J.W. & Benfey, P.N. (2004) A broad competence to respond to SHORT ROOT revealed by tissue-specific ectopic expression. *Development*, 131(12), 2817–2826. Available from: <https://doi.org/10.1242/dev.01144>
- Shiono, K. & Taira, S. (2020) Imaging of multiple plant hormones in roots of rice (*Oryza sativa*) using nanoparticle-assisted laser desorption/ionization mass spectrometry. *Journal of Agricultural and Food Chemistry*, 68(24), 6770–6775. Available from: <https://doi.org/10.1021/acs.jafc.0c00749>
- Shtatrnikova, V.Y., Kudryakova, N.V., Kudoyarova, G.R., Korobova, A.V., Akhlyarova, G.R., Danilova, M.N. et al. (2015) Effects of nitrate and ammonium on growth of *Arabidopsis thaliana* plants transformed with the ARR5::GUS construct and a role for cytokinins in suppression of disturbances induced by the

- presence of ammonium. *Russian Journal of Plant Physiology*, 62(6), 741–752. Available from: <https://doi.org/10.1134/S1021443715060151>
- Smit, M.E., Llavata-Peris, C.I., Roosjen, M., van Beijnum, H., Novikova, D., Levitsky, V. et al. (2020) Specification and regulation of vascular tissue identity in the *Arabidopsis* embryo. *Development*, 147(8), dev186130. Available from: <https://doi.org/10.1242/dev.186130>
- Steyn, Q.J. & Schultz, E.A. (2003) The FORKED genes are essential for distal vein meeting in *Arabidopsis*. *Development*, 130(19), 4695–4708. Available from: <https://doi.org/10.1242/dev.00689>
- Sun, L., Di, D., Li, G., Kronzucker, H.J. & Shi, W. (2017) Spatio-temporal dynamics in global rice gene expression (*Oryza sativa* L.) in response to high ammonium stress. *Journal of Plant Physiology*, 212, 94–104. Available from: <https://doi.org/10.1016/j.jplph.2017.02.006>
- Sun, L., Di, D.W., Li, G., Kronzucker, H.J., Wu, X. & Shi, W. (2020) Endogenous ABA alleviates rice ammonium toxicity by reducing ROS and free ammonium via regulation of the SAPK9–bZIP20 pathway. *Journal of Experimental Botany*, 71(15), 4562–4577. Available from: <https://doi.org/10.1093/jxb/eraa076>
- Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. (2011) REVIGO summarizes and visualizes long lists of gene ontology terms. *PLoS One*, 6(7), e21800. Available from: <https://doi.org/10.1371/journal.pone.0021800>
- Takei, K., Ueda, N., Aoki, K., Kuromori, T., Hirayama, T., Shinozaki, K. et al. (2004) AtIPT3 is a key determinant of nitrate-dependent cytokinin biosynthesis in *Arabidopsis*. *Plant and Cell Physiology*, 45(8), 1053–1062. Available from: <https://doi.org/10.1093/pcp/pch119>
- Tamura, W., Hidaka, Y., Tabuchi, M., Kojima, S., Hayakawa, T., Sato, T. et al. (2010) Reverse genetics approach to characterize a function of NADH-glutamate synthase1 in rice plants. *Amino Acids*, 39(4), 1003–1012. Available from: <https://doi.org/10.1007/s00726-010-0531-5>
- Tan, W., Zhang, D., Zhou, H., Zheng, T., Yin, Y. & Lin, H. (2018) Transcription factor HAT1 is a substrate of SnRK2.3 kinase and negatively regulates ABA synthesis and signaling in *Arabidopsis* responding to drought. *PLoS Genetics*, 14(4), e1007336. Available from: <https://doi.org/10.1371/journal.pgen.1007336>
- Tian, T., Liu, Y., Yan, H., You, Q., Yi, X., Du, Z. et al. (2017) agriGO v2.0: a GO analysis toolkit for the agricultural community, 2017 update. *Nucleic Acids Research*, 45(W1), W122–W129. Available from: <https://doi.org/10.1093/nar/gkx382>
- Walker, L., Boddington, C., Jenkins, D., Wang, Y., Grønland, J.T., Hulsmans, J. et al. (2017) Changes in gene expression in space and time orchestrate environmentally mediated shaping of root architecture. *The Plant Cell*, 29(10), 2393–2412. Available from: <https://doi.org/10.1105/tpc.16.00961>
- Wang, L. & Ruan, Y.-L. (2013) Regulation of cell division and expansion by sugar and auxin signaling. *Frontiers in Plant Science*, 4. Available from: <https://doi.org/10.3389/fpls.2013.00163>
- Wang, S., Chang, Y., Guo, J., Zeng, Q., Ellis, B.E. & Chen, J.G. (2011) *Arabidopsis* ovate family proteins, a novel transcriptional repressor family, control multiple aspects of plant growth and development. *PLoS One*, 6(8), e23896. Available from: <https://doi.org/10.1371/journal.pone.0023896>
- Ward, J.M., Cufir, C.A., Denzel, M.A. & Neff, M.M. (2005) The Dof transcription factor OBP3 modulates phytochrome and cryptochrome signaling in *Arabidopsis*. *The Plant Cell*, 17(2), 475–485. Available from: <https://doi.org/10.1105/tpc.104.027722>
- Warren, C.R. & Adams, M.A. (2002) Possible causes of slow growth of nitrate-supplied *Pinus pinaster*. *Canadian Journal of Forest Research*, 32(4), 569–580. Available from: <https://doi.org/10.1139/x01-225>
- Werner, T., Motyka, V., Laucou, V., Smets, R., Van Onckelen, H. & Schmülling, T. (2003) Cytokinin-deficient transgenic *Arabidopsis* plants show multiple developmental alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem activity. *The Plant Cell*, 15(11), 2532–2550. Available from: <https://doi.org/10.1105/tpc.014928>
- Willemssen, V., Bauch, M., Bennett, T., Campilho, A., Wolkenfelt, H., Xu, J. et al. (2008) The NAC domain transcription factors FEZ and SOMBRERO control the orientation of cell division plane in *Arabidopsis* root stem cells. *Developmental Cell*, 15(6), 913–922. Available from: <https://doi.org/10.1016/j.devcel.2008.09.019>
- Xu, J., Hofhuis, H., Heidstra, R., Sauer, M., Friml, J. & Scheres, B. (2006) A molecular framework for plant regeneration. *Science*, 311(5759), 385–388. Available from: <https://doi.org/10.1126/science.1121790>
- Youssef, C., Bizet, F., Bastien, R., Legland, D., Bogeat-Triboulot, M.B. & Hummel, I. (2018) Quantitative dissection of variations in root growth rate: a matter of cell proliferation or of cell expansion? *Journal of Experimental Botany*, 69(21), 5157–5168. Available from: <https://doi.org/10.1093/jxb/ery272>
- Zhang, D., Ye, H., Guo, H., Johnson, A., Zhang, M., Lin, H. et al. (2014) Transcription factor HAT1 is phosphorylated by BIN2 kinase and mediates brassinosteroid repressed gene expression in *Arabidopsis*. *The Plant Journal*, 77(1), 59–70. Available from: <https://doi.org/10.1111/tpj.12368>
- Zhang, S. & Wang, X. (2017) One new kind of phytohormonal signaling integrator: up-and-coming GASA family genes. *Plant Signaling & Behavior*, 12(2), e1226453. Available from: <https://doi.org/10.1080/15592324.2016.1226453>
- Zhang, W., Swarup, R., Bennett, M., Schaller, G.E. & Kieber, J.J. (2013) Cytokinin induces cell division in the quiescent center of the *Arabidopsis* root apical meristem. *Current Biology*, 23(20), 1979–1989. Available from: <https://doi.org/10.1016/j.cub.2013.08.008>
- Zhou, W., Wei, L., Xu, J., Zhai, Q., Jiang, H., Chen, R. et al. (2010) *Arabidopsis* tyrosylprotein sulfotransferase acts in the auxin/PLETHORA pathway in regulating postembryonic maintenance of the root stem cell niche. *The Plant Cell*, 22(11), 3692–3709. Available from: <https://doi.org/10.1105/tpc.110.075721>
- Zou, N., Li, B., Chen, H., Su, Y., Kronzucker, H.J., Xiong, L. et al. (2013) GSA-1/ARG1 protects root gravitropism in *Arabidopsis* under ammonium stress. *New Phytologist*, 200(1), 97–111. Available from: <https://doi.org/10.1111/nph.12365>
- Zúñiga-Mayo, V.M., Marsch-Martínez, N. & de Folter, S. (2012) The class II HD-ZIP JAIBA gene is involved in meristematic activity and important for gynoecium and fruit development in *Arabidopsis*. *Plant Signaling & Behavior*, 7(11), 1501–1503. Available from: <https://doi.org/10.4161/psb.21901>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Ortigosa, F., Lobato-Fernández, C., Shikano, H., Ávila, C., Taira, S., Cánovas, F.M. et al. (2021) Ammonium regulates the development of pine roots through hormonal crosstalk and differential expression of transcription factors in the apex. *Plant, Cell & Environment*, 1–21. <https://doi.org/10.1111/pce.14214>