

Ferlizza E., Isani G., Dondi F., Andreani G., Vasylyeva K., Bellei E., Almeida A.M., Matzapetakis M. 2020. Urinary proteome and metabolome in dogs (*Canis lupus familiaris*): The effect of chronic kidney disease. Journal of Proteomics 222: 103795. <https://doi.org/10.1016/j.jprot.2020.103795>

# Urinary proteome and metabolome in dogs (*Canis lupus familiaris*): The effect of chronic kidney disease

Author links open overlay

panelEneaFerlizza<sup>a</sup>Glorialsani<sup>b</sup>FrancescoDondi<sup>b</sup>GiuliaAndreani<sup>b</sup>KaterinaVasylyeva<sup>b</sup>ElisaBellei<sup>c</sup>André M.Almeida<sup>d</sup>ManolisMatzapetakis<sup>e</sup>

[Show more](#)

[Add to Mendeley](#)

[Share](#)

[Cite](#)

<https://doi.org/10.1016/j.jprot.2020.103795>[Get rights and content](#)

## Highlights

- 

Urinary proteome and metabolome were studied in healthy and CKD dogs.

- 

Proteomics highlighted a decrease of uromodulin and an increase of albumin.

- 

<sup>1</sup>H NMR evidenced 17 metabolites significantly different between healthy and CKD dogs.

-

Proteomics and metabolomics successfully suggested putative biomarkers for CKD.

## Abstract


Chronic kidney disease (CKD) is a progressive and irreversible disease. Although urine is an ideal biological sample for proteomics and metabolomics studies, sensitive and specific biomarkers are currently lacking in dogs. This study characterised dog urine proteome and metabolome aiming to identify and possibly quantify putative biomarkers of CKD in dogs. Twenty-two healthy dogs and 28 dogs with spontaneous CKD were selected and urine samples were collected. Urinary proteome was separated by SDS-PAGE and analysed by mass spectrometry, while urinary metabolome was analysed in protein-depleted samples by 1D  $^1\text{H}$  NMR spectra. The most abundant proteins in urine samples from healthy dogs were uromodulin, albumin and, in entire male dogs, arginine esterase. In urine samples from CKD dogs, the concentrations of uromodulin and albumin were significantly lower and higher, respectively, than in healthy dogs. In addition, these samples were characterised by a more complex protein pattern indicating mixed glomerular (protein bands  $\geq 65$  kDa) and tubular (protein bands  $< 65$  kDa) proteinuria. Urine spectra acquired by NMR allowed the identification of 86 metabolites in healthy dogs, belonging to 49 different pathways mainly involved in amino acid metabolism, purine and aminoacyl-tRNA biosynthesis or tricarboxylic acid cycle. Seventeen metabolites showed significantly different concentrations when comparing healthy and CKD dogs. In particular, carnosine, trigonelline, and cis-aconitate, might be suggested as putative biomarkers of CKD in dogs.

## Significance

Urine is an ideal biological sample, however few proteomics and metabolomics studies investigated this fluid in dogs and in the context of CKD (chronic kidney disease). In this research, applying a multi-omics approach, new insights were gained regarding the molecular changes triggered by this disease in canine urinary proteome and metabolome. In particular, the involvement of the tubular component was

highlighted, suggesting uromodulin, trigonelline and carnosine as possible biomarkers of CKD in dogs.

## Graphical abstract



1. [Download : Download high-res image \(348KB\)](#)
  2. [Download : Download full-size image](#)
- [Previous](#) article in issue
  - [Next](#) article in issue

## 1. Introduction

Chronic kidney disease (CKD) is a progressive and irreversible disease characterised by the presence of structural or functional abnormalities in one or both kidneys over a period of three months or longer [1]. CKD is one of the most common renal diseases in dogs with an estimated prevalence varying from 0.5 to 3.64% depending on the inclusion criteria of the cases [1], [2], [3]. Early diagnosis of CKD may hinder the disease progression and improve patient quality of life. International Renal Interest Society (IRIS) guidelines for staging and treatment of CKD help clinicians to correctly classify patients and establish the best therapies [4]. Nonetheless, sensitive and specific biomarkers for early detection and monitoring of CKD in dogs are currently lacking. The gold standard to evaluate the renal function is the determination of the glomerular filtration rate (GFR); however, this value does not provide information on CKD aetiology and the available methods for its estimation are difficult to be applied in the routine clinical practice [5,6]. Renal biopsy is considered the gold standard for determining the type of renal damage, but it is an invasive

procedure and not always feasible [7]. Therefore, the assessment of the kidney function is currently based on conventional blood (serum creatinine or urea) and urine (proteinuria and specific gravity) clinicopathological variables, whose alterations are usual findings of CKD, but have limitations when used as early indicators of the disease [7]. For these reasons, other sensitive and specific biomarkers measurable in non- or minimally invasive biological samples are required in clinical practice to identify early renal damage in dogs.

Over the last years, significant efforts have been made in veterinary medicine to apply proteomics to search for new biomarkers or for validating detection methods for proteins already considered as potential early indicators of kidney disease in dogs and cats [[8], [9], [10], [11], [12], [13], [14], [15], [16], [17]]. However, proteins are only some of the molecular species present in urine and a broader approach with the aid of metabolomics can offer additional clinical information.

Metabolomics enables the assessment of a broad range of endogenous and exogenous small molecular mass metabolites, potentially useful to investigate the physiologic status and the pathogenesis of the diseases, and to discover new biomarkers of altered biochemical pathways [[18], [19], [20], [21]]. Metabolites are in general not specific for a single metabolic pathway and in most cases different biochemical reactions contribute to the production of the same metabolite; this peculiarity offers the opportunity to obtain a more comprehensive insight into the complexity of a biological sample. In human medicine, metabolomics was extensively applied to urine to analyse the healthy metabolome [22] and to search for small molecules as potential biomarkers of different diseases, such as immune-mediated inflammatory diseases [23], different cancers [24], [25], [26], and renal diseases [19],[27], [28], [29], [30]]. However, in veterinary medicine, the application of metabolomics techniques to urine is still limited [[31], [32], [33], [34], [35]].

Owing to the metabolic and protein complexity of urine, the aim of this work was to combine the analytical power of proteomics and metabolomics to obtain a more comprehensive characterisation of the urine in healthy dogs and to compare it with the urine of CKD patients with our ultimate goal to suggest new biomarkers of CKD in the canine species.

## **2. Materials and methods**

## 2.1. Animal selection and sample collection

The present study was performed on urine samples collected at the Veterinary Teaching Hospital of the University of Bologna from owned dogs. The dogs were divided into two experimental groups and specimens considered as biological replicates. Upon arrival, all dogs were subjected to physical examination and routine laboratory tests, including complete blood count, serum chemistry and complete urinalysis with urine protein to creatinine ratio (UPC).

Blood samples were collected by venipuncture using a vacuum collection system (Vacutest Kima, Arzergrande, Italy) after at least a 12-h fasting period. Blood samples were processed within one hour after collection. Serum samples were collected in tubes with clot activator (Vacutest Kima, Arzergrande, Italy), centrifuged at 3000 g for 10 min and analysed in an automated chemistry analyser (AU 480, Olympus/Beckman Coulter, Atlanta, GE, USA).

Urine samples were collected by ultrasound-guided cystocentesis. All urine specimens were processed on a routine basis and evaluated in our laboratory within 24 h after collection. In particular, urinalysis consisted in macroscopic examination, urine specific gravity (USG) measured by manual refractometer (American Optical, Buffalo, New York), urine dipstick test (Combur10Test, Roche Diagnostic, Mannheim, Germany) applied on an automated reader (Urisys 1100, Roche Diagnostic, Mannheim, Germany) and microscopic sediment evaluation. Urine sediment was obtained after centrifugation at 500 g for 10 min. Urine supernatants were immediately analysed (dipstick examination), divided in aliquots and stored at  $-80\text{ }^{\circ}\text{C}$  for the subsequent proteomics and metabolomics analysis. Urine chemistry was performed on a refrigerated ( $+4\text{ }^{\circ}\text{C}$ ) aliquot if performed within 24 h after the sample processing or on an aliquot kept frozen at  $-20\text{ }^{\circ}\text{C}$  for a maximum of 7 days. Dogs were considered healthy or diseased on the basis of history, clinical signs and the results of the above-mentioned routine laboratory tests. The control group included 22 healthy dogs presented at the hospital as blood donors. The 22 healthy dogs were 10 males (3 castrated) and 12 females (7 spayed) with an average age of  $37 \pm 20$  months. Mixed-breed dogs were 7/22 (32%), while purebred dogs were 15/22 (68%) (3 Galgo Espanol, 3 Labrador Retriever, 2 Australian Shepherd, 1 Great Dane,

1 Border Collie, 1 Flat-Coated Retriever, 1 German Shepherd, 1 Miniature Pinscher, 1 Boxer, 1 Weimaraner). The diseased group included 28 dogs affected by naturally occurring CKD. The 28 CKD dogs were 14 males (5 castrated) and 14 females (9 spayed) with a mean age of  $111 \pm 61$  months. Mixed-breed dogs were 10/28 (36%), while purebred dogs were 18/28 (64%) (3 Jack Russell Terrier, 2 Chinese Shar-Pei, 2 Labrador Retriever, 2 Cavalier King Charles spaniel, 1 Beagle, 1 Bernese mountain dog, 1 Boxer, 1 German Shepherd, 1 English Springer spaniel, 1 American Staffordshire terrier, 1 Bull terrier, 1 German shorthaired pointer, 1 Lagotto Romagnolo). The diagnosis of CKD was based on history, clinical signs, clinicopathological and imaging results, according to the literature [3,4]. In particular, the presence of clinical findings, abdominal imaging results and (a) persistent pathologic renal proteinuria based on the UPC (UPC > 0.5), assessed and confirmed over a one-month period, and/or (b) serum creatinine (sCrea) concentration  $\geq 1.40$  mg/dL and/or (c) urine specific gravity (USG) <1.030 were considered diagnostic. The IRIS CKD guidelines were used to subsequently stage CKD dogs [4]. Basing on serum creatinine, 8 dogs were classified with CKD stage 1, 6 with stage 2, 9 with stage 3 and 5 with stage 4. On the basis of UPC, 4 dogs were non-proteinuric (UPC < 0.2), 6 dogs were borderline proteinuric (UPC 0.2-0.5) and 18 were proteinuric (UPC > 0.5).

The study was conducted according to the EU Directive 2010/63/EU for animal experiments and approved by the Institutional Scientific Ethical Committee of the University of Bologna for animal testing.

## **2.2. Urine protein to creatinine ratio**

Five mL of urine were collected from each animal by ultrasound-guided cystocentesis. After centrifugation at 500 g for 10 min, urine total proteins and creatinine were measured using commercial kits (Urinary/CSF Protein, OSR6170, and Creatinine OSR6178, Olympus/Beckman Coulter, Atlanta, GE, USA) on an automated chemistry analyser (AU 480, Olympus/Beckman Coulter, Atlanta, GE, USA). The UPC was calculated with the following formula: UPC = urine protein (mg/dL)/urine creatinine (mg/dL).

## **2.3. SDS-PAGE and protein identification**

Urine proteins were separated using an electrophoresis system (NuPAGE, Thermo Fisher Scientific, Waltham, MA, USA) as previously described [8,36]. Briefly, three to five  $\mu\text{g}$  of protein were loaded on 4–12% polyacrylamide gel in MOPS buffer with SDS (Thermo Fisher Scientific, Waltham, MA, USA). The gels were stained with Coomassie brilliant blue (PageBlu protein staining solution; Thermo Fisher Scientific, Waltham, MA, USA) compatible with mass spectrometry analysis. After staining, each gel was digitalised (ChemidocMP, BioRad, Hercules, California, USA) and the pherograms were obtained using a commercial software (ImageLab, BioRad, Hercules, California, USA). The bands at 100, 67 and 18 kDa were cut and identified by electrospray ionization quadrupole time-of-flight mass spectrometry (ESI-Q-TOF/MS) as previously reported [8,36].

To quantify the bands at 100 kDa and 67 kDa, on each sample, one  $\mu\text{g}$  of protein, obtained from a solution containing 1  $\mu\text{g}/\mu\text{L}$  of lactate dehydrogenase (LDH), (Sigma-Aldrich/Merck KGaA, Darmstadt, Germany) was added as internal standard of quantity. The ImageLab software estimated the volume of each protein band based on pixel density within the band boundaries in the digital image. The volume of the band of interest was then compared to the internal standard (LDH) of the corresponding lane and the concentration was calculated as

follows:  $X\text{mg/dL} = V_{\text{band}}/V_{\text{LDH}}/\mu\text{L}_{\text{sample}} * 100$ .

X = concentration of the protein at 100 kDa or at 67 kDa.

$V_{\text{band}}$  = volume of the band at 100 kDa, or at 67 kDa determined by the software.

$V_{\text{LDH}}$  = volume of the band of the internal standard (LDH) determined by the software.

$\mu\text{L}_{\text{sample}}$  =  $\mu\text{L}$  of the sample loaded in the gel.

Subsequently, the respective ratios with urine creatinine (uromodulin [mg/dL]: creatinine [mg/dL], UMC; albumin [mg/dL]: creatinine [mg/dL], UAC) were calculated.

#### 2.4. NMR sample preparation

Urine metabolites were extracted for NMR as follows: 500  $\mu\text{L}$  of urine supernatants were mixed with 550  $\mu\text{L}$  of chloroform and 550  $\mu\text{L}$  of methanol, vortexed for 1 min, left to rest for 15 min at +4 °C and centrifuged at 12000 g for 15 min at room temperature. Nine hundred  $\mu\text{L}$  of the upper phase (urine/methanol) were dried in a

vacuum centrifuge (SpeedVac, Thermo Fischer Scientific, Waltham, MA, USA) overnight at 30 °C. The resulting pellets were suspended with 200 µL of phosphate buffer (PB, 240 mM pH 7.4 in D<sub>2</sub>O with trimethylsilylpropanoic acid [TSP] and sodium azide [NaN<sub>3</sub>]) and 400 µL of D<sub>2</sub>O to a final concentration of 80 mM PB, 0.087 mM TSP and 0.022% (v/v) NaN<sub>3</sub>. Samples were vortexed for 1 min, centrifuged at 12,000 g for 1 min and 560 µL transferred into a 5 mm NMR tubes.

## 2.5. NMR acquisition

NMR spectroscopy was conducted on an 800 MHz spectrometer with a triple resonance HCN Z-gradient probe, at 298 K (Bruker AvanceII+, Ettlingen, Germany). Acquisition and processing were carried out using standard software (Topsin 3.2, Bruker Biospin, Billerica, MA, USA). One dimensional <sup>1</sup>H NMR spectra with Carr-Purcell-Meiboom-Gill (CPMG) filter to attenuate signals from macromolecules were acquired using a standard vendor pulse sequence (cpmgpr1d). Spectra were acquired at 25 °C, with a 20 ppm spectral width, spin lock duration of 78.72 ms, presaturation for 4 s using 20 µW and acquisition time of 2 s. A total of 16 dummy scans and 128 scans were acquired for each sample. All spectra were processed with an exponential window function with 1 Hz line broadening and automated phasing and baseline correction. For the chemometric analysis, the processed data were further processed in the “nmrprocflow” platform [37]. Bins were obtained using manually curated, intelligent binning after referencing, baseline correction, water signal removal and peak alignment. For selected samples, additional homonuclear and heteronuclear spectra (<sup>1</sup>H *J*-resolved, <sup>1</sup>H—<sup>1</sup>H COSY, and <sup>1</sup>H—<sup>13</sup>C HSQC) were also collected to assist with compound identification.

## 2.6. Metabolite annotation and identification

The bins obtained from the “nmrprocflow” platform [37] were annotated with the help of database assisted spectral decomposition using commercial software (Chenomx 8.2 NMR Suit, Edmonton, Alberta, Canada) and the internal reference library (Version 10) as well as the Biological Magnetic Resonance Data Bank (BMRB, <http://www.bmrb.wisc.edu>) reference spectra for compounds absent in the internal reference library. Buckets were attributed to multiple metabolites where peaks



were found to overlap. Pathway analysis module of a free web-based analytical platform (Metaboanalyst 4.0, [www.metaboanalyst.ca](http://www.metaboanalyst.ca)), that used the high-quality Kyoto encyclopaedia of genes and genomes (KEGG) metabolic pathways as the backend knowledgebase, was used to search for the metabolic pathways.

## 2.7. Statistical analysis

Serum and urine chemistry data were analysed with statistical software (R version 3.4.4). Normal distribution was tested graphically and by Shapiro-Wilk normality test, and data were expressed as mean  $\pm$  standard deviation (SD) or median (range; minimum – maximum value) if normally or non-normally distributed, respectively. Variables were compared between healthy ( $N = 22$ ) and CKD ( $N = 28$ ) dogs using the Student *t*-test or the Mann-Whitney *U* test depending on their distribution, assuming  $P < .05$  as a significant probability. The Kruskal-Wallis rank sum test was applied to evaluate differences among healthy and CKD stages (stages 1–4, basing on serum creatinine and according to the IRIS guidelines [4]) and adjusted *P*-values lower than 0.05 were considered statistically significant.

For metabolomics statistical analysis, processed spectra were aligned, baseline corrected and divided into 397 variable width spectral regions or ‘buckets’ with the intensity of each bucket divided by the bucket width. To identify the signals differentially present in the two groups, the buckets were loaded into a web-based platform (Metaboanalyst 4.0, [www.metaboanalyst.ca](http://www.metaboanalyst.ca)) which uses the R package of statistical computing software [38]. For multivariate analysis, buckets were scaled by auto-scaling (mean-centred and divided by the standard deviation of each variable) while, for univariate analysis, and in order to remove the influences attributed to muscle mass and urine concentration, the bucket intensities were normalised to the peak of creatinine (bucket 3.0360 ppm). Both univariate and multivariate statistics were employed. *t*-test and fold change analysis were used to identify the buckets with differential presence, while the list was supplemented with the use of unsupervised principal components analysis (PCA) and supervised partial least squares discriminant analysis (PLS-DA). Both PCA and PLS-DA can identify signals (buckets) whose importance becomes significant via correlated variance. In addition, PCA provides a global view of the differentiability between the two experimental conditions and the

groups of observables that are mostly responsible. In contrast, PLS-DA, since it is a supervised method, highlights the variables most responsible for the differences between groups as previously used in other metabolomics approaches [39,40]. The small sample size that is typical in such studies and the inherent large number of variables obtained may affect the consistency of the multivariate analysis used. To evaluate the consistency of the results, the software performs a number of tests and reports the parameters  $Q^2$  and  $R^2$  as quality parameters of the models.  $Q^2$  indicates the predictive ability of the model, while  $R^2$  is the indicator of the suitability of the fit. For PLSDA  $Q^2 > 0.6$  were selected as acceptable models. Variable importance in projection (VIP) scores greater than 1 and  $t$ -test with a  $P$  value  $<.05$  were used to identify metabolites as differentially expressed.

### 3. Results

#### 3.1. Clinical data

Mean clinical data, serum and urine biochemistry of healthy and CKD dogs are reported in [Table 1](#), while the results for each dog are reported in Supplement Table 1.

Table 1. Clinical data for healthy and CKD dogs. Data are reported as mean  $\pm$  SD or median (range) depending on normal or non-normal distribution, respectively.

| Signalment                 | Healthy (N = 22) | CKD (N = 28)  |                                 |           | P       |
|----------------------------|------------------|---------------|---------------------------------|-----------|---------|
| Age in months              | 37 $\pm$ 20      | 112 $\pm$ 61  |                                 |           | <0.0001 |
| Female n (entire/neutered) | 12 (5/7)         | 14 (5/9)      |                                 |           |         |
| Male n (entire/neutered)   | 10 (7/3)         | 14 (9/5)      |                                 |           |         |
| <b>Serum biochemistry</b>  | <b>Healthy</b>   | <b>CKD</b>    | <b>N (%) CKD &lt; / &gt; RI</b> | <b>RI</b> |         |
| Total Proteins (g/dL)      | 6.4 $\pm$ 0.4    | 6.0 $\pm$ 0.8 | 5 (17.9) < / 1 (3.6) >          | 5.6–7.3   | 0.109   |
| Albumin (g/dL)             | 3.4 $\pm$ 0.3    | 3.0 (1.1–3.8) | 12 (42.9) <                     | 2.8–3.9   | <0.0001 |
| Creatinine (mg/dL)         | 1.1 $\pm$ 0.2    | 2.0 (0.6–9.8) | 1 (3.6) < / 20 (71.4) >         | 0.8–1.4   | <0.0001 |
| Urea (mg/dL)               | 33 $\pm$ 8       | 110 (17–519)  | 22 (78.6) >                     | 17–48     | <0.0001 |

| Signalment                | Healthy (N = 22)    | CKD (N = 28)        |                                 |                      | P       |
|---------------------------|---------------------|---------------------|---------------------------------|----------------------|---------|
| Phosphorus (mg/dL)        | 4.6 ± 0.9           | 4.9 (2.6–14.1)      | 11 (39.3) >                     | 2.7–5.4              | 0.056   |
| <b>Urine biochemistry</b> | <b>Healthy</b>      | <b>CKD</b>          | <b>N (%) CKD &lt; / &gt; RI</b> | <b>RI</b>            |         |
| UPC                       | 0.07 (0.04–0.19)    | 0.78 (0.09–12.8)    | 18 (64.3) >                     | < 0.5                | <0.0001 |
| USG                       | 1.052 (1.034–1.064) | 1.014 (1.006–1.062) | 27 (96.4) <                     | > 1.030 <sup>a</sup> | <0.0001 |
| <b>IRIS Stage</b>         |                     | <b>N (%)</b>        |                                 | <b>RI</b>            |         |
| I                         |                     | 8 (28.6)            |                                 | < 1.4                |         |
| II                        |                     | 6 (21.4)            |                                 | 1.4–2.0              |         |
| III                       |                     | 9 (32.1)            |                                 | 2.1–5.0              |         |
| IV                        |                     | 5 (17.8)            |                                 | > 5.0                |         |

RI, reference intervals; N, number of samples; UPC, urine protein to creatinine ratio; USG, urine specific gravity;


a

Considered as adequate USG in dogs.

CKD dogs were significantly older ( $P < .0001$ ), had significantly higher concentration of serum creatinine ( $P < .0001$ ), urea ( $P < .0001$ ) and UPC ( $P < .0001$ ), while USG ( $P < .0001$ ) was significantly lower than in the healthy dogs. CKD patients were also staged according to serum creatinine concentration following IRIS guidelines [4] and the differences of UPC and USG were evaluated. USG was significantly lower in each CKD stage group than in the healthy dogs ( $P < .01$ ), and samples classified as CKD stage 1 had higher USG than those classified as Stage 3 ( $P = .016$ ) and 4 ( $P = .007$ ). UPC was significantly higher in each CKD stage group than in healthy dogs ( $P < .05$ ), however, no significant differences were found among CKD stages.

### 3.2. SDS-PAGE proteomics analysis

Representative gels and pherograms from healthy and CKD dog urines are reported in [Fig. 1](#).



1. [Download : Download high-res image \(557KB\)](#)
2. [Download : Download full-size image](#)

Fig. 1. Representative SDS-PAGE gels of urine samples from healthy and CKD dogs. Black continuous box indicates uromodulin (103 kDa); black dotted box indicates albumin (67 kDa); black dashed box indicates the internal standard of quantity (1 µg); black dashed and dotted box indicates arginine esterase (18 kDa). M, male; MC, male castrated; F, female; FS, female spayed. LMM, low molecular mass (kDa < 67 kDa); HMM high molecular mass (kDa > 67 kDa). S1–4 under each lane indicate the CKD stage of the patient according to IRIS guideline. NP (non-proteinuric, UPC < 0.2), BP (borderline proteinuric, UPC 0.2–0.5) or P (proteinuric, UPC > 0.5) under each lane indicate the classification of proteinuria according to IRIS guideline.

Urine samples from the healthy group presented similar profiles characterised by the presence of three most abundant bands at apparent molecular mass (MM) of 103, 80 and 67 kDa, respectively. The bands at 103 and 67 kDa were identified by mass spectrometry as uromodulin and albumin, respectively ([Table 2](#)). Moreover, most of the samples presented other three to five low abundance bands at apparent MM between 55 and 14 kDa and two bands at MM < 14 kDa. In addition, urine samples from entire males presented other two evident bands at apparent MM of 18 and 12 kDa. The band at 18 kDa was identified as arginine esterase ([Table 2](#)).

Table 2. Proteins identified in dog urine by mass spectrometry.

| Protein name      | Protein entry name <sup>a</sup> | MM (kDa) <sup>b</sup> | Score <sup>c</sup> | Pept <sup>d</sup> | Sign Pept <sup>e</sup> | Seq <sup>f</sup> | Sign seq <sup>g</sup> |
|-------------------|---------------------------------|-----------------------|--------------------|-------------------|------------------------|------------------|-----------------------|
| Uromodulin        | UROM_CANFA                      | 73                    | 2298               | 138               | 113                    | 15               | 13                    |
| Albumin           | ALBU_CANFA                      | 69                    | 5802               | 470               | 321                    | 44               | 39                    |
| Arginine esterase | ESTA_CANFA                      | 29                    | 532                | 111               | 52                     | 10               | 9                     |

a

Protein entry name from UniProt knowledge database.

b

Theoretical protein molecular mass.

c

The highest scores obtained with Mascot search engine.

d

Peptides: total number of peptides matching the identified proteins.

e

Significant peptides: total number of significant peptides matching the identified proteins.

f

Sequence: total number of distinct sequences matching the identified proteins.

g

Significant sequences: total number of significant distinct sequences matching the identified proteins.

CKD samples presented different and more variable electrophoretic profiles. The disappearance of uromodulin and/or the increase of intensity of albumin and of the band at 80 kDa were clearly evident in all the analysed samples. The increase in number and intensity of the bands at high (>67 kDa) and low (<67 kDa) MM was also evidenced. Particularly, two samples presented an increase in number and intensity of

the bands at high (>67 kDa) MM only, nine samples showed an increase in number and intensity of the bands at low (<67 kDa) MM only, while the remaining 17 samples presented an increase in number and intensity of the bands at both high and low MM. Additionally, in 12 samples ([Fig. 1](#); Lanes 2, 4, 6, 9) was evidenced a band at 21 kDa that was not present in healthy samples.


Concentrations of uromodulin and albumin and their ratio with creatinine (UMC and UAC) are reported in [Table 3](#). Urine samples from healthy dogs presented a low amount of albumin ( $3.1 \pm 1.4$  mg/dL) and a high amount of uromodulin ( $11.9 \pm 2.3$  mg/dL). CKD dogs presented a significantly higher concentration of albumin ( $P = .0025$ ) and UAC value ( $P = .0002$ ) and a significantly lower concentration of uromodulin ( $P < .0001$ ) and UMC value ( $P = .0044$ ), compared to healthy animals.

Table 3. Data for albumin and uromodulin quantification by SDS-PAGE. Data are reported as mean  $\pm$  SD or median (range) depending on normal or non-normal distribution, respectively.

|                           | Healthy           | CKD                 | <i>P</i> |
|---------------------------|-------------------|---------------------|----------|
| <b>Albumin (mg/dL)</b>    | $3.1 \pm 1.4$     | 26.6 (1.4–228.9)    | 0.0025   |
| <b>UAC</b>                | $0.010 \pm 0.007$ | 0.213 (0.028–1.395) | 0.0002   |
| <b>Uromodulin (mg/dL)</b> | $11.9 \pm 2.3$    | 0 (0–5.1)           | <0.0001  |
| <b>UMC</b>                | $0.038 \pm 0.012$ | 0 (0–0.044)         | 0.0044   |

### 3.3. Metabolites annotation and identification





Representative NMR spectra from healthy and CKD dog urine samples are reported in [Fig. 2](#).



1. [Download : Download high-res image \(219KB\)](#)
2. [Download : Download full-size image](#)

Fig. 2. Representative NMR spectra of urine samples collected from healthy and CKD dogs.

An overview of the NMR spectra of samples from healthy dogs evidenced similar profiles, while the urine from CKD patients showed more variable spectra and differences in metabolite abundance. From the 397 buckets, 86 metabolites were identified in healthy samples, with different biological functions and belonging to different pathways. An entire spectrum of the urine of a healthy dog with the assigned metabolites is reported in [Fig. 3](#). The five most abundant metabolites were creatinine, urea, taurine, lactate and 1-methylnicotinamide, while the list of all the identified metabolites is reported in [Table 4](#). After MetaboAnalyst pathway analysis, metabolites were shown as belonging to 49 different pathways, and 23 of these pathways were represented by at least 3 different metabolites. The most represented pathways are mainly involved in amino acid metabolism, purine and aminoacyl-tRNA biosynthesis and tricarboxylic acid cycle ([Table 5](#)). In particular, 10 metabolites belonged to glycine, serine and threonine metabolism and aminoacyl-tRNA biosynthesis, while 8 metabolites were involved in phenylalanine metabolism and purine metabolism.



1. [Download](#) : [Download high-res image \(265KB\)](#)



## 2. Download : Download full-size image

Fig. 3. Representative spectrum of urine from an healthy dog. For a better visualisation, the spectrum has been divided into four parts. a) From 0.0 to 2.8 ppm; b) from 2.4 to 4.9 ppm; c) from 4.6 to 7.0 ppm; d) from 7.0 to 10.0 ppm. The reported metabolites are: 1 valine; 2 fucose; 3 lactate; 4 alanine; 5 acetate; 6 N6-acetyllysine; 7 *N*-acetylglycine; 8 acetylcysteine; 9 succinate; 10 pyridoxamine; 11 citrate; 12 dimethylamine; 13 methylguanidine; 14 trimethylamine; 15 *N,N*-dimethylglycine; 16 creatine; 17 creatinine; 18 choline; 19 phosphorylcholine; 20 carnitine; 21 taurine, trimethylamine N-oxyde, betaine; 22 taurine; 23 trans-aconitate; 24 3-hydroxyphenilacetate; 25 3-methylxantine; 26 2-hydroxyphenilacetate; 27 glycine; 28 *N*-phenilacetylglicine; 29 7-methylxantine; 30 creatine, creatine phosphate, glycolate; 31 pseudouridine; 32 trigonelline; 33 1-methylnicotinamide; 34 allantoin; 35 cis-aconitate; 36 urea; 37 xanthosine; 38 cytosine; 39 urocanate; 40 tyramine, tyrosine; 41 1-methylhistidine; 42 histidine; 43 3-indoxylsulphate; 44 tyramine; 45 hippurate; 46 hypoxanthine.

Table 4. Assigned metabolites in the urine of healthy dogs.

| Query                      | HMDB        | PubChem | KEGG   |
|----------------------------|-------------|---------|--------|
| 1,7-Dimethylxanthine       | HMDB0001860 | 4687    | C13747 |
| 1-Methyladenosine          | HMDB0003331 | 27476   | C02494 |
| 1-Methylguanine            | HMDB0003282 | 70315   | C04152 |
| 1-Methylhistidine          | HMDB0000001 | 92105   | C01152 |
| 1-Methylnicotinamide       | HMDB0000699 | 457     | C02918 |
| 2-Furoylglycine            | HMDB0000439 | 21863   | NA     |
| 2-Hydroxybutyric acid      | HMDB0000008 | 11266   | C05984 |
| 2-Hydroxyphenylacetic acid | HMDB0000669 | 11970   | C05852 |
| 2-Ketobutyric acid         | HMDB0000005 | 58      | C00109 |
| 2-Methylglutaric acid      | HMDB0000422 | 12046   | NA     |
| 3-Aminoisobutyric acid     | HMDB0003911 | 64956   | C05145 |
| 3-Hydroxyphenylacetic acid | HMDB0000440 | 12122   | C05593 |

| Query                      | HMDB        | PubChem  | KEGG   |
|----------------------------|-------------|----------|--------|
| 3-Indoxylsulfic acid       | HMDB0000682 | 10258    | NA     |
| 3-Methyl-2-oxovaleric acid | HMDB0000491 | 47       | C03465 |
| 3-Methylglutaric acid      | HMDB0000752 | 12284    | NA     |
| 3-Methylxanthine           | HMDB0001886 | 70639    | C16357 |
| 4-Aminohippuric acid       | HMDB0001867 | 2148     | NA     |
| 4-Hydroxybenzoic acid      | HMDB0000500 | 135      | C00156 |
| 4-Hydroxyphenylacetic acid | HMDB0000020 | 127      | C00642 |
| 4-Pyridoxic acid           | HMDB0000017 | 6723     | C00847 |
| 7-Methyladenine            | HMDB0011614 | 71593    | C02241 |
| 7-Methylxanthine           | HMDB0001991 | 68374    | C16353 |
| Acetic acid                | HMDB0000042 | 176      | C00033 |
| Acetylcysteine             | HMDB0001890 | 12035    | C06809 |
| Adenosine                  | HMDB0000050 | 60961    | C00212 |
| Alanine                    | HMDB0000161 | 5950     | C00041 |
| Allantoin                  | HMDB0000462 | 204      | C01551 |
| Arabinitol                 | HMDB0001851 | 439255   | C00532 |
| Ascorbic acid              | HMDB0000044 | 54670067 | C00072 |
| Betaine                    | HMDB0000043 | 247      | C00719 |
| Carnitine                  | HMDB0000062 | 2724480  | C00318 |
| Choline                    | HMDB0000097 | 305      | C00114 |
| cis-Aconitic acid          | HMDB0000072 | 643757   | C00417 |

| Query                | HMDB        | PubChem | KEGG   |
|----------------------|-------------|---------|--------|
| Citric acid          | HMDB0000094 | 311     | C00158 |
| Creatine             | HMDB0000064 | 586     | C00300 |
| Creatine phosphate   | HMDB0001511 | 587     | C02305 |
| Creatinine           | HMDB0000562 | 588     | C00791 |
| Cytosine             | HMDB0000630 | 597     | C00380 |
| Dimethylamine        | HMDB0000087 | 674     | C00543 |
| Ferulic acid         | HMDB0000954 | 445858  | C01494 |
| Formic acid          | HMDB0000142 | 284     | C00058 |
| Fucose               | HMDB0000174 | 17106   | C01019 |
| Galactonic acid      | HMDB0000565 | 128869  | C00880 |
| Galactose            | HMDB0000143 | 439357  | C00984 |
| Glucaric acid        | HMDB0000663 | 33037   | C00818 |
| Glucuronic acid      | HMDB0000127 | 444791  | C00191 |
| Glycine              | HMDB0000123 | 750     | C00037 |
| Glycolic acid        | HMDB0000115 | 757     | C00160 |
| Glyoxylic acid       | HMDB0000119 | 760     | C00048 |
| Hippuric acid        | HMDB0000714 | 464     | C01586 |
| Histidine            | HMDB0000177 | 6274    | C00135 |
| Hypoxanthine         | HMDB0000157 | 790     | C00262 |
| 3-Methylhistidine    | HMDB0000479 | 64969   | C01152 |
| Indole-3-lactic acid | HMDB0000671 | 92904   | C02043 |

| Query                         | HMDB        | PubChem | KEGG   |
|-------------------------------|-------------|---------|--------|
| Isobutyric acid               | HMDB0001873 | 6590    | C02632 |
| Isoleucine                    | HMDB0000172 | 6306    | C00407 |
| Kynurenic acid                | HMDB0000715 | 3845    | C01717 |
| Lactic acid                   | HMDB0000190 | 107689  | C00186 |
| Lysine                        | HMDB0000182 | 5962    | C00047 |
| Mannitol                      | HMDB0000765 | 6251    | C00392 |
| Methylguanidine               | HMDB0001522 | 10111   | C02294 |
| <i>N,N</i> -Dimethylglycine   | HMDB0000092 | 673     | C01026 |
| N6-Acetyllysine               | HMDB0000206 | 92832   | C02727 |
| N-Acetylglycine               | HMDB0000532 | 10972   | NA     |
| <i>N</i> -Phenylacetylglycine | HMDB0000821 | 68144   | C05598 |
| Oxoglutaric acid              | HMDB0000208 | 51      | C00026 |
| Phosphorylcholine             | HMDB0001565 | 1014    | C00588 |
| Pseudouridine                 | HMDB0000767 | 15047   | C02067 |
| Pyridoxamine                  | HMDB0001431 | 1052    | C00534 |
| Serine                        | HMDB0000187 | 5951    | C00065 |
| Succinic acid                 | HMDB0000254 | 1110    | C00042 |
| Taurine                       | HMDB0000251 | 1123    | C00245 |
| Threonine                     | HMDB0000167 | 6288    | C00188 |
| trans-Aconitic acid           | HMDB0000958 | 444212  | C02341 |
| Trigonelline                  | HMDB0000875 | 5570    | C01004 |

| Query                  | HMDB        | PubChem | KEGG   |
|------------------------|-------------|---------|--------|
| Trimethylamine         | HMDB0000906 | 1146    | C00565 |
| Trimethylamine N-oxide | HMDB0000925 | 1145    | C01104 |
| Tryptophan             | HMDB0000929 | 6305    | C00078 |
| Tyramine               | HMDB0000306 | 5610    | C00483 |
| Tyrosine               | HMDB0000158 | 6057    | C00082 |
| Uracil                 | HMDB0000300 | 1174    | C00106 |
| Urea                   | HMDB0000294 | 1176    | C00086 |
| Urocanic acid          | HMDB0000301 | 736715  | C00785 |
| Valine                 | HMDB0000883 | 6287    | C00183 |
| Xanthine               | HMDB0000292 | 1188    | C00385 |
| Xanthosine             | HMDB0000299 | 64959   | C01762 |
| Xanthurenic acid       | HMDB0000881 | 5699    | C02470 |

Table 5. Significant pathways obtained by the pathway analysis module of MetaboAnalyst.

| Pathway                                  | Total <sup>a</sup> | Hits <sup>b</sup> | Raw p    | Metabolites   |
|--|--------------------|-------------------|----------|---|
| Glycine, serine and threonine metabolism | 48                 | 10                | 1.94E-06 | Serine; Choline; Betaine; Dimethylglycine; Glycine; Threonine; Creatine; 2-Ketobutyric acid; Glyoxylic acid; Tryptophan   |
| Phenylalanine metabolism                 | 45                 | 8                 | 7.59E-05 | Hippuric acid; N-Phenylacetyl-glycine; Succinic acid; 2-Hydroxyphenylacetic acid; 4-Hydroxybenzoic acid; 4-Hydroxyphenylacetic acid; Tyrosine; 3-Hydroxyphenylacetic acid |
| Aminoacyl-tRNA biosynthesis              | 75                 | 10                | 0.00012  | Histidine; Glycine; Serine; Valine; Alanine; Lysine; Isoleucine; Threonine; Tryptophan; Tyrosine  |
| Caffeine metabolism                      | 21                 | 6                 | 0.000452 | 1,7-Dimethylxanthine; 3-Methylxanthine; 7-Methylxanthine; Xanthosine; Xanthine; Glyoxylic acid  |

| Pathway                                     | Total <sup>a</sup> | Hits <sup>b</sup> | Raw p    | Metabolites  |
|---|--------------------|-------------------|----------|--|
| Methane metabolism                          | 34                 | 6                 | 0.000666 | Glycine; Formic acid; Trimethylamine; Trimethylamine N-oxide; Dimethylamine; Serine;                         |
| Glyoxylate and dicarboxylate metabolism     | 50                 | 7                 | 0.001004 | cis-Aconitic acid; Glyoxylic acid; Oxoglutaric acid; Formic acid; Glycolic acid; Citric acid; Succinic acid; |
| Nitrogen metabolism                         | 39                 | 6                 | 0.001418 | Tyrosine; Tryptophan; Taurine; Histidine; Glycine; Formic acid;  |
| Citrate cycle (TCA cycle)                   | 20                 | 4                 | 0.00349  | Succinic acid; Oxoglutaric acid; cis-Aconitic acid; Citric acid;   |
| Propanoate metabolism                       | 35                 | 5                 | 0.005027 | 2-Ketobutyric acid; Succinic acid; Lactic acid; 2-Hydroxybutyric acid; Valine;                               |
| Valine, leucine and isoleucine biosynthesis | 27                 | 4                 | 0.010619 | Threonine; Valine; Isoleucine; 2-Ketobutyric acid;   |
| Taurine and hypotaurine metabolism          | 20                 | 3                 | 0.025942 | Taurine; Alanine; Acetic acid;   |
| Purine metabolism                           | 92                 | 8                 | 0.029144 | Xanthine; Adenosine; Xanthosine; Hypoxanthine; Urea; Glyoxylic acid; Glycine; Allantoin                      |
| Alanine, aspartate and glutamate metabolism | 24                 | 3                 | 0.041947 | Alanine; Oxoglutaric acid; Succinic acid;  |
| Pyrimidine metabolism                       | 60                 | 5                 | 0.044772 | Cytosine; Uracil; Pseudouridine; Urea; 3-Aminoisobutanoic acid;  |


a

Total metabolites belonging to the pathway as reported by the pathway analysis module of MetaboAnalyst.

b

Metabolites assigned in urine of healthy dogs belonging to the pathway as obtained by the pathway analysis module of MetaboAnalyst.

By univariate *t*-test, 83 buckets resulted significantly different between healthy and CKD dog urine samples. Unsupervised multivariate analysis (PCA) was able to distinguish between healthy and CKD dogs (Fig. 4). The supervised multivariate analysis using PLS-DA (Fig. 4, Table 6) indicated that the optimal model comprised 5 components ( $R^2 = 0.99$ ,  $Q^2 = 0.74$ ), but also the model with only one component had reasonable predictive value ( $R^2 = 0.73$ ,  $Q^2 = 0.62$ ). Both univariate and multivariate analysis were used to identify the differentially abundant metabolites. Of the 83 significantly different buckets, 21 were assigned to 17 metabolites (Table 6). The metabolites showing the highest increase in CKD samples were carnosine, 7-methylxanthine and cis-aconitic acid, while the metabolites showing the most evident decrease were trigonelline and urocanic acid.



1. [Download : Download high-res image \(342KB\)](#)
2. [Download : Download full-size image](#)

Fig. 4. a) Principal component analysis (PCA) score plot of healthy (crosses and dark grey circle) and CKD (triangles and faint grey circle) urine samples. b) Partial Least Square – Discriminant Analysis (PLS-DA) distribution plot of healthy (crosses and dark grey circle) and CKD (triangles and faint grey circle) urine samples. c) Variable Importance in Projection (VIP) scores for the 25 most influential buckets of PLS-DA. Table 6. Metabolites showing significant differences between healthy and CKD dogs.

| Bucket  | Metabolite | Fold change CKD/Healthy | VIP score <sup>a</sup> | P value |
|---------|------------|-------------------------|------------------------|---------|
| B6_9876 | Carnosine  | 3.15                    | 1.922                  | 0.001   |

| Bucket  | Metabolite             | Fold change CKD/Healthy | VIP score <sup>a</sup> | P value |
|---------|------------------------|-------------------------|------------------------|---------|
| B3_9190 | 7-Methylxanthine       | 2.94                    | 1.444                  | 0.037   |
| B5_6610 | cis-Aconitic acid      | 2.67                    | 1.754                  | 0.014   |
| B2_7085 | Dimethylamine          | 1.86                    | 1.512                  | 0.017   |
| B2_8135 | Methylguanidine        | 1.80                    | 1.415                  | 0.025   |
| B7_8490 | Kynurenic acid         | 1.77                    | 1.333                  | 0.045   |
| B5_8415 | Xanthosine             | 1.72                    | 2.054                  | 0.002   |
| B4_2825 | Pseudouridine          | 1.70                    | 1.886                  | 0.002   |
| B7_6681 | Pseudouridine          | 1.59                    | 2.294                  | 0.000   |
| B5_3745 | Allantoin              | 1.47                    | 2.051                  | 0.007   |
| B7_3740 | Urocanic acid          | 0.49                    | 1.323                  | 0.032   |
| B0_9355 | 2-Hydroxybutyrric acid | 0.48                    | 1.343                  | 0.005   |
| B1_0360 | Valine                 | 0.44                    | 1.583                  | 0.010   |
| B7_7874 | 4-Hydroxybenzoic acid  | 0.40                    | 1.278                  | 0.042   |
| B7_1303 | Ferulic acid           | 0.34                    | 1.566                  | 0.009   |
| B8_1155 | 7-Methyladenine        | 0.32                    | 1.360                  | 0.030   |
| B7_7217 | Indole-3-lactic acid   | 0.26                    | 1.450                  | 0.018   |
| B6_3648 | Ferulic acid           | 0.26                    | 1.590                  | 0.009   |
| B6_3739 | Urocanic acid          | 0.21                    | 1.713                  | 0.005   |
| B8_8262 | Trigonelline           | 0.15                    | 1.284                  | 0.043   |
| B9_1121 | Trigonelline           | 0.10                    | 1.345                  | 0.034   |

a



## 4. Discussion

The aim of the present research was to characterise the urinary proteome and metabolome in healthy dogs and to compare it with that of urine collected from CKD patients, to suggest biomarkers of the disease that would be useful in veterinary medicine.

In the present study, SDS-PAGE allowed the separation of the urinary proteins based on their molecular mass, giving information about the localisation of the nephronal damage. Most urine samples (17/28) of CKD dogs analysed in this study had protein bands at both high and low MM, indicating a mixed glomerular and tubular pattern. It is generally recognised that the renal proteinuria with an UPC > 2 is strongly indicative of glomerular involvement [41,42]. Our data support this evidence, as the electrophoretic profiles of the seven urine samples with an UPC > 2 were characterised by protein bands with high MM. However, in all these samples, bands with low MM were also present, suggesting a concomitant tubular damage. Other authors reported a tubular impairment in dogs with UPC > 2 [42,43]. On the other hand, in our study, 7 of the 21 samples with UPC < 2 indicated also a glomerular involvement and hence the evaluation of proteinuria by UPC could lead to misinterpretation regarding the nephronal origin of the proteinuria, as previously suggested by other authors [41,43], [44], [45], [46].

In the present study, 8 dogs with early stages of CKD (I and II; serum creatinine < 2.1 mg/dL), classified as non-proteinuric (UPC < 0.2) or borderline proteinuric (UPC 0.2–0.5), showed altered electrophoretic profiles with the decrease of uromodulin and the increase in number and intensity of low MM bands. Chacar et al., [43] also reported the prevalence of tubular pattern in urine samples of dogs with early stages of CKD. On the other hand, out of 14 dogs affected by CKD at advanced IRIS stages (III and IV; serum creatinine > 2.1 mg/dL), 10 patients had a mixed profile, while 4 dogs presented a clear tubular pattern, with absent or mild glomerular involvement. Tubular epithelium seems to be more susceptible to ongoing stress and dysregulation promoting interstitial inflammation and fibrosis [47]. Therefore, it can be hypothesised that, in general, dogs with CKD in the initial phases (serum creatinine

<2.1 mg/dL and UPC < 0.5) might present a prevalent tubulointerstitial involvement followed by a gradual glomerular impairment leading to an increase of albumin and higher MM proteins in urine. In this complex scenario, the analysis of qualitative proteinuria could be essential to better characterise the kidney damage and the nephronal involvement.

In addition to the evaluation of the electrophoretic protein profiles, SDS-PAGE allowed the quantification of urinary uromodulin and albumin. In urine samples of healthy dogs, the abundance of uromodulin associated with the low concentration of albumin is confirmatory of data previously reported by other authors [43,48], [49], [50]. In our study, the quantification of these two proteins, followed by UMC and UAC calculation, represents an additional step for their clinical use. In fact, uromodulin and albumin are known markers of renal dysfunction or damage, in particular of CKD [51,52]. Raila et al., [50] reported a decrease of uromodulin in azotaemic and proteinuric dogs affected by renal disease and, despite the different method used for protein quantification (western blot), UMC values determined in the healthy dogs were comparable to our results. In addition, Chacar et al., [43] quantified uromodulin by western blotting and reported a decrease of uromodulin only in the late stages of CKD (IRIS 3–4), suggesting this protein as a marker of CKD progression rather than of early diagnosis. Differently, in the present study, the decrease of uromodulin was observed by SDS-PAGE already in stage 1 non-proteinuric CKD dogs, suggesting uromodulin as a promising and early biomarker of renal dysfunction in dogs.

Urinary albumin concentration is low in healthy dogs and an increase occurs in the presence of renal involvement [11,53]. Accordingly, in the present study, healthy dogs have low values of albumin and UAC, in the range of those reported by other authors [11,54], [55], [56], [57]. Different authors determined albuminuria in dogs affected by a variety of diseases and conditions, including CKD [11,53], [54], [55], [56],[58], [59], [60], [61], [62]. However, despite the clinical importance of albumin quantification in urine, the reference intervals for albuminuria are still lacking for dogs and should be the aim of further research.

Finally, the presence of arginine esterase in urine of entire male dogs was also evidenced and needs to be considered to correctly interpret urine electrophoretic profile and to exclude false tubular involvement as previously reported [15,42].

The second part of the study focused on the application of NMR to characterise the urinary metabolome of healthy dogs and to evaluate the differences with CKD patients. As most CKD samples contained high protein concentrations, to avoid interferences on NMR spectra and possible false positives, a further step in sample preparation was added by precipitating the proteins. This step allowed the enrichment of urine metabolome, improving the quality of the spectra and the identification of a higher number of metabolites. Moreover, since the high repeatability of NMR metabolomics is well known [20,21], no technical replicates were analysed and only biological replicates were considered. From the corresponding spectra, 86 metabolites were identified in healthy samples, a number higher than those previously reported in dog urine by other authors [32,33,63,64] and producing, so far and to the best of our knowledge, one of the most complete dataset of canine urinary metabolome. Most of these metabolites are of endogenous origin, while others, like ferulic acid, are of exogenous or mixed origin. Most of the identified metabolites were previously reported in urine of healthy or diseased dogs [32,33,65], in human urine [19,22] and also in feline urine [31]. The majority of these metabolites is involved in amino acids metabolism, purine and pyrimidine metabolism, tricarboxylic acid cycle and methane metabolism.

Nine metabolites were significantly increased in urine of CKD dogs. Carnosine, a dipeptide composed by alanine and histidine acting as an antioxidant scavenger, showed the most evident increase. This molecule is filtered by the glomerulus and then reabsorbed at the level of the proximal tubule by the proton-coupled oligotransporter PEPT2 [66]. It has been recently reported that the kidney has an intrinsic carnosine metabolism with carnosine synthase and carnosinase 1 activity in the glomeruli and tubular cells [67,68]. In CKD dogs, the increased urinary excretion of carnosine may reflect an oxidative stress suffered by the kidney, a condition hypothesised also by other authors in obese dogs [32,65]. Moreover, since carnosine is present at high concentrations in muscle tissues, and muscle weakness and atrophy are common findings in CKD patients, the increase of this molecule in urine may also reflect an increased muscle catabolism [65]. Finally, as a causative event, a damage of the epithelium of the proximal tubule might also be hypothesised, leading to impaired reabsorption of carnosine; this hypothesis is supported by the decrease of uromodulin evidenced by SDS-PAGE and by the increase of cis-aconitic acid in urine of CKD

dogs. Cis-aconitic acid, an intermediate in the tricarboxylic acid cycle, was observed in the urine of type 2 diabetic human patients. In fact, increased excretion of this metabolite reflects local effects on tubular transport in the kidneys [69]. Therefore, a damage of the tubular epithelium might determine an inefficient reabsorption leading to an increased concentration of urinary cis-aconitic acid and carnosine.

Regarding other urinary metabolites increased in urine of CKD dogs, xanthosine, allantoin, and 7-methylxanthine are of interest. These metabolites belong to the complex pathways of purine metabolism; in particular, during purine catabolism, the nucleoside xanthosine is transformed into xanthine, which in turn is oxidized to uric acid by uricase. In humans, uric acid is the end product of purine catabolism, while in dogs an additional reaction transforms this metabolite into allantoin. In humans, some of these metabolites were suggested as possible markers of diabetic nephropathy [70], end stage renal disease [71] or other kidney disorders [72], while an increase of allantoin and xanthine to creatinine ratios were previously reported in urine of dogs affected by CKD [73]. Despite the possible influence of medications, such as allopurinol or diuretics received by two CKD dogs included in the present study, that could have affected purine metabolism, these data show evidence that CKD is associated with alterations in urinary concentrations of purine metabolites, and thus, this issue deserves more attention in further research.

Three additional metabolites increased in urine of CKD dogs. They were methylguanidine (MG), kynurenic acid (KnA) and dimethylamine (DA). These molecules are well known uremic toxins that accumulate in serum and urine due to the impairment of renal function [74,75]. MG derives from creatinine and is often detected in serum and urine of uremic human patients [76,77]. MG was detected also in serum of uremic dogs and was shown to increase in urine of dogs affected by transitional cell carcinoma [33,78]. In the present study, 14 samples were collected from dogs at advanced CKD stages (serum creatinine > 2.1 mg/dL; IRIS 3 and 4). Therefore, the increase of MG in urine of CKD dogs might be considered in further studies as a possible biomarker of advanced CKD stages. KnA is a key inflammatory metabolite of the tryptophan catabolic pathway: the degradation of tryptophan occurs through the formation of kynurenine, which in turn can be transformed into KnA and other related metabolites. Kidneys are involved in tryptophan metabolism either eliminating the catabolites or producing the enzymes involved in tryptophan

metabolism. In case of renal failure, these metabolites, which are physiologically excreted in urine, accumulate in the blood, contributing to uremia. Accordingly, the study of Rhee et al., [79] reported that serum levels of KnA increased with CKD development and severity. Moreover, increased KnA urinary excretion was associated with adverse clinical outcomes in critically ill patients with acute kidney injury [80] and four tryptophan metabolites, including urinary KnA, were reported to be associated with an estimated glomerular filtration rate (eGFR) decline and with oxidative stress after eight years follow-up [81,82].

Eight metabolites were significantly reduced in urine of CKD patients and the most consistent decrease was evident for trigonelline, which can be obtained from the diet, or alternatively produced as a niacin-derived metabolite. Proximal tubule epithelia synthesize NAD from precursors taken up from urine and an excess of metabolites of the biosynthetic pathway, including trigonelline, is normally secreted in urine. In case of tubular damage, a reduced/absent absorption of nicotinamide or nicotinic acid occurs leading to a reduced/absent trigonelline secretion. Accordingly, in a mice model of acute kidney injury, trigonelline removal from urine was reported as a consequence of tubular damage [[83], [84], [85]].

Significant decrease was observed also for urocanic, indole-3-lactic and ferulic acids. The two first metabolites derive from hepatic histidine and tryptophan catabolism, respectively. In particular, histidine can be converted to histamine, 3-methylhistidine or urocanic acid by different pathways, while indole-3-lactic acid is obtained through the reduction of indolepyruvic acid derived by oxidative deamination of tryptophan. Finally, ferulic acid is a phenolic acid widely distributed in plants that can be absorbed by the small intestine and excreted through the urine. All these metabolites can be found in plasma and urine [22,31,86]. Serum indole-3-lactic acid was recently associated to eGFR in human CKD patients [87], but, to the best of our knowledge, no information is available in the literature on the decrease of these metabolites in the urine of CKD patients. Further studies are therefore needed to clarify their role as possible biomarkers.

This study presents some limitations. The first one is related to the different age between healthy and diseased dogs. Since CKD is a disease of older animals, and adult/old dogs are usually presented to the Veterinary Teaching Hospital due to pathologic conditions, it was not possible to collect samples from age-matched

controls. However, none of the different metabolites identified between healthy and CKD dogs were reported by Wang et al., [88] as affected by age in healthy dogs. Therefore, despite a possible age effect on urine metabolome cannot be completely excluded, we hypothesise that the effect of CKD was more consistent than the effect of the age. In addition, numerous dog breeds were included in the two study groups; therefore, breed-specific differences or effects on the urinary proteome and metabolome could not be determined in the present study. Secondly, the limited number of CKD samples did not allow to highlight significant differences among CKD stages for both proteomics and metabolomics results and it was not possible to highlight clear trends in biomarkers as the disease worsens. The final limitation relates to the absence of technical replicates for the evaluation of the robustness of our data. The technical evaluation of the performance of NMR applied to the dog urine was out of the scope of the present research, especially since the high repeatability of NMR metabolomics is well-known [20,21].

## 5. Conclusions

The integrated application of proteomics and metabolomics on urine samples yielded new insight into the molecular complexity of urine in healthy dogs and highlighted biochemical changes in response to CKD. SDS-PAGE evidenced the involvement of the tubular compartment with the decrease of uromodulin and the presence of low MM bands also in non-proteinuric and non-azotaemic dogs and could be considered a useful and complementary diagnostic tool for clinical pathologists, clinicians and researchers working in veterinary nephrology and urology.

NMR metabolomics was successfully applied to canine urinary samples allowing the identification of 86 metabolites. Of these, 17 showed significant differences in CKD dogs. In particular, the increase of carnosine and cis-aconitic acid and the decrease of trigonelline are indicative of the tubular involvement, adding further evidence to the results of SDS-PAGE. Additional studies are needed to clarify the molecular mechanisms underlying the pathophysiology of CKD and to confirm the role of the discovered metabolites as biomarkers of this disease in dogs. In particular, increasing the number of urine samples collected from dogs affected by all stages of CKD should

be the focus of future research to confirm early biomarkers and highlight trends as the disease worsens.

## **Funding**

EF was supported by a study grant from the Fondazione ItPA Onlus (2017, Rome, Italy). GI was supported by a grant from University of Bologna (RFO). The NMR data were acquired at CERMAX, ITQB-NOVA, Oeiras, Portugal with equipment funded by FCT, project AAC 01/SAICT/2016 and ONEIDA (LISBOA-01-0145-FEDER-016417) which is co-funded by FEEI - “Fundos Europeus Estruturais e de Investimento” from “Programa Operacional Regional Lisboa 2020” and by national funds from FCT.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## **Author contribution**

EF processed and analysed samples for proteomics and metabolomics, analysed and interpreted data and wrote the manuscript. GI designed and supervised the study, interpreted data, wrote and critically revised the manuscript. FD supervised the recruitment of the patients, interpreted data and critically revised the manuscript. GA interpreted data and critically revised the manuscript. KV recruited the patients, collected samples, performed urine, blood and serum biochemistry and critically revised the manuscript. EB performed protein mass identification and critically revised the manuscript. AMA designed and supervised the study and critically revised the manuscript. MM acquired and analysed metabolomics data and critically revised the manuscript. All authors read and approved the final manuscript.

## **Declaration of Competing Interest**

None

## **Appendix A. Supplementary data**

[Download : Download spreadsheet \(20KB\)](#)

Supplementary Table 1. Clinical data, serum and urine biochemistry of each dog included in the study.

## References

[1]

D.J. Polzin

**Chronic kidney disease in small animals**

Vet. Clin. North Am. Small Anim. Pract., 41 (2011), pp. 15-30, [10.1016/j.cvsm.2010.09.004](https://doi.org/10.1016/j.cvsm.2010.09.004)

[ArticleDownload](#) [PDFView Record in Scopus](#)[Google Scholar](#)

[2]

D.G. O'Neill, J. Elliott, D.B. Church, P.D. Mcgreavy, P.C. Thomson, D.C. Brodbelt

**Chronic kidney disease in dogs in UK veterinary practices: prevalence, risk factors, and survival**

J. Vet. Intern. Med., 27 (2013), pp. 814-821, [10.1111/jvim.12090](https://doi.org/10.1111/jvim.12090)

[CrossRefView Record in Scopus](#)[Google Scholar](#)

[3]

J.W. Bartges

**Chronic kidney disease in dogs and cats**

Vet. Clin. North Am. Small Anim. Pract., 42 (2012), pp. 669-692, [10.1016/j.cvsm.2012.04.008](https://doi.org/10.1016/j.cvsm.2012.04.008)

[ArticleDownload](#) [PDFView Record in Scopus](#)[Google Scholar](#)

[4]

International Renal Interest Society

**IRIS Staging of CKD (modified 2019)**

(2019), pp. 1-8

[View Record in Scopus](#)[Google Scholar](#)

[5]

S. Kovarikova

**Indirect markers of glomerular filtration rate in dogs and cats: A review**

Vet. Med. (Praha), 63 (2018), pp. 395-412, [10.17221/77/2017-VETMED](https://doi.org/10.17221/77/2017-VETMED)

[CrossRefView Record in Scopus](#)[Google Scholar](#)

[6]



P. Pocar, P. Scarpa, A. Berrini, P. Cagnardi, R. Rizzi, V. Borromeo

**Diagnostic potential of simplified methods for measuring glomerular filtration rate to detect chronic kidney disease in dogs**

J. Vet. Intern. Med., 33 (2019), pp. 2105-2116, [10.1111/jvim.15573](https://doi.org/10.1111/jvim.15573)

[CrossRefView Record in ScopusGoogle Scholar](#)

[7]

J.A. Hokamp, M.B. Nabity

**Renal biomarkers in domestic species**

Vet. Clin. Pathol., 45 (2016), pp. 28-56, [10.1111/vcp.12333](https://doi.org/10.1111/vcp.12333)

[CrossRefView Record in ScopusGoogle Scholar](#)

[8]

E. Ferlizza, A. Campos, A. Neagu, A. Cuoghi, E. Bellei, E. Monari, F. Dondi, A.M. Almeida, G. Isani

**The effect of chronic kidney disease on the urine proteome in the domestic cat (*Felis catus*)**

Vet. J., 204 (2015), pp. 73-81, [10.1016/j.tvjl.2015.01.023](https://doi.org/10.1016/j.tvjl.2015.01.023)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

[9]

E. Ferlizza, F. Dondi, G. Andreani, D. Bucci, J. Archer, G. Isani

**Validation of an electrophoretic method to detect albuminuria in cats**

J. Feline Med. Surg., 19 (2017), pp. 860-868, [10.1177/1098612X16664112](https://doi.org/10.1177/1098612X16664112)

[View Record in ScopusGoogle Scholar](#)

[10]

I. van Hoek, E. Meyer, L. Duchateau, K. Peremans, P. Smets, S. Daminet

**Retinol-binding protein in serum and urine of hyperthyroid cats before and after treatment with radioiodine**

J. Vet. Intern. Med., 23 (2009), pp. 1031-1037, [10.1111/j.1939-1676.2009.0364.x](https://doi.org/10.1111/j.1939-1676.2009.0364.x)

[CrossRefView Record in ScopusGoogle Scholar](#)

[11]

P.M.Y. Smets, E. Meyer, B.E.J. Maddens, L. Duchateau, S. Daminet

**Urinary markers in healthy young and aged dogs and dogs with chronic kidney disease**

J. Vet. Intern. Med., 24 (2010), pp. 65-72, [10.1111/j.1939-1676.2009.0426.x](https://doi.org/10.1111/j.1939-1676.2009.0426.x)

[CrossRefView Record in ScopusGoogle Scholar](#)

[12]

M.B. Nabity, G.E. Lees, L.J. Dangott, R. Cianciolo, J.S. Suchodolski, J.M. Steiner

**Proteomic analysis of urine from male dogs during early stages of tubulointerstitial injury in a canine model of progressive glomerular disease**

Vet. Clin. Pathol., 40 (2011), pp. 222-236, [10.1111/j.1939-165X.2011.00307.x](https://doi.org/10.1111/j.1939-165X.2011.00307.x)

[View Record in ScopusGoogle Scholar](#)

[13]

T.L. Williams, J. Archer

**Evaluation of urinary biomarkers for azotemic chronic kidney disease in cats**

J. Small Anim. Pract., 57 (2015), pp. 122-129, [10.1111/jsap.12439](https://doi.org/10.1111/jsap.12439)

[View Record in ScopusGoogle Scholar](#)

[  
1  
4  
]

G. Isani, E. Ferlizza, J.E. Nally

**Proteomic research in urine and other fluids**

A.M. de Almeida, P.D. Eckersall, I. Miller (Eds.), Proteomics Domestic Animals: From Farm to System Biology, Springer, Cham (2018), pp. 121-147, [10.1007/978-3-319-69682-9\\_7](https://doi.org/10.1007/978-3-319-69682-9_7)

[CrossRefView Record in ScopusGoogle Scholar](#)

[  
1  
5  
]

I. Miller, A. Preßlmayer-Hartler, R. Wait, K. Hummel, C. Sensi, I. Eberini, E. Razzazi-Fazeli, E. Gianazza

**In between — proteomics of dog biological fluids**

J. Proteome, 106 (2014), pp. 30-45, [10.1016/j.jprot.2014.04.016](https://doi.org/10.1016/j.jprot.2014.04.016)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

J. Kuleš, P. Bilić, B. Beer Ljubić, J. Gotić, M. Crnogaj, M. Brkljačić, V. Mrljak

**Glomerular and tubular kidney damage markers in canine babesiosis caused by *Babesia canis***

Ticks Tick. Borne. Dis., 9 (2018), pp. 1508-1517, [10.1016/j.ttbdis.2018.07.012](https://doi.org/10.1016/j.ttbdis.2018.07.012)

[ArticleDownload](#) [PDFView](#) [Record in ScopusGoogle Scholar](#)

L. Pelander, V. Brunchault, B. Buffin-

Meyer, J. Klein, B. Breuil, P. Züribig, P. Magalhães, W. Mullen, J. Elliott, H. Syme, J.P. Schanstra, J. Häggström, I. Ljungvall

**Urinary peptidome analyses for the diagnosis of chronic kidney disease in dogs**

Vet. J., 249 (2019), pp. 73-79, [10.1016/j.tvjl.2019.05.010](https://doi.org/10.1016/j.tvjl.2019.05.010)

[ArticleDownload](#) [PDFView](#) [Record in ScopusGoogle Scholar](#)

I.F. Duarte, S.O. Diaz, A.M. Gil

**NMR metabolomics of human blood and urine in disease research**

J. Pharm. Biomed. Anal., 93 (2014), pp. 17-26, [10.1016/j.jpba.2013.09.025](https://doi.org/10.1016/j.jpba.2013.09.025)

[ArticleDownload](#) [PDFView](#) [Record in ScopusGoogle Scholar](#)

R.A. Silva, T.C.S. Pereira, A.R. Souza, P.R. Ribeiro

**<sup>1</sup>H NMR-based metabolite profiling for biomarker identification**

Clin. Chim. Acta, 502 (2020), pp. 269-279, [10.1016/j.cca.2019.11.015](https://doi.org/10.1016/j.cca.2019.11.015)

[ArticleDownload](#) [PDFView](#) [Record in ScopusGoogle Scholar](#)

J.L. Markley, R. Brüschweiler, A.S. Edison, H.R. Eghbalnia, R. Powers, D. Raftery, D.S. Wishart

**The future of NMR-based metabolomics**

Curr. Opin. Biotechnol., 43 (2017), pp. 34-40, [10.1016/j.copbio.2016.08.001](https://doi.org/10.1016/j.copbio.2016.08.001)

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)[Google Scholar](#)

P.G. Takis, V. Ghini, L. Tenori, P. Turano, C. Luchinat

**Uniqueness of the NMR approach to metabolomics**

Trends Anal. Chem., 120 (2019), p. 115300, [10.1016/j.trac.2018.10.036](https://doi.org/10.1016/j.trac.2018.10.036)

[ArticleDownload](#) [PDFView](#) [Record in Scopus](#)[Google Scholar](#)

S. Bouatra, F. Aziat, R. Mandal, A.C. Guo, M.R. Wilson, C. Knox, T.C. Bjorndahl, R. Krishnamurthy, F. Saleem, P. Liu, Z.T. Dame, J. Poelzer, J. Huynh, F.S. Yallou, N. Psychogios, E. Dong, R. Bogumil, C. Roehring, D.S. Wishart

**The human urine metabolome**

PLoS One, 8 (2013), Article e73076, [10.1371/journal.pone.0073076](https://doi.org/10.1371/journal.pone.0073076)

[CrossRefView](#) [Record in Scopus](#)[Google Scholar](#)

A. Julià, M. Vinaixa, E. Domènech, A. Fernández-Nebro, J.D. Cañete, C. Ferrándiz, J. Tornero, J.P. Gisbert, P. Nos, A.G. Casbas, L. Puig, I. González-Álvaro, J.A. Pinto-Tasende, R. Blanco, M.A. Rodríguez, A. Beltran, X. Correig, S. Marsal, E. Fernández, R. Sanmartí, J. Gratacós, V.M. Martínez-Taboada, F. Gomollón, E. Daudén, J. Maymó, R. Queiró, F.J.L. Longo, E. Garcia-Planella, J.L.S. Carazo, M. Alperi-López, C. Montilla, J.J. Pérez-Venegas, B. Fernández-Gutiérrez, J.L. Mendoza, J.L. López Estebaranz, À. Olivé, J.C. Torre-Alonso, M. Barreiro-de Acosta, D.M. Ramírez, H. Corominas, S. Muñoz-Fernández, J.L. Andreu, F. Muñoz, P. de la Cueva, A. Erra, C.M. González, M.Á. Aguirre-Zamorano, M. Vera, F. Vanaclocha, D. Roig, P. Vela, C. Saro, E. Herrera, P. Zarco, J.M. Nolla, M. E

steve, J.L. Marenco de la Fuente, J.M. Pego-Reigosa, V. García-Sánchez, J. Panés, E. Fonseca, F. Blanco, J. Rodríguez-Moreno, P. Carreira, J. Ramírez, G. Avila, L. Codó, J.L. Gelpí, A.C. García-Montero, N. Palau, M. López-Lasanta, R. Tortosa, A. Alonso

**Urine metabolome profiling of immune-mediated inflammatory diseases**

BMC Med., 14 (2016), p. 133, [10.1186/s12916-016-0681-8](https://doi.org/10.1186/s12916-016-0681-8)

[View Record in ScopusGoogle Scholar](#)

E.A. Mathé, A.D. Patterson, M. Haznadar, S.K. Manna, K.W. Krausz, E.D. Bowman, P.G. Shields, J. R. Idle, P.B. Smith, K. Anami, D.G. Kazandjian, E. Hatzakis, F.J. Gonzalez, C.C. Harris

**Non-invasive urinary metabolomic profiling identifies diagnostic and prognostic markers in lung cancer**

Cancer Res., 74 (2014), pp. 3259-3270, [10.1158/0008-5472.CAN-14-0109](https://doi.org/10.1158/0008-5472.CAN-14-0109)

[View Record in ScopusGoogle Scholar](#)

C.L. Silva, A. Olival, R. Perestrelo, P. Silva, H. Tomás, J.S. Câmara

**Untargeted urinary<sup>1</sup>H NMR-based metabolomic pattern as a potential platform in breast cancer detection**

Metabolites., 9 (2019), p. 269, [10.3390/metabo9110269](https://doi.org/10.3390/metabo9110269)

[CrossRefView Record in ScopusGoogle Scholar](#)

E.R. Kim, H.N. Kwon, H. Nam, J.J. Kim, S. Park, Y.H. Kim

**Urine-NMR metabolomics for screening of advanced colorectal adenoma and early stage colorectal cancer**

Sci. Rep., 9 (2019), p. 4786, [10.1038/s41598-019-41216-y](https://doi.org/10.1038/s41598-019-41216-y)

[View Record in ScopusGoogle Scholar](#)

M. Martin-Lorenzo, L. Gonzalez-Calero, A. Ramos-Barron, M.D. Sanchez-Niño, C. Gomez-Alamillo, J.M. García-Segura, A. Ortiz, M. Arias, F. Vivanco, G. Alvarez-Llamas

**Urine metabolomics insight into acute kidney injury point to oxidative stress disruptions in energy generation and H<sub>2</sub>S availability**

J. Mol. Med., 95 (2017), pp. 1399-1409, [10.1007/s00109-017-1594-5](https://doi.org/10.1007/s00109-017-1594-5)

[CrossRefView Record in ScopusGoogle Scholar](#)

M. Posada-Ayala, I. Zubiri, M. Martin-Lorenzo, A. Sanz-Maroto, D. Molero, L. Gonzalez-Calero, B. Fernandez-Fernandez, F. De La Cuesta, C.M. Laborde, M.G. Barderas, A. Ortiz, F. Vivanco, G. Alvarez-Llamas

**Identification of a urine metabolomic signature in patients with advanced-stage chronic kidney disease**

Kidney Int., 85 (2014), pp. 103-111, [10.1038/ki.2013.328](https://doi.org/10.1038/ki.2013.328)

[ArticleDownload PDFCrossRefView Record in ScopusGoogle Scholar](#)

M. Breit, K.M. Weinberger

**Metabolic biomarkers for chronic kidney disease**

Arch. Biochem. Biophys., 589 (2016), pp. 62-80, [10.1016/j.abb.2015.07.018](https://doi.org/10.1016/j.abb.2015.07.018)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

S. Hallan, M. Afkarian, L.R. Zelnick, B. Kestenbaum, S. Sharma, R. Saito, M. Darshi, G. Barding, D. Raftery, W. Ju, M. Kretzler, K. Sharma, I.H. de Boer

**Metabolomics and gene expression analysis reveal down-regulation of the citric acid (TCA) Cycle in non-diabetic CKD patients**

EBioMedicine., 26 (2017), pp. 68-77, [10.1016/j.ebiom.2017.10.027](https://doi.org/10.1016/j.ebiom.2017.10.027)

[ArticleDownload](#) [PDFView Record in Scopus](#)[Google Scholar](#)

S.-M. Rivera-Vélez, N.F. Villarino

**Feline urine metabolomic signature: characterization of low-molecular-weight substances in urine from domestic cats**

J. Feline Med. Surg., 20 (2017), pp. 155-163, [10.1177/1098612X17701010](https://doi.org/10.1177/1098612X17701010)

[Google Scholar](#)

J. Soder, R. Hagman, J. Dicksved, S. Lindase, K. Malmlof, P. Agback, A. Moazzami, K. Hoglund, S. Wernersson

**The urine metabolome differs between lean and overweight Labrador Retriever dogs during a feed-challenge**

PLoS One, 12 (2017), Article e0180086, [10.1371/journal.pone.0180086](https://doi.org/10.1371/journal.pone.0180086)

[CrossRefView Record in Scopus](#)[Google Scholar](#)

J. Zhang, S. Wei, L. Liu, G.A. Nagana Gowda, P. Bonney, J. Stewart, D.W. Knapp, D. Raftery

**NMR-based metabolomics study of canine bladder cancer**

Biochim. Biophys. Acta, 1822 (2012), pp. 1807-1814, [10.1016/j.bbadis.2012.08.001](https://doi.org/10.1016/j.bbadis.2012.08.001)

[ArticleDownload](#) [PDFView Record in Scopus](#)[Google Scholar](#)

Y.A. Lawrence, B.C. Guard, J.M. Steiner, J.S. Suchodolski, J.A. Lidbury

**Untargeted metabolomic profiling of urine from healthy dogs and dogs with chronic hepatic disease**

PLoS One, 14 (2019), Article e0217797, [10.1371/journal.pone.0217797](https://doi.org/10.1371/journal.pone.0217797)

[CrossRefView Record in ScopusGoogle Scholar](#)

C. Zhu, V. Faillace, F. Laus, M. Bazzano, L. Laghi

**Characterization of trotter horses urine metabolome by means of proton nuclear magnetic resonance spectroscopy**

Metabolomics., 14 (2018), p. 106, [10.1007/s11306-018-1403-3](https://doi.org/10.1007/s11306-018-1403-3)

[View Record in ScopusGoogle Scholar](#)

G. Isani, E. Ferlizza, A. Cuoghi, E. Bellei, E. Monari, B. Bianchin Butina, C. Castagnetti

**Identification of the most abundant proteins in equine amniotic fluid by a proteomic approach**

Anim. Reprod. Sci., 174 (2016), pp. 150-160, [10.1016/j.anireprosci.2016.10.003](https://doi.org/10.1016/j.anireprosci.2016.10.003)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

D. Jacob, C. Deborde, M. Lefebvre, M. Maucourt, A. Moing

**NMRProcFlow: a graphical and interactive tool dedicated to 1D spectra processing for NMR-based metabolomics**

Metabolomics., 13 (2017), p. 36, [10.1007/s11306-017-1178-y](https://doi.org/10.1007/s11306-017-1178-y)

[View Record in ScopusGoogle Scholar](#)



J. Xia, D.S. Wishart

**Using MetaboAnalyst 3.0 for comprehensive metabolomics data analysis**

Curr. Protoc. Bioinforma, 55 (2016), [10.1002/cpbi.11](https://doi.org/10.1002/cpbi.11)

[Google Scholar](#)

M. Palma, L.E. Hernández-Castellano, N. Castro, A. Arguëllo, J. Capote, M. Matzapetakis, A.M. de Almeida

**NMR-metabolomics profiling of mammary gland secretory tissue and milk serum in two goat breeds with different levels of tolerance to seasonal weight loss**

Mol. BioSyst., 12 (2016), pp. 2094-2107, [10.1039/C5MB00851D](https://doi.org/10.1039/C5MB00851D)

[View Record in Scopus](#)[Google Scholar](#)

E. Szymańska, E. Saccenti, A.K. Smilde, J.A. Westerhuis

**Double-check: validation of diagnostic statistics for PLS-DA models in metabolomics studies**

Metabolomics., 8 (2012), pp. 3-16, [10.1007/s11306-011-0330-3](https://doi.org/10.1007/s11306-011-0330-3)

[CrossRefView Record in Scopus](#)[Google Scholar](#)

J.A. Hokamp, R.E. Cianciolo, M. Boggess, G.E. Lees, S.L. Benali, M. Kovarsky, M.B. Nabity

**Correlation of urine and serum biomarkers with renal damage and survival in dogs with naturally occurring proteinuric chronic kidney disease**

J. Vet. Intern. Med., 30 (2016), pp. 591-601, [10.1111/jvim.13832](https://doi.org/10.1111/jvim.13832)

[CrossRefView Record in Scopus](#)[Google Scholar](#)

J.A. Hokamp, S.A. Leidy, I. Gaynanova, R.E. Cianciolo, M.B. Nabity

**Correlation of electrophoretic urine protein banding patterns with severity of renal damage in dogs with proteinuric chronic kidney disease**

Vet. Clin. Pathol., 47 (2018), pp. 425-434, [10.1111/vcp.12648](https://doi.org/10.1111/vcp.12648)

[CrossRefView Record in ScopusGoogle Scholar](#)

F. Chacar, M. Kogika, T.R. Sanches, D. Caragelasco, C. Martorelli, C. Rodrigues, J.M.C. Capcha, D. Chew, L. Andrade

**Urinary Tamm-Horsfall protein, albumin, vitamin D-binding protein, and retinol-binding protein as early biomarkers of chronic kidney disease in dogs**

Physiol. Rep., 5 (2017), p. e13262, [10.14814/phy2.13262](https://doi.org/10.14814/phy2.13262)

[CrossRefView Record in ScopusGoogle Scholar](#)

R. Cianciolo, J. Hokamp, M. Nabity

**Advances in the evaluation of canine renal disease**

Vet. J., 215 (2016), pp. 21-29, [10.1016/j.tvjl.2016.04.012](https://doi.org/10.1016/j.tvjl.2016.04.012)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

O. Jaturakan, M. Vanichwatanaramlouk, A. Kornkaew, M. Trisiroj

**SDS-PAGE electrophoresis for urinary protein analysis in dogs with chronic kidney disease and urinary tract infection**

Thai J. Vet. Med., 43 (2013), pp. 75-83

[View Record in ScopusGoogle Scholar](#)

D.S. Caragelasco, M.M. Kogika, C.R. Martorelli, K.K. Kanayama, D.M.N. Simões

**Urine protein electrophoresis study in dogs with pituitary dependent hyperadrenocorticism during therapy with trilostane**

Pesqui. Vet. Bras., 37 (2017), pp. 734-740, [10.1590/s0100-736x2017000700014](https://doi.org/10.1590/s0100-736x2017000700014)

[CrossRefView Record in ScopusGoogle Scholar](#)

G. Segev, M.B. Nabity, G.F. Grauer, S. Brown, D.J. Polzin, L.D. Cowgill, C. Langston, A.M. van Dongen, J. Elliott

**Is progressive chronic kidney disease a slow acute kidney injury?**

Vet. Clin. North Am. Small Anim. Pract., 46 (2016), pp. 995-1013, [10.1016/j.cvsm.2016.06.001](https://doi.org/10.1016/j.cvsm.2016.06.001)

[Google Scholar](#)

J. Raila, H. Aupperle, G. Raila

**Renal pathology and urinary protein excretion in a 14-month-old Bernese mountain dog with chronic renal failure**

J. Vet. Med A., 54 (2007), pp. 131-135, [10.1111/j.1439-0442.2007.00919.x](https://doi.org/10.1111/j.1439-0442.2007.00919.x)

[CrossRefView Record in ScopusGoogle Scholar](#)

J. Raila, L. Brunnberg

**Influence of kidney function on urinary excretion of albumin and retinol-binding protein in dogs with naturally occurring renal disease**

Am. J. Vet. Res., 71 (2010), pp. 1387-1394, [10.2460/ajvr.71.11.1387](https://doi.org/10.2460/ajvr.71.11.1387)

[CrossRefView Record in ScopusGoogle Scholar](#)

J. Raila, F.J. Schweigert, B. Kohn

**Relationship between urinary Tamm-Horsfall protein excretion and renal function in dogs with naturally occurring renal disease**

Vet. Clin. Pathol., 43 (2014), pp. 261-265, [10.1111/vcp.12143](https://doi.org/10.1111/vcp.12143)

[CrossRefView Record in ScopusGoogle Scholar](#)

O. Devuyst, E. Olinger, L. Rampoldi

**Uromodulin: from physiology to rare and complex kidney disorders**

Nat. Rev. Nephrol., 13 (2017), pp. 525-544, [10.1038/nrneph.2017.101](https://doi.org/10.1038/nrneph.2017.101)

[CrossRefView Record in ScopusGoogle Scholar](#)

A. Okubo, A. Nakashima, S. Doi, T. Doi, T. Ueno, K. Maeda, R. Tamura, K. Yamane, T. Masaki

**High-normal albuminuria is strongly associated with incident chronic kidney disease in a nondiabetic population with normal range of albuminuria and normal kidney function**

Clin. Exp. Nephrol. (2020), [10.1007/s10157-019-01842-2](https://doi.org/10.1007/s10157-019-01842-2)

[Google Scholar](#)

J. Martinez, C. Kellogg, M.C. Iazbik, C.G. Couto, B.M. Pressler, T.M. Hoepf, M.J. Radin

**The renin-angiotensin-aldosterone system in Greyhounds and non-Greyhound dogs**

J. Vet. Intern. Med., 31 (2017), pp. 988-993, [10.1111/jvim.14720](https://doi.org/10.1111/jvim.14720)

[CrossRefView Record in ScopusGoogle Scholar](#)

A. Bacic, M.M. Kogika, K.C. Barbaro, C.S. Iuamoto, D.M.N. Simões, M.L. Santoro

**Evaluation of albuminuria and its relationship with blood pressure in dogs with chronic kidney disease**

Vet. Clin. Pathol., 39 (2010), pp. 203-209, [10.1111/j.1939-165X.2009.00207.x](https://doi.org/10.1111/j.1939-165X.2009.00207.x)

[View Record in ScopusGoogle Scholar](#)

C.Z. Cavalcante, M.M. Kogika, A. Bacic, M.L. Santoro, S.I. Miyashiro, J.P. Sault, M.K. Oyafuso, D.M. Simões

**Evaluation of albuminuria and electrophoresis of urinary proteins from dogs with hyperadrenocorticism and relation with systemic arterial pressure**

Pesqui. Vet. Bras., 33 (2013), pp. 1354-1363, [10.1590/S0100-736X2013001100013](https://doi.org/10.1590/S0100-736X2013001100013)

[Google Scholar](#)

P.M.Y. Smets, H.P. Lefebvre, B.P. Meij, S. Croubels, E. Meyer, I. Van de Maele, S. Daminet

**Long-term follow-up of renal function in dogs after treatment for ACTH-dependent hyperadrenocorticism**

J. Vet. Intern. Med., 26 (2012), pp. 565-574, [10.1111/j.1939-1676.2012.00915.x](https://doi.org/10.1111/j.1939-1676.2012.00915.x)

[CrossRefView Record in ScopusGoogle Scholar](#)

O.L. Ioannis, S. Nectarios, C. Jose Joaquin, T. Konstantina, R. Timoleon, P. Zoe

**The prognostic value of microalbuminuria in puppies with canine parvoviral enteritis**

Acta Vet. Brno., 69 (2019), pp. 116-122, [10.2478/acve-2019-0008](https://doi.org/10.2478/acve-2019-0008)

[View Record in ScopusGoogle Scholar](#)

H. Schaefer, B. Kohn, F.J. Schweigert, J. Raila

**Quantitative and qualitative urine protein excretion in dogs with severe inflammatory response syndrome**

J. Vet. Intern. Med., 25 (2011), pp. 1292-1297, [10.1111/j.1939-1676.2011.00829.x](https://doi.org/10.1111/j.1939-1676.2011.00829.x)

[CrossRefView Record in ScopusGoogle Scholar](#)

S.L. Vaden, C.A. Turman, T.L. Harris, S.L. Marks

**The prevalence of albuminuria in dogs and cats in an ICU or recovering from anesthesia**

J. Vet. Emerg. Crit. Care, 20 (2010), pp. 479-487, [10.1111/j.1476-4431.2010.00584.x](https://doi.org/10.1111/j.1476-4431.2010.00584.x)

[CrossRefView Record in ScopusGoogle Scholar](#)

Y.H. Lien, T.Y. Hsiang, H.P. Huang

**Associations among systemic blood pressure, microalbuminuria and albuminuria in dogs affected with pituitary- and adrenal-dependent hyperadrenocorticism**

Acta Vet. Scand., 52 (2010), p. 61, [10.1186/1751-0147-52-61](https://doi.org/10.1186/1751-0147-52-61)

[View Record in ScopusGoogle Scholar](#)

J. Whittemore, B. Marcum, D. Mawby, M. Coleman, T. Hacket, L. Lappin

**Associations among albuminuria, C-reactive protein concentrations, survival predictor index scores, and survival in 78 critically ill dogs**

J. Vet. Intern. Med., 25 (2011), pp. 818-824

[CrossRefView Record in ScopusGoogle Scholar](#)

R. Troia, M. Gruarin, C. Grisetti, F. Serafini, L. Magna, E. Monari, M. Giunti, F. Dondi

**Fractional excretion of electrolytes in volume-responsive and intrinsic acute kidney injury in dogs: diagnostic and prognostic implications**

J. Vet. Intern. Med., 32 (2018), pp. 1372-1382, [10.1111/jvim.15146](#)

[CrossRefView Record in ScopusGoogle Scholar](#)

M.R. Viant, C. Ludwig, S. Rhodes, U.L. Günther, D. Allaway

**Validation of a urine metabolome fingerprint in dog for phenotypic classification**

Metabolomics., 3 (2007), pp. 453-463, [10.1007/s11306-007-0092-0](#)

[CrossRefView Record in ScopusGoogle Scholar](#)

G. Carlos, F.P. dos Santos, P.E. Fröhlich

**Canine metabolomics advances**

Metabolomics., 16 (2020), p. 16, [10.1007/s11306-020-1638-7](#)

[View Record in ScopusGoogle Scholar](#)

G.M. Forster, J. Stockman, N. Noyes, A.L. Heuberger, C.D. Broeckling, C.M. Bantle, E.P. Ryan

**A comparative study of serum biochemistry, metabolome and microbiome parameters of clinically healthy, normal weight, overweight, and obese companion dogs**

Top. Companion Anim. Med., 33 (2018), pp. 126-135, [10.1053/j.tcam.2018.08.003](#)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

D. Jappar, Y. Hu, R.F. Keep, D.E. Smith

**Transport mechanisms of carnosine in SKPT cells: contribution of apical and basolateral membrane transporters**

Pharm. Res., 26 (2009), pp. 172-181, [10.1007/s11095-008-9726-9](#)

[CrossRefView Record in ScopusGoogle Scholar](#)

V. Peters, C.Q.F. Klessens, H.J. Baelde, B. Singler, K.A.M. Veraar, A. Zutinic, J. Drozak, J. Zschocke, C.P. Schmitt, E. De Heer

**Intrinsic carnosine metabolism in the human kidney**

Amino Acids, 47 (2015), pp. 2541-2550, [10.1007/s00726-015-2045-7](https://doi.org/10.1007/s00726-015-2045-7)

[CrossRefView Record in ScopusGoogle Scholar](#)

K. Kilis-Pstrusinska

**Carnosine and kidney diseases: what we currently know?**

Curr. Med. Chem., 20 (2020), [10.2174/0929867326666190730130024](https://doi.org/10.2174/0929867326666190730130024)

[Google Scholar](#)

M. Urpi-Sarda, E. Almanza-Aguilera, R. Llorach, R. Vázquez-Fresno, R. Estruch, D. Corella, J.V. Sorli, F. Carmona, A. Sanchez-Pla, J. Salas-Salvadó, C. Andres-Lacueva

**Non-targeted metabolomic biomarkers and metabolotypes of type 2 diabetes: A cross-sectional study of PREDIMED trial participants**

Diabetes Metab., 45 (2019), pp. 167-174, [10.1016/j.diabet.2018.02.006](https://doi.org/10.1016/j.diabet.2018.02.006)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

C.J. Chen, W.L. Liao, C.T. Chang, Y.N. Lin, F.J. Tsai

**Identification of urinary metabolite biomarkers of type 2 diabetes nephropathy using an untargeted metabolomic approach**

J. Proteome Res., 17 (2018), pp. 3997-4007, [10.1021/acs.jproteome.8b00644](https://doi.org/10.1021/acs.jproteome.8b00644)

[CrossRefView Record in ScopusGoogle Scholar](#)

D.Q. Chen, G. Cao, H. Chen, D. Liu, W. Su, X.Y. Yu, N.D. Vaziri, X.H. Liu, X. Bai, L. Zhang, Y.Y. Zhao

**Gene and protein expressions and metabolomics exhibit activated redox signaling and wnt/ $\beta$ -catenin pathway are associated with metabolite dysfunction in patients with chronic kidney disease**

Redox Biol., 12 (2017), pp. 505-521, [10.1016/j.redox.2017.03.017](https://doi.org/10.1016/j.redox.2017.03.017)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

R.E. Silva, J.L. Baldim, D.A. Chagas-Paula, M.G. Soares, J.H.G. Lago, R.V. Gonçalves, R.D. Novaes

**Predictive metabolomic signatures of end-stage renal disease: A multivariate analysis of population-based data**

Biochimie., 152 (2018), pp. 14-30, [10.1016/j.biochi.2018.06.009](https://doi.org/10.1016/j.biochi.2018.06.009)

[ArticleDownload](#) [PDFView Record in ScopusGoogle Scholar](#)

C.M. Rivara, C.R. Johnson, J.P. Lulich, C.A. Osborne, M. Murtaugh

**The effect of disease on the urinary purine metabolite concentrations in dogs**

Vet. Rec., 173 (2013), p. 219, [10.1136/vr.101237](https://doi.org/10.1136/vr.101237)

[CrossRefView Record in ScopusGoogle Scholar](#)

F. Durantou, G. Cohen, R. De Smet, M. Rodriguez, J. Jankowski, R. Vanholder, A. Argiles

**Normal and pathologic concentrations of uremic toxins**

J. Am. Soc. Nephrol., 23 (2012), pp. 1258-1270, [10.1681/ASN.2011121175](https://doi.org/10.1681/ASN.2011121175)

[View Record in ScopusGoogle Scholar](#)

M.A. Risso, S. Sallustio, V. Sueiro, V. Bertoni, H. Gonzalez-Torres, C.G. Musso

**The importance of tubular function in chronic kidney disease**

Int. J. Nephrol. Renovasc. Dis., 12 (2019), pp. 257-262, [10.2147/IJNRD.S216673](https://doi.org/10.2147/IJNRD.S216673)

[CrossRefView Record in ScopusGoogle Scholar](#)

K. Ienaga, K. Nakamura, T. Fujisawa, Y. Fukunaga, H. Nihei, M. Narita, Y. Tomino, T. Sanaka, K. Aoyagi, K. Nakano, H. Koide

**Urinary excretion of creatol, an *in vivo* biomarker of hydroxyl radical, in patients with chronic renal failure**

Ren. Fail., 29 (2007), pp. 279-283, [10.1080/08860220701219863](https://doi.org/10.1080/08860220701219863)

[CrossRefView Record in ScopusGoogle Scholar](#)

N. Neiryneck, R. Vanholder, E. Schepers, S. Eloit, A. Pletinck, G. Glorieux

**An update on uremic toxins**

Int. Urol. Nephrol., 45 (2013), pp. 139-150, [10.1007/s11255-012-0258-1](https://doi.org/10.1007/s11255-012-0258-1)



[CrossRefView Record in ScopusGoogle Scholar](#)

A.M. Bosco, B.F.M. Almeida, P.P. Pereira, D.B. dos Santos, J.S. Neto, W.L. Ferreira, P.C. Ciarlini

**The uremic toxin methylguanidine increases the oxidative metabolism and accelerates the apoptosis of canine neutrophils**

Vet. Immunol. Immunopathol., 185 (2017), pp. 14-19, [10.1016/j.vetimm.2017.01.006](#)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

E.P. Rhee, C.B. Clish, A. Ghorbani, M.G. Larson, S. Elmariah, E. McCabe, Q. Yang, S. Cheng, K. Pierce, A. Deik, A.L. Souza, L. Farrell, C. Domos, R.W. Yeh, I. Palacios, K. Rosenfield, R.S. Vasan, J.C. Florez, T.J. Wang, C.S. Fox, R.E. Gerszten

**A combined epidemiologic and metabolomic approach improves CKD prediction**

J. Am. Soc. Nephrol., 24 (2013), pp. 1330-1338, [10.1681/ASN.2012101006](#)

[View Record in ScopusGoogle Scholar](#)

F. Aregger, D.E. Uehlinger, G. Fusch, A. Bahunjic, R. Pschowski, M. Walter, J.C. Schefold

**Increased urinary excretion of kynurenic acid is associated with non-recovery from acute kidney injury in critically ill patients**

BMC Nephrol., 19 (2018), p. 44, [10.1186/s12882-018-0841-5](#)

[View Record in ScopusGoogle Scholar](#)

C. Barrios, T.D. Spector, C. Menni

**Blood, urine and faecal metabolite profiles in the study of adult renal disease**

Arch. Biochem. Biophys., 589 (2016), pp. 81-92, [10.1016/j.abb.2015.10.006](#)

[ArticleDownload PDFView Record in ScopusGoogle Scholar](#)

A.L. Saucedo, M.M. Perales-Quintana, D. Paniagua-Vega, C. Sanchez-Martinez, P. Cordero-Perez, N.W. Minsky

**Chronic kidney disease and the search for new biomarkers for early diagnosis**

Curr. Med. Chem., 25 (2018), pp. 3719-3747, [10.2174/0929867325666180307110908](#)

[CrossRefView Record in ScopusGoogle Scholar](#)

T. Chihanga, Q. Ma, J.D. Nicholson, H.N. Ruby, R.E. Edelman, P. Devarajan, M.A. Kennedy

**NMR spectroscopy and electron microscopy identification of metabolic and ultrastructural changes to the kidney following ischemia-reperfusion injury**

Am. J. Physiol. Physiol., 314 (2018), pp. 154-166, [10.1152/ajprenal.00363.2017](https://doi.org/10.1152/ajprenal.00363.2017)

[Google Scholar](#)

M.A. Hanifa, M. Skott, R.G. Maltesen, B.S. Rasmussen, S. Nielsen, J. Frøkiær, T. Ring, R. Wimmer

**Tissue, urine and blood metabolite signatures of chronic kidney disease in the 5/6 nephrectomy rat model**

Metabolomics., 15 (2019), p. 112, [10.1007/s11306-019-1569-3](https://doi.org/10.1007/s11306-019-1569-3)

[View Record in Scopus](#)[Google Scholar](#)

A.J. Won, S. Kim, Y.G. Kim, K.B. Kim, W.S. Choi, S. Kacew, K.S. Kim, J.H. Jung, B.M. Lee, S. Kim, H.S. Kim

**Discovery of urinary metabolomic biomarkers for early detection of acute kidney injury**

Mol. BioSyst., 12 (2016), pp. 133-144, [10.1039/c5mb00492f](https://doi.org/10.1039/c5mb00492f)

[View Record in Scopus](#)[Google Scholar](#)

Y. He, W. Su, T. Chen, X. Zeng, Z. Yan, J. Guo, W. Yang, H. Wu

**Identification of prototype compounds and derived metabolites of naoxintong capsule in beagle dog urine and feces by UFLC-Q-TOF-MS/MS**

J. Pharm. Biomed. Anal., 176 (2019), p. 112806, [10.1016/j.jpba.2019.112806](https://doi.org/10.1016/j.jpba.2019.112806)

[ArticleDownload](#) [PDFView Record in Scopus](#)[Google Scholar](#)

S.M. Titan, G. Venturini, K. Padilha, G. Tavares, R. Zatz, I. Bensenor, P.A. Lotufo, E.P. Rhee, R.I. Thadhani, A.C. Pereira

**Metabolites related to eGFR: evaluation of candidate molecules for GFR estimation using untargeted metabolomics**

Clin. Chim. Acta, 489 (2019), pp. 242-248, [10.1016/j.cca.2018.08.037](https://doi.org/10.1016/j.cca.2018.08.037)

[ArticleDownload](#) [PDFView Record in Scopus](#)[Google Scholar](#)

Y. Wang, D. Lawler, B. Larson, Z. Ramadan, S. Kochhar, E. Holmes, J.K. Nicholson

**Metabonomic investigations of aging and caloric restriction in a life-long dog study**

J. Proteome Res., 6 (2007), pp. 1846-1854, [10.1021/pr060685n](https://doi.org/10.1021/pr060685n)

[CrossRefView Record in ScopusGoogle Scholar](#)