

# Transcription Factors in the Development and Function of Group 2 Innate Lymphoid Cells

著者	Ebihara Takashi, Taniuchi Ichiro
journal or publication title	International Journal of Molecular Sciences
volume	20
number	6
page range	1377
year	2019
出版者	MDPI
関連リンク	<a href="https://doi.org/10.3390/ijms20061377">https://doi.org/10.3390/ijms20061377</a>
著作権等	(C) 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license ( <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> ).
URL	<a href="http://hdl.handle.net/10295/00005782">http://hdl.handle.net/10295/00005782</a>

doi: 10.3390/ijms20061377



Review

# Transcription Factors in the Development and Function of Group 2 Innate Lymphoid Cells

Takashi Ebihara <sup>\*,†</sup>  and Ichiro Taniuchi

Laboratory for Transcriptional Regulation, RIKEN Center for Integrative Medical Sciences (IMS),  
1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama 230-0045, Japan; ichiro.taniuchi@riken.jp

\* Correspondence: takashi.ebihara@riken.jp; Tel.: +81-45-503-7065

† Present address: Department of Medical Biology, Akita University School of Medicine, 1-1-1 Hondo,  
Akita 010-8543, Japan; tebihara@med.akita-u.ac.jp.

Received: 28 February 2019; Accepted: 18 March 2019; Published: 19 March 2019



**Abstract:** Group 2 innate lymphoid cells (ILC2s) are tissue-resident cells and are a major source of innate  $T_H2$  cytokine secretion upon allergen exposure or parasitic-worm infection. Accumulating studies have revealed that transcription factors, including GATA-3, Bcl11b, Gfi1, ROR $\alpha$ , and Ets-1, play a role in ILC2 differentiation. Recent reports have further revealed that the characteristics and functions of ILC2 are influenced by the physiological state of the tissues. Specifically, the type of inflammation strongly affects the ILC2 phenotype in tissues. Inhibitory ILC2s, memory-like ILC2s, and ex-ILC2s with ILC1 features acquire their characteristic properties following exposure to their specific inflammatory environment. We have recently reported a new ILC2 population, designated as exhausted-like ILC2s, which emerges after a severe allergic inflammation. Exhausted-like ILC2s are featured with low reactivity and high expression of inhibitory receptors. Therefore, for a more comprehensive understanding of ILC2 function and differentiation, we review the recent knowledge of transcriptional regulation of ILC2 differentiation and discuss the roles of the Runx transcription factor in controlling the emergence of exhausted-like ILC2s. The concept of exhausted-like ILC2s sheds a light on a new aspect of ILC2 biology in allergic diseases.

**Keywords:** group 2 innate lymphoid cells; transcription factor; exhausted-like ILC2; Runx

## 1. Introduction

Innate lymphoid cells (ILCs) are lymphocytes responsible for innate cytotoxicity or helper functions [1–3]. While ILCs with cytotoxic capability are conventional natural killer (NK) cells which do not have tissue-resident properties, ILCs with helper functions generally reside in the tissue, provoke immune responses against pathogens, and maintain mucosal integrity [4,5]. Helper ILCs are unable to directly sense pathogenic antigens because they do not express antigen-specific receptors such as T cell receptor and B cell receptor. Instead, helper ILCs are generally stimulated by cytokines released from damaged-epithelial cells, phagocytes, and dendritic cells [6–11]. Based on cytokine production and requirements of transcription factors, helper ILCs are classified into three distinct subsets; ILC1s, ILC2s, and ILC3s which mirror  $T_H1$ ,  $T_H2$ , and  $T_H17$ , respectively [3,12]. ILC1s secrete interferon  $\gamma$  (IFN $\gamma$ ) and require T-bet for their differentiation [13]. T-bet is also important for the final maturation of conventional NK cells, which is highly dependent also on eomesodermin (Eomes) [14]. ILC2s are characterized by the production of the  $T_H2$  cytokines interleukin 4 (IL-4), IL-5, IL-13, and granulocyte-macrophage colony-stimulating factor (GM-CSF). All of these expressions are positively regulated by the transcription factor GATA binding protein 3 (GATA-3) [15]. Both RAR-related orphan receptor gamma t (ROR $\gamma$ t) and aryl hydrocarbon receptor (AhR) are required for the differentiation of ILC3s as well as for the production of IL-17 or IL-22 by ILC3s [16–18]. A substantial number of ILC1s

are found in the liver, intestinal intraepithelial layer, intestinal lamina propria, adipose tissue, and the skin [13,19–21]. ILC2s are preferentially localized in the lung, skin, adipose tissue, fat-associated lymphoid clusters, and the intestinal lamina propria [7,8,11,22]. ILC3s are abundant in the intestinal lamina propria, skin, and secondary lymphoid organs, such as Peyer's patches and mesenteric lymph nodes [16,18,23].

All ILC subsets develop from common lymphoid progenitor cells (CLPs) [24]. During the first commitment process from CLPs to the ILC lineage, early innate lymphoid progenitor cells (EILPs) are generated [25]. EILPs give rise to conventional NK cells and all helper ILC subsets. EILPs are marked by the expression of T cell factor 1 (TCF-1), thymocyte selection associated high mobility group box (Tox), and nuclear factor, interleukin 3 regulated (NFIL3) [25–27]. Progenitors specific to helper ILCs are Id2<sup>+</sup> common helper innate lymphoid progenitors (CHILPs) that lose the developmental potential for conventional NK cells [13]. Half of all CHILPs express promyelocytic leukemia zinc finger (PLZF) and are designated as innate lymphoid cell precursors (ILCPs) [10,24]. ILCPs can differentiate into helper ILCs except for a subpopulation of ILC3s, lymphoid tissue inducer-like (Lti-like) cells.

ILC2s have been extensively studied in the research field of allergy or parasitic worm infection [9]. Physiological roles of ILC2s are also implicated in metabolic homeostasis and virus infection [22,28,29]. Steady state ILC2s in tissues constitutively produce IL-5, which recruits eosinophils to the tissues [30]. Eosinophils are one of the essential components of allergy, as well as cytotoxic effector cells against parasitic worms. When mucosal tissues are injured by allergens or worms, IL-33, IL-25, and thymic stromal lymphopoietin (TSLP) are released from the damaged epithelial cells [31]. ILCs sense these signal cytokines by their cognate receptors and produce a variety of T<sub>H</sub>2 cytokines, including IL-5, IL-9, and IL-13. IL-9 is necessary for the maintenance of ILC2 activity in an autocrine manner during the early phase of an allergic reaction [32]. IL-13 from ILC2s induces chemokine CCL17 expression from dendritic cells to attract T<sub>H</sub>2 cells, resulting in enhanced T<sub>H</sub>2 responses [33]. ILC2s are an early source of amphiregulin, which is a ligand of the epidermal growth factor receptor and plays a critical role in the repair of damaged epithelial cells [28]. Therefore, ILC2s are involved in the initiation of allergy as well as tissue-repair.


The functional state of ILC2s is influenced by environmental cues. ILC2s acquire ILC1-like properties and produce IFN $\gamma$  when mice are infected with viruses and bacteria or are exposed to cigarette smoke to induce chronic obstructive pulmonary-like disease [34,35]. IL-1b, IL-12, and IL-18 are involved in the phenotypic conversion from ILC2s to ILC1-like cells. ILC2s also acquire "memory-like" signatures [36]. Once ILC2s are stimulated with IL-33, the activated ILC2s live long in the tissue and produce increased levels of T<sub>H</sub>2 cytokines upon a second challenge. Systemic IL-25 treatment and helminth infection induce inflammatory ILC2s (iILC2s), which are characterized by IL-17 production [37]. iILC2s migrate from the site of inflammation to the systemic tissues through lymphatics in an S1P-dependent manner [38]. Despite such migratory capacity of iILC2s, the majority of ILC2s reside in the tissue during and after helminth infection [4].

Transcription factors specifically required for ILC2 differentiation have been studied. Previous studies demonstrated the importance of transcription factors, such as GATA-3, B-Cell lymphoma/leukaemia 11B (Bcl11b), growth factor independent 1 transcription repressor (Gfi-1), RAR related orphan receptor A (ROR $\alpha$ ), and v-ets erythroblastosis virus E26 oncogene homolog 1 (Ets-1), for ILC2 differentiation [7,15,39–46]. ILC2s change their characteristics in response to environments. The mechanism by which transcription factor networks dictate phenotypical changes of ILC2s remains to be determined. For a comprehensive understanding of physiological roles of ILC2s in the allergic pathology, we summarize the current understanding of the transcriptional control of ILC2 differentiation and function, followed by our recent findings regarding how Runx-related transcription factor (Runx) proteins regulate the functional state of ILC2.

## 2. Transcription Factors Involved in the Development of Group 2 Innate Lymphoid Cells (ILC2s)

### 2.1. GATA-3

GATA-3 is necessary for the differentiation of CHILPs, which specifically give rise to all helper ILC subsets (Figure 1) [13]. While ILC1s and ILC3s express GATA-3, the expression level of GATA-3 in ILC2s is significantly higher compared to other ILCs. GATA-3 binds the gene loci which are critical for ILC2 activity. The genes positively regulated by GATA-3 include *Il5*, *Il13*, *Areg* encoding amphiregulin, *Il1rl1* encoding IL-33 receptor ST2, *Il9*, and *Il2ra* [15]. The conditional deletion of GATA-3 in mature ILC2s leads to reduced cytokine production and hypo proliferation [15,39,40]. Therefore, GATA-3 governs ILC2 identity, differentiation, homeostasis, and function.



**Figure 1.** Group 2 innate lymphoid cells (ILC2) differentiation and transcription factors. GATA-3 is expressed in early innate lymphoid progenitor cell (EILP), and its expression is maintained in all ILC subsets. Ets-1 up-regulates Id2 in common helper innate lymphoid progenitors (CHILP). Bcl11b-expressing progenitors in the CHILP population give rise to ILC2 (green blocked arrow). Bcl11b positively regulates ROR $\alpha$  and Gfi-1 both of which are indispensable for ILC2 differentiation (black blocked arrow from Bcl11b). Gfi-1 induces GATA-3 for ILC2 specification (black blocked arrow from Gfi-1). Runx proteins repress GATA-3 activity by antagonistic binding (red T-bar).

### 2.2. Bcl11b

Bcl11b is the earliest marker for the ILC2-lineage. The Bcl11b-expressing cells among the CHILP population specifically develop into ILC2s (Figure 1) [41]. Bcl11b ablation in hematopoietic stem cells impairs ILC2 differentiation in the bone marrow and peripheral tissues. Bcl11b expression in ILC subsets is controversial. While Yu et al. suggested that Bcl11b expression among ILC subsets was restricted to ILC2s [43], and Walker et al. showed that ILC1s and ILC3s expressed Bcl11b [42]. Conditional deletion of Bcl11b in ILC2s results in the generation of ILC2s with ILC3 gene signatures [41]. Thus, Bcl11b-deficient ILC2s have a reduction in GATA-3, Gfi-1, and ROR $\alpha$  expression, and an increase in sex determining region Y-box 4 (Sox4), ROR $\gamma$ t, and AhR expression. Bcl11b is associated with the *Gfi1* and *Ahr* gene loci, but not the *Gata3* or *Rorc* gene loci, suggesting that Gfi1 and AhR expression may be directly regulated by Bcl11b. Enhanced Gfi-1 expression in Bcl11b-deficient ILC2s restores GATA-3 expression and down-regulates ROR $\gamma$ t expression. Therefore, Bcl11b positively regulates GATA-3 expression and negatively regulates ROR $\gamma$ t expression through Gfi-1 induction.

### 2.3. *Gfi-1*

*Gfi-1* expression in ILC2 is relatively higher than other ILCs. *Gfi-1* is required for ILC2 differentiation and optimal  $T_H2$  cytokine responses to IL-33 [44]. *Gfi-1* positively regulates the expression of *Gata3* and *Il1rl1* (Figure 1). Deletion of *Gfi-1* causes aberrant IL-17a production by ILC2s due to dysregulated expression of ROR $\gamma$ t [44]. Increased ROR $\gamma$ t expression in *Gfi-1*-deficient ILC2s may be explained by increased expression of Sox4, which is known to induce ROR $\gamma$ t expression. Thus, *Gfi-1* suppresses the machinery for the ILC3 lineage and maintains the effector state of ILC2s in allergic diseases.

### 2.4. *ROR $\alpha$*

Although ROR $\alpha$  is expressed in CHILPs and all ILC subsets, it is specifically necessary for ILC2 differentiation among ILC subsets (Figure 1) [7,45,47]. A ROR $\alpha$ -dependent ILC2 differentiation mechanism has not been described yet. Differential interaction of ROR $\alpha$  with the gene loci in ILC2s and other ILCs should be examined to understand ROR $\alpha$  function in ILC2 differentiation and function. ROR $\alpha$ -deficient staggerer mice have been used to observe physiological effects resulting from ILC2 deficiency [7,45]. However, deletion of ROR $\alpha$  dampens regulatory T cell (Treg) and ILC3 function, but not their differentiation [48,49]. These data suggest that there might be other unknown immunological defects in the ROR $\alpha$ -deficient staggerer mice and the related mice with conditional deletion of ROR $\alpha$ .

### 2.5. *Ets1*

*Ets-1* induces *Id2* which is critical for differentiation of ILC progenitors, CHILPs [46]. However, for unknown reasons, ILC2s without *Ets-1* are reduced only in the bone marrow, but not in the lung. *Ets-1* is required for ILC2s to expand and produce IL-5 and IL-13 in response to IL-33. In contrast, IL-6 and IL-9 production are up-regulated in *Ets-1*-deficient ILC2s cultured with IL-33. It remains to be elucidated how *Ets-1* regulates the ILC2 fitness and function.

## 3. Runx Proteins and Immune Cells

### 3.1. Global Effects of Runx Proteins in Immune Cells

Runx proteins are a family of transcription factors necessary for many biological processes ranging from differentiation, function, proliferation, tumorigenesis, and cellular identity [50,51]. There are three mammal members in the Runx protein family: Runx1, Runx2, and Runx3. All Runx proteins require heterodimer formation with Subunit b of core binding factor (Cbf $\beta$ ) to exert their function as transcription factors. All three Runx genes generate two transcripts from distal (P1) and proximal (P2) promoters [52]. These two transcripts generate Runx protein isoforms with distinct N-terminal sequences. Utilization of the promoters differs among cell types. *Runx1* and *Runx3* genes are expressed in many types of immune cells [50]. Runx1 is required for the emergence of hematopoietic stem cells from hemogenic endothelium. Loss of Runx1 causes embryonic lethality due to bleeding in the brain caused by the absence of platelets [53,54]. Runx1 is also involved in the early thymocyte differentiation and development of invariant natural killer T (iNKT) cells, regulatory T cells and  $T_H17$  cells [51,55–58]. Runx3 is a critical transcription factor for the differentiation of CD8 $^+$  T cells and  $T_H1$  cells [56,59]. Runx3 is crucial to induce effector genes such as *Ifng*, *Gzmb*, *Eomes*, and to repress *Il4* gene in  $T_H1$  cells. *Runx3* transcript from the P1 promoter is correlated with protein expression level in T cells because the P2-*Runx3* transcript utilizes IRES-mediated translation [52,60]. Runx2 is required for the differentiation of osteoblasts and bone formation [61,62]. Runx2 expression in hematopoietic cells is rather restricted to plasmacytoid dendritic cells [63]. Runx2-deficient plasmacytoid dendritic cells have deficits in migrating to the peripheral tissues.

### 3.2. *Runx3 Is Required for the Differentiation of ILC1s and ILC3s*

Helper ILCs and conventional NK cells preferentially express *Runx3* mRNA among all *Runx* family genes [64,65]. However, ILC subsets are characterized by the levels of *Runx3* P1 transcripts and *Runx3* protein expression. High *Runx3* protein expression is observed in NK cells, which are innate counterparts of CD8<sup>+</sup>T cells, and in ILC1s, innate counterparts of T<sub>H</sub>1 cells. ILC3s, innate counterpart for T<sub>H</sub>17 cells, express an intermediate level of *Runx3*, while *Runx1* is likely to be a major *Runx* protein to drive T<sub>H</sub>17 program. *Runx3* expression is very low in ILC2s, innate lymphoid cells. Ablation of *Runx3* in all hematopoietic cells leads to a reduction of NK cells, ILC1s, and ILC3s but not ILC2s. *Runx3* is necessary for optimal IFN $\gamma$  production by ILC1s. Ablation of *Cbfb* or a combined loss of both *Runx1* and *Runx3* result in a dramatic reduction of NK cells and ILC1s, suggesting that *Runx1* compensates for the loss of *Runx3* in these cells. ILC1s fall into an apoptotic state in the absence of *Runx* proteins, which is a result of low expression of anti-apoptotic *Bcl2*. For the development of ILC3s, *Runx3* induces expression of an ILC3 master regulator gene, *ROR $\gamma$ t* mainly by direct binding to the intronic enhancer region of the *ROR $\gamma$ t* [66]. *Ahr* is also required for full production of IL-17 and IL-22 together with *ROR $\gamma$ t* [17]. *Runx3* deletion impairs expression of *Ahr* in ILC3s. Upon a half dosage of *ROR $\gamma$ t* expression caused by heterozygous gene knock-out, *Ahr* expression is reduced by half in the ILC3s. Thus, *Runx3* regulates *ROR $\gamma$ t* expression and its downstream target, *Ahr*.

Lymphoid tissue-inducer cells (Lti) cells are indispensable for secondary lymphoid organogenesis and are considered as a subpopulation of ILC3s because Lti cells are characterized by and are dependent on *ROR $\gamma$ t* expression. Lti cells appear in the embryonic intestine at embryonic day 12. An early study clarified the importance of *Runx1* for Lti cell differentiation via *ROR $\gamma$ t* induction in the embryos [67]. Although *ROR $\gamma$ t* expression in Lti-cells is reduced by *Runx3* deficiency, the number of Lti cells in the embryonic intestine is normal in the absence of *Runx3* [64]. These data suggest that there might be distinct functions of *Runx* proteins for the generation of Lti cells.

*Runx* proteins also regulate early ILC differentiation. *Runx1* and *Runx3* expression in ILC progenitors increases as CLPs give rise to CHILPs and ILCPs (Figure 1) [64]. Expression of *Runx1* and *Runx3* reaches the highest level at an ILCP stage in the course of ILC differentiation from CLPs to ILC subsets. *Runx* proteins are essential to induce *PLZF* expression, which is the marker for ILCPs, by direct interaction with an enhancer of the *Zbtb16* locus encoding *PLZF* [68]. ILC1s and ILC3s down-regulate *Runx1* expression but maintain intermediate to high levels of *Runx3* expression [64]. In contrast, down-regulation of *Runx3* occurs in ILC2 progenitor cells and ILC2s which express more *Runx1* than other ILCs. Therefore, the levels of *Runx3* expression specifies ILC lineages.

## 4. The Function of *Runx* Proteins in ILC2s


### 4.1. *Runx Proteins Prevent Steady-State ILC2s from Overactivation*

*Runx3* deletion is not enough for obvious phenotypic changes in ILC2s [64,69]. ILC2s lacking *Runx3* normally populate the peripheral tissues and are functionally intact because *Runx1* is also expressed and compensate *Runx3* function in ILC2s. When the function of all *Runx* proteins is abrogated by ablation of *Cbfb*, a binding partner of *Runx* proteins, in the ILC2s, the ILC2s are normally present in the tissues but exhibit an activated phenotype even in the steady-state condition. Killer cell lectin-like receptor G1 (KLRG1) is a known marker for activated ILC2s, and *Thy1* is down-regulated in the ILC2s when they are stimulated by IL-25. Interestingly, loss of *Runx* function in ILC2s induces high KLRG1 expression and low *Thy1* expression with unleashed IL-5 production. As ILC2s are a major source of IL-5, which recruits eosinophils to the lung, eosinophils in the bronchoalveolar space are increased by overproduction of IL-5 from the activated ILC2s lacking *Runx* function. The infiltration of eosinophils to the bronchoalveolar space seems to occur at subclinical levels because lung epithelial cells are not significantly damaged by the adoptive transfer of the *Cbfb*-deficient ILC2s.

The activated ILC2 phenotype caused by *Cbfb* deficiency is reminiscent of those observed in the T cells lacking *Cbfb* function. T<sub>H</sub>2 skewing occurs in CD4 T cells in the absence of *Cbfb*, resulting in

asthma-like symptoms in the lung [59]. Runx/Cbfb complexes associate and antagonize GATA-3, a master regulator of TH2 cells, in T cells [70]. Runx/Cbfb complexes also suppress the expression of IL-4, which is a critical TH2 cytokine for type 2 immunity, through binding to a silencer region in the *Il4* gene locus [59]. Therefore, the balance between Runx and GATA-3 dictates the functional state of TH2 cells.

ILC2 activity at steady state is also determined by the balancing action of Runx proteins against GATA-3 (Figure 2). While GATA-3 overexpression increases IL-5 production in ILC2s, Runx3 cancels such effects in ILC2s [69]. In addition to IL-5, transcriptome analysis demonstrates that many genes regulated by GATA-3 are inversely regulated by Runx proteins. In steady-state ILC2s, IL-5 production is increased by the absence of Cbfb, presumably via antagonizing GATA-3 function. However, we cannot deny the possibility that Runx/Cbfb complexes work as a direct repressor for the *Il5* gene as is the case for *Il4*, because Cbfb associates with several genomic regions around the *Il5* gene in ILC2s. Deletion of either Runx1 or Runx3 alone does not induce IL-5 overproduction from ILC2s. However, when both Runx1 and Runx3 are deleted, ILC2s show the activated phenotype, suggesting that the redundant function of Runx proteins restrains the basic activity of ILC2s in steady-state.



**Figure 2.** Runx/Cbfb complexes differentially regulate ILC2 effector function at steady state and in severe inflammation. In the steady state of the lung, Runx/Cbfb complexes suppress ILC2 activity through antagonistic binding to GATA-3 which positively regulates ILC2 effector function including constitutive IL-5 production (left red T-bar). During severe allergic airway inflammation, Runx/Cbfb complexes are essential for ILC2s to respond to IL-33 and maintain their activity. Runx/Cbfb complexes inhibit the emergence of exhausted-like ILC2s which are normally rare at the site of severe inflammation (right red T-bar). Red dots: IL-5. Blue dots: IL-13.

#### 4.2. Runx Proteins Inhibit the Emergence of Exhausted-Like ILC2s during Allergic Inflammation

ILC2s are activated by IL-33, IL-25, TSLP, TNF superfamily ligand TL1A (TL1A), and neuropeptides such as vasoactive intestinal polypeptide (VIP), neuromedin U and, calcitonin gene-related peptide [6,11,30,31,71–73]. Furthermore, IL-33 is the most potent ILC2 stimulator in vitro and in vivo and plays an important role in the pathogenesis of allergy. Interestingly, Cbfb-deficient ILC2s have defects in ILC2 cytokine production and proliferation during in vitro responses to IL-33 [69]. Expression of *Il5*, *Il9*, *Il13*, *Csf2* encoding GM-CSF, and *Areg* encoding amphiregulin upon IL-33 stimulation is all downregulated in Cbfb-deficient ILC2s. In addition, the expression of many molecules critical for ILC2 activity is dampened in the Cbfb-deficient ILC2s. ILC2s have machinery for extending immune responses upon their activation. For instance, IL-9 produced by ILC2s binds IL-9 receptor on ILC2s and stimulates their expansion [32]. Inducible costimulatory ligand (ICOSL) induced on the activated ILC2s contributes to their survival and cytokine production by receiving


ICOS signals [74]. Activated ILC2s express a variety of neuropeptide receptors including *Vipr2*, a VIP receptor, and *Nmur1*, a neuromedin receptor for expansion and cytokine production of ILC2s during allergy [30,71–73]. Surprisingly, expression of *Il9*, *Il9r*, *Icos*, *Vipr2*, and *Nmur1* in activated ILC2 are all impaired by *Cbfb*-deficiency [69]. Furthermore, ILC2s lacking *Cbfb* show up-regulated expressions of a series of T-cell exhaustion marker genes, such as *Tigit*, *Il10*, *Prdm1*, *Ctla4*, and *Lag3*. It has been documented that T cells acquire a state of dysfunction, called T cell exhaustion, after chronic exposure to cancer and viruses as a result of continuous antigen stimulation [75–77]. Exhausted T cells are characterized by impaired ability of proliferation, cytokine production and, effector function after re-stimulation. Expression of a variety of inhibitory receptors such as programmed cell death 1 (PD-1), T cell immunoreceptor with Ig and ITIM domains (*Tigit*), and lymphocyte Activation Gene-3 (*Lag3*) are highly induced on exhausted T cells. We, therefore, define hyporesponsive ILC2s with increased expression of T cell exhaustion markers as “exhausted-like” ILC2s.

Since both *Tigit* and IL-10 are highly expressed in *Cbfb*-deficient ILC2s, these two molecules could serve as good markers to define ILC2s with hyporesponsive characteristics. Indeed, a low number of hyporesponsive *Tigit*<sup>+</sup>IL-10<sup>+</sup> ILC2s emerge at the site of severe allergic inflammation even in *wild-type* mice [69]. Severe allergy is known to induce IL-10-producing ILC2s, which are associated with decreased eosinophil recruitment in the lung [26]. *Tigit* expression is restricted to high IL-10-producers in the activated ILC2s [69]. The low reactive *Tigit*<sup>+</sup>IL-10<sup>+</sup> ILC2s also express high levels of ILC2 activation markers such as PD-1, KLRG1, and glucocorticoid-induced tumor necrosis factor receptor (GITR). When mice are intranasally administered with papain, a proteinase allergen, the bronchoalveolar space is the most inflamed site in the lung. The *Tigit*<sup>+</sup>IL-10<sup>+</sup> ILC2s with low reactivity can be found only in the bronchoalveolar space, but not in the lung during severe subacute allergy induced by high doses of papain. The absence of *Cbfb* in ILC2s enhances the generation of the *Tigit*<sup>+</sup>IL-10<sup>+</sup> ILC2s and ameliorates allergic inflammation induced by papain inhalation in the subacute phase. However, when mice are chronically inoculated with papain for a month, increased low reactive *Tigit*<sup>+</sup>IL-10<sup>+</sup> ILC2s from *Cbfb* deficiency do not reduce the degree of chronic inflammation. If mice have the papain treatment interval for recovery after the first course of challenges, the emergence of the low reactive *Tigit*<sup>+</sup>IL-10<sup>+</sup> ILC2s enhanced by *Cbfb* deficiency is associated with attenuation of allergic inflammation induced by second challenges of papain treatment. Therefore, the presence of exhausted-like *Tigit*<sup>+</sup>IL-10<sup>+</sup> ILC2s is likely to reduce acute deterioration of chronic allergy.

Our data clearly show that pre-activated *Cbfb*-deficient ILC2s are hyporesponsive to IL-33 stimulation *in vitro* and *in vivo* [69]. If enhanced GATA-3 activation by the absence of *Cbfb* is involved in the generation of exhausted-like ILC2s, GATA-3 overexpression alone would lead to the emergence of exhausted-like ILC2s. However, ectopic expression of GATA-3 conversely elicits increased ILC2 reactivity upon IL-33 stimulation *in vitro* and *in vivo*. Of note, *Runx3* overexpression over GATA-3 expression down-modulates ILC2 responsiveness, suggesting that the reactivity of ILC2s to IL33 stimulation is regulated by the balance between GATA-3 and *Runx3* even in the activation phase (Figure 3). Conversely, transcription factors regulate gene expression through association with regulatory genomic regions such as promoters, enhancers, or repressors. Chromatin immunoprecipitation sequencing data suggest that the *Cbfb* binding pattern in ILC2s activated with IL-33 differs from that in ILC2s without IL-33 stimulation. Interestingly, *Cbfb*-binding peaks induced specifically upon IL-33 stimulation are enriched in regions nearby the signature genes for exhausted-like ILC2s. For example, the *Il5*, *Il13*, *Nmur1*, *Vipr2*, *Il10*, *Tigit*, *Prdm1*, *Lag3*, and *Ctla4* gene loci are marked by *Cbfb*-bindings specific to IL-33 stimulation. These *Cbfb*-binding peaks are not bound by GATA-3, suggesting that regulation of the activity of those possible regulatory regions by *Runx/Cbfb* is likely to be independent of GATA-3. Furthermore, *Runx/Cbfb* complexes bind the presumed enhancers of ILC2 functional gene loci marked by H3K27 acetylation in the IL-33-stimulated ILC2s (Figure 3). *Cbfb* binding peaks overlapping with H3K27 acetylation peaks are found around the *Il5*, *Il9*, *Il13*, *Areg*, and *Nmur1* gene loci. A decrease of H3K27 tri-methylation at the *Il10* gene by the absence of *Cbfb* suggests that *Cbfb* could repress the *Il10* locus through epigenetic mechanisms



(Figure 3). Therefore, Runx/Cbf $\beta$  complexes directly contribute to the phenotypical gene expression of exhausted like ILC2s.



**Figure 3.** Transcriptional regulation by Runx/Cbf $\beta$  complexes in ILC2s. Runx/Cbf $\beta$  complexes antagonize GATA-3 (red cross) and suppress the ILC2 activity irrespective of allergic inflammation in a dose dependent manner. The presence of Runx/Cbf $\beta$  complexes is indispensable for ILC2s not to acquire a gene expression profile of exhausted-like ILC2s. Runx/Cbf $\beta$  complexes function as activators for gene expression of ILC2 cytokines and receptors, and as repressors for inhibitory molecules, such as IL-10.

## 5. Conclusions

We reviewed how transcription factors regulate ILC2 differentiation and function. A new mechanism by which Runx proteins confer sustained ILC2 reactivity during allergic inflammation was identified. Exhausted-like ILC2s are a small population of over-activated ILC2s expressing inhibitory molecules. PD-1 is a good example of an inhibitory receptor that is induced on activated ILC2s and downmodulate ILC2 activity against worm infection [78]. However, PD-1 on ILC2s could be an activation marker rather than exhaustion marker, because PD-1<sup>+</sup> ILC2s generally produce more T<sub>H</sub>2 cytokines than PD-1<sup>-</sup> ILC2s. ILC2s produce IL-10 when the cells are exposed to severe allergic inflammation. However, higher T<sub>H</sub>2 cytokine production by IL-10-producing ILC2s suggests that IL-10-producing ILC2s are likely to be in more activated than non-IL-10 producing ILC2s [26]. We revealed that Tigit<sup>+</sup>IL-10<sup>+</sup> ILC2s have exhausted-like features at the site of severe inflammation. Exhausted-like ILC2s are very rare in the mouse model of subacute allergy as long as Runx proteins are functional. Whether exhausted-like ILC2s are present in humans remains elusive. However, if they are present, exhausted-like ILC2s might be accumulated in patients with chronic allergy and contribute to constraining allergic responses. Thus, exhausted-like ILC2s may serve as beneficial cells to those patients and have the therapeutic potency to be designated a new target for allergic disorders. Clinical trials are warranted to clarify the physiological effects of exhausted-like ILC2s in chronic allergy.

**Author Contributions:** T.E and I.T. discuss the concept and wrote this review article.

**Funding:** This research was funded by “Kibou Project 2016” Startup Support for Young Researchers in Immunology (T.E.), JSPS KAKENHI grant number 18H02647 (T.E.), Riken (Incentive Research Project, FY2017) (T.E.), Takeda Science Foundation (T.E.), Mochida Memorial Foundation for Medical and Pharmaceutical Research (T.E.), and Suzuken Memorial Foundation (T.E.).

**Conflicts of Interest:** The authors declare no conflict of interest.

## Abbreviations

GATA-3	GATA binding protein 3
Bcl11b	B-Cell lymphoma/leukaemia 11B
Gfi1	Growth factor independent 1 transcription repressor
ROR $\alpha$	RAR related orphan receptor A
Ets-1	v-ets erythroblastosis virus E26 oncogene homolog 1
Cbfb	Subunit b of core binding factor
Runx	Runt-related transcription factor
Tox	Thymocyte selection associated high mobility group box
NFIL3	Nuclear factor, interleukin 3 regulated
PLZF	Promyelocytic leukemia zinc finger
T-bet	T-box-containing protein expressed in T cells
Eomes	Eomesodermin
ROR $\gamma$ t	RAR-related orphan receptor gamma t
AhR	Aryl hydrocarbon receptor
TCF-1	T cell factor 1
ILC	Innate lymphoid cell
Sox4	Sex determining region Y-box 4
T <sub>H</sub>	Helper T cell
CLP	Common lymphoid progenitor
EILP	Early innate lymphoid progenitor
CHILP	Common helper innate lymphoid progenitors
ILCP	Innate lymphoid cell precursors
Lti	Lymphoid tissue inducer
IFN $\gamma$	Interferon gamma
IL	Interleukin
GM-CSF	Granulocyte-macrophage colony-stimulating factor
TSLP	Thymic stromal lymphopoietin
TL1A	TNF superfamily ligand TL1A
Vip	Vasoactive intestinal peptide
PD-1	Programmed cell death 1
Tigit	T cell immunoreceptor with Ig and ITIM domains
ICOS	Inducible costimulator
Lag3	Lymphocyte activation gene-3
KLRG1	Killer cell lectin-like receptor G1
GITR	Glucocorticoid-induced tumor necrosis factor receptor
ICOSL	Inducible costimulatory ligand
CCL17	CC chemokine ligand 17

## References

1. Ebbo, M.; Crinier, A.; Vely, F.; Vivier, E. Innate lymphoid cells: major players in inflammatory diseases. *Nat. Rev. Immunol.* **2017**, *17*, 665–678. [[CrossRef](#)] [[PubMed](#)]
2. Sonnenberg, G.F.; Artis, D. Innate lymphoid cells in the initiation, regulation and resolution of inflammation. *Nat. Med.* **2015**, *21*, 698–708. [[CrossRef](#)]
3. Vivier, E.; Artis, D.; Colonna, M.; Diefenbach, A.; di Santo, J.P.; Eberl, G.; Koyasu, S.; Locksley, R.M.; McKenzie, A.N.J.; Mebius, R.E.; et al. Innate Lymphoid Cells: 10 Years On. *Cell* **2018**, *174*, 1054–1066. [[CrossRef](#)] [[PubMed](#)]

4. Gasteiger, G.; Fan, X.; Dikiy, S.; Lee, S.Y.; Rudensky, A.Y. Tissue residency of innate lymphoid cells in lymphoid and nonlymphoid organs. *Science* **2015**, *350*, 981–985. [[CrossRef](#)] [[PubMed](#)]
5. Klose, C.S.; Artis, D. Innate lymphoid cells as regulators of immunity, inflammation and tissue homeostasis. *Nat. Immunol.* **2016**, *17*, 765–774. [[CrossRef](#)] [[PubMed](#)]
6. Von Moltke, J.; Ji, M.; Liang, H.E.; Locksley, R.M. Tuft-cell-derived IL-25 regulates an intestinal ILC2-epithelial response circuit. *Nature* **2016**, *529*, 221–225. [[CrossRef](#)] [[PubMed](#)]
7. Neill, D.R.; Wong, S.H.; Bellosi, A.; Flynn, R.J.; Daly, M.; Langford, T.K.; Bucks, C.; Kane, C.M.; Fallon, P.G.; Pannell, R.; et al. Nuocytes represent a new innate effector leukocyte that mediates type-2 immunity. *Nature* **2010**, *464*, 1367–1370. [[CrossRef](#)] [[PubMed](#)]
8. Moro, K.; Yamada, T.; Tanabe, M.; Takeuchi, T.; Ikawa, T.; Kawamoto, H.; Furusawa, J.; Ohtani, M.; Fujii, H.; Koyasu, S. Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)/Sca-1(+) lymphoid cells. *Nature* **2010**, *463*, 540–544. [[CrossRef](#)]
9. Kim, B.S.; Artis, D. Group 2 innate lymphoid cells in health and disease. *Cold Spring Harb. Perspect. Biol.* **2015**, *7*. [[CrossRef](#)]
10. Longman, R.S.; Diehl, G.E.; Victorio, D.A.; Huh, J.R.; Galan, C.; Miraldi, E.R.; Swaminath, A.; Bonneau, R.; Scherl, E.J.; Littman, D.R. CX(3)CR1(+) mononuclear phagocytes support colitis-associated innate lymphoid cell production of IL-22. *J. Exp. Med.* **2014**, *211*, 1571–1583. [[CrossRef](#)]
11. Kim, B.S.; Siracusa, M.C.; Saenz, S.A.; Noti, M.; Monticelli, L.A.; Sonnenberg, G.F.; Hepworth, M.R.; van Voorhees, A.S.; Comeau, M.R.; Artis, D. TSLP elicits IL-33-independent innate lymphoid cell responses to promote skin inflammation. *Sci. Transl. Med.* **2013**, *5*. [[CrossRef](#)] [[PubMed](#)]
12. Zook, E.C.; Kee, B.L. Development of innate lymphoid cells. *Nat. Immunol.* **2016**, *17*, 775–782. [[CrossRef](#)] [[PubMed](#)]
13. Klose, C.S.; Flach, M.; Mohle, L.; Rogell, L.; Hoyler, T.; Ebert, K.; Fabiunke, C.; Pfeifer, D.; Sexl, V.; Fonseca-Pereira, D.; et al. Differentiation of type 1 ILCs from a common progenitor to all helper-like innate lymphoid cell lineages. *Cell* **2014**, *157*, 340–356. [[CrossRef](#)] [[PubMed](#)]
14. Daussy, C.; Faure, F.; Mayol, K.; Viel, S.; Gasteiger, G.; Charrier, E.; Bienvenu, J.; Henry, T.; Debien, E.; Hasan, U.A.; et al. T-bet and Eomes instruct the development of two distinct natural killer cell lineages in the liver and in the bone marrow. *J. Exp. Med.* **2014**, *211*, 563–577. [[CrossRef](#)] [[PubMed](#)]
15. Yagi, R.; Zhong, C.; Northrup, D.L.; Yu, F.; Bouladoux, N.; Spencer, S.; Hu, G.; Barron, L.; Sharma, S.; Nakayama, T.; et al. The transcription factor GATA3 is critical for the development of all IL-7R $\alpha$ -expressing innate lymphoid cells. *Immunity* **2014**, *40*, 378–388. [[CrossRef](#)] [[PubMed](#)]
16. Lee, J.S.; Cella, M.; McDonald, K.G.; Garlanda, C.; Kennedy, G.D.; Nukaya, M.; Mantovani, A.; Kopan, R.; Bradfield, C.A.; Newberry, R.D.; et al. AHR drives the development of gut ILC22 cells and postnatal lymphoid tissues via pathways dependent on and independent of Notch. *Nat. Immunol.* **2012**, *13*, 144–151. [[CrossRef](#)]
17. Kiss, E.A.; Diefenbach, A. Role of the Aryl Hydrocarbon Receptor in Controlling Maintenance and Functional Programs of ROR $\gamma$ mat(+) Innate Lymphoid Cells and Intraepithelial Lymphocytes. *Front. Immunol.* **2012**, *3*, 124. [[CrossRef](#)]
18. Sawa, S.; Cherrier, M.; Lochner, M.; Satoh-Takayama, N.; Fehling, H.J.; Langa, F.; di Santo, J.P.; Eberl, G. Lineage relationship analysis of ROR $\gamma$ mat+ innate lymphoid cells. *Science* **2010**, *330*, 665–669. [[CrossRef](#)]
19. O’Sullivan, T.E.; Rapp, M.; Fan, X.; Weizman, O.E.; Bhardwaj, P.; Adams, N.M.; Walzer, T.; Dannenberg, A.J.; Sun, J.C. Adipose-Resident Group 1 Innate Lymphoid Cells Promote Obesity-Associated Insulin Resistance. *Immunity* **2016**, *45*, 428–441. [[CrossRef](#)]
20. Sojka, D.K.; Plougastel-Douglas, B.; Yang, L.; Pak-Wittel, M.A.; Artyomov, M.N.; Ivanova, Y.; Zhong, C.; Chase, J.M.; Rothman, P.B.; Yu, J.; et al. Tissue-resident natural killer (NK) cells are cell lineages distinct from thymic and conventional splenic NK cells. *eLife* **2014**, *3*, e01659. [[CrossRef](#)] [[PubMed](#)]
21. Fuchs, A.; Vermi, W.; Lee, J.S.; Lonardi, S.; Gilfillan, S.; Newberry, R.D.; Cella, M.; Colonna, M. Intraepithelial type 1 innate lymphoid cells are a unique subset of IL-12- and IL-15-responsive IFN- $\gamma$ -producing cells. *Immunity* **2013**, *38*, 769–781. [[CrossRef](#)] [[PubMed](#)]

22. Lee, M.W.; Odegaard, J.I.; Mukundan, L.; Qiu, Y.; Molofsky, A.B.; Nussbaum, J.C.; Yun, K.; Locksley, R.M.; Chawla, A. Activated type 2 innate lymphoid cells regulate beige fat biogenesis. *Cell* **2015**, *160*, 74–87. [[CrossRef](#)] [[PubMed](#)]
23. Pantelyushin, S.; Haak, S.; Ingold, B.; Kulig, P.; Heppner, F.L.; Navarini, A.A.; Becher, B. Ror $\gamma$ mat<sup>+</sup> innate lymphocytes and gammadelta T cells initiate psoriasiform plaque formation in mice. *J. Clin. Investig.* **2012**, *122*, 2252–2256. [[CrossRef](#)] [[PubMed](#)]
24. Constantinides, M.G.; McDonald, B.D.; Verhoef, P.A.; Bendelac, A. A committed precursor to innate lymphoid cells. *Nature* **2014**, *508*, 397–401. [[CrossRef](#)] [[PubMed](#)]
25. Yang, Q.; Li, F.; Harly, C.; Xing, S.; Ye, L.; Xia, X.; Wang, H.; Wang, X.; Yu, S.; Zhou, X.; et al. TCF-1 upregulation identifies early innate lymphoid progenitors in the bone marrow. *Nat. Immunol.* **2015**, *16*, 1044–1050. [[CrossRef](#)] [[PubMed](#)]
26. Seehus, C.R.; Kadavallore, A.; Torre, B.; Yeckes, A.R.; Wang, Y.; Tang, J.; Kaye, J. Alternative activation generates IL-10 producing type 2 innate lymphoid cells. *Nat. Commun.* **2017**, *8*, 1900. [[CrossRef](#)]
27. Yu, X.; Wang, Y.; Deng, M.; Li, Y.; Ruhn, K.A.; Zhang, C.C.; Hooper, L.V. The basic leucine zipper transcription factor NFIL3 directs the development of a common innate lymphoid cell precursor. *eLife* **2014**, *3*. [[CrossRef](#)] [[PubMed](#)]
28. Monticelli, L.A.; Sonnenberg, G.F.; Abt, M.C.; Alenghat, T.; Ziegler, C.G.; Doering, T.A.; Angelosanto, J.M.; Laidlaw, B.J.; Yang, C.Y.; Sathaliyawala, T.; et al. Innate lymphoid cells promote lung-tissue homeostasis after infection with influenza virus. *Nat. Immunol.* **2011**, *12*, 1045–1054. [[CrossRef](#)]
29. Chang, Y.J.; Kim, H.Y.; Albacker, L.A.; Baumgarth, N.; McKenzie, A.N.; Smith, D.E.; Dekruff, R.H.; Umetsu, D.T. Innate lymphoid cells mediate influenza-induced airway hyper-reactivity independently of adaptive immunity. *Nat. Immunol.* **2011**, *12*, 631–638. [[CrossRef](#)]
30. Nussbaum, J.C.; van Dyken, S.J.; von Moltke, J.; Cheng, L.E.; Mohapatra, A.; Molofsky, A.B.; Thornton, E.E.; Krummel, M.F.; Chawla, A.; Liang, H.E.; et al. Type 2 innate lymphoid cells control eosinophil homeostasis. *Nature* **2013**, *502*, 245–248. [[CrossRef](#)]
31. Halim, T.Y. Group 2 innate lymphoid cells in disease. *Int. Immunol.* **2016**, *28*, 13–22. [[CrossRef](#)]
32. Wilhelm, C.; Hirota, K.; Stieglitz, B.; van Snick, J.; Tolaini, M.; Lahl, K.; Sparwasser, T.; Helmby, H.; Stockinger, B. An IL-9 fate reporter demonstrates the induction of an innate IL-9 response in lung inflammation. *Nat. Immunol.* **2011**, *12*, 1071–1077. [[CrossRef](#)]
33. Halim, T.Y.; Hwang, Y.Y.; Scanlon, S.T.; Zaghoulani, H.; Garbi, N.; Fallon, P.G.; McKenzie, A.N. Group 2 innate lymphoid cells license dendritic cells to potentiate memory T<sub>H</sub>2 cell responses. *Nat. Immunol.* **2016**, *17*, 57–64. [[CrossRef](#)]
34. Silver, J.S.; Kearley, J.; Copenhaver, A.M.; Sanden, C.; Mori, M.; Yu, L.; Pritchard, G.H.; Berlin, A.A.; Hunter, C.A.; Bowler, R.; et al. Inflammatory triggers associated with exacerbations of COPD orchestrate plasticity of group 2 innate lymphoid cells in the lungs. *Nat. Immunol.* **2016**, *17*, 626–635. [[CrossRef](#)]
35. Ohne, Y.; Silver, J.S.; Thompson-Snipes, L.; Collet, M.A.; Blanck, J.P.; Cantarel, B.L.; Copenhaver, A.M.; Humbles, A.A.; Liu, Y.J. IL-1 is a critical regulator of group 2 innate lymphoid cell function and plasticity. *Nat. Immunol.* **2016**, *17*, 646–655. [[CrossRef](#)]
36. Martinez-Gonzalez, I.; Matha, L.; Steer, C.A.; Ghaedi, M.; Poon, G.F.; Takei, F. Allergen-Experienced Group 2 Innate Lymphoid Cells Acquire Memory-like Properties and Enhance Allergic Lung Inflammation. *Immunity* **2016**, *45*, 198–208. [[CrossRef](#)]
37. Huang, Y.; Guo, L.; Qiu, J.; Chen, X.; Hu-Li, J.; Siebenlist, U.; Williamson, P.R.; Urban, J.F., Jr.; Paul, W.E. IL-25-responsive, lineage-negative KLRG1(hi) cells are multipotential ‘inflammatory’ type 2 innate lymphoid cells. *Nat. Immunol.* **2015**, *16*, 161–169. [[CrossRef](#)]
38. Huang, Y.; Mao, K.; Chen, X.; Sun, M.A.; Kawabe, T.; Li, W.; Usher, N.; Zhu, J.; Urban, J.F., Jr.; Paul, W.E.; et al. S1P-dependent interorgan trafficking of group 2 innate lymphoid cells supports host defense. *Science* **2018**, *359*, 114–119. [[CrossRef](#)]
39. Klein Wolterink, R.G.; Serafini, N.; van Nimwegen, M.; Vosshenrich, C.A.; de Bruijn, M.J.; Fonseca Pereira, D.; Veiga Fernandes, H.; Hendriks, R.W.; di Santo, J.P. Essential, dose-dependent role for the transcription factor Gata3 in the development of IL-5<sup>+</sup> and IL-13<sup>+</sup> type 2 innate lymphoid cells. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 10240–10245. [[CrossRef](#)]

40. Hoyler, T.; Klose, C.S.; Souabni, A.; Turqueti-Neves, A.; Pfeifer, D.; Rawlins, E.L.; Voehringer, D.; Busslinger, M.; Diefenbach, A. The transcription factor GATA-3 controls cell fate and maintenance of type 2 innate lymphoid cells. *Immunity* **2012**, *37*, 634–648. [[CrossRef](#)]
41. Califano, D.; Cho, J.J.; Uddin, M.N.; Lorentsen, K.J.; Yang, Q.; Bhandoola, A.; Li, H.; Avram, D. Transcription Factor Bcl11b Controls Identity and Function of Mature Type 2 Innate Lymphoid Cells. *Immunity* **2015**, *43*, 354–368. [[CrossRef](#)]
42. Walker, J.A.; Oliphant, C.J.; Englezakis, A.; Yu, Y.; Clare, S.; Rodewald, H.R.; Belz, G.; Liu, P.; Fallon, P.G.; McKenzie, A.N. Bcl11b is essential for group 2 innate lymphoid cell development. *J. Exp. Med.* **2015**, *212*, 875–882. [[CrossRef](#)]
43. Yu, Y.; Wang, C.; Clare, S.; Wang, J.; Lee, S.C.; Brandt, C.; Burke, S.; Lu, L.; He, D.; Jenkins, N.A.; et al. The transcription factor Bcl11b is specifically expressed in group 2 innate lymphoid cells and is essential for their development. *J. Exp. Med.* **2015**, *212*, 865–874. [[CrossRef](#)]
44. Spooner, C.J.; Lesch, J.; Yan, D.; Khan, A.A.; Abbas, A.; Ramirez-Carrozzi, V.; Zhou, M.; Soriano, R.; Eastham-Anderson, J.; Diehl, L.; et al. Specification of type 2 innate lymphocytes by the transcriptional determinant Gfi1. *Nat. Immunol.* **2013**, *14*, 1229–1236. [[CrossRef](#)]
45. Halim, T.Y.; MacLaren, A.; Romanish, M.T.; Gold, M.J.; McNagny, K.M.; Takei, F. Retinoic-acid-receptor-related orphan nuclear receptor alpha is required for natural helper cell development and allergic inflammation. *Immunity* **2012**, *37*, 463–474. [[CrossRef](#)]
46. Zook, E.C.; Ramirez, K.; Guo, X.; van der Voort, G.; Sigvardsson, M.; Svensson, E.C.; Fu, Y.X.; Kee, B.L. The ETS1 transcription factor is required for the development and cytokine-induced expansion of ILC2. *J. Exp. Med.* **2016**, *213*, 687–696. [[CrossRef](#)]
47. Robinette, M.L.; Fuchs, A.; Cortez, V.S.; Lee, J.S.; Wang, Y.; Durum, S.K.; Gilfillan, S.; Colonna, M.; Immunological Genome, C. Transcriptional programs define molecular characteristics of innate lymphoid cell classes and subsets. *Nat. Immunol.* **2015**, *16*, 306–317. [[CrossRef](#)]
48. Malhotra, N.; Leyva-Castillo, J.M.; Jadhav, U.; Barreiro, O.; Kam, C.; O'Neill, N.K.; Meylan, F.; Chambon, P.; von Andrian, U.H.; Siegel, R.M.; et al. RORalpha-expressing T regulatory cells restrain allergic skin inflammation. *Sci. Immunol.* **2018**, *3*. [[CrossRef](#)]
49. Lo, B.C.; Gold, M.J.; Hughes, M.R.; Antignano, F.; Valdez, Y.; Zaph, C.; Harder, K.W.; McNagny, K.M. The orphan nuclear receptor ROR alpha and group 3 innate lymphoid cells drive fibrosis in a mouse model of Crohn's disease. *Sci. Immunol.* **2016**, *1*. [[CrossRef](#)]
50. Voon, D.C.; Hor, Y.T.; Ito, Y. The RUNX complex: reaching beyond haematopoiesis into immunity. *Immunology* **2015**, *146*, 523–536. [[CrossRef](#)]
51. Collins, A.; Littman, D.R.; Taniuchi, I. RUNX proteins in transcription factor networks that regulate T-cell lineage choice. *Nat. Rev. Immunol.* **2009**, *9*, 106–115. [[CrossRef](#)]
52. Bone, K.R.; Gruper, Y.; Goldenberg, D.; Levanon, D.; Groner, Y. Translation regulation of Runx3. *Blood Cells Mol. Dis.* **2010**, *45*, 112–116. [[CrossRef](#)]
53. Okuda, T.; van Deursen, J.; Hiebert, S.W.; Grosveld, G.; Downing, J.R. AML1, the target of multiple chromosomal translocations in human leukemia, is essential for normal fetal liver hematopoiesis. *Cell* **1996**, *84*, 321–330. [[CrossRef](#)]
54. Wang, Q.; Stacy, T.; Miller, J.D.; Lewis, A.F.; Gu, T.L.; Huang, X.; Bushweller, J.H.; Bories, J.C.; Alt, F.W.; Ryan, G.; et al. The CBFbeta subunit is essential for CBFalpha2 (AML1) function in vivo. *Cell* **1996**, *87*, 697–708. [[CrossRef](#)]
55. Taniuchi, I.; Osato, M.; Egawa, T.; Sunshine, M.J.; Bae, S.C.; Komori, T.; Ito, Y.; Littman, D.R. Differential requirements for Runx proteins in CD4 repression and epigenetic silencing during T lymphocyte development. *Cell* **2002**, *111*, 621–633. [[CrossRef](#)]
56. Egawa, T.; Tillman, R.E.; Naoe, Y.; Taniuchi, I.; Littman, D.R. The role of the Runx transcription factors in thymocyte differentiation and in homeostasis of naive T cells. *J. Exp. Med.* **2007**, *204*, 1945–1957. [[CrossRef](#)]
57. Kitoh, A.; Ono, M.; Naoe, Y.; Ohkura, N.; Yamaguchi, T.; Yaguchi, H.; Kitabayashi, I.; Tsukada, T.; Nomura, T.; Miyachi, Y.; et al. Indispensable role of the Runx1-Cbfbeta transcription complex for in vivo-suppressive function of FoxP3+ regulatory T cells. *Immunity* **2009**, *31*, 609–620. [[CrossRef](#)]

58. Zhang, F.; Meng, G.; Strober, W. Interactions among the transcription factors Runx1, ROR $\gamma$  and Foxp3 regulate the differentiation of interleukin 17-producing T cells. *Nat. Immunol.* **2008**, *9*, 1297–1306. [[CrossRef](#)]
59. Naoe, Y.; Setoguchi, R.; Akiyama, K.; Muroi, S.; Kuroda, M.; Hatam, F.; Littman, D.R.; Taniuchi, I. Repression of interleukin-4 in T helper type 1 cells by Runx/Cbfbeta binding to the Il4 silencer. *J. Exp. Med.* **2007**, *204*, 1749–1755. [[CrossRef](#)]
60. Kim, B.; Sasaki, Y.; Egawa, T. Restriction of Nonpermissive RUNX3 Protein Expression in T Lymphocytes by the Kozak Sequence. *J. Immunol.* **2015**, *195*, 1517–1523. [[CrossRef](#)]
61. Komori, T.; Yagi, H.; Nomura, S.; Yamaguchi, A.; Sasaki, K.; Deguchi, K.; Shimizu, Y.; Bronson, R.T.; Gao, Y.H.; Inada, M.; et al. Targeted disruption of Cbfa1 results in a complete lack of bone formation owing to maturational arrest of osteoblasts. *Cell* **1997**, *89*, 755–764. [[CrossRef](#)]
62. Otto, F.; Thornell, A.P.; Crompton, T.; Denzel, A.; Gilmour, K.C.; Rosewell, I.R.; Stamp, G.W.; Beddington, R.S.; Mundlos, S.; Olsen, B.R.; et al. Cbfa1, a candidate gene for cleidocranial dysplasia syndrome, is essential for osteoblast differentiation and bone development. *Cell* **1997**, *89*, 765–771. [[CrossRef](#)]
63. Sawai, C.M.; Sisirak, V.; Ghosh, H.S.; Hou, E.Z.; Ceribelli, M.; Staudt, L.M.; Reizis, B. Transcription factor Runx2 controls the development and migration of plasmacytoid dendritic cells. *J. Exp. Med.* **2013**, *210*, 2151–2159. [[CrossRef](#)]
64. Ebihara, T.; Song, C.; Ryu, S.H.; Plougastel-Douglas, B.; Yang, L.; Levanon, D.; Groner, Y.; Bern, M.D.; Stappenbeck, T.S.; Colonna, M.; et al. Runx3 specifies lineage commitment of innate lymphoid cells. *Nat. Immunol.* **2015**, *16*, 1124–1133. [[CrossRef](#)]
65. Levanon, D.; Negreanu, V.; Lotem, J.; Bone, K.R.; Brenner, O.; Leshkowitz, D.; Groner, Y. Transcription factor Runx3 regulates interleukin-15-dependent natural killer cell activation. *Mol. Cell. Biol.* **2014**, *34*, 1158–1169. [[CrossRef](#)]
66. Tenno, M.; Kojo, S.; Lawir, D.F.; Hess, I.; Shiroguchi, K.; Ebihara, T.; Endo, T.A.; Muroi, S.; Satoh, R.; Kawamoto, H.; et al. Cbfbeta2 controls differentiation of and confers homing capacity to prethymic progenitors. *J. Exp. Med.* **2018**, *215*, 595–610. [[CrossRef](#)]
67. Tachibana, M.; Tenno, M.; Tezuka, C.; Sugiyama, M.; Yoshida, H.; Taniuchi, I. Runx1/Cbfbeta2 complexes are required for lymphoid tissue inducer cell differentiation at two developmental stages. *J. Immunol.* **2011**, *186*, 1450–1457. [[CrossRef](#)]
68. Mao, A.P.; Ishizuka, I.E.; Kasal, D.N.; Mandal, M.; Bendelac, A. A shared Runx1-bound Zbtb16 enhancer directs innate and innate-like lymphoid lineage development. *Nat. Commun.* **2017**, *8*, 863. [[CrossRef](#)]
69. Miyamoto, C.; Kojo, S.; Yamashita, M.; Moro, K.; Lacaud, G.; Shiroguchi, K.; Taniuchi, I.; Ebihara, T. Runx/Cbfbeta complexes protect group 2 innate lymphoid cells from exhausted-like hyporesponsiveness during allergic airway inflammation. *Nat. Commun.* **2019**, *10*, 447. [[CrossRef](#)]
70. Yagi, R.; Junttila, I.S.; Wei, G.; Urban, J.F., Jr.; Zhao, K.; Paul, W.E.; Zhu, J. The transcription factor GATA3 actively represses RUNX3 protein-regulated production of interferon-gamma. *Immunity* **2010**, *32*, 507–517. [[CrossRef](#)]
71. Wallrapp, A.; Riesenfeld, S.J.; Burkett, P.R.; Abdunour, R.E.; Nyman, J.; Dionne, D.; Hofree, M.; Cuoco, M.S.; Rodman, C.; Farouq, D.; et al. The neuropeptide NMU amplifies ILC2-driven allergic lung inflammation. *Nature* **2017**, *549*, 351–356. [[CrossRef](#)]
72. Cardoso, V.; Chesne, J.; Ribeiro, H.; Garcia-Cassani, B.; Carvalho, T.; Bouchery, T.; Shah, K.; Barbosa-Morais, N.L.; Harris, N.; Veiga-Fernandes, H. Neuronal regulation of type 2 innate lymphoid cells via neuromedin U. *Nature* **2017**, *549*, 277–281. [[CrossRef](#)]
73. Klose, C.S.N.; Mahlakoiv, T.; Moeller, J.B.; Rankin, L.C.; Flamar, A.L.; Kabata, H.; Monticelli, L.A.; Moriyama, S.; Putzel, G.G.; Rakhilin, N.; et al. The neuropeptide neuromedin U stimulates innate lymphoid cells and type 2 inflammation. *Nature* **2017**, *549*, 282–286. [[CrossRef](#)]
74. Maazi, H.; Patel, N.; Sankaranarayanan, I.; Suzuki, Y.; Rigas, D.; Soroosh, P.; Freeman, G.J.; Sharpe, A.H.; Akbari, O. ICOS:ICOS-ligand interaction is required for type 2 innate lymphoid cell function, homeostasis, and induction of airway hyperreactivity. *Immunity* **2015**, *42*, 538–551. [[CrossRef](#)]
75. Wherry, E.J.; Kurachi, M. Molecular and cellular insights into T cell exhaustion. *Nat. Rev. Immunol.* **2015**, *15*, 486–499. [[CrossRef](#)]

76. Crespo, J.; Sun, H.; Welling, T.H.; Tian, Z.; Zou, W. T cell anergy, exhaustion, senescence, and stemness in the tumor microenvironment. *Curr. Opin. Immunol.* **2013**, *25*, 214–221. [[CrossRef](#)]
77. Yi, J.S.; Cox, M.A.; Zajac, A.J. T-cell exhaustion: characteristics, causes and conversion. *Immunology* **2010**, *129*, 474–481. [[CrossRef](#)]
78. Taylor, S.; Huang, Y.; Mallett, G.; Stathopoulou, C.; Felizardo, T.C.; Sun, M.A.; Martin, E.L.; Zhu, N.; Woodward, E.L.; Elias, M.S.; et al. PD-1 regulates KLRG1(+) group 2 innate lymphoid cells. *J. Exp. Med.* **2017**, *214*, 1663–1678. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).