

1 **Rescue behaviour in a social bird: removal of sticky ‘bird-catcher tree’**
2 **seeds by group members**

3

4 Martijn Hammers^{1,2*} & Lyanne Brouwer³

5

6 ¹ Behavioural and Physiological Ecology, GELIFES, University of Groningen, P.O. Box
7 11103, 9700 CC, Groningen, The Netherlands

8 ² Nature Seychelles, Roche Caiman, Mahé, Republic of Seychelles

9 ³ Division of Evolution & Ecology, Research School of Biology, Australian National
10 University, Canberra, 0200 ACT, Australia*

11

12 Correspondence: m.hammers@rug.nl

13

14 Short title: Rescue behaviour in a social bird

15

16 Keywords: *Acrocephalus sechellensis*, cooperative breeding, reciprocity, rescue behaviour,
17 *Pisonia grandis*

18

19

20 **Summary**

21 Rescue behaviour is a special form of cooperation in which a rescuer exhibits behaviours
22 directed towards averting a threat to an endangered individual, thereby potentially putting
23 itself at risk. Although rescue behaviour has been well-documented in experimental studies
24 on rats and ants, published cases in other non-human animals are rare. Here, we report
25 observations of rescue behaviour in the cooperatively breeding Seychelles warbler

26 (*Acrocephalus sechellensis*). In this species, individuals sometimes become entangled in seed
27 clusters of ‘bird-catcher trees’ (*Pisonia grandis*). Just one or a few of these sticky seeds can
28 prevent Seychelles warblers to fly and may lead to mortality. In four cases, individuals were
29 observed displaying behaviour aimed at removing sticky seeds from the feathers of an
30 entangled individual belonging to their group. Intriguingly, the rescuing individuals engaged
31 in this behaviour despite potentially risking entanglement. To our knowledge, this is the first
32 recorded case of rescue behaviour in birds.

33

34 **Introduction**

35 The question how and why individuals cooperate and engage in seemingly altruistic
36 behaviour has received much attention in the past decades and has been listed as one of the
37 ‘important 125 questions’ in science (Pennisi 2005). Rescue behaviour is a special form of
38 cooperative behaviour in which a rescuer exhibits behaviours directed towards averting a
39 threat to an endangered individual, thereby potentially putting itself at risk. From an
40 evolutionary perspective, such behaviour is intriguing because it is likely to be costly and not
41 necessarily beneficial for the individual displaying the behaviour in the short term.

42 Nowbahari & Hollis (2010) proposed a four-point definition of rescue behaviour in
43 order to stimulate research and to aid separating rescue behaviour from other forms of
44 cooperative behaviour: First, an individual must be in danger and likely to suffer physical
45 harm if the hazard is not eliminated. Second, the rescue behaviour is, or may be, costly for
46 the rescuer (i.e. the rescuer risks endangerment). Third, the rescuers’ action is appropriate for
47 the type of distress of the victim, independent of the outcome of the rescue event (i.e. the
48 rescue event is not necessarily successful). Finally, the rescuer gains no direct rewards in
49 exchange for the rescue action (e.g. food, mating opportunity), but may indirectly benefit

50 (e.g. future rewards, improve survival and reproduction of family members) (see also Hollis
51 & Nowbahari 2013a).

52 Although common in humans, rescue behaviour has only been described in a few
53 other animal taxa. The most well-known examples are experiments on restrained ants and
54 rats, where individuals were shown to exhibit behaviours aimed at rescuing a restrained
55 individual (Bartal et al. 2011; Hollis & Nowbahari 2013b). In the peer-reviewed literature,
56 observations from vertebrates in the wild are rare and mostly anecdotal. For example,
57 Chimpanzees (*Pan troglodytes*) have been reported to save group members that were attacked
58 by a leopard (*Panthera pardus*) (Boesch 1991) and white-faced capuchin monkeys (*Cebus*
59 *capucinus*) have been observed saving group members during an attack by another group
60 (Vogel & Fuentes-Jimenez 2009). Further, banded mongooses (*Mungos mungo*) have been
61 observed rescuing a group member from an attack by a martial eagle (*Polemaetus bellicosus*)
62 (Rood 1983) and humpback whales (*Megaptera novaeangliae*) rescue conspecifics and other
63 species from attacks by killer whales (*Orcinus orca*) (Pitman et al. 2017). However, in these
64 examples, individuals are not rescued from an inanimate trap as in experiments on rats and
65 ants, but from a predator. Therefore, it is unclear whether such cases represent rescue
66 behaviour as defined above or can be better described as cooperative self-defence against
67 predators attacking group members (see Barash 1976). It is likely that rescue behaviour
68 occurs in more group-living animals (Nowbahari & Hollis 2010; Hollis & Nowbahari 2013a),
69 but as far as we are aware rescue behaviour has not been reported in any birds yet.

70 Here, we report four cases of Seychelles warblers (*Acrocephalus sechellensis*)
71 attempting to rescue group members. In this species, individuals sometimes become
72 entangled in seed clusters of the tree *Pisonia grandis* (Figure 1). Just one or a few of the
73 sticky seeds of this ‘bird-catcher tree’ can prevent a Seychelles warbler from flying, and
74 frequently lead to mortality. In four cases, individuals were observed picking and pulling at

75 sticky seeds that were attached to the feathers of another group member. We discuss why this
76 behaviour qualifies as rescue behaviour and discuss its potential adaptive significance.

77

78 **Methods**

79 The Seychelles warbler is a small passerine (Figure 1) that is currently confined to five small
80 islands in Seychelles. In the population on the 27 ha Cousin Island (4°19'48"S 55°39'48"E),
81 which has been the subject of intensive study since 1985, *ca* 320 individuals occur in *ca* 115
82 territories (Komdeur 1992; Hammers et al. 2015). Seychelles warblers are cooperative
83 breeders, with groups defending their territory year-round. Fifty percent of territories contain
84 one to four additional independent (i.e. non-juvenile) subordinates in addition to the dominant
85 breeding pair (Brouwer et al. 2006; Kingma et al. 2016). Around 50% of these subordinates
86 show helping behaviour during the breeding season (e.g. territorial defence, nestbuilding,
87 offspring care) (Komdeur 1994a). Due to the absence of predators and the relatively benign
88 environment, extrinsic mortality is probably lower than in most other passerines (Hammers et
89 al. 2015). Indeed, Seychelles warblers show high annual survival (84% in adults, 61% in
90 juveniles; Brouwer et al. 2006) and the maximum recorded lifespan is 19 years (M.
91 Hammers; unpublished data). Seychelles warblers are insectivorous and glean the vast
92 majority of their food from leaves (Komdeur 1991). On Cousin Island, native *Pisonia grandis*
93 is the most common tree species and important for Seychelles warblers in terms of food and
94 nesting sites. Importantly, Seychelles warblers glean insects from *Pisonia* leaves, but do not
95 eat *Pisonia* seeds. *Pisonia* seed clusters contain from 12 to over 200 seeds, which become
96 extremely sticky when they are ripe and fall from the tree (Burger 2005). The seeds easily
97 attach to bird feathers and are very difficult to remove (Figure 1). While predominantly
98 ground-nesting seabirds (their long-distance seed dispersal vector; Walker 1991) become
99 entangled in *Pisonia* seeds, passerines might also risk entanglement (Burger 2005), especially

100 when they spend some time on the ground (where most ripe seeds are located). Seychelles
101 warblers spend most of their time foraging in the canopy where the risk of entanglement is
102 probably low, but may be exposed to sticky *Pisonia* seeds on the ground when collecting
103 nesting material (mainly dominant female breeders), or during territorial fights (both sexes).
104 Depending on the bird species, just a few seeds are sufficient to prevent an individual flying
105 and may cause mortality, which is likely a negative side effect of selection for extreme
106 stickiness of the seeds to facilitate dispersal (Burger 2005).

107 In 1999–2015, in each year during the main breeding season (June to September), and
108 in some years during the minor breeding season (December-March), Seychelles warblers
109 were recorded in all territories across the island (total = 21,781 resightings; $1,361 \pm 181$
110 (mean \pm SE) resightings per year [no observations were entered in 2000]) and regular mist-
111 netting sessions were performed (total = 3,517 catches; 207 ± 20 (mean \pm SE) catches per
112 year). Records of birds entangled in *Pisonia* seeds and rescue behaviour were collected
113 opportunistically (i.e. recorded whenever encountered).

114

115 **Results**

116 For Seychelles warblers, who spend most of their time foraging in the canopy, the risk of
117 entanglement in *Pisonia* seeds is generally low. Between 1999 and 2015, 35 individuals (17
118 dominants: 11 ♀, 6 ♂; 12 subordinates: 5 ♀, 7 ♂; 3 fledglings: 3 ♀, 0 ♂; 3 unringed individuals:
119 unknown sexes) were observed to have *Pisonia* seeds stuck to their feathers, ranging from
120 one or a few (<5) seeds (14 individuals) to individuals being completely covered in seeds
121 (typically >10 seeds) or with a seed cluster attached (21 individuals). Dominant female
122 breeders were almost twice as likely to become entangled as males, possibly because mainly
123 dominant females engage in nestbuilding behaviour (Komdeur 1991). At least 60% (N=21) of
124 entangled individuals showed difficulty flying or could not fly at the time of the observation.

125 Seychelles warblers observed to be entangled in *Pisonia* had a high risk of mortality, with
126 44% (N=14 out of 32 ringed individuals) of individuals not surviving after the current
127 breeding season. The rate of mortality is significantly higher than the previously recorded
128 population average of 8% mortality over a six-month period for adults (binomial test: $P <$
129 0.001) and the 19.5% mortality recorded for juveniles (binomial test: $P = 0.002$; Brouwer et
130 al. 2006). Thirteen of the 35 individuals with *Pisonia* seeds (37%) were caught by hand
131 (which was possible because these individuals had many seeds attached to their feathers and
132 could not fly) and the seeds were removed. Six of the 13 hand-caught individuals (46%)
133 survived until at least six months later. Three of the 35 individuals with *Pisonia* seeds were
134 caught using mist-nets and one of these three individuals survived. These three individuals
135 had 1-2 seeds attached to their wing or tail and were able to fly. The remaining eleven ringed
136 individuals that survived until at least six months later and were not caught and treated were
137 able to clear the *Pisonia* seeds in a natural way, as they were not observed with seeds in their
138 feathers during later observations.

139 While one individual was observed to remove seeds (with difficulty) from its own
140 feathers, we recorded four cases where other individuals helped removing the seeds and all
141 these individuals survived (i.e. in four of the eleven cases (36%) where individuals got rid of
142 the seeds naturally and survived until at least six months later). In these cases (twice in May
143 2004, July 2009, August 2009), the victim was entangled, flight performance was impaired
144 and the bird made alarm calls. One individual had seeds attached to both wings, whereas the
145 other three individuals were entangled in a seed cluster containing several seeds. The
146 rescuing individuals tried to remove seeds by picking and pulling at them. Although we could
147 not establish with certainty whether seeds were successfully removed, none of the individuals
148 had *Pisonia* seeds attached to their feathers on the next occasion that they were observed (i.e.
149 4, 8, 8, and 40 days later, respectively). The rescuer and the victim always belonged to the

150 same group, but had alternative social and genetic relationships: 1) a dominant male breeder
151 helping a subordinate female (father and daughter); 2) a dominant female breeder helping a
152 dominant male breeder (partners); 3) a subordinate male helping a dominant female breeder
153 (son to mother); 4) a dominant female breeder helping a male subordinate (not genetically
154 related, but the subordinate later became a helper in the same territory).

155

156 **Discussion**

157 We observed rare rescuing behaviour in group-living wild birds, in which a group member
158 showed behaviour aimed at removing sticky seeds from an endangered individual's feathers.
159 The seed removal behaviour fulfils all four criteria for rescue behaviour proposed by
160 Nowbahari & Hollis (2010) and Hollis & Nowbahari (2013). First, the individuals caught in
161 *Pisonia* seeds were clearly in distress and likely to have died if the seeds had not been
162 removed. Currently, we lack detailed information about the sources of extrinsic mortality in
163 the Seychelles warbler and future research should investigate whether *Pisonia* entanglement
164 contributes significantly to mortality in this species. The alarm calls produced by the victims
165 are perhaps 'calls-for-help' to alert other individuals that help is needed. Indeed, in this
166 species, individuals often produce alarm calls to alert and recruit group members, for
167 example when a nest predator approaches the nest or when a conspecific intruder is detected
168 in the territory. The observed behaviour is unlikely to be an extension of typical allopreening
169 (i.e. an individual preening another individual) behaviour, since, despite conducting
170 thousands of hours of field observations, we have not observed allopreening behaviour in this
171 species. Second, although we have not observed rescuers becoming entangled in seed
172 clusters, it appears likely that individuals that help other individuals to remove seeds may put
173 themselves at risk of also becoming entangled. This potential risk of entanglement, and the
174 associated high risk of mortality, may make this behaviour potentially much more costly than

175 other cooperative behaviours regularly observed in this species (e.g. food provisioning and
176 territorial defence). Third, picking and pulling at the seeds is appropriate behaviour to help
177 the victim. Finally, except perhaps in the case of the dominant female breeder helping her
178 partner, warblers do not appear to benefit directly from saving group members, as this does
179 not yield immediate improved access to reproduction or food. However, improving the
180 survival of group members is likely to yield indirect fitness benefits, including maintaining
181 the future reproductive success of related individuals (Brouwer et al. 2012). Interestingly, all
182 four observed cases of rescue behaviour were between members of the opposite sex.
183 Mortality of the entangled bird would have decreased the indirect fitness benefits of the
184 rescuer, because either a new breeder (most likely less related or less experienced) would
185 have taken up the breeding vacancy or the group would have lost a (future) helper (Komdeur
186 1994b). Apart from these indirect fitness benefits, in long-lived species like the Seychelles
187 warbler, where social bonds within groups may persist for several years, it is likely that
188 favours are returned later, an example of reciprocal altruism (e.g. Rutte & Taborsky 2007;
189 Freidin et al. 2015); rescue behaviour may therefore be adaptive in this long-lived
190 cooperatively breeding bird.

191 There is an ongoing discussion on whether rescue behaviour can be used as evidence
192 for behaviour driven by empathic emotions in non-human animals. Decety et al. (2016)
193 highlighted in a recent review that there is good evidence for basic forms of empathy in non-
194 human animals and that empathic behaviour probably mediates social behaviour, but also that
195 many debates originate from using different definitions of empathy. The authors of a study on
196 rescue behaviour in rats argued that rescue can be interpreted as behaviour driven by empathy
197 (i.e. individuals responding to the needs of others; Bartal et al. 2011), while others have
198 argued that this conclusion is premature and that the results can be interpreted differently
199 (e.g. Vasconcelos et al. 2012; Silberberg et al. 2014). For example, individuals may not show

200 rescue behaviour in order to remove a threat, but merely to stop the distress signals of a
201 distressed animal, or to re-establish a social connection arising from isolation of the rescuer.
202 Unfortunately, our observations do not allow us to conclude if empathy plays a role, but we
203 hope that these observations encourage further study on the causes and consequences of
204 seemingly altruistic rescue behaviour in non-human animals.

205

206 **Acknowledgments**

207 We thank Nature Seychelles and the Cousin Island staff for providing accommodation and
208 facilities during our visits and the Department of Environment and the Seychelles Bureau
209 of Standards for permission for fieldwork and sampling. We thank many people for their
210 invaluable contributions to fieldwork, lab work and database management. We thank Terry
211 Burke, Owen Howison, Sjouke Kingma and two anonymous reviewers for helpful
212 comments on the manuscript. The Seychelles warbler research project is coordinated by
213 Jan Komdeur, David S. Richardson, Terry Burke and Hannah Dugdale and supported by
214 grants from NERC and the Netherlands Organisation for Scientific Research (NWO). MH
215 is supported by a NWO VENI fellowship (863.15.020), LB is supported by an Australian
216 Research Council DECRA fellowship (DE130100174).

217

218 **References**

- 219 Barash, D. P. (1976). Mobbing behavior by crows: the effect of the "crow-in-distress"
220 model." *The Condor* 78, 120-120.
- 221 Bartal, I. B. A., Decety, J., & Mason, P. (2011). Empathy and pro-social behaviour in rats.
222 *Science*, 334, 1427-1430.
- 223 Boesch, C. (1991). The effects of leopard predation on grouping patterns in forest
224 chimpanzees. *Behaviour*, 117, 220-241.

225 Brouwer, L., Richardson, D. S., Eikenaar, C., & Komdeur, J. (2006). The role of group size
226 and environmental factors on survival in a cooperatively breeding tropical passerine. *Journal*
227 *of Animal Ecology*, *75*, 1321-1329.

228 Brouwer, L., Richardson, D. S., & Komdeur, J. (2012). Helpers at the nest improve late-life
229 offspring performance: evidence from a long-term study and a cross-foster experiment. *PloS*
230 *ONE*, *7*, e33167.

231 Burger, A. E. (2005). Dispersal and germination of seeds of *Pisonia grandis*, an Indo-Pacific
232 tropical tree associated with insular seabird colonies. *Journal of Tropical Ecology*, *21*, 263-
233 271.

234 Decety, J., Bartal, I. B. A., Uzefovsky, F., & Knafo-Noam, A. (2016). Empathy as a driver of
235 prosocial behaviour: highly conserved neurobehavioural mechanisms across
236 species. *Philosophical Transactions of the Royal Society B*, *371*, 20150077.

237 Freidin, E., Carballo, F., & Bentosela, M. (2015). Direct reciprocity in animals: The roles of
238 bonding and affective processes. *International Journal of Psychology*.
239 doi:10.1002/ijop.12215

240 Hammers, M., Kingma, S. A., Bebbington, K., van de Crommenacker, J., Spurgin, L. G.,
241 Richardson, D. S., Burke, T., Dugdale, H. L., & Komdeur, J. (2015). Senescence in the wild:
242 Insights from a long-term study on Seychelles warblers. *Experimental gerontology*, *71*, 69-
243 79.

244 Kingma, S. A., Bebbington, K., Hammers, M., Richardson, D. S., & Komdeur, J. (2016).
245 Delayed dispersal and the costs and benefits of different routes to independent breeding in a
246 cooperatively breeding bird. *Evolution*, *70*, 2595-2610.

247 Hollis, K. L., & Nowbahari, E. (2013a). Toward a behavioral ecology of rescue behavior.
248 *Evolutionary Psychology*, *11*, 147470491301100311.

249 Hollis, K. L., & Nowbahari, E. (2013b). A comparative analysis of precision rescue
250 behaviour in sand-dwelling ants. *Animal Behaviour*, 85, 537-544.

251 Komdeur, J. (1991). *Cooperative breeding in the Seychelles warbler* (Doctoral dissertation,
252 University of Cambridge).

253 Komdeur, J. (1992). Importance of habitat saturation and territory quality for evolution of
254 cooperative breeding in the Seychelles warbler. *Nature*, 358, 493-495.

255 Komdeur, J. (1994a). Experimental evidence for helping and hindering by previous offspring
256 in the cooperative-breeding Seychelles warbler *Acrocephalus sechellensis*. *Behavioral*
257 *Ecology and Sociobiology*, 34, 175-186.

258 Komdeur, J. (1994b) The effect of kinship on helping in the cooperative breeding Seychelles
259 warbler (*Acrocephalus sechellensis*). *Proceedings of the Royal Society of London B:*
260 *Biological Sciences* 256: 47-52.

261 Nowbahari, E., & Hollis, K. L. (2010). Rescue behavior: Distinguishing between rescue,
262 cooperation and other forms of altruistic behavior. *Communicative & integrative biology*, 3,
263 77-79.

264 Pennisi, E. (2005). How did cooperative behavior evolve? *Science*, 309, 93-93.

265 Pitman, R. L., Deecke, V. B., Gabriele, C. M., Srinivasan, M., Black, N., Denking, J.,
266 Durban, J. W., Mathews, E. A., Matkin, D. R., Neilson, J. L., Schulman-Janiger, A.,
267 Shearwater, D. Stap, P., & Ternullo, R. (2017). Humpback whales interfering when mammal-
268 eating killer whales attack other species: Mobbing behavior and interspecific
269 altruism? *Marine Mammal Science*, 33, 7-58.

270 Rood, J. P. (1983). Banded mongoose rescues pack member from eagle. *Animal*
271 *Behaviour*, 31, 1261-1262.

272 Rutte, C., & Taborsky, M. (2007). Generalized reciprocity in rats. *PLoS Biology*, 5, e196.

273 Silberberg, A., Allouch, C., Sandfort, S., Kearns, D., Karpel, H., & Slotnick, B. (2014).
274 Desire for social contact, not empathy, may explain “rescue” behavior in rats. *Animal*
275 *cognition*, 17, 609-618.

276 Vasconcelos, M., Hollis, K., Nowbahari, E., & Kacelnik, A. (2012). Pro-sociality without
277 empathy. *Biology letters*, 8(6), 910-912.

278 Vogel, E. R., & Fuentes-Jiménez, A. (2006). Rescue behavior in white-faced capuchin
279 monkeys during an intergroup attack: support for the infanticide avoidance
280 hypothesis. *American journal of primatology*, 68, 1012-1016.

281 Walker, T.A. (1991). *Pisonia* islands of the Great Barrier Reef I. The distribution, abundance
282 and dispersal by seabirds of *Pisonia grandis*. *Atoll Research Bulletin*, 350, 1-23.

283 **Figures**

284 Figure 1. A Seychelles warbler entangled in a seed cluster of *Pisonia grandis*. The seeds are
285 extremely sticky and easily attach to feathers. This individual (a dominant male) was unable
286 to fly and was attacked by skinks. It was captured by hand and the seeds were removed, after
287 which it survived for another six years. Picture by Martijn Hammers.



288