

Access to dialysis services: A systematic mapping review based on geographical information systems

Benyamin Hoseini,^{1,2} Nasser Bagheri,³ Behzad Kiani,² Amirabbas Azizi,⁴ Hamed Tabesh,² Mahmood Tara²

¹*Department of Health Information Technology, Neyshabur University of Medical Sciences, Neyshabur, Iran;* ²*Department of Medical Informatics, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran;* ³*Department of Health Services Research and Policy, College of Health and Medicine, Biology and Environment, Australian National University, Canberra, ACT, Australia;* ⁴*Department of Health Information Technology, School of Allied Health Professions, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran*

Correspondence: Mahmood Tara, Department of Medical Informatics, Faculty of Medicine, Mashhad University of Medical Sciences, Park Square, Mashhad, Iran
Tel.: +985138002428 - Fax: +985138002445.
E-mail: smtara@gmail.com

Key words: Access; Dialysis; Geographical information systems; Floating catchment; Renal.

Contributions: NB designed the study; BH searched for the literature. Then, the EndNote X5 (Thomson Reuters, New York, NY, USA) reference management software was applied to aggregate all the search strategy returns. Two reviewers (BH and BK) screened the exported articles in EndNote and compared them with the inclusion and exclusion criteria independently. Any disagreement was resolved by another reviewer (MT), who was also responsible for the supervision of the research. Each study was assessed independently by two reviewers for the eligibility criteria. The full text of the qualified studies was retrieved and saved. The eligible papers were read, tagged and summarized by one reviewer (BH) and then verified by the second reviewer (MT). Quality scores were assigned by two reviewers (BH and AA) and verified by the third reviewer (HT). BH wrote the draft of the paper. BK and BH reviewed the paper and made revisions.

Acknowledgements: the authors would like to thank the authors of all the studies they reviewed.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: the work was supported by a grant [941344] from Mashhad University of Medical Sciences Research Council.

Received for publication: 16 April 2017.

Revision received: 28 November 2017.

Accepted for publication: 28 November 2017.

©Copyright B. Hoseini et al., 2018
Licensee PAGEPress, Italy
Geospatial Health 2018; 13:577
doi:10.4081/gh.2018.577

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Abstract

Equitable access to healthcare services constitutes one of the leading priorities of healthcare provision and *access to dialysis services* (ADS) has an essential impact on patients depending on renal dialysis. The many existing GIS-based ADS evaluations include various spatial and non-spatial factors affecting ADS. We systematically mapped and reviewed the available literature with reference to this area identifying gaps in current GIS-based ADS measurements and developing recommendations for future studies. A three-step, systematic mapping review of the available GIS-related evidence in PubMed, Embase, Web of science, Scopus, Science Direct and IEEE Xplore was performed in May 2016 and the information collected updated October 2017 by two independent selection processes. The quality of the studies was assessed using an informal, mixed-approach scoring system. Out of 1119 literature references identified, 36 were identified and used for final review after removal of duplicates, study screenings and applying inclusion/exclusion criteria. Given the contents of the selected studies, three study groups were identified and 41 factors with potential effects on ADS determined. These studies mainly addressed the potential and/or spatial aspects of ADS. Our systematic mapping review of the evidence revealed that current GIS-based measures of ADS tend to calculate potential ADS instead of a realized one. It was also noted that listed factors affecting ADS were mainly non-spatial bringing forth the hypothesis that designing an integrated ADS index could possibly produce better ADS score than those currently advocated. Some primary and secondary research suggestions are made and a list of recommendations offered.

Introduction

Although equitable *access to healthcare services* (AHS) is a leading public health priority, the importance of AHS and the attention received from the policy managers differ from one disease to another. *Access to dialysis services* (ADS) is a crucial necessity for patients who have to travel to a dialysis facility three times a week (Mactier, 2007; Stephens et al., 2013). Thus, poorly developed ADS lead to poor health outcomes such as increased morbidity and mortality (Moist et al., 2008; Diamant et al., 2010; Rucker et al., 2011; Thompson et al., 2012).

A vital aspect of AHS is the ease with which they can be



accessed and used when needed (McLafferty, 2003). AHS has five main dimensions: accessibility, availability, accommodation, affordability and acceptability (Levesque *et al.*, 2013; Russell *et al.*, 2013; Saurman, 2016). The accessibility and availability dimensions are usually related to geographical factors and are therefore labelled *spatial accessibility* (Mao and Nekorchuk, 2013), while non-geographic-dependent dimensions such as affordability, accommodation and acceptability are considered non-spatial (Guagliardo, 2004). Additionally, AHS can be divided into two broad categories: potential AHS and realized AHS. The former is simply defined as the presence of enabling healthcare, while the latter is the actual use of its services (Andersen, 1995).

Geographic information systems (GISs) enable researchers in the healthcare field to apply combined spatial accessibility measures for the inspection of the equitability of resource allocation. Various studies have demonstrated the application of GIS tools which are useful for calculation and visualization of accessibility scores (Guagliardo, 2004; Yang *et al.*, 2006; Matsumoto *et al.*, 2012; Kiani *et al.*, 2017). Stephens *et al.* (2013) performed a GIS-based measure of ADS which included travel impedance to dialysis facilities as an indicator. A recent study performed in Iran by Kiani *et al.* (2017) revealed the importance of spatial accessibility to dialysis services and showed that this variable is strongly underestimated when potential travel time is calculated (Kiani *et al.*, 2017). They developed a comprehensive measure of revealed accessibility that includes travel time and some other spatial and also non-spatial factors into one indicator framework. However, their index does not include facility capacity. Matsumoto *et al.* (2012) designed an algorithm embedded in a GIS-based measure of ADS which demonstrates that patients cannot always be accepted by the nearest dialysis facility due to limited capacity. Another study performed in the US emphasized the acceptability dimension and revealed that dialysis patients may have a short travel time to one dialysis facility but might decide to go to another centre due to ethnic disparity (Saunders *et al.*, 2014). Some studies, mainly conducted in Britain (Roderick *et al.*, 1999; Christie *et al.*, 2005; White *et al.*, 2006; Judge *et al.*, 2012), calculated deprivation as an important ADS indicator which includes some major non-spatial factors. Deprivation can also potentially affect *need of dialysis services* (NDS) (Roderick *et al.*, 1999; Thomas, 2005; Yang *et al.*, 2006; Judge *et al.*, 2012). NDS is defined as the number of patients in an area who need dialysis (Yang *et al.*, 2006).

The diversity of GIS-based ADS measures include various indicators with spatial as well as non-spatial confounding factors that may confuse researchers and policymakers. A mapping exercise could potentially lead to a more conclusive index producing better scores than currently used. GIS-based methods are inherently spatial; some of them, such as the *two-step floating catchment area* (2SFCA), demonstrated the capability of integrating both spatial and non-spatial AHS factors into one framework (Bagheri *et al.*, 2008; McGrail and Humphreys, 2009; McGrail and Humphreys, 2015). Yang *et al.* (2006) provide a 2SFCA platform that integrates only dialysis patients and dialysis machines within a 30-minutes potential travel time catchment. Although this represents a certain progress, to the best of our knowledge, no study so far has made an attempt to develop a truly integrated GIS-based ADS index. In an effort to do so, we decided to focus on the gaps in the current approach to GIS-based ADS measures through a systematic review of the available literature. In addition, we aimed to develop recommendations and a list of factors affecting ADS that would improve the research on the measuring ADS based on GIS.

Materials and Methods

A systematic mapping review of the available GIS-based literature was performed. It aimed to describe the extent of the study on a particular topic and to identify knowledge gaps in the study base, where further primary and/or secondary studies are needed (Grant and Booth, 2009).

Search strategy

The scientific literature was explored with regard to relevant communications on ADS and the use of GIS. It was done in May 2016 and the information collected updated in October 2017. We included the following electronic databases: PubMed, Web of Science, Scopus, ScienceDirect, EMBASE and IEEE Xplore. Initially, the databases were systematically searched using a variation of access concept (access, accessibility, availability, affordability, acceptability, accommodation, utilization, deprivation, disparity and equity) in connection with spatial terms (geographic information, GIS, geomapping, location-allocation, and spatial analysis) and dialysis (dialysis, haemodialysis, renal and kidney). To combine the search terms within each category, we utilized the disjunction *OR*, and to combine categories we utilized the conjunction *AND*.

In order to identify additional relevant information related to ADS and GIS, the reference lists obtained were surveyed manually. To boost our search strategy further, we also looked for the so called gray literature, *i.e.* reports, standards, manuals and guidelines on the topic using general search engines such as Google. No date or study design limitation was imposed in any of the research steps described. The complete search strategy is available upon request.

Eligibility criteria

After the literature search had been completed, the EndNote X5 (Thomson Reuters, New York, NY, USA) reference management software was applied to aggregate all search returns. The articles were then screened and each study assessed independently for eligibility by two of the authors in different combinations. A study was considered eligible for inclusion if it included assessment of ADS as the primary or secondary outcome, while it was excluded if i) it had not clearly reported and calculated ADS indicators; ii) it lacked a methodological description of the measurement of ADS or its indicators; or iii) it consisted of a letter to the editor, an editorial, general comments, a position paper or it was an unstructured paper.

The full text of the qualified studies was read, tagged and summarized by one author and verified by one other author. A brief flow diagram of the strategy is depicted in Figure 1.

Data extraction

Studies deemed eligible for review underwent data extraction. For each paper, essential data items related to ADS measurement were extracted and fitted into a form with a choice of headings, such as ADS indicator; Factor(s) affecting the indicator; Method of measurement; Primary outcome measures; Secondary outcome measures; and Study design. Additional properties, such as conclusive comments and suggested measurement intervals, were also recorded when available and applicable.

Quality assessment

Since systematic mapping reviews mainly aim to describe the state of the art of a particular topic, it is desirable to include communication types of a range as wide as possible. Due to the high diversity of study types in our review, this necessitated an informal quality assessment that was performed by classifying the literature by type of study. We used a mixed-approach scoring system as applied by Azizi *et al.* (2016) under similar circumstances (Azizi *et al.*, 2016). Quality scores were assigned by two authors and verified by a third author. The weight of the literature was assigned according to their study design by the quality scoring system. In this approach, papers such as reviews and *randomized controlled trials* (RCTs) achieved the highest score (score 4) while score one was the lowest score. In this approach, the gray literature was given score one as formal or expert consensus regarding quality score. The summary of the quality assessment approach is outlined in Table 1.

Data analysis

Following mapping review methods, a tabular method for the synthesis of qualitative research findings was used. Also, the quantity and quality of the literature were characterized by study design and other key features, and need for some primary or secondary research was identified.

Results

Out of 1119 literature items collected, 76 were deemed eligible for further full-text review. After reviewing the full-text studies for final content match, 36 were selected for the review. Further details pertaining to the included studies are shown in Figure 1.

Characteristics of included studies

Three groups of studies were identified. The first group included eight cross-sectional studies addressing ADS and treating it as a primary outcome. The focus of these communications was the design of a GIS-based model intended for gauging the degree of ADS. They mostly calculated potential travel time (Roderick *et al.*, 1999; Christie *et al.*, 2005; White *et al.*, 2006; Yang *et al.*, 2006; Judge *et al.*, 2012; Matsumoto *et al.*, 2012; Stephens *et al.*, 2013) or facility capacity (White *et al.*, 2006; Yang *et al.*, 2006; Matsumoto *et al.*, 2012) as the key indicators of spatial accessibility. Some papers in this group also considered deprivation (Roderick *et al.*, 1999; Christie *et al.*, 2005; White *et al.*, 2006; Judge *et al.*, 2012) as an important, non-spatial indicator of ADS. One study designed a measure to calculate actual travel time revealed significant effects on the travel time of other non-spatial factors such as sex, income level, caregivers, transportation mode, education level, ethnicity and personal vehicle ownership. It demonstrated the large gap between potential travel time and actual travel time (Kiani *et al.*, 2017). Another study showed that ignoring facility capacity and accounting only for travel time when evaluating ADS may result in misleading conclusions (Matsumoto *et al.*, 2012). Further details pertaining to this group are outlined in Table 2.

The second group of studies (10 peer-reviewed papers) considered gauging ADS as a secondary outcome, while their primary outcome measures focused on the association between ADS and health-related outcomes (Tonelli *et al.*, 2007; Moist *et al.*, 2008; Diamant *et al.*, 2010; Rucker *et al.*, 2011; Thompson *et al.*, 2012;

Miller *et al.*, 2014) determining the relationship between ADS and prevalence rates of dialysis patients (Kashima *et al.*, 2012) or designing models to locate dialysis facilities (Ayyalasomayajula *et al.*, 2011; Salgado *et al.*, 2011; Faruque *et al.*, 2012). Almost all of them calculated potential travel impedance as an indicator of ADS based on GIS.

Finally, the third group of 16 studies considered factors affecting ADS. Among them, were ten peer-reviewed articles (Smith *et al.*, 1985; Tonelli *et al.*, 2006; Hall *et al.*, 2008; Prakash *et al.*, 2010; Matsumoto *et al.*, 2013; Omrani-Khoo *et al.*, 2013; Rodriguez *et al.*, 2013; Plantinga *et al.*, 2014; Saunders *et al.*, 2014; Kiani *et al.*, 2017), one case series (Tshamba *et al.*, 2014), one proceeding (Richard *et al.*, 2009), and four gray literature items (Maheswaran *et al.*, 2003; Mactier, 2007; Levinson, 2011; Amy Martin, 2013).

Considering the contents of the total number of GIS-based studies discussed here and the literature extracted from their comprehensive reference lists, 41 factors affecting ADS were determined (Table 3). Travel impedance, especially travel time, were the main indicators of spatial accessibility, the rest were mostly non-spatial. Some studies emphasize that factors such as ethnicity or the patient's health insurance status also affect ADS (Kashima *et al.*, 2012; Thompson *et al.*, 2012; Saunders *et al.*, 2014), while other studies imply that they do not affect ADS (White *et al.*, 2006; Matsumoto *et al.*, 2012). The factors we found in our literature search and their frequency are given in Table 3.




Figure 1. PRISMA Flowchart of included and excluded studies.

Table 1. Classification according to evidence quality.

Type of evidence	Score
RCT*, meta-analysis, systematic review	4
Case-control, cohort studies, quasi-experimental studies	3
Observational studies (case reports, case series)	2
Formal/expert consensus	1

*Randomized controlled trial



Discussion

Our systematic mapping review of the evidence revealed that current GIS-based measures of ADS tend to calculate potential ADS instead of a realized one (Table 2). Also, we found no study

including both spatial and non-spatial dimensions of ADS into one framework that could produce a more realistic score than current attempts in this direction. However, in a recent study performed in Iran, Kiani *et al.* (2017) developed an integrated measure of ADS by calculating a reasonable measure of actual travel time in contrast to previous reports that mainly focus on estimated potential

Table 2. List of studies aiming at a GIS-based measure of access to dialysis service (ADS): an extraction table.

Reference, country	Setting and study participant	Method employed	Indicators discussed	Component	Comment
Roderick <i>et al.</i> , 1999 England, UK	Renal facilities Number of patients: 5,715	Multilevel modelling using Poisson regression based on patient postcodes matched to each census ward Dependent variable: the number of accepted patients (used as access indicator) Independent effects modelled: 1) age and sex; 2) census ward need factors and supply factors; and 3) district health authority level effects	Travel impedance Service availability Jarman index ^a Townsend index ^b Carstairs index ^c DoE Index ^d	Spatial accessibility Deprivation	All deprivation indices integrated as a customized index for best representing the deprivation component Emphasis on deprivation as the key indicator of both ADS ^e and NDS ^f but authors did not incorporate the components into one ADS ^e framework Emphasis on impact of diabetes, hypertension and socio-economic factors on NDS ^f but authors did not incorporate them in the model Paper implies that age and gender affect ADSe and ND ^{Sf} and matches ADSe indicators for these factors
Christie <i>et al.</i> , 2005 Wales, UK	Patient registers and main and satellite dialysis facilities Population: 2.4 million Number of patients: 1,514	Calculated a fifth of small area deprivation using the Townsend index, three bands of travel time using GIS ^g , RRT ^h prevalence, and comparing patients and population accessibility between catchments, and presenting estimates of RRT ^h prevalence in the population by a fifth of small area deprivation	Travel time, Townsend index	Spatial accessibility Deprivation	Travel time calculated as an indicator of spatial accessibility and deprivation calculated to compare RRT ^h prevalence between areas. The components were not incorporated into one ADSe framework Potential travel time calculated instead of actual travel time, and estimates of RRT ^h prevalence were lower in more deprived areas
White <i>et al.</i> , 2006 Wales, UK Wales	Main dialysis facilities Population: 2.9 million Number of patients: 2,173 (RRT ^h), 821 (HD ⁱ), 342 (PDK)	UK Renal Registry, Welsh population census data and WIMD ⁱ 2005 analyzed using GIS ^g patient location; dialysis facilities geocoded and travel time calculated. Facility capacity calculated using the number of dialysis consoles assuming two shifts of dialysis per day and each patient received thrice a week. Linear regression analysis of data based on a 7% increase in the number of dialysis patients per year	Travel time, Facility capacity, WIMD ⁱ	Spatial accessibility Deprivation	Travel time and facility capacity calculated as indicators of spatial accessibility separately, and the WIMD ⁱ calculated to compare catchments prevalence It was estimated there would be 1,400 patients on haemodialysis in Wales by 2014 Emphasis on contributing factors led to some socioeconomic factors being addressed in the WIMD ⁱ but some others, such as age, were not addressed
Yang <i>et al.</i> , 2006 USA	Dialysis facilities (44), census block data and dialysis consoles (867) Number of patients: 3,722	Two GISg-based methods, 2SFCA ^j and kernel density applied considering the number of dialysis consoles in each dialysis facility as the supply and the number of patients as the demand with a 30-minute threshold for travel time. Demand estimate based on the population in each census tract multiplied by 0.12%	Travel time, Facility capacity	Spatial accessibility	Both methods integrated two indicators to calculate spatial accessibility. The implied revealed need most likely the same as the potential need when analyzing current ADS ^e . The study implied that diabetes, hypertension, age, gender and race may influence the ND ^{Sf} but this variable was not adjusted for them

Continued on the next page.

travel time (Christie *et al.*, 2005; White *et al.*, 2006; Yang *et al.*, 2006; Judge *et al.*, 2012; Matsumoto *et al.*, 2012; Stephens *et al.*, 2013) However, the results are still far from a truly realized ADS index since the key ADS indicator of facility capacity generally remains ignored.

Matsumoto *et al.* (2012) compared two GIS-based ADS mea-

asures, the capacity-distance model and the distance model, and found the former to be more realistic than the latter. In the capacity-distance model, which addresses both travel time and facility capacity, patients were forced to travel further due to capacity limitations at closer centres. Furthermore, an American study noted that proximity to dialysis services does not directly translate into

Table 2. Continued from the previous page.

Reference, country	Setting and study participant	Method employed	Indicators discussed	Component	Comment
Matsumoto <i>et al.</i> , 2012 Japan	Dialysis facilities (98) Number of patients: 7,374	Travel time calculated by GIS ^g in two models: the distance model (travel time to nearest facility) and the capacity-distance model (both travel time and facility capacity). The latter allowed facilities to accept the nearest patient (based on travel time) until capacity is full and then remit patients to the second-nearest facility run through facilities until all patients are accepted	Travel time, Facility capacity	Spatial accessibility	The distance model addresses only travel time to calculate spatial accessibility, while the capacity-distance model uses two ADS ^e indicators jointly to calculate spatial accessibility Potential travel time calculated instead of real travel time; no model considering non-spatial factors indicating a potential gap between calculated accessibility and real access, which is more than just the equity of travel time
Judge <i>et al.</i> , 2012 UK	Renal Registry and England, UK local authority districts (354)	Multilevel Poisson regression models constructed separately for incidence and prevalence. Travel times and dialysis facilities catchment areas estimated by GIS ^g ; small area estimates of RRT ^h prevalence produced for all 354 local districts.	Travel time, IMDm	Spatial accessibility Deprivation	Components not integrated into one ADS ^e framework Deprivation adjusted for socio-demographic differences but without adjusting is directly relative to NDS ^f Potential travel time calculated instead of real travel time
Stephens <i>et al.</i> , 2013 USA	Centers for Medicare and Medicaid services (5,007) Number of patients: 332,117	Three data points estimated: (1) patients' location; (2) location of dialysis facility currently serving patient; and (3) location of replacement facility. GIS ^g -based calculation of travel impedance between them Included calculations: 1. Travel impedance from patient location to current facility; 2. Travel impedance from patient location to replacement facility; 3. Incremental travel impedance for travel to replacement facility compared with travel to current facility	Travel time	Spatial accessibility	Actual data not obtained from each patient's original dialysis facility. As it was assumed that patients utilize the closest location, potential accessibility was calculated Indication that facility capacity is a key indicator of spatial accessibility to dialysis facilities, but calculation of spatial accessibility not integrated with travel time. Non-spatial factors emphasized although not integrated with the spatial accessibility component to calculate a more realistic ADS ^e
Kiani <i>et al.</i> , 2017 Iran	Dialysis facilities (6) Number of patients: 168	Data for AAT ⁿ and other spatial or non-spatial variables gathered via a semi-structured questionnaire for patients. Univariate analysis and univariate general linear model used to identify AAT ⁿ associated factors Driving time and distance calculated using Google maps; AAT of rural and urban analyzed separately	Travel impedance	Realized spatial accessibility	Non-spatial factors (sex, income level, caregivers, transportation mode, education level, ethnicity and personal vehicle ownership) influenced the revealed access identified in rural and urban groups but spatial factors were identified only in the former AAT ⁿ calculated but some other important ADS ^e indicators such as facility capacity was ignored

^aA scoring system developed by the British general practitioner Brian Jarman (b. 1933) for the level of social deprivation; ^{ba} measure of material deprivation within a population; ^can index of deprivation used in spatial epidemiology to identify socio-economic confounding; ^dan index of urban poverty published by the UK Department for Environment, Food and Rural Affairs designed to assess relative levels of deprivation in local authorities; ^eAccess to dialysis services; ^fNeed of dialysis services; ^gGeographical information systems; ^hRenal replacement therapy; ⁱWelsh index of multiple deprivation; ^jHemodialysis; ^kPeritoneal dialysis; ^lTwo-step floating catchment area; ^mIndex of multiple deprivation; ⁿAAT actual access time.



access owing to potential racial segregation (Saunders *et al.*, 2014). However, both these studies calculated potential travel time. Meanwhile, many British studies (Roderick *et al.*, 1999; Christie *et al.*, 2005; White *et al.*, 2006; Judge *et al.*, 2012) emphasize the importance of non-spatial factors, *e.g.*, Kiani's *et al.* study (2017), considering deprivation as one of the main ADS indicators. Indeed, controversies in this field show a gap between calculated and real ADS. Although this study did not find any research integrating spatial and non-spatial factors into a GIS-embedded model for measuring ADS, various GIS-oriented studies join some spatial ADS indicators together to measure spatial accessibility (Yang *et al.*, 2006; Matsumoto *et al.*, 2012). For instance, Yang *et al.* compared two GIS-based methods, the 2SFCA approach and the kernel density method, in a case study on renal dialysis facilities in Chicago, USA. In this study, based on the main spatial ADS indicators travel time and facility capacity, the 2SFCA method produced better accessibility ratios overall (Yang *et al.*, 2006). Although, this work only integrated spatial ADS dimensions, it has provided a platform that has been successfully used in other contexts, especially in primary healthcare where an overall measure including both spatial and non-spatial factors of access is included (Wang and Luo, 2005; Bagheri *et al.*, 2008; McGrail and Humphreys, 2009; McGrail and Humphreys, 2015). For example, Bagheri *et al.* (2008) developed an integrated *access to primary healthcare* (APH) index which combined spatial accessibility, calculated by 2SFCA method, and a need index into one framework, while McGrail and Humphreys (2009, 2015) improved the 2SFCA method by introducing a concept based on three key components, *i.e.* spatial accessibility, population health needs and mobility.

The progress covered by this review, as well as the gaps revealed, raises the hypothesis that *an integrated ADS index should*

calculate access more realistically than current GIS-based measures. However, further research examining whether this hypothesis is correct or not is needed. This mapping review provides some evidence-based recommendations that may help researchers and policymakers perform a primary study assessing this hypothesis. Three components of the 2SFCA platform should be addressed in order to construct an integrated ADS index, *i.e.* spatial accessibility, mobility and NDS.

Spatial accessibility must take into account both accessibility and availability. Travel time, discussed by most current studies, should be calculated as an indicator of accessibility with a threshold of 30 minutes as haemodialysis guidelines recommended (Mactier, 2007). However, actual travel time as proposed by Kiani *et al.* (Kiani *et al.*, 2017) should be used together with facility capacity, expressed as the number of dialysis machines (supply) to the number of patients (demand) in each facility (Yang *et al.*, 2006; Matsumoto *et al.*, 2012) that seems to be the key availability indicator. It is as important as the travel time and should be incorporated into the 2SFCA framework with an appropriate threshold to construct catchment areas more realistically. Patients need dialysis thrice a week according to current haemodialysis guidelines (Mactier, 2007), which means that each machine can serve up to four patients per week (two patients on even days and two patients on odd days) leading to a supply-to-demand ratio threshold of $\frac{1}{4}$. If, regardless of this threshold, all catchments are constructed with a radius of 30 minutes travel time, some of them might include patients more than their facility capacity.

Mobility is defined as the population's ability to overcome distance (Bisht *et al.*, 2010). Taking the relative population size of those aged either <18 years or >75 years as the measure of reduced personal mobility, McGrail *et al.* (McGrail and Humphreys, 2009)

Table 3. Factors affecting access to dialysis services (ADS) elicited from literature.

Factors/Variables	Rate	Factors/Variables	Rate
Patient age	14	Population density	2
Patient gender	11	Province need	2
Patient ethnicity	13	Nephrology beds	3
Patient's education status	6	Location of the patient's nephrologist	1
Marital status	1	Patient's familiarity with facility	1
Patient's income	9	Routine procedures at the facility	1
Patient's employment status	5	Availability of transportation	3
Patient's disability	1	Car ownership	23
Patient mobility	4	Road quality linking home and facility	4
Patient's comorbidities	1	Transportation mode	3
Patient's health insurance status	7	Travel cost	17
Home owner	1	Travel speed	1
Socioeconomic status of neighbourhood	1	Travel time	3
Location of dialysis patient's residence	12	Travel distance	1
Language status	1	Time of the day for dialysis visit	1
Number of dialysis facilities	2	Degree of family support	3
Facility type (satellite or in-center)	4	Public assistance	1
Dialysis type needed*	2	Availability of human resources	1
Facility capacity	10	Deprivation	10
Number of dialysis consoles per facility	9	Climate conditions	1
Cost of care	2		

*haemodialysis (HD) or peritoneal dialysis (PD).

point out that the three indicators, *i.e.* households without a car, individuals with reduced mobility and public transport availability, measure different aspects of mobility and that correlations between them are small. Considering this nature of mobility, it seems useful to apply these three indicators when calculating mobility in the ADS context.

Revealed NDS (the actual demand) is most likely to be the same as the potential NDS when the current ADS is calculated (Yang *et al.*, 2006), but future ADS calculations will differ because the number of patients then is unclear, in particular as the prevalence of end-stage renal disease (ESRD) is increasing (Chadban *et al.*, 2003; Eggers, 2011). An easy way to estimate NDS is by multiplying the annual growth rate of demand at current demand (White *et al.*, 2006; Yang *et al.*, 2006). But an adjustment is needed if there is an increase in diabetes which is already anticipated (Roderick *et al.*, 1999; Yang *et al.*, 2006) and hypertension (Roderick *et al.*, 1999; Yang *et al.*, 2006), the two principal causes of ESRD. In addition, deprivation (Roderick *et al.*, 1999; Thomas, 2005; Judge *et al.*, 2012), age (Roderick *et al.*, 1999; Thomas, 2005), gender (Roderick *et al.*, 1999; Yang *et al.*, 2006) and ethnicity (Roderick *et al.*, 1996; Judge *et al.*, 2012) may influence the NDS differently in the future.

To our knowledge, this is the first systematic mapping review of the available literature aimed at identifying gaps in current GIS-based ADS measures and developing evidence-based recommendations. However, by limiting the search strategy by specifying it for the GIS category and developing a list of factors affecting ADS elicited, we may have lost evidence in studies exclusively focused on exploring factors affecting ADS. This would be true, even though we had a comprehensive list referencing the literature elicited from our systematic mapping review. However, this was secondary outcome measure of the study, and we suggest a systematic review with a wider scope that could list all factors affecting ADS. Although we did not systematically focus on seeking factors affecting NDS, we think that our findings related to estimating NDS are appropriate and enough. Moreover, we found that hospitalization rate (Rucker *et al.*, 2011) and mortality rate (Rucker *et al.*, 2011; Thompson *et al.*, 2012) are negatively associated with ADS, facts that can be used for validation of the integrated index of ADS, a proper validation of this index remains open for future study. Finally, even though we highlighted the absence of evidence-based recommendations incorporating indicators related to the acceptability dimension in an integrated ADS index, we could not alleviate this weakness of current GIS-based measures of ADS, which needs further research.

Conclusions

Current GIS-based measures of ADS tend to calculate potential ADS instead of a realized one and there is a need to examine whether an integrated index of ADS can calculate a realistic score. Listed factors affecting ADS are mainly non-spatial encouraging the design of an integrated ADS index produce better ADS score than those currently advocated. The mapping review strongly suggests exploring the hypothesis that a combined index of ADS including most dimensions of ADS can be developed and produce a better ADS score than current available. The 2SFCA method may be capable of providing a platform for this aim as the study recommended and researchers and policymakers are encouraged to examine and validate this hypothesis.

References

- Amy Martin KJB, Nathan Hale, Janice C. Probst, 2013. Dialysis availability in rural America, South Carolina Rural Health Research Center.
- Andersen RM, 1995. Revisiting the behavioral model and access to medical care: does it matter? *J Health Soc Behav* 36:1-10.
- Ayyalasomayajula B, Wiebe N, Hemmelgarn BR, Bello A, Manns B, Klarenbach S, Tonelli M, 2011. A novel technique to optimize facility locations of new nephrology services for remote areas. *Clin J Am Soc Nephrol* 6:2157-64.
- Azizi A, Aboutorabi R, Mazloum-Khorasani Z, Hoseini B, Tara M, 2016. Diabetic personal health record: a systematic review article. *Iran J Publ Health* 45:1388-98.
- Bagheri N, Benwell GL, PGDipSci M, 2008. Modelling Accessibility to primary health care using a spatial accessibility index and a need index. *Hawai 'I J Pub Health* 1:14.
- Bisht SS, Mishra V, Fuloria S, 2010. Measuring accessibility for inclusive development: a census based index. *Soc Indicators Res* 98:167-81.
- Chadban SJ, Briganti EM, Kerr PG, Dunstan DW, Welborn TA, Zimmet PZ, Atkins RC, 2003. Prevalence of kidney damage in Australian adults: The AusDiab kidney study. *J Am Soc Nephrol* 14:S131-8.
- Christie S, Morgan G, Heaven M, Sandifer Q, Woerden H, 2005. Analysis of renal service provision in south and mid Wales. *Publ Health* 119:738-42.
- Diamant MJ, Harwood L, Movva S, Wilson B, Stitt L, Lindsay RM, Moist LM, 2010. A comparison of quality of life and travel-related factors between in-center and satellite-based hemodialysis patients. *Clin J Am Soc Nephrol* 5:268-74.
- Eggers PW, 2011. Has the incidence of end-stage renal disease in the USA and other countries stabilized? *Curr Opin Nephrol Hypertens* 20:241-5.
- Faruque LI, Ayyalasomayajula B, Pelletier R, Klarenbach S, Hemmelgarn BR, Tonelli M, 2012. Spatial analysis to locate new clinics for diabetic kidney patients in the underserved communities in Alberta. *Nephrol Dial Transplant* 27:4102-9.
- Grant MJ, Booth A, 2009. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J* 26:91-108.
- Guagliardo MF, 2004. Spatial accessibility of primary care: concepts, methods and challenges. *Int J Health Geogr* 3:3.
- Hall YN, O'Hare AM, Young BA, Boyko EJ, Chertow GM, 2008. Neighborhood poverty and kidney transplantation among US asians and pacific islanders with end-stage renal disease. *Am J Transplant* 8:2402-9.
- Judge A, Caskey FJ, Welton NJ, Ansell D, Tomson CR, Roderick PJ, Ben-Shlomo Y, 2012. Inequalities in rates of renal replacement therapy in England: does it matter who you are or where you live? *Nephrol Dial Transplant* 27:1598-607.
- Kashima S, Matsumoto M, Ogawa T, Eboshida A, Takeuchi K, 2012. The impact of travel time on geographic distribution of dialysis patients. *PLoS One* 7:e47753.
- Kiani B, Bagheri N, Tara A, Hoseini B, Tabesh H, Tara M, 2017. Revealed access to haemodialysis facilities in northeastern Iran: Factors that matter in rural and urban areas. *Geospat Health* 12:237-44.
- Kiani B, Bagheri N, Tara A, Hoseini B, Tara M, 2017. Haemodialysis services in the northeastern region of Iran. *Geospat Health* 12:130-6.



- Levesque JF, Harris MF, Russell G, 2013. Patient-centred access to health care: conceptualising access at the interface of health systems and populations. *Int J Equity Health* 12:18.
- Levinson DR, 2011. Access to kidney dialysis services at indian health service and tribal facilities. Department of Health and Human Services USA.
- Mactier R, 2007. Clinical practice guidelines module 2: haemodialysis. UK Renal Association.
- Maheswaran R, Payne N, Meechan D, Burden RP, Fryers PR, Wight J, Hutchinson A, 2003. Socioeconomic deprivation, travel distance, and renal replacement therapy in the Trent Region, United Kingdom 2000: an ecological study. *J Epidemiol Commun Health* 57:523-4.
- Mao L, Nekorchuk D, 2013. Measuring spatial accessibility to healthcare for populations with multiple transportation modes. *Health Place* 24:115-22.
- Matsumoto M, Kashima S, Ogawa T, Takeuchi K, 2013. Do rural and remote areas really have limited accessibility to health care? Geographic analysis of dialysis patients in Hiroshima, Japan. *Rural Remote Health* 13:2507.
- Matsumoto M, Ogawa T, Kashima S, Takeuchi K, 2012. The impact of rural hospital closures on equity of commuting time for haemodialysis patients: simulation analysis using the capacity-distance model. *Int J Health Geogr* 11:28.
- McGrail MR, Humphreys JS, 2009. The index of rural access: an innovative integrated approach for measuring primary care access. *BMC Health Serv Res* 9:124.
- McGrail MR, Humphreys JS, 2015. Spatial access disparities to primary health care in rural and remote Australia. *Geospat Health* 10:358.
- McLafferty SL, 2003. GIS and health care. *Annu Rev Public Health* 24:25-42.
- Miller LM, Vercaigne LM, Moist L, Lok CE, Tangri N, Komenda P, Rigatto C, Mojica J, Sood MM, 2014. The association between geographic proximity to a dialysis facility and use of dialysis catheters. *BMC Nephrol* 15:40.
- Moist LM, Bragg-Gresham JL, Pisoni RL, Saran R, Akiba T, Jacobson SH, Fukuhara S, Mapes DL, Rayner HC, Saito A, Port FK, 2008. Travel time to dialysis as a predictor of health-related quality of life, adherence, and mortality: the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Am J Kidney Dis* 51:641-50.
- Omrani-Khoo H, Lotfi F, Safari H, Zargar Balaye Jame S, Moghri J, Shafii M, 2013. Equity in distribution of health care resources; assessment of need and access, using three practical indicators. *Iran J Public Health* 42:1299-308.
- Plantinga LC, Kim M, Goetz M, Kleinbaum DG, McClellan W, Patzer RE, 2014. Pre-end-stage renal disease care not associated with dialysis facility neighborhood poverty in the United States. *Am J Nephrol* 39, 50-8.
- Prakash S, Rodriguez RA, Austin PC, Saskin R, Fernandez A, Moist LM, O'Hare AM, 2010. Racial composition of residential areas associates with access to pre-ESRD nephrology care. *J Am Soc Nephrol* 21:1192-9.
- Richard JB, Aldigier JC, Le Mignot L, Glaudet F, Ben Said M, Landais P, 2009. Equity of accessibility to dialysis facilities. *Stud Health Technol Inform* 150:777-81.
- Roderick P, Clements S, Stone N, Martin D, Diamond I, 1999. What determines geographical variation in rates of acceptance onto renal replacement therapy in England? *J Health Serv Res Policy* 4:139-46.
- Roderick PJ, Raleigh VS, Hallam L, Mallick NP, 1996. The need and demand for renal replacement therapy in ethnic minorities in England. *J Epidemiol Commun Health* 50:334-9.
- Rodriguez RA, Hotchkiss JR, O'Hare AM, 2013. Geographic information systems and chronic kidney disease: racial disparities, rural residence and forecasting. *J Nephrol* 26:3-15.
- Rucker D, Hemmelgarn BR, Lin M, Manns BJ, Klarenbach SW, Ayyalasomayajula B, James MT, Bello A, Gordon D, Jindal KK, Tonelli M, 2011. Quality of care and mortality are worse in chronic kidney disease patients living in remote areas. *Kidney Int* 79:210-7.
- Russell DJ, Humphreys JS, Ward B, Chisholm M, Buykx P, McGrail M, Wakerman J, 2013. Helping policy-makers address rural health access problems. *Aust J Rural Health* 21:61-71.
- Salgado TM, Moles R, Benrimoj SI, Fernandez-Llimos F, 2011. Designing a model to minimize inequities in hemodialysis facilities distribution. *Geospat Health* 6:5-12.
- Saunders MR, Lee H, Maene C, Schuble T, Cagney KA, 2014. Proximity does not equal access: racial disparities in access to high quality dialysis facilities. *J Racial Ethn Health Disparities* 1:291-9.
- Saurman E, 2016. Improving access: modifying Penchansky and Thomas's theory of access. *J Health Serv Res Policy* 21:36-9.
- Smith MD, Robson AM, Woodward RS, Michelman JE, Valerius TJ, Hong BA, 1985. Geographic access to health care services: the case of maintenance hemodialysis. *Am J Kidney Dis* 5:19-26.
- Stephens JM, Brotherton S, Dunning SC, Emerson LC, Gilbertson DT, Harrison DJ, Kochevar JJ, McClellan AC, McClellan WM, Wan S, Gitlin M, 2013. Geographic disparities in patient travel for dialysis in the United States. *J Rural Health* 29:339-48.
- Thomas M, 2005. Deprivation and dialysis: pathways to kidney failure in Australian Aborigines. *Adv Chronic Kidney Dis* 12:84-7.
- Thompson S, Gill J, Wang X, Padwal R, Pelletier R, Bello A, Klarenbach S, Tonelli M, 2012. Higher mortality among remote compared to rural or urban dwelling hemodialysis patients in the United States. *Kidney Int* 82:352-9.
- Tonelli M, Hemmelgarn B, Kim AK, Bertazzon S, Klarenbach S, Manns B, Wiebe N, Culleton B, Gill JS, 2006. Association between residence location and likelihood of kidney transplantation in Aboriginal patients treated with dialysis in Canada. *Kidney Int* 70:924-30.
- Tonelli M, Manns B, Culleton B, Klarenbach S, Hemmelgarn B, Wiebe N, Gill JS, 2007. Association between proximity to the attending nephrologist and mortality among patients receiving hemodialysis. *Cmaj* 177:1039-44.
- Tshamba HM, Van Caillie D, Nawej FN, Kapend FM, Kaj FM, Yav GD, Nawej PT, 2014. Risk of death and the economic accessibility at the dialysis therapy for the renal insufficient patients in Lubumbashi city, Democratic Republic of Congo. *Pan Afr Med J* 19:61.
- Wang F, Luo W, 2005. Assessing spatial and nonspatial factors for healthcare access: towards an integrated approach to defining health professional shortage areas. *Health Place* 11:131-46.
- White P, James V, Ansell D, Lodhi V, Donovan KL, 2006. Equity of access to dialysis facilities in Wales. *Qjm* 99:445-52.
- Yang DH, Goerge R, Mullner R, 2006. Comparing GIS-based methods of measuring spatial accessibility to health services. *J Med Syst* 30:23-32.