



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XIX International Grassland Congress

---

## Test of a Conversion Equation to Increase the Accuracy of Microhistological Analysis of Herbivore Diets in the Pantanal

S. A. Santos  
*EMBRAPA, Brazil*

C. Costa  
*Universidade Estadual Paulista, Brazil*

A. Pott  
*EMBRAPA, Brazil*

J. M. Alvarez  
*EMBRAPA, Brazil*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/19/8/8>

This collection is currently under construction.

The XIX International Grassland Congress took place in São Pedro, São Paulo, Brazil from February 11 through February 21, 2001.

Proceedings published by Fundacao de Estudos Agrarios Luiz de Queiroz

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

**TEST OF A CONVERSION EQUATION TO INCREASE THE ACCURACY OF  
MICROHISTOLOGICAL ANALYSIS OF HERBIVORE DIETS IN THE  
PANTANAL**

S.A.Santos<sup>1</sup>, C. Costa<sup>2</sup>, A. Pott<sup>1</sup>, and J. M. Alvarez<sup>1</sup>

<sup>1</sup>Embrapa Pantanal, C.P. 109, Corumbá, MS, Brasil, 79320-900, asantos@cpap.embrapa.br  
<sup>2</sup> FMVZ-UNESP, Distrito Rubião Júnior, Botucatu, SP, 18618-000.

**Abstract**

Forty forage species samples were collected in September 1999, including grass, sedges, forbs, shrubs and trees. Seven hand-compounded mixtures of known botanical composition and dry weight were prepared. Five slides of each mixture were made and 20 frequency observations were recorded per slide. Two procedures were used to determine percent composition on a dry weight basis and their values were converted to correct the proportion of the unidentifiable fragments amongst species. Estimated and actual values differed significantly ( $P < 0.05$ ) in 31.7%, 34.1%, 12.2% and 12.2% of the species for frequency addition (FA), density conversion (DC), frequency addition converted (FAC) and density conversion converted (DCC), respectively. The average similarity values between estimated and observed mixture were 90.2, 86.0, 94.2 and 93.3 for FA, DC, FAC and DCC, respectively. In conclusion, food habits by microscopic analysis may be described using any of the procedures tested because they rank important forage species for herbivores.

**Keywords:** Native pastures, botanical composition.

## Introduction

The fecal microhistological technique based on the frequency of occurrence of plant fragments on a microscope field (presence or absence) has become one of the most popular methods of determining food habits of large herbivores (Mcinnis *et al.*, 1983). However, this method presents several limitations. Several studies have evaluated its accuracy (Sparks and Malechek, 1968; Holechek and Gross, 1982; Mcinnis *et al.*, 1983; Norbury, 1988). Known mixtures may be used by all technicians to evaluate their accuracy and provide factors for correction if certain species are over or under estimated (Holechek and Gross, 1982). One of the limitations refers to the fact that the relation between identifiable epidermal tissue and unidentifiable tissue is not similar for all species, overestimating or underestimating some of them. Norbury (1988) elaborated a conversion equation based on the proportion of identifiable tissues for each species. This equation may provide an accurate analysis of herbivore diets.

The objective of this study is to evaluate the use of this conversion equation for microhistological analysis of herbivore diets in the Pantanal. For this purpose two procedures are used for calculating the dry weight composition: addition frequency and frequency converted to density.

## Material and Methods

Forty species were selected for experimental use, based on preference and abundance. Cattle preferred parts were harvested and kept in a refrigerator until processing. Species used in the diets included grasses: *Axonopus purpusii*, *Panicum laxum*, *Paspalidium paludivagum*, *Leersia hexandra*, *Hymenachne amplexicaulis*, *Reimarochloa*

*brasiliensis*, *Paspalum plicatulum*, *Andropogon hypogynus*, *Mesosetum chaseae*, *Andropogon bicornis*, *Sorghastrum setosum*, *Elyonurus muticus*, *Andropogon selloanus*, *Setaria geniculata*, *Axonopus paraguayensis* and *Loudetia flammida*; sedges: *Eleocharis acutangula*, *Rhynchospora trispicata*, *Rhynchospora tenuis*, *Cyperus haspan*, *Eleocharis minima* and *Eleocharis interstincta*; forbs: *Diodia kuntzei*, *Richardia grandiflora*, *Psittacanthus calyculatus*, *Sebastiania hispida*, *Bidens gardneri*, *Thalia geniculata*, *Melochia simplex*, *Aeschnomone fluminensis*, *Hydrolea spinosa*, *Desmodium barbatum* and *Hyptis brevipes* ; shrubs: *Annona dioica*, *Smilax fluminensis*, *Arrabidaea brachypoda*, *Doliocarpus dentatus* and *Banisteriopsis pubipetala*; trees: *Curatella americana* and *Scheelea phalerata*.

Seven mixtures were hand-compounded to represent simulated diets. Five slides of each species and each mixture were prepared based on the methodology of Sparks and Malechek (1968). Only one observer analyzed both mixture and reference slides using Nikon binocular microscopes at 100 magnification. Drawings, photographs, microscope slides and keys were developed for separating these species on the basis of epidermal and cellular characteristics. Twenty frequency observations were recorded per slide to insure high repeatability among slides. Two procedures were used to calculate the percentage of dry weight. In the first procedure the number of identifiable fragments of each species in all fields was divided by the total number of identifiable fragments of all species in all fields (Holechek and Gross, 1982). Procedure two involved the conversion of frequency to density described by Fracker and Brischle (1944).

The proportion of identifiable epidermal tissue was determined for each species by examining the reference slides of chopped samples of individual species. A conversion equation was calculated as discussed by Norbury (1988) for both procedures:

$$\text{Converted point frequency} = [f_i/f(\text{ID}_i)] / \sum_{i=1}^n [f_i * f(\text{ID}_i)]$$

Where  $f_i$  = point frequency of epidermis for species  $i$  in the mixture,  $f(\text{ID}_i)$  = point frequency of identifiable epidermis for species  $i$  on the reference slide, and  $n$  = total number of species in the mixture.

Similarity indices were calculated using Kulczyński's formula (Oosting, 1956) to show the similarity between estimated and actual dry weight composition. The estimated percentage of each species in each mixture was compared to the actual percentage using a t-test for paired comparisons.

## **Results and Discussion**

Estimated and actual values differed significantly ( $P < 0.05$ ) in 31.7 %, 34.1%, 12.2% and 12.2% of the species for frequency addition, density conversion, frequency addition converted and density conversion converted, respectively (Table 1). The similarity values between estimated and observed diets for the two procedures and their converted point frequencies suggest that the test of accuracy applied to the two procedures increased the estimated value, especially for the density conversion procedure (Table 2). Without the use of converted point frequency, the frequency addition procedure provided the most accurate evaluation of dry weight composition for almost all mixtures. Similar results were found by Holeček and Gross (1982) and Alipayo *et al.* (1992). Barker (1986) suggests that, to make the use of correction factors worth, the accuracy should be at least doubled. According to Holeček and Gross (1982) the frequency addition procedure reduces overestimation of species with easily identifiable fragments while the density procedure either magnifies or reduces it depending on whether the frequency of the overestimated

species is low or high in relation to other species in the mixture. These cases were also observed in this study as in *A. purpussi* (mixture 1), in *H. amplexicaulis* (mixture 2), *P. plicatulum* (mixture 3) and *D. dentatus* (mixture 7). These species constitute major mixture components. The species estimated values were reduced by frequency addition and magnified by density procedure. However, the use of the test of conversion increased these values. The results indicated that food habits by microscopic analysis of fecal material may be described using any of the procedures tested because they rank forage species important for herbivores. Based on these data, the frequency addition procedure with converted values gives better accuracy for calculating the percentage of dry weight. However, as the use of correction factors did not greatly improve the accuracy, its application could not be justified.

### References

- Alipayo, D., Valdez R., Holechek J.L. and Cardenas M.** (1992). Evaluation of microhistological analysis for determining ruminant diet botanical composition. *J. Range. Manage.*, **45**:148-152.
- Barker, R.D.** (1986). A technique to simplify herbivore diet analysis. *Aust. Wildl. Res.*, **13**:569-573.
- Fracker, S.B. and Brischle J.A.** (1944). Measuring the local distribution of *Ribes*. *Ecology*, **25**:283-303.
- Holechek, J.L. and Gross, B.D.** (1982). Evaluation of different calculation procedures for microhistological analysis. *J. Range. Manage.*, **35**:721-723.
- Mcinnis, M.L., Vavra M. and Krueger W.C.** (1983). A comparison of four methods used to determine the diets of large herbivores. *J. Range. Manage.*, **36**:302-306.

**Norbury, G.L.** (1988). Microscopic analysis of herbivore diets – a problem and a solution. Aust. Wildl. Res., **15**:51-57.

**Oosting, H.J.** (1956). The study of plant communities. W.H.Freeman and Company, San Francisco. 440 p.

**Sparks, D.R. and Malechek J. C.** (1968). Estimating percentage dry weight in diets using a microscope technique. J. Range. Manage., **21**:264-265.

**Table 1-** The percentage mean ( $\pm$ sd) of dry weight of seven hand-compounded mixtures, using two methods (fia and fib) and converted frequency.

Mixtures	Actual % dry weight	% dry weight using frequency (fia)	% dry weight using frequency to density (fib)	F (IDi) <sup>1</sup> reference slide (%)	Converted point frequency using fia	Converted point frequency using fib
<b>Mixture 1</b>						
<i>A. purpusii</i>	41.8	39.5 $\pm$ 1.5*	50.1 $\pm$ 4.1*	86.0	36.3 $\pm$ 1.6*	46.8 $\pm$ 4.5
<i>P. laxum</i>	21.2	23.4 $\pm$ 2.8	20.7 $\pm$ 3.1	81.0	23.4 $\pm$ 3.5	20.3 $\pm$ 3.1
<i>E. acutangula</i>	20.9	18.3 $\pm$ 4.6	15.3 $\pm$ 5.7	70.0	20.5 $\pm$ 5.3	17.5 $\pm$ 6.2
<i>L. hexandra</i>	8.5	11.5 $\pm$ 2.8	8.6 $\pm$ 2.4	87.0	10.4 $\pm$ 2.5	7.8 $\pm$ 1.5
<i>P. paludivagum</i>	7.6	7.3 $\pm$ 1.1	5.1 $\pm$ 0.9*	56.0	10.0 $\pm$ 1.5*	8.2 $\pm$ 1.5
<b>Mixture 2</b>						
<i>H. amplexicaulis</i>	26.8	31.2 $\pm$ 4.1*	44.8 $\pm$ 7.7*	84.0	27.8 $\pm$ 4.1	33.4 $\pm$ 6.8
<i>D. kuntzei</i>	26.8	22.7 $\pm$ 3.5*	21.5 $\pm$ 3.4	45.0	28.2 $\pm$ 2.7	27.8 $\pm$ 3.8
<i>Reimarochloa sp</i>	20.3	20.6 $\pm$ 4.4	20.1 $\pm$ 5.4	55.0	21.6 $\pm$ 4.8	21.3 $\pm$ 5.8
<i>R. tenuis</i>	13.8	13.6 $\pm$ 3.6	12.3 $\pm$ 3.7	62.0	12.9 $\pm$ 3.4	11.7 $\pm$ 3.6
<i>R. trispicata</i>	12.3	12.5 $\pm$ 3.3	11.2 $\pm$ 3.8	57.0	12.6 $\pm$ 8.0	11.5 $\pm$ 3.6
<b>Mixture 3</b>						
<i>P. plicatum</i>	27.5	38.5 $\pm$ 4.1*	44.8 $\pm$ 7.7*	84.0	27.8 $\pm$ 4.1	33.4 $\pm$ 6.8
<i>A. hypogynus</i>	22.3	15.1 $\pm$ 3.7*	13.1 $\pm$ 3.1	44.0	20.6 $\pm$ 3.8	18.8 $\pm$ 3.9
<i>M. chaseae</i>	22.0	20.8 $\pm$ 5.2	19.6 $\pm$ 5.3	67.0	18.6 $\pm$ 3.5	18.3 $\pm$ 4.7
<i>A. bicornis</i>	12.3	8.3 $\pm$ 1.8*	7.4 $\pm$ 2.1*	37.0	13.6 $\pm$ 2.5	12.5 $\pm$ 3.4
<i>C. haspan</i>	10.7	7.6 $\pm$ 1.7*	6.1 $\pm$ 1.7	40.0	11.4 $\pm$ 2.7	9.6 $\pm$ 2.8
<i>R. grandiflora</i>	5.2	10.3 $\pm$ 2.6*	8.7 $\pm$ 2.8*	74.0	8.4 $\pm$ 2.5*	7.4 $\pm$ 2.2*
<b>Mixture 4</b>						
<i>S. setosum</i>	24.4	15.5 $\pm$ 4.7*	15.0 $\pm$ 5.8*	50.0	19.9 $\pm$ 4.8	19.3 $\pm$ 6.0
<i>E. muticus</i>	19.9	19.0 $\pm$ 3.4	20.2 $\pm$ 6.5	80.0	15.7 $\pm$ 3.5	19.1 $\pm$ 3.8
<i>A. selloanus</i>	11.7	9.4 $\pm$ 2.5	7.8 $\pm$ 2.1	42.0	14.4 $\pm$ 3.1	12.2 $\pm$ 2.7
<i>A. dioica</i>	10.6	15.5 $\pm$ 5.7	15.1 $\pm$ 6.1	90.0	11.1 $\pm$ 4.1	11.3 $\pm$ 5.3
<i>C. americana</i>	9.6	16.8 $\pm$ 1.3*	16.4 $\pm$ 1.0*	85.0	12.3 $\pm$ 1.3*	13.4 $\pm$ 1.2*
<i>S. hispida</i>	9.0	5.7 $\pm$ 1.6*	4.7 $\pm$ 1.7*	56.0	7.6 $\pm$ 2.9	5.6 $\pm$ 1.9*
<i>P. calyculatus</i>	8.8	12.9 $\pm$ 4.2	11.8 $\pm$ 4.7	82.0	10.1 $\pm$ 3.8	10.9 $\pm$ 3.7
<i>B. gardneri</i>	6.0	7.9 $\pm$ 2.9	6.5 $\pm$ 2.6	51.0	9.8 $\pm$ 3.0	8.2 $\pm$ 2.6
<b>Mixture 5</b>						
<i>T. geniculata</i>	22.8	22.5 $\pm$ 3.2	25.9 $\pm$ 5.0	75.0	18.0 $\pm$ 2.9*	21.0 $\pm$ 4.3
<i>M. simplex</i>	21.5	16.5 $\pm$ 4.4	17.1 $\pm$ 6.0	53.0	18.7 $\pm$ 4.6	19.3 $\pm$ 6.2
<i>E. minima</i>	11.1	14.0 $\pm$ 1.8*	13.8 $\pm$ 2.1*	80.0	10.5 $\pm$ 1.6	10.6 $\pm$ 1.8
<i>A. fluminensis</i>	11.0	7.0 $\pm$ 1.4*	6.1 $\pm$ 1.4*	40.0	10.5 $\pm$ 2.1	9.3 $\pm$ 2.0
<i>R. trispicata</i>	9.8	12.4 $\pm$ 1.0*	11.8 $\pm$ 1.2*	57.0	13.0 $\pm$ 0.8*	12.4 $\pm$ 4.0*
<i>E. interstincta</i>	8.5	8.9 $\pm$ 2.7	8.1 $\pm$ 2.8	40.0	13.2 $\pm$ 3.9	12.2 $\pm$ 4.0
<i>S. geniculata</i>	8.1	10.9 $\pm$ 3.5	10.3 $\pm$ 4.0	82.0	7.8 $\pm$ 2.8	7.7 $\pm$ 3.2
<i>H. spinosa</i>	7.2	7.8 $\pm$ 2.4	6.9 $\pm$ 2.3	56.0	8.1 $\pm$ 2.7	7.5 $\pm$ 2.6
<b>Mixture 6</b>						
<i>D. barbatum</i>	23.5	25.2 $\pm$ 2.7	26.0 $\pm$ 4.4	69.0	23.9 $\pm$ 2.6	25.6 $\pm$ 4.0
<i>S. phalerata</i>	22.1	28.9 $\pm$ 5.6*	32.1 $\pm$ 9.1*	95.0	20.2 $\pm$ 4.1	23.0 $\pm$ 7.1
<i>A. paraguayensis</i>	20.5	8.8 $\pm$ 1.4*	7.0 $\pm$ 1.1*	30.0	19.5 $\pm$ 2.8	16.0 $\pm$ 2.4*
<i>S. fluminensis</i>	17.7	20.1 $\pm$ 4.3	19.2 $\pm$ 5.6	71.0	18.7 $\pm$ 4.0	18.6 $\pm$ 5.4
<i>A. brachypoda</i>	16.2	16.9 $\pm$ 4.9	15.5 $\pm$ 6.1	62.0	17.7 $\pm$ 4.9	16.8 $\pm$ 6.0
<b>Mixture 7</b>						
<i>D. dentatus</i>	41.4	45.8 $\pm$ 1.5*	60.7 $\pm$ 2.5*	85.0	36.9 $\pm$ 1.8*	51.6 $\pm$ 2.7*
<i>B. pubipetala</i>	25.0	20.9 $\pm$ 3.7*	15.8 $\pm$ 4.0*	55.0	26.1 $\pm$ 4.5	20.7 $\pm$ 5.0
<i>L. flammida</i>	20.7	17.7 $\pm$ 4.0	12.8 $\pm$ 3.3*	55.0	22.1 $\pm$ 4.6	16.8 $\pm$ 4.0*
<i>H. brevipes</i>	12.9	16.0 $\pm$ 3.8	11.3 $\pm$ 3.0	75.0	14.8 $\pm$ 3.5	10.9 $\pm$ 3.2

1- f(IDi) = point frequency of identifiable epidermis for species i on the reference slide

\* Observed value differs significantly (P<0.05) from the actual value, by paired t-test



**Table 2** - The similarity index between actual and estimated mixture mean percentage in two procedures (frequency addition and frequency converted to density) and their converted point frequencies.

Mixtures	Frequency addition (fia)	Frequency converted to density (fib)	Fia converted	Fib converted
1	94.8	91.4	94.1	95.0
2	95.7	91.9	97.0	97.1
3	84.5	79.1	94.9	91.7
4	84.6	82.4	89.9	90.7
5	90.7	90.0	91.0	93.4
6	88.3	85.8	97.1	95.5
7	92.9	81.3	95.5	89.8
<b>Mean</b>	<b>90.2</b>	<b>86.0</b>	<b>94.2</b>	<b>93.3</b>