



## Changes in Pasture Growth Rate Due to Fertiliser and Grazing Management

P. Quigley  
*Agriculture Victoria, Australia*

D. Chapman  
*University of Melbourne, Australia*

J. Lamb  
*Agriculture Victoria, Australia*

G. Kearney  
*Agriculture Victoria, Australia*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

Digital part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

Commons document is available at <https://uknowledge.uky.edu/igc/19/8/11>

This collection is currently under construction.

The XIX International Grassland Congress took place in São Pedro, São Paulo, Brazil from February 11 through February 21, 2001.

Proceedings published by Fundacao de Estudos Agrarios Luiz de Queiroz

## **CHANGES IN PASTURE GROWTH RATE DUE TO FERTILISER AND GRAZING MANAGEMENT**

<sup>1</sup>P. Quigley, <sup>2</sup>D. Chapman, <sup>1</sup>J. Lamb and <sup>1</sup>G. Kearney

<sup>1</sup> Agriculture Victoria, Pastoral and Veterinary Institute, Private Bag 105, Hamilton, Victoria, 3300, Australia

<sup>2</sup> Institute of Land and Food Resources, The University of Melbourne, Victoria, 3010, Australia

### **Abstract**

It is imperative that sheep production systems in southern Australia continue to be refined so producers remain financially viable but at the same time the environment is not degraded. As part of a national thrust for development and promotion of better production systems, one research site has been established in Victoria where pastures and animal production are measured together with water and nutrient movement. Results for pasture growth rates over two years are presented here and will be used to develop best industry practice at completion of the project.

**Keywords:** Phalaris, subterranean clover, phosphorus, stocking rate, sheep

### **Introduction**

There is an urgent need for many livestock producers in southern Australia to increase the efficiency of production in order to remain financial viable. A contributing factor to this pressure has been the decline in the quality of pasture over large tracts of this region

during the last three decades (Quigley *et al.* 1992, Kemp and Dowling 1991). Causes have included reduced fertiliser inputs, reduction in resowing, over-grazing in summer and autumn, and under-grazing in spring (Mason and Andrew 1998). Perennial ryegrass (*Lolium perenne* L.) pasture and animal production can be substantially increased by optimising rates of applied phosphorus fertilisers in this climatic region (Cayley *et al.* 1998, Cayley and Kearney 2000). The additional benefits from using improved pasture species together with appropriate rates of phosphorus fertiliser have now been demonstrated for pastures based on perennial ryegrass and phalaris (*Phalaris aquatica* L.) mixtures (Saul *et al.* 1993) or for ryegrass alone (Waller *et al.* 1999). Other research has shown that phalaris-based pastures have potential for higher productivity when some form of controlled grazing is implemented (Lodge 1997). In Australia, however, farm pastures are normally continuously grazed (Cook *et al.* 1978). This paper reports preliminary data for growth rates of pastures subjected to different grazing management treatments. The project was designed to develop best industry practice for sheep production from pasture in the temperate zone of southern Australia.

### **Material and Methods**

The site for this experiment (37° 24' S, 141° 55' E) formed part of a national network of sites, each examining different tactical strategies for improving animal production, and at the same time maintaining environmental sustainability. The 20 ha site was sown to phalaris cv. Australian and subterranean clover (*Trifolium subterraneum* L.) cv. Trikkala in 1994. In July 1997, five grazing management treatments were replicated three times in plots 1.5 to 2.25 ha each (Table 1). These were stocked with Merino ewes. Those in the 4-paddock rotation were moved to fresh pasture every 2 weeks, except in spring when they were moved weekly. Ewes on the variable rotations were moved every 2 to 12 days, depending on pasture growth, except from lambing (August) to weaning (November) when they were set-stocked. Ewes

were weighed every 4 to 6 weeks, and stocking rates were adjusted several times per year to achieve similar liveweights across all treatments.

The low and high P treatments received 8 and 30 kg of phosphorus respectively, applied as superphosphate in autumn each year. The nitrogen treatment had 50 kg N/ha applied in March and 50 kg N/ha applied in late July.

Pasture growth was measured monthly during the 1998 and 1999 growing seasons using 20 or 24 enclosure cages per plot (Cayley and Bird 1996). Herbage mass was determined using a falling-plate pasture meter (manufactured by Arborline Pty Ltd, Hamilton, Victoria). The meter was calibrated using 12 herbage cuts, each time measurements were made.

A linear mixed model was fitted using ASREML (Gilmour *et al.* 1997) to test for the effects of treatment on pasture growth rate interacted with a cubic smoothing spline of time.

## **Results and Discussion**

The level of P on the set-stocked treatments (A and B) did not affect pasture growth rates. It is likely that, after 2 years from starting this experiment, there has been insufficient difference in soil available phosphorus to lead to differences in pasture growth rates. Measurement for an additional two growing seasons, with continuing differential fertiliser application, will assist in clarifying this conclusion.

There were no significant differences among any of the rotational-grazed treatments (C, D, and E), therefore only treatment E is shown in Fig. 1.

A combination of nitrogen fertiliser and rotational grazing (E) significantly ( $P<0.05$ ) increased growth rates above set-stocked treatments (A and B) (Fig 1). The set-stocked, low P treatment (A) had significantly ( $P<0.05$ ) lower growth rates than all the rotation treatments (C, D and E) which also had high P. There was a significant curvature trend between years

( $P < 0.05$ ) but no significant interaction between year and management treatment. These differences in growth have been attributed to differences in phalaris tiller size but not differences in tiller density (Cullen *et al.* 2000).

In 1998, pasture growth commenced following rainfall in April (77 mm monthly total) but conditions after this were generally dry until further heavy rain in June (93 mm monthly total), by which time mean daily temperature was  $7.6^{\circ}\text{C}$  and limiting pasture growth rates. In 1999, growth commenced in March (54 mm monthly total) with additional good rainfall in May (99 mm monthly total). Although combined winter and spring rainfall was higher in 1998 (439 mm) than 1999 (299 mm), maximum growth rates were approximately 57 kg DM/ha/day and 67 kg DM/ha/day in 1998 and 1999, respectively. Thompson *et al.* (2001) have shown that sheep production from this site was substantially improved in response to these changes in pasture production.

Considerable progress is being made in development of better management systems that increase financial viability of meat and wool production in temperate rainfall regions of Australia. As a consequence of the national program, progressive landholders are now beginning to implement these management systems, with suitable adaptations to fit their needs.

### **Acknowledgments**

Our thanks go to Meat and Livestock Australia, Land and Water Resources Research and Development Corporation and the Department of Natural Resources and Environment for funding; Robert and Marilyn Lyons for access to land and in-kind support.

### **References**

**Cayley J.W.D. and Bird P.R.** (1996). Techniques for Measuring Pastures (Pastoral and Veterinary Institute, Hamilton, Victoria).

**Cayley J.W.D., Hannah M.C., Kearney G.A. and Clark S.G.** (1998). Effects of phosphorus fertiliser and rate of stocking on the seasonal pasture production of perennial ryegrass-subterranean clover pasture. *Aust. J. Agric. Res.* **49**: 233-248.

**Cayley J.W.D. and Kearney G.A.** (2000). Profitable use of phosphorus fertiliser for temperate pastoral Australia. *Asian-Aust. J. Anim. Sci.* **13**: 191-194.

**Cook S.J., Lazenby A. and Blair G.** (1978). Pasture degeneration. II The importance of superphosphate, nitrogen and grazing management. *Aust. J. Agric. Res.* **29**: 19-29.

**Cullen B., Quigley P. and Chapman D.** (2000). Sustainable grazing systems for western Victoria. 3. How does rotational; grazing increase phalaris production? *Proc. 41st Ann. Conf. Vic. Grassland Soc.* (In press).

**Gilmour A.R., Thompson R., Cullis B.R. and Welham S.J.** (1997). ASREML. NSW Agriculture.

**Kemp D.R. and Dowling P.M.** (1991). Species distribution within improved pastures over central N.S.W. in relation to rainfall and altitude. *Aust. J. Agric. Res.* **42**: 647-659.

**Lodge G.M.** (1997). Phalaris. Pages 8-9 in R.D. FitzGerald and G.M. Lodge, eds. *Grazing management of temperate pasture: literature reviews and grazing guidelines for major species.* NSW Agric. Tech. Bull. 47.

**Mason W. and Andrew M.** (1998). Sustainable Grazing Systems (SGS) – developing a national experiment. *Proc. 9th Aust. Agron. Conf., Wagga Wagga* pp. 314-317.

**Quigley P.E., Ward G.N. and Morgan, T.** (1992). Botanical composition of pasture in south-western Victoria. *Proc. 6th Aust. Agron. Conf., Armidale* p. 533.

**Saul G.R., Jowett D., Morgan T., Noble P. and Borg D.** (1993). Improved pasture species, fertiliser and pasture use to increase the productivity of wool sheep in south-western Victoria. *Proc. XVII Int. Grassld Cong., New Zealand* pp. 1289-1292.


**Thompson A.N., Borg D, Graham J.F., Lamb J. and Saul G.R.** (2001). Rotational grazing increases wool and lamb production from phalaris-subterranean clover pastures in south eastern Australia. Proc. XIX Int. Grassld Cong., Brazil (In press).

**Waller R.A., Sale P.W.G., Saul G.R., Quigley P.E. and Kearney G.A.** (1999). Tactical versus continuous stocking for persistence of perennial ryegrass (*Lolium perenne* L.) in pastures grazed by sheep in south-western Victoria. Aust. J. Exp. Agric. **39**: 265-274.

**Table 1** - Treatments and details for the experiment.

Description	Potential stocking rate (DSE/ha) <sup>(a)</sup>	Target Olsen P (mg/kg)	Paddock area (ha)
A Set stock, low P <sup>(b)</sup>	12	4 – 6	2.25
B Set stock, high P <sup>(c)</sup>	16	12 – 14	1.5
C Simple rotation, high P	20	12 - 14	1.5
D Intensive rotation, high P	21	12 - 14	1.5
E Intensive rotation, high P + N <sup>(d)</sup>	24	12 - 14	1.5

<sup>(a)</sup> dry sheep equivalent based on a 45 kg wether    <sup>(b)</sup> Approximately 8 kg P/ha/year;    <sup>(c)</sup> Approximately 30 kg P/ha/year;    <sup>(d)</sup> 100 kg N/ha/year



**Figure 1** - The effects of three management treatments on pasture growth rates in 1998 and 1999.