



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXIV International Grassland Congress /
XI International Rangeland Congress

Title Evaluation of Yields and Nutritive Composition of Dual Purpose Sweet Potato Vine Cultivars for Forage Use

R. S. Kenana

Kenya Agricultural Research and Livestock Organization, Kenya

A. I. Hoka

Kenya Agricultural Research and Livestock Organization, Kenya

N. Ondabu

Kenya Agricultural Research and Livestock Organization, Kenya

F. Ondabu

Kenya Agricultural Research and Livestock Organization, Kenya

J. Mercy

Kenya Agricultural Research and Livestock Organization, Kenya

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/24/2/27>

This collection is currently under construction.

The XXIV International Grassland Congress / XI International Rangeland Congress (Sustainable Use of Grassland and Rangeland Resources for Improved Livelihoods) takes place virtually from October 25 through October 29, 2021.

Proceedings edited by the National Organizing Committee of 2021 IGC/IRC Congress

Published by the Kenya Agricultural and Livestock Research Organization

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Title Evaluation of Yields and Nutritive Composition of Dual Purpose Sweet Potato Vine Cultivars for Forage Use

Kenana R.S., Hoka A.I, Ondabu N., Ondabu F. and Mercy J.
KALRO-National Beef Research Institute-Lanet
P.O. Box 3840, Nakuru, Kenya.

Key words: *Cultivar; digestible energy; Feed; fodder; Protein.*

Abstract

The sweet potato vine (SPV) is a widely grown but underutilized feed resource for livestock. The vines have desirable characteristics suitable for fodder production due to high contents of protein and digestible energy. The objective of this study was to evaluate the yield and nutritive composition of five dual purpose cultivars based on vine production for forage use. The experimental design was arranged in randomized complete block design where six cultivars of sweet potato were established in plots of 2.5m x 2.5m in three replicates. The cultivars used were; Kenspot 1, 2, 3, 4, and 5. An improved forage type cultivar (Wagbolige) served as a control. Data was subjected to analysis of variance (ANOVA) using a general linear model (SAS, 2002). Results showed that there was no significant difference ($p < 0.05$) in dry matter (DM) among the cultivars. Wagbolige (control) was much superior in yield ($p < 0.05$) compared to the others cultivars. There was no significant difference ($p < 0.05$) in yield between Kenspot 1, 3, 4 and 5. Kenspot 2 had the lowest yields. Kenspot 1 had the highest CP ($p < 0.05$) compared with the other 5 cultivars. However, there was no significant difference ($p < 0.05$) in CP between Kenspot 5, Kenspot 2 and Wagbolige. Kenspot 3 had the lowest CP content among the cultivars. Energy was highest in Kenspot 1 and lowest in Kenspot 2. NDF and ADF were highest in Kenspot 2 and lowest in Kenspot 4. There was no significant difference ($p < 0.05$) in Ca, P, K and Mg. This study concluded that the five dual purpose cultivars demonstrated excellent potential in terms of quality for forage use. Biomass production however, was the main limitation.

Introduction

The increasing population and diminishing land sizes, has reduced the land available for forage production, as the little land available is used for growing food crops (FAO, 2006). This has necessitated the need to identify multipurpose crops which can be used both as human food and livestock feeds. In recent years, sweet potato (*Ipomoea batatas*) has elicited a lot of interest from various research organizations and governments due to its adaptability to semi-arid areas and the possibility of being used both as human food and livestock feed. However, there is limited information on its contribution to livestock production in sub-Saharan Africa and as a result its potential as livestock feed has not been fully exploited (Peter, 2008).

Studies conducted at the International Potato Centre (CIP) in 2008 revealed that farmers preferred the dual purpose varieties because they have enough tubers for human consumption as well as fodder for livestock. The dual purpose cultivars are particularly preferred because harvesting can be done throughout the growing season. In Kenya about 59.2 thousand ha of land is grown with sweet potato annually with a production of 12.8t/ha (FAO, 2012). However, the amount of vines produced alongside has not been documented. Sweet potato vines (SPVs) can play a significant role as a partial replacement for other forages and pastures in the nutrition of dairy cows, goats and pigs in East Africa (Peters, 2008). A study conducted in Uganda to determine the potential of sweet potato vine-based diets as partial milk substitute (PMS) for dairy calves reduced the amount of milk consumed per calf by 120 litres over the 70-day period (Taabu *et al.*, 2016).

In recent years several sweet potato varieties were released for various agro-ecological zones at Kenya Agricultural and Livestock Research Organization (KALRO), Foods crop Research Institute in Njoro. These included cultivars; Kenspot 1, Kenspot 2, Kenspot 3, Kenspot 4, Kenspot 5 among others. However, studies on biomass and nutritional value for livestock are missing and this study is a step towards filling this knowledge gap

Methods and Study Site

This study was carried out at KALRO-Lanet situated in Nakuru County. The center lies between longitude 36° 09' E and latitude 00° 18' S at an altitude of 1920 m above sea level. The center occupies 1418 hectares of land within two Agro ecological zones (AEZs), where 20% of the land lies within AEZ III and 80% in AEZ IV (Jaetzol *et al.*, 2006). The region has a bimodal rainfall season with the (long rains) starting from March to June and the (short rains) falling from November to December. The annual rainfall ranges between 600-1000 mm and temperatures between 10°C and 30°C (Jaetzol *et al.*, 2006). Soil pH ranges between 5.5- 6.5. Soils

are deep sandy loam with good water holding classified as humic nitosols under Food and Agricultural Organization of the United Nations (FAO) classification.

Cultivation of Forage Sweet Potato

The land was prepared into a fine seed bed. The land was divided into 18 plots of 2.5m x2.5m with a 1-meter-wide path way between the plots. A randomized complete block design (RCBD) was used to layout the experiment. Six cultivars of SPV namely Kenspot 1, Kenspot 2, Kenspot 3, Kenspot 4, Kenspot 5 and Wagabolige were planted randomly in three replicates. Wagabolige which is an improved forage type sweet potato cultivar (Ondabu et al., 2007) was chosen as the control cultivar. The cultivars were planted on their respective plots using 50 cm long cuttings, in holes dug on the flat ground in rows 60 cm apart and the plants were spaced 30 cm apart within rows. A base fertilizer dressing with Di-ammonium phosphate was done at planting and top dressing at 45 days after planting. The fertilizer was applied according to Wielemaker and Boxem (1982) during planting at the rate of 54 and 20 kg ha⁻¹ of nitrogen (N) and phosphorus (P), respectively; and after establishment (45 days after planting), the blocks were cut back and top dressed with 52 kg ha⁻¹N. Potassium was not applied as soils in Lanet area were known to contain adequate potash to meet the requirements for normal growth of forage sweet potato cultivars (Wielemaker and Boxem, 1982). These blocks were kept clean by regular hand weeding and the subsequent re-growth was harvested at 120 days

Results

Yield and Nutrient Composition

The yield and composition of the cultivars are shown in Table 1. Wagabolige, the control cultivar, apparently produced the highest biomass quantity ($p<0.05$) followed by Kenspot 3, Kenspot 4, Kenspot 1, Kenspot 5 and Kenspot 2 respectively. Kenspot 4 had the highest dry matter (DM) though not significant ($p<0.05$) from the other cultivars, while Kenspot 3 had the lowest. There was no major significant difference in energy among the cultivar though Kenspot 1 had the highest while Kenspot 2 had the lowest. Crude fibre (NDF and ADF) was highest ($p<0.05$) in Kenspot 2 and lowest in Kenspot 1. Crude protein (CP) was highest ($p<0.05$) in Kenspot 1 and lowest in Kenspot 3. There was no significant difference ($p<0.05$) in Ca, P, K and Mg among the cultivars.

Table 1 Yield and chemical composition of six cultivars of sweet potato vines

Cultivar	Yield t/ha	DM%	Crude Protein (%)	Energy (Kcal)	NDF %	ADF %	Ca %	P %	K %	Mg %
Kenspot 1	29.4 ^{cb}	13.02	23.70 ^a	2084.05	40.15 ^c	21.50	1.29	0.36	4.60	0.39
Kenspot 2	16.4 ^d	12.93	19.23 ^c	1757.09	46.11 ^a	28.78 ^a	1.26	0.34	4.79	0.39
Kenspot 3	31.7 ^b	12.92	16.84 ^d	1793.62	42.28 ^b	24.96 ^{cb}	1.23	0.31	4.73	0.39
Kenspot 4	29.2 ^{cb}	13.73	21.82 ^b	1911.84	42.85 ^b	23.29 ^d	1.19	0.37	5.33	0.31
Kenspot 5	19.5 ^{cd}	13.50	19.78 ^c	1785.06	43.02 ^b	25.46 ^b	1.38	0.38	5.61	0.36
Wagabolige	64.5 ^a	13.12	19.60 ^c	1846.22	42.94 ^b	24.40 ^c	1.29	0.33	4.95	0.36
Average	32.1	13.2	20.2	1863.1	42.9	24.7	1.27	0.35	4.97	0.37
CV%	13.3	3.6	2.5	0.1	0.68	1.1	11.1	13.7	9.9	12.5

a, b, c, d: means with same superscripts in the column are not significantly different ($P>0.05$). DM: dry matter, CP: crude protein, NDF: neutral detergent fibre, ADF: acid detergent fibre, Ca: calcium, P: phosphorous, K: Potassium and Mg: Magnesium

Discussion

Yield and Nutrient Composition

The forage type sweet potatoes (SPV) are generally high in biomass production. This presented itself very well in this study. Cultivar (Wagabolige) an improved forage type (Ondabu *et al.*, 2007) produced the highest yields (8.5ton/ha). This is more than double the highest among the dual purpose cultivars (4.1ton/ha). The six cultivars had a CP range of 16% to 23%. This agrees with the range given by Dung (2001). All the cultivars were generally of high nutritional quality. They had much higher CP more than the 80 g CP/kg DM, below which forages are considered low in quality (Irungu *et al.*, 2016; Ondiek *et al.*, 2000; Leng, 1990) and hence may not limit microbial activity in the rumen (Van Soest, 1994). The NDF was also below the 600 g/kg DM usually considered as the threshold for ruminants (Irungu *et al.*, 2016; Meissner *et al.*, 1991). The low NDF was consistent with the general observation of lower NDF in non-grass forages (Minson, 1990). The six cultivars had adequate fibre, measured as NDF and defined as total cell wall content which is essential for rumination, saliva flow, rumen buffering and health of the rumen wall (Fox *et al.*, 1992). Furthermore, the high energy will enable the animals to obtain adequate ME required for incorporating nitrogen into microbial protein (Preston and Leng, 1987; Muia, 2000; Irungu *et al.* 2007). The chemical composition of the six forage sweet potato cultivars was in agreement with available literature on forage sweet potato (Snijders *et al.*, 1992;

Kiragu and Tamminga, 1997; Larbi *et al.*, 2007). The DM was in agreement with values reported by Chhay *et al.* (2007), Olorunnisomo (2007) and Kebede *et al.* (2008). The CP values were also within those reported by Giang *et al.* (2004), Lam and Ledin (2004) and Kiragu *et al.* (2007). The fibre content (NDF and ADF) were all in the range reported by Olorunnisomo, (2007) and Kebede *et al.* (2008). However, their values were lower than those reported in the current study and those by Snijders *et al.* (1992), Kariuki *et al.* (1998) and Kiragu *et al.* (2007). These higher values may be due to harvesting age, variety used and a site effect (Karachi, 1982a, b; Irungu *et al.*, 2000). The low NDF in forage sweet potato cultivars was consistent with the general observation of lower NDF in non-grass fodders (Minson, 1990). Nonetheless, all forage sweet potato cultivars in the current study had higher NDF than 150 g/kg DM recommended by Strasia and Gill (1990) as being suitable for growing ruminants. Furthermore, these cultivars contained lower than 600 g NDF/kg DM, beyond which a feed is classified as poor quality (Meissner *et al.*, 1991). Acid detergent fiber is associated with the digestibility of a feed, thus a feed with high ADF is less digestible than that of low ADF and a less digestible feed reduces the intake by animals.

Conclusions

All cultivars were similar in terms of DM, energy and minerals (Ca, P, K and Mg). These cultivars however, differed in yields, CP, NDF, and ADF. The six cultivars had superior feed qualities for livestock as they contained high CP and less fibre compared to common Kenyan forages. They were also high in energy, which classified them as high quality forages. However, the dry matter was low and might not be sufficient to meet the dry matter requirement of the animals if fed solely. They are, therefore, recommended to livestock farmers as high quality forage supplements. Additional trials are needed to determine their performance on livestock.

Acknowledgements

I wish to acknowledge my employer Kenya Agricultural and Livestock Research Organization (KALRO) and the Institute Director, KALRO-BRI-Lanet for logistical support

References

- [AOAC. 1995. Official Method of Analysis. 15th Edn. Association of Official Analytical Chemists, Washington, DC., USA.
- AOAC (Association of Official Analytical Chemists) (1998). Official Methods of Analysis. 16th edition, 4th revision. AOAC, Gaithersburg, MD.
- Chhay, T., Borin, K., Phiny, C. (2007). A note on the effect of fresh mulberry leaves, fresh sweet potato vines or mixture of both foliages on intake, digestibility and N retention of growing pigs given a basal diet of broken rice. *Livestock Research for Rural Development*. Volume 19, article 136
- Dung, N. N. X. (2001). Evaluation of green plants and by-products from the Mekong Delta with emphasis on fibre utilization by pigs (Doctoral thesis). Swedish University of Agricultural Sciences: Uppsala, Sweden.
- Food and Agriculture Organization (FAO). 2012. Food and Agriculture Organization Statistical database for agriculture. Crops and products domain. <http://faostat.fao.org/> (Accessed 19, August 2013).
- Giang, H.H., Ly, L.V. Ogle, B. (2004). Evaluation of ensiling methods to preserve Sweet potato roots and vines as pig feed. *Livestock Research for Rural Development*. Vol. 1 article 45.
- Irungu, J.W., Kirubi, D., Nangayo, F., Ndolo, P.J., Munga, T., Wafula, J.S., Macharia, C. and Maina, D. (2000). *Agronomic and virological comparison of sweet potato varieties in different agro-ecological zones in Kenya*. Proceedings of the 7th KARI Biennial Scientific Conference, 13–17 November 2000, Nairobi, Kenya, pp. 413–421.
- Jaetzold, R., Schmidt, H., Hornetz, B. and Shisanya, C. (2006). *The Farm Management Handbook of Kenya*. Central Kenya. Volume II, 2nd edition. Ministry of Agriculture, Kenya.
- Kaitho, R.J. (1997). *Nutritive value of browse as protein supplements to poor quality roughages*. PhD thesis. Wageningen University, Netherlands.
- Karachi, M.K. (1982a). The performance of sweet potato in Western Kenya: I. Effect of nitrogen and phosphorus combination on yield. *E. Afr. Agric. For. J.*47, 55–59.
- Karachi, M.K. (1982b). The performance of sweet potato in Western Kenya: II. Yield of 31 cultivars. *E. Afr. Agric. For. J.*47, 60–67.
- Karachi, M.K. and Dzwela, D.H. (1990). The potential of sweet potato as dual purpose crop in semi-arid Crop and livestock system in Kenya. *PANESA/ARNAB, ILCA*, Adis Ababa. pp. 518–532.
- Kariuki, J.N. (1998). *The potential of improving napier grass under smallholder dairy farmer's conditions in Kenya*. PhD thesis. Wageningen University, Netherlands.
- Kariuki, J.N., Gitau, G.K., Gachui, C.K., Tamminga, S., Van Bruchem, J., Muia, J.M.K. and Irungu, K.R.G. (1998). Effect of feeding Napier grass, lucerne and sweet potato vines as sole diets to dairy heifers on nutrient intake, weight gain and rumen degradation. *Liv. Prod.Sci.*55, 13–20.
- Kebede, T., Lemma, T., Tadesse, E. and Guru, M. (2008). Effect of level of substitution of sweet potato vines for concentrate on body weight gain and carcass characteristic of browsing Arsi Bale goats. *JCAB* 2, 36–42.
- Kiragu, J.W. and Tamminga, S. (1997). *Utilization of sweet potato vines as a roughage source in the diet of early weaned Friesian calves*. Proceedings of the 5th KARI Scientific Conference, 14–16 October, 1996, Nairobi, Kenya. pp. 562–579.

- Kiragu, J.W., Mitaru, B.N. and Irungu, K.R.G. (2007). Supplementing Napier grass with different proportions of sweet potato vines on growth and rumen development. *Paper presented at Joint Tanzania Society of Animal Production and Veterinary Association of Tanzania*, 29 November–1 December 2007, Arusha, Tanzania.
- Lam, V. and Ledin, I. (2004). Effect of feeding different proportions of sweet potato vines (*Ipomoea batatas*) and *Sesbania gradiflorain* in the diet on feed intake and growth of goats. *Livestock Research for Rural Development*. Vol. 16, article 77
- Larbi, A., Etela, I., Nwokocho, H.N., Oji, U.I., Anyanwu, N.J., Gbaraneh, L.D., Anioke, S.C., Balogun, R.O. and Muhammad, I.R. (2007). Fodder and tuber yields and fodder quality of sweet potato cultivars at different maturity stages in West African humid forest and Savanna zones. *Anim. Feed Sci. and Techn.* 135, 126–138.
- Leng, R.A. (1990). Factors affecting the utilization of poor quality forages by ruminants particularly under tropical conditions. *Nutr. Res. Reviews*. 3, 277–303.
- Meissner, H.H., Koster, H.H., Nieuwuodt, S.H. and Coertze, R.J. (1991). Effect of energy supplementation on intake and digestion of early and mid-season rye grass and panicum/smuts finger hay. *S. Afr. J. Anim. Sci.* 21, 33–42.
- Minson, D.J. (1990). Forage in Ruminant Nutrition. *Academic Press*, London.
- Muia, J.M.K. (2000). Use of Napier grass to improve smallholder milk production in Kenya. PhD thesis, Wageningen University, Netherlands.
- Olorunnisomo, O.A. (2007). Yield and quality of sweet potato forage pruned at different intervals for West African dwarf sheep. *Livestock Research for Rural Development*. Volume 19, article 36.
- Ondabu, N., Irungu, K.R.G., Tarus, S. Ontiti, S. (2007). *Sweet Potato Vine selection programme at Lanet*. Proceedings of the 9th Triennial Symposium of the International Society for Tropical Root Crops Africa Branch, Mombasa, Kenya, 1–5 November 2004, pp. 490–494.
- Peters, D. (2008). Assessment of the potential of sweet potato as livestock feed in East Africa: Rwanda, Uganda, and Kenya. Report presented to the International Potato Center (CIP).
- Snijders, P.J.M., Muia, J. and Kariuki, J.N. (1992). Yield and quality of sweet potato vines harvested at different stages, Research Report, Naivasha, Kenya.
- Strasia, C.A., Gill, D.R. (1990). Formulating feedlot diets. Great Plains Beef Cattle Hand book. Animal Science Research Report. Agricultural Experimental Station, Oklahoma State University. Factsheet.
- Taabu H L, Ndyomugenyeni E K, Mutetikka D, Ebong C 2016: Effect of feeding sweet potato vine-based diets as partial milk substitutes for dairy calves in Uganda. *Livestock Research for Rural Development*. Volume 28, Article 18.
- Van Soest, P.J. (1994). Nutritional Ecology of the Ruminant, 2nd edn. Cornell University Press, Ithaca, NY.
- Van Soest, P.J., Robertson, J.B., Lewis, B.A. (1991). Methods for dietary fibre, neutral detergent fibre, and non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.*, 74, 3583–3597.
- Van Soest, P J and Robertson J B 1985 Analysis of forages and fibrous foods. AS 613 Manual. Department of Animal Science, Cornell University, Ithaca, pp 105-106.
- Wielemaker, W.G. and Boxem, H.W. (1982). Soils of the Kisii Area. Ministry of Agriculture. Kenya Soil Survey, Nairobi, Kenya.