



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XIX International Grassland Congress

---

## Urine Calcium But Not Plasma Calcium or Urine Hydroxyproline Is Increased by a Systemic Acidosis in the Dairy Cow

J. R. Roche

*Dairying Research Corporation, New Zealand*

Dawn E. Dalley

*Agriculture Victoria Ellinbank, Australia*

F. O'Mara

*University College Dublin, Ireland*

E. S. Kolver

*Dairying Research Corporation, New Zealand*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/19/19/7>

This collection is currently under construction.

The XIX International Grassland Congress took place in São Pedro, São Paulo, Brazil from February 11 through February 21, 2001.

Proceedings published by Fundacao de Estudos Agrarios Luiz de Queiroz

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

**URINE CALCIUM BUT NOT PLASMA CALCIUM OR URINE HYDROXYPROLINE  
IS INCREASED BY A SYSTEMIC ACIDOSIS IN THE DAIRY COW**

J.R. Roche<sup>1</sup>, D.E. Dalley<sup>2</sup>, F. O'Mara<sup>3</sup> and E.S. Kolver<sup>1</sup>

<sup>1</sup>Dairying Research Corporation, Private Bag 3123, Hamilton, New Zealand.

<sup>2</sup>Agriculture Victoria Ellinbank, RMB 2460, Ellinbank, Victoria 3821, Australia.

<sup>3</sup>Department of Animal Science and Production, University College Dublin, Ireland.

**Abstract**

Eight non-lactating, pregnant Holstein-Friesian cows were allocated to two treatments and individually offered diets differing in dietary cation-anion difference. Decreasing the dietary cation-anion difference reduced the urine pH within hours of anionic salt supplementation. Plasma calcium concentration was unaffected by dietary cation-anion difference but urine calcium concentration was significantly increased within 10 days of including anionic salts in the diet. Faecal calcium concentration was significantly reduced, indicating increased calcium absorption. Dietary calcium concentration or dietary cation-anion difference did not significantly affect urinary hydroxyproline.

**Keywords:** Dairy cows, non-lactating, pasture, dietary cation-anion difference.

**Introduction**

Some researchers have measured an increase in plasma and urine calcium concentration at calving when a reduced dietary cation-anion difference (DCAD) was fed (Block, 1984). In contrast Roche (1999) found no increase in plasma calcium during a decrease in DCAD, although pre-calving urinary calcium concentration increased

exponentially ( $r^2 = 0.95$ ). Schonewille *et al.* (1994) found that increased absorption of calcium only accounted for 60% of the increased excretion of calcium. The source of the additional calcium in urine remains unclear.

Some researchers (Block, 1984; Goff *et al.*, 1991) have measured an increase in urinary hydroxyproline as a result of feeding a low DCAD and have concluded that the reduction in blood pH increased bone resorption. However, Schonewille *et al.* (1994) and Roche (1999) found no increase in urinary hydroxyproline when blood pH was reduced and Van Mosel *et al.* (1994) claimed that the extra calcium excreted was due to decreased bone accretion.

The objective of the work reported here was to test the effects of a reduction in systemic pH on the excretion of calcium and hydroxyproline by the cow.

### **Materials and Methods**

Eight non-lactating, pregnant, multiparous, rumen-fistulated Holstein-Friesian cows were allocated, in a randomized block design, to two treatments on the basis of age ( $5.75 \pm 0.44$  years) and live weight ( $584 \pm 52$  kg). Treatments differed in DCAD; the high treatment (High) had a DCAD of  $+24 \text{ mEq } 100\text{g}^{-1}$  and the low treatment (Low) had a DCAD of  $-20 \text{ mEq } 100\text{g}^{-1}$ , achieved by the addition of magnesium chloride ( $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ ) and ammonium chloride ( $\text{NH}_4\text{Cl}$ ). DCAD was calculated using the equation of Tucker *et al.* (1992).

#### *Management and feeds*

Animal measurements taken for four days (days -3 to 0) prior to the beginning of the experiment were used as a covariate. Cows in both treatments were fed the same base feed of 7kg DM of hay and 3 kg DM of barley. Cows on Low were supplemented with  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$  and  $\text{NH}_4\text{Cl}$ , through the rumen fistula, at 09.00 and 15.00 h. Following the adaptation period

(Days 1-14) cows were individually fed indoors for seven days (days 15-21) during which individual DM intake and total urine and faecal measurements were made. Barley (1.5 kg cow<sup>-1</sup>) was offered for 20 minutes at 09.00 and 15.00 h after which the cows had access to hay until 21.30 h. On day 16 the cows were fitted with faecal and urine separation equipment as described by Grainger (1982). The calcium balance period extended from day 17 to 21.

### *Measurements*

A midstream urine sample was collected from each cow following manual stimulation on day -3 to 16. Urine pH was determined and two sub-samples were frozen awaiting calcium and creatinine analysis.

Samples of all feeds were analyzed for macro minerals using x-ray spectroscopy (Norrish and Hutton, 1977) and DCAD calculated.

During the Ca balance period total faecal and urine outputs were taken for faecal DM and calcium concentration, and urinary pH and calcium, hydroxyproline and creatinine concentration. Blood was collected from each cow daily, centrifuged for 10 minutes at 1120 g and the plasma removed.

Faecal, urine and plasma calcium concentrations were determined on a Perkin Elmer 372 atomic absorption spectrophotometer at 422.7 nm against a series of calcium standards. Urine hydroxyproline was determined by a method developed by Parekh and Jung (1970) using a micro plate reader (Biorad 550, USA). Urine creatinine was determined by a method modified from Bartels *et al.* (1972), using an autoanalyser (Boehringer Mannheim Hitachi 911).

### *Statistical analysis*

All data were analyzed by analysis of variance for a randomized block design using the statistical procedures of Genstat V (1997), with cows as the experimental unit.

## Results and Discussion

The lower DCAD caused a significant reduction ( $P < 0.001$ ) in the pH of excreted urine within a couple of hours of anionic salt supplementation (Figure 1). Urine pH on the Low treatment continued to decline for 2 days, reached a plateau and on day 6 increased from 5.98 to 7.83. On day 9, urine pH began to decline again to a trough of approximately 5.5 at day 19. Figure 1 also shows a concomitant rise in the corrected urinary calcium concentration (CUCa) as urine pH declined after day 9, which is consistent with the results of Vagg and Payne (1970) and Roche (1999).

There was no significant difference between treatments in calcium intake or plasma calcium concentrations (Table 1). The reduced DCAD caused a decrease in urine pH, an increase in urinary calcium excretion ( $P < 0.001$ ) and a reduction in faecal calcium excretion ( $P < 0.05$ ) suggesting calcium absorption was increased.

Hydroxyproline excretion in urine, an indicator of bone resorption (Robins, 1994), was not affected by treatment (Table 1). These results may indicate that when DCAD was reduced bone resorption was not responsible for the additional calcium excretion unaccounted for by increased absorption. Alternatively, they question the effectiveness of urinary hydroxyproline as a marker, specific to bone resorption, when used in the periparturient period. As hydroxyproline indicates any catabolism of collagen including the release of collagen by the uterus its use as a specific marker for collagen released during bone resorption is probably not accurate. The direct measure of bone vicissitude by Van Mosel *et al.* (1994) who reported a decrease in bone accretion during systemic acidosis but found no effect on bone resorption supports this theory.

A DCAD of  $-20 \text{ mEq } 100\text{g}^{-1}$  caused a significant depression in urine pH and an increase in urine calcium: creatinine ratio. Plasma calcium and urine hydroxyproline were unaffected by DCAD. Urine hydroxyproline is probably not a reliable indicator of bone resorption in the periparturient period.

### References

- Bartels, H., Boemer, M., and Heirli, C.** (1972) Serum creatinine determination without deproteinization. *Clinica Chimica. Acta.* **37**:193-197.
- Block, E.** (1984) Manipulating dietary anions and cations for prepartum dairy cows to reduce incidence of milk fever. *Journal of Dairy Science.* **67**:2939-2948.
- Genstat V Committee** (1997) Genstat 5, Release 4.1. Reference Manual. Oxford University Press, Oxford, United Kingdom.
- Goff, J.P., Horst, R.L., Mueller, F.J., Miller, J.K., Kiess, G.A. and Dowlen, H.H.** (1991) Addition of chloride to a prepartal diet high in cations increases 1,25-dihydroxyvitamin D response to hypocalcaemia preventing milk fever. *Journal of Dairy Science.* **74**:3863-3871.
- Grainger, C.** (1982) Some effects of genotype on the conversion of pasture to milk by Friesian cows. Ph.D. Dissertation. Massey University, New Zealand.
- Norrish, K. and Hutton, J.T.** (1977) Plant analyses by x-ray spectrometry. 1. Low atomic number elements, sodium to calcium. *X-Ray Spectrometry.* **6**:6-11.
- Parekh, A.C., and Jung, D. H.** (1970) An improved method for determination of total hydroxyproline in urine. *Biochemical Medicine.* **4**:446-456.
- Robins, S.P.** (1994) Biochemical markers for assessing skeletal growth. *European Journal of Clinical Nutrition.* **48**:199-209.

**Roche, J.R.** (1999) Dietary Cation-Anion Difference for Pasture-fed cows. Ph.D. Dissertation. University College Dublin, Ireland.

**Schonewille, J.T., Van't Klooster, A.T., Dirkswager, A. and Bayen, A.** (1994) Stimulatory effect of an anion (chloride)-rich ration on apparent calcium absorption in dairy cows. *Livestock Production Science*. **40**:233-240.

**Tucker, W.B., Hogue, J.F., Waterman, D.F., Swenson, T.S., Xin, Z., Hemken, R.W., Jackson, J.A., Adams, G.D. and Spicer, L.J.** (1992) Sulphur should be included when calculating the dietary cation-anion balance of diets for lactating dairy cows. *Animal Science Research Report*, Oklahoma Research Station. pp.141-150.

**Vagg, M.J. and Payne, J.M.** (1970) The effect of ammonium chloride induced acidosis on calcium metabolism in ruminants. *British Veterinary Journal*. **126**:530-537.


**Van Mosel M., Wouterse H.S. and Van't Klooster A.T.** (1994) Effects of reducing dietary  $([Na + K]-[Cl+S])$  on bone in dairy cows at parturition. *Research in Veterinary Science*, **56**, 270-276.

**Table 1** - Effect of High or Low DCAD on the net calcium balance of the non-lactating dairy cow.

	High	Low	SEM
Calcium Intake (g day <sup>-1</sup> )	27.6	28.4	1.66
Urine Calcium (g day <sup>-1</sup> )	0.8 <sup>a</sup>	4.7 <sup>b</sup>	0.59
Faecal Calcium (g day <sup>-1</sup> )	46.0 <sup>b</sup>	32.8 <sup>a</sup>	3.7
Plasma Calcium (mg l <sup>-1</sup> )	100.2	97.0	1.56
Urinary Hydroxyproline (mg day <sup>-1</sup> )	0.0087	0.0077	0.00081

<sup>a, b</sup> Means in a row with different superscripts differ significantly





**Figure 1** - Effect of dietary cation-anion difference on the urine pH (◆ and ■) and urine Ca: creatinine ratio (◇ and □) of High and Low DCAD, respectively