

# National Cancer Institute Biospecimen Evidence-Based Practices: Harmonizing Procedures for Nucleic Acid Extraction from Formalin-Fixed, Paraffin-Embedded Tissue

Sarah R. Greytak,<sup>1</sup> Kelly B. Engel,<sup>2</sup> Erik Zmuda,<sup>3</sup> Esmeralda Casas-Silva,<sup>4</sup> Ping Guan,<sup>4</sup> Katherine A. Hoadley,<sup>5</sup> Andrew J. Mungall,<sup>6</sup> David A. Wheeler,<sup>7</sup> Harsha V. Doddapaneni,<sup>7</sup> and Helen M. Moore<sup>4</sup>

## Introduction

VARIABLE AND SUBOPTIMAL biospecimen handling practices have been identified as impediments to biomarker discovery,<sup>1,2</sup> including predictive biomarkers for oncology<sup>3</sup> indicating a clear and present need for evidence-based, standardized practices. The United States and international efforts have been launched to better understand and mitigate variability during the preanalytical phase and decrease associated effects by promoting harmonization of procedures both within and across institutions. The Biorepositories and Biospecimen Research Branch (BBRB) of the United States National Cancer Institute (NCI) publishes best practice documents aimed at improving the quality of data generated from human biospecimens (<https://biospecimens.cancer.gov/bestpractices/overview.asp>) and sponsors research initiatives (<https://biospecimens.cancer.gov/programs/default.asp>) and the Biospecimen Research Database (BRD; <http://biospecimens.cancer.gov/brd>) to better understand thresholds and effects of individual preanalytical factors in biospecimen handling. The BRD allows users to query both a curated literature repository and standard operating procedure (SOP) library for a specific preservative, diagnosis, analyte, or preanalytical factor. The BRD incorporates information from international efforts, including those by the International Society for Biological and Environmental Repositories (ISBER) and the European Union-sponsored SPI-DIA program (standardization and improvement of generic preanalytical tools and procedures for *in vitro* diagnostics; [www.spidia.eu](http://www.spidia.eu)), among others.

To facilitate the implementation of evidence-based practices in biospecimen handling, BBRB has developed a document series termed Biospecimen Evidence-Based Practices

(BEBP), which contains step-by-step procedural guidelines derived from peer-reviewed primary research articles and expert experience.<sup>4</sup> The aim of the BEBP series is to promote a practical level of standardization and improve overall biospecimen quality and data reproducibility by specifying both optimal methods and suitable alternatives, while merging published evidence with practical knowledge of experts in the field. The intent of the BEBP is not to serve as a SOP, but to facilitate the development of evidence-based SOPs by individual laboratories.

The present BEBP focuses on nucleic acid extraction from formalin-fixed, paraffin-embedded (FFPE) tissue biospecimens (see Supplementary Data; Supplementary Data are available online at [www.liebertpub.com/bio](http://www.liebertpub.com/bio)). Such biospecimens are being increasingly utilized in genomic research, and it has become clear over the past decade that variable and suboptimal FFPE biospecimen collection and processing practices can alter the quantity and/or quality of extracted DNA and RNA.<sup>5</sup> Concordance between the molecular data generated with FFPE and snap-frozen biospecimens varies widely among reports, with correlations ranging from weak to very strong for the same analytical method.<sup>6</sup> While lack of concordance may be partially attributable to differences in biospecimen handling during FFPE processing, available evidence suggests that the extraction method can compound or mitigate effects introduced during biospecimen collection, processing, and storage, thus affecting the suitability of samples for downstream analysis.<sup>7-11</sup> When deciding on a nucleic acid extraction procedure, it is paramount to consider artifacts that may have been introduced during formalin fixation and processing, such as nucleic acid fragmentation,<sup>12</sup> nucleic acid-protein crosslinking,<sup>13,14</sup> denaturation,<sup>15,16</sup> and

<sup>1</sup>Kelly Government Solutions, Rockville, Maryland.

<sup>2</sup>Preferred Solutions Group, Arlington, Virginia.

<sup>3</sup>Cytogenetics/Molecular Genetics Laboratory at Nationwide Children's Hospital, Columbus, Ohio.

<sup>4</sup>Biorepositories and Biospecimen Research Branch, Cancer Diagnosis Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute, Bethesda, Maryland.

<sup>5</sup>Department of Genetics, Lineberger Comprehensive Cancer Center, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina.

<sup>6</sup>Canada's Michael Smith Genome Sciences Center, BC Cancer Agency, Vancouver, Canada.

<sup>7</sup>Human Genome Sequencing Center, Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas.

additions of methylol groups to nucleic acids.<sup>15,17</sup> Importantly, many such formalin-induced modifications to DNA and RNA are reversible<sup>13,18–21</sup>; for example, the addition of a demodification step as well as optimization of multiple steps during the extraction procedure can attenuate effects introduced during fixation and processing. However, FFPE-specific optimization steps are interdependent and must be considered collectively when developing a strategy for extraction. While several commercial extraction kits are marketed as being tailored for FFPE biospecimens, experts contributing to the BEBP advised validating any kit for the intended tissue type, processing regime, and analytical platform before implementation in experimental studies. Validation includes experiments to confirm the feasibility and accuracy of the intended analytical use, the reproducible performance of the kit, and the robustness to processing regimes.

## Materials and Methods

The BEBP for nucleic acid extraction from FFPE biospecimens was built upon a framework of key procedural steps (or preanalytical factors) during DNA and RNA extraction from FFPE tissue biospecimens that was generated through an initial survey of the literature and available SOPs. These preanalytical factors were used to guide targeted searches of the National Library of Medicine (NLM)’s

PubMed database (pubmed.gov) and NCI’s BRD. Published evidence was compiled regarding how each step or extraction-specific preanalytical factor affects nucleic acid yield, quality, integrity, or downstream analysis. The list of potential pre-analytical factors was expanded through crossreferencing and pertinent articles were curated for inclusion in the BRD. Once a draft document was available for review, experts were invited to participate on a review panel based on their expertise and publication history in the field. Feedback based on the knowledge, experiences, and opinions of participating experts was collected primarily through electronic correspondence both upon review of the original document and in response to detailed questions. Additional input was captured during a teleconference after experts had reviewed a complete draft.

## Results

The final expert-vetted BEBP document contains step-by-step procedural guidelines for the extraction of nucleic acids from FFPE tissue (Supplementary Data). Recommendations outlined within the procedural guideline section are a culmination of the summarized literature, protocols, and details from commercial extraction kits, published guidelines by the Centers for Disease Control and Prevention and the Clinical & Laboratory Standards Institute Laboratory, and guidance from the expert panel. As more than one method was often

TABLE 1. DNA QUALITY GUIDELINES BY ANALYTICAL PLATFORM

<i>Application</i>	<i>Criteria employed</i>	<i>Reference</i>
NGS	Monitor yield after library preparation and mean insert size as predictors for characterization success. Age of sample is not a predictor of success, as fixation technique plays a greater role.	BROAD Institute ( <a href="http://genomics.broadinstitute.org/data-sheets/DTS_FFPE_4-2017.pdf">http://genomics.broadinstitute.org/data-sheets/DTS_FFPE_4-2017.pdf</a> )
	A minimum of 100 ng DNA is used for library construction. DNA integrity is not assessed by gel electrophoresis. Library metrics are used to determine pass/fail status before sequencing. Successful libraries should have the majority of library fragments between 300 and 600 bp in size with a minimum yield of 15 µL at 3 nM.	Dr. Andrew Mungall BC Cancer ( <a href="http://www.bcgsc.ca/services/sequencing-libraries-faq">www.bcgsc.ca/services/sequencing-libraries-faq</a> )
	Sample intake QC-minimum DNA integrity (>200 bp) and absence of protein contamination evaluated by E-Gel.	Dr. Harsha Doddapaneni and Dr. David Wheeler (Baylor College of Medicine Human Genome Sequencing Center)
	Library construction yields should be >300 ng with fragments between 200 and 800 bp when using 100 ng input (manual preparation) or 250 ng (robotic preparation).	
	Post-library capture should have >10 nM yield and devoid of primer dimers.	
	150 ng double-stranded DNA, amplification of 100 bp product; ΔCt <2 using the FFPE QC Kit >6% amplifiable copies; Input adjusted based on PCR amplification of TBP or FTH1	Personal communication Dr. Betsou (IBBL, Luxembourg) Sah et al. <sup>22</sup>
PCR	ΔCt <1.55 real-time PCR-based Illumina FFPE QC Kit	Serizawa et al. <sup>23</sup> Wang et al. <sup>24</sup>
aCGH	Comparative assessment of differentially sized GAPDH PCR amplicons: 100, 236, 299, 411, bp visualized by HPLC	
	Amplification of a 200 bp fragment of GAPDH from 100 ng DNA	van Beers et al. <sup>25</sup>
	Amplification of 200 bp product; >2 µg DNA	Personal Communication Dr. Betsou (IBBL, Luxembourg)

aCGH, array comparative genomic hybridization; FFPE, formalin-fixed paraffin-embedded; FTH1, ferritin heavy polypeptide 1; GAPDH, glyceraldehyde 3-phosphate dehydrogenase; HPLC, high-performance liquid chromatography; NGS, next-generation sequencing; PCR, polymerase chain reaction; TBP, tata-binding protein.



using the BEBP should prove a valuable tool in improving the quality and reproducibility of molecular data generated with FFPE biospecimens. This document can also be found on BBRB's website (<http://biospecimens.cancer.gov>).

## Acknowledgments

The authors thank Dr. Fay Betsou (IBBL, Luxembourg), Dr. Jakob Hedegaard (MOMA, Denmark), Dr. William Mathieson (IBBL, Luxembourg), and Dr. Geraldine Thomas (Imperial College, England) for their participation on the expert panel and their insightful recommendations.

This work was funded by the National Cancer Institute's Biorepositories and Biospecimen Research Branch.

## Author Disclosure Statement

No conflicting financial interests exist.

## References

1. Poste G. Bring on the biomarkers. *Nature* 2011;469:156–157.
2. Taube SE, Clark GM, Dancey JE, et al. A perspective on challenges and issues in biomarker development and drug and biomarker codevelopment. *J Natl Cancer Inst* 2009;101:1453–1463.
3. de Gramont A, Watson S, Ellis LM, et al. Pragmatic issues in biomarker evaluation for targeted therapies in cancer. *Nat Rev Clin Oncol* 2015;12:197–212.
4. Engel KB, Vaught J, Moore HM. National Cancer Institute Biospecimen Evidence-Based Practices: A novel approach to pre-analytical standardization. *Biopreserv Biobank* 2014;12:148–150.
5. Bass BP, Engel KB, Greytak SR, et al. A review of pre-analytical factors affecting molecular, protein, and morphological analysis of formalin-fixed, paraffin-embedded tissue. *Arch Pathol Lab Med* 2014;138:1520–1530.
6. Greytak SR, Engel KB, Bass BP, et al. Accuracy of molecular data generated with FFPE biospecimens: Lessons from the literature. *Cancer Res* 2015;75:1541–1547.
7. Turashvili G, Yang W, McKinney S, et al. Nucleic acid quantity and quality from paraffin blocks: Defining optimal fixation, processing and DNA/RNA extraction techniques. *Exp Mol Pathol* 2012;92:33–43.
8. Funabashi KS, Barcelos D, Visoná I, et al. DNA extraction and molecular analysis of non-tumoral liver, spleen, and brain from autopsy samples: The effect of formalin fixation and paraffin embedding. *Pathol Res Pract* 2012;208:584–591.
9. Kalmár A, Péterfia B, Hollósi P, et al. Bisulfite-based DNA methylation analysis from recent and archived formalin-fixed, paraffin embedded colorectal tissue samples. *Pathol Oncol Res* 2015;21:1149–1156.
10. Tourmier B, Chapusot C, Courcet E, et al. Why do results conflict regarding the prognostic value of the methylation status in colon cancers? The role of the preservation method. *BMC Cancer* 2012;12:12.
11. Ludyga N, Grünwald B, Azimzadeh O, et al. Nucleic acids from long-term preserved FFPE tissues are suitable for downstream analyses. *Virchows Arch* 2012;460:131–140.
12. Pavelic J, Gall-Troselj K, Bosnar MH, et al. PCR amplification of DNA from archival specimens. A methodological approach. *Neoplasma* 1996;43:75–81.
13. Jackson V. Studies on histone organization in the nucleosome using formaldehyde as a reversible cross-linking agent. *Cell* 1978;15:945–954.
14. Brutlag D, Schlehuber C, Bonner J. Properties of formaldehyde-treated nucleohistone. *Biochemistry* 1969;8:3214–3218.
15. McGhee JD, von Hippel PH. Formaldehyde as a probe of DNA structure. 4. Mechanism of the initial reaction of Formaldehyde with DNA. *Biochemistry* 1977;16:3276–3293.
16. Karlsen F, Kalantari M, Chitemerere M, et al. Modifications of human and viral deoxyribonucleic acid by formaldehyde fixation. *Lab Invest* 1994;71:604–611.
17. Masuda N, Ohnishi T, Kawamoto S, et al. Analysis of chemical modification of RNA from formalin-fixed samples and optimization of molecular biology applications for such samples. *Nucleic Acids Res* 1999;27:4436–4443.
18. Oberli A, Popovici V, Delorenzi M, et al. Expression profiling with RNA from formalin-fixed, paraffin-embedded material. *BMC Med Genomics* 2008;1:9.
19. Hamatani K, Eguchi H, Takahashi K, et al. Improved RT-PCR amplification for molecular analyses with long-term preserved formalin-fixed, paraffin-embedded tissue specimens. *J Histochem Cytochem* 2006;54:773–780.
20. Jackson V. Formaldehyde cross-linking for studying nucleosomal dynamics. *Methods* 1999;17:125–139.
21. Shi SR, Cote RJ, Wu L, et al. DNA extraction from archival formalin-fixed, paraffin-embedded tissue sections based on the antigen retrieval principle: Heating under the influence of pH. *J Histochem Cytochem* 2002;50:1005–1011.
22. Sah S, Chen L, Houghton J, et al. Functional DNA quantification guides accurate next-generation sequencing mutation detection in formalin-fixed, paraffin-embedded tumor biopsies. *Genome Med* 2013;5:77.
23. Serizawa M, Yokota T, Hosokawa A, et al. The efficacy of uracil DNA glycosylase pretreatment in amplicon-based massively parallel sequencing with DNA extracted from archived formalin-fixed paraffin-embedded esophageal cancer tissues. *Cancer Genet* 2015;208:415–427.
24. Wang F, Wang L, Briggs C, et al. DNA degradation test predicts success in whole-genome amplification from diverse clinical samples. *J Mol Diagn* 2007;9:441–451.
25. van Beers EH, Joosse SA, Ligtenberg MJ, et al. A multiplex PCR predictor for aCGH success of FFPE samples. *Br J Cancer* 2006;94:333–337.
26. Penland SK, Keku TO, Torrice C, et al. RNA expression analysis of formalin-fixed paraffin-embedded tumors. *Lab Invest* 2007;87:383–391.
27. Roberts L, Bowers J, Sensinger K, et al. Identification of methods for use of formalin-fixed, paraffin-embedded tissue samples in RNA expression profiling. *Genomics* 2009;94:341–348.
28. Duenwald S, Zhou M, Wang Y, et al. Development of a microarray platform for FFPET profiling: Application to the classification of human tumors. *J Transl Med* 2009;7:65.
29. Abramovitz M, Ordanic-Kodani M, Wang Y, et al. Optimization of RNA extraction from FFPE tissues for expression profiling in the DASL assay. *BioTechniques* 2008;44:417–423.

Address correspondence to:

*Helen M. Moore, PhD*

*Biorepositories and Biospecimen Research Branch*

*Cancer Diagnosis Program*

*Division of Cancer Treatment and Diagnosis*

*National Cancer Institute*

*9609 Medical Center Drive, Rm 3W422, MSC 9728*

*Bethesda, MD 20892*

*E-mail: moorehe@mail.nih.gov*