

Rowan University

## Rowan Digital Works

---

Stratford Campus Research Day

25th Annual Research Day

---

May 6th, 12:00 AM

### Lipoxin A 4 (LxA 4 ) Promotes Reduction and Antibiotic Efficacy Against *Pseudomonas aeruginosa* Biofilm

Julianne M. Thornton  
*Rowan University*

Jean Walker  
*Rowan University*

Prem Y.K. Sundarasivarao  
*Rowan University*

Bernd Spur  
*Rowan University*

Ana Rodriguez  
*Rowan University*

Follow this and additional works at: [https://rdw.rowan.edu/stratford\\_research\\_day](https://rdw.rowan.edu/stratford_research_day)

See next page for additional authors

 Part of the [Bacterial Infections and Mycoses Commons](#), [Bacteriology Commons](#), [Biological Phenomena](#), [Cell Phenomena](#), and [Immunity Commons](#), [Medical Cell Biology Commons](#), and the [Medicinal and Pharmaceutical Chemistry Commons](#)

Let us know how access to this document benefits you - share your thoughts on our [feedback form](#).

---

Thornton, Julianne M.; Walker, Jean; Sundarasivarao, Prem Y.K.; Spur, Bernd; Rodriguez, Ana; and Yin, Kingsley, "Lipoxin A 4 (LxA 4 ) Promotes Reduction and Antibiotic Efficacy Against *Pseudomonas aeruginosa* Biofilm" (2021). *Stratford Campus Research Day*. 52.  
[https://rdw.rowan.edu/stratford\\_research\\_day/2021/may6/52](https://rdw.rowan.edu/stratford_research_day/2021/may6/52)

This Poster is brought to you for free and open access by the Conferences, Events, and Symposia at Rowan Digital Works. It has been accepted for inclusion in Stratford Campus Research Day by an authorized administrator of Rowan Digital Works.

---

**Author(s)**

Julianne M. Thornton, Jean Walker, Prem Y.K. Sundarasivarao, Bernd Spur, Ana Rodriguez, and Kingsley Yin



Julianne M. Thornton, Jean Walker, Prem Y. K. Sundarasivarao, Bernd Spur, Ana Rodriguez, Kingsley Yin

Rowan University School of Osteopathic Medicine, Stratford, NJ, USA


## Abstract

*Pseudomonas aeruginosa* (*P. aeruginosa*) is an opportunistic bacterium commonly found in wound infections and airways of cystic fibrosis patients. *P. aeruginosa* readily forms biofilms which can reduce the efficacy of antibiotics used to eradicate the pathogen. We have previously shown that a Specialized Pro-resolving Mediator (SPM), Lipoxin A<sub>4</sub> (LxA<sub>4</sub>) is a quorum sensing inhibitor which can reduce *P. aeruginosa* virulence. In this study, we examined the direct actions of LxA<sub>4</sub> and RvD<sub>2</sub> on *P. aeruginosa* biofilm formation and virulence gene expression. The influence of LxA<sub>4</sub> on antibiotic efficacy and the combined effects on biofilm formation were also investigated. LxA<sub>4</sub> and RvD<sub>2</sub> reduced *P. aeruginosa* biofilm formation and virulence gene expression. LxA<sub>4</sub> increased ciprofloxacin inhibition on biofilm formation but did not affect ciprofloxacin's action on non-adherent bacteria. On the other hand, LxA<sub>4</sub> increased bacterial killing action of imipenem but did not affect imipenem's action on biofilm. We also found that LxA<sub>4</sub> can increase ciprofloxacin's bacterial killing ability in established biofilm. Together these results suggest that LxA<sub>4</sub> has direct effects on *P. aeruginosa* biofilm formation and can increase antibiotic efficacy directly.

## Specialized Pro-Resolving Mediators


Fig. 1. Chemical structures of specialized pro-resolving mediators RvD<sub>1</sub>, RvD<sub>2</sub>, RvD<sub>5</sub>, and LxA<sub>4</sub>.

All three resolvins are 22C chains; RvD<sub>1</sub> has hydroxyl groups at C7, 8 and 17; RvD<sub>2</sub> has hydroxyl groups at C7, 16 and 17; and RvD<sub>5</sub> has hydroxyl groups at C7 and 17. LxA<sub>4</sub> is a 20C chain with hydroxyl groups on C5, 6 and 15.



## LxA<sub>4</sub> and RvD<sub>2</sub> Reduce Two Quorum-sensing Genes' Expression


Fig. 4. Effects of SPMs on *Pseudomonas aeruginosa* quorum-sensing virulence gene expression. (a) 1/ $\Delta$ CT of two virulence genes' expression when treated with 1 nM LxA<sub>4</sub>. LxA<sub>4</sub> significantly reduced both genes' expression at 4.5 hr. (b) 1/ $\Delta$ CT of two virulence genes' expression when treated with 1 nM RvD<sub>2</sub>. RvD<sub>2</sub> significantly reduced *pqsA* gene expression at 4.5 hr. Data are mean  $\pm$  s.e.m., with p-values determined by unpaired t-test comparing untreated control to treatment group of same timepoint. \*  $p < 0.05$ , \*\*  $p < 0.01$ . n = 3 independent experiments.



## Hypotheses: How LxA<sub>4</sub> Can Assist Antibiotics Against Biofilms

**Biofilm Formation:** LxA<sub>4</sub> antagonizes the LasR receptor, blocking the quorum-sensing auto-inducer 3-OC12 HSL, reducing biofilm formation. Antibiotics (ABX) are then able to access the bacteria and kill them.

**Established Biofilm:** LxA<sub>4</sub>, as a lipid, can integrate with the biofilm matrix, weakening its integrity. This provides an opportunity for antibiotics (ABX) to access the bacteria within the biofilm and kill them.



## LxA<sub>4</sub> Enhances Ciprofloxacin Effects on Biofilm Formation But Has No Effect on Planktonic Cell Growth





Fig. 5. LxA<sub>4</sub> treatment can aid the efficacy of the antibiotic ciprofloxacin in reducing biofilm formation. Cultures were treated in a 96-well plate overnight with multiple concentrations of ciprofloxacin and LxA<sub>4</sub>. The cultures and apical biofilm were washed away, and the remaining apical biofilm rings were stained with 0.1% crystal violet. Absorbance was measured at 600 nm. Some cultures were recovered from the 96-well plate, diluted and spread on tryptic soy agar plates to incubate overnight at 37°C. Colonies were counted the next day. (a) LxA<sub>4</sub> does not have an effect when combined with ciprofloxacin on non-adherent cell growth, determined by colony forming units (CFUs). (b) When combined with LxA<sub>4</sub>, biofilm formation is significantly reduced in ciprofloxacin treatments at and above bactericidal doses (0.5 µg/mL). CFU data are mean  $\pm$  s.e.m. of percent change from control adjusted to zero. Biofilm data are mean  $\pm$  s.e.m. percent of control. \*\* =  $p < 0.01$ ; CFUs n = 4 independent experiments; biofilm n = 6 (1 nM) and n = 3 (10 nM) independent experiments.


## LxA<sub>4</sub> Does Not Affect Imipenem Efficacy Against Biofilm Formation But Enhances Suppression of Planktonic Growth

Fig. 6. LxA<sub>4</sub> in combination with imipenem significantly decreased non-adherent cell growth but not biofilm formation. Cultures were treated in a 96-well plate overnight with multiple concentrations of imipenem and LxA<sub>4</sub>. The cultures and apical biofilm were washed away, and the remaining apical biofilm rings were stained with 0.1% crystal violet. Absorbance was measured at 600 nm. Some cultures were recovered from a 96-well plate, diluted and spread on tryptic soy agar plates to incubate overnight at 37°C. Colonies were counted the next day. (a) LxA<sub>4</sub> (10 nM) in combination with imipenem (0.1 µg/mL) significantly decreased non-adherent cell growth. (b) LxA<sub>4</sub> does not significantly affect the efficacy of imipenem against biofilm formation. These data combined with the ciprofloxacin data (Fig. 4) suggests the synergistic action of LxA<sub>4</sub> with antibiotics is dependent on the class of antibiotics and their mechanism of action. CFU data are mean  $\pm$  s.e.m. of percent change from control adjusted to zero. Biofilm data are mean  $\pm$  s.e.m. percent of control. \* =  $p < 0.05$ ; CFUs n = 4 independent experiments; biofilm n = 3 (LxA<sub>4</sub> 1 nM) and n = 4 (LxA<sub>4</sub> 10 nM) independent experiments.



## LxA<sub>4</sub> Can Increase the Efficacy of Ciprofloxacin Against Bacteria of an Established Biofilm

Fig. 7. Effects of LxA<sub>4</sub> on ciprofloxacin action in pre-formed biofilms. (a) LxA<sub>4</sub> (1 nM) significantly increased the efficacy of ciprofloxacin (0.5 µg/mL) to reduce the amount of metabolically active bacterial cells associated with the biofilm, determined by MTT assay. (b) LxA<sub>4</sub> does not significantly alter the efficacy of ciprofloxacin to reduce biofilm biomass, determined by crystal violet staining. This suggests that LxA<sub>4</sub> treatment can aid the ciprofloxacin in accessing the bacteria associated with a pre-formed biofilm. Viability data are mean  $\pm$  s.e.m. of percent change from control adjusted to zero. Biofilm data are mean  $\pm$  s.e.m. percent of control. \* =  $p < 0.05$ ; viability n = 5 independent experiments; biomass n = 5 independent experiments.



## Conclusions

In summary, our results show that LxA<sub>4</sub> and RvD<sub>2</sub> can directly reduce *P. aeruginosa* biofilm formation. LxA<sub>4</sub> and RvD<sub>2</sub> can also downregulate virulence gene expression. LxA<sub>4</sub> enhances the efficacy of antibiotics directly against *P. aeruginosa* biofilm formation and bacterial cells within existing biofilm. These studies also provide evidence that further investigation into the antimicrobial mechanisms of RvD<sub>2</sub> is warranted. The results suggest that there is relative selectivity in SPM inhibition of biofilm formation.

## Future Directions


- SPM treatment of *P. aeruginosa* to evaluate the effects of SPMs on quorum-sensing gene expression and regulation
- 2-hit treatments of SPMs  $\pm$  antibiotics on established biofilms to assess the effects of SPMs on antibiotic ability to kill biofilm-associated persister cells
- Pre-treatment of THP-1 monocytes with SPMs against established biofilm
- Co-incubation of THP-1 monocytes with SPMs against established biofilm to assess the effects of SPMs on immune cell ability to phagocytose biofilm

## Acknowledgments

We thank Rowan University Graduate School of Biomedical Sciences, New Jersey Health Foundation and the NIH (RO1 A128202)

## RvD<sub>2</sub> and LxA<sub>4</sub> Reduce *P. aeruginosa* Biofilm Formation

Fig. 2. RvD<sub>2</sub> and LxA<sub>4</sub> reduce *P. aeruginosa* biofilm formation. (a) In a 96-well plate, cultures were treated with multiple concentrations of each SPM and incubated at 37°C overnight. The cultures and apical biofilm were washed away, and the remaining apical biofilm rings were stained with 0.1% crystal violet. Absorbance was measured at 600 nm. RvD<sub>2</sub> significantly reduced biofilm formation at all concentrations, with treatment at 0.1 nM less effective than at 1 nM or 10 nM. LxA<sub>4</sub> significantly reduced biofilm formation, though 0.1 nM and 10 nM were less effective than 1 nM. (b) Effects of PPR2 receptor peptide agonist on *P. aeruginosa* biofilm formation. Cultures were treated with multiple concentrations of the peptide agonist WKYMVm. Peptide agonist WKYMVm did not significantly affect the formation of biofilm at either concentration tested. Data are mean  $\pm$  s.e.m. of percent change from control adjusted to zero. \* =  $p < 0.05$ ; \*\* =  $p < 0.01$ ; n = 3 independent experiments for all treatments.



## Non-adherent cell growth of *P. aeruginosa* was unaffected by LxA<sub>4</sub> or RvD<sub>2</sub>




Fig. 3. Non-adherent cell growth of *P. aeruginosa* was unaffected by LxA<sub>4</sub> or RvD<sub>2</sub>. SPMs were added to *P. aeruginosa* cultures in 96-well plates and incubated with orbital shaking at 37°C in a microplate reader for 15 hr. Absorbance was measured every 10 min. Neither LxA<sub>4</sub> (a) nor RvD<sub>2</sub> (b) showed any significant effect on non-adherent cell growth in minimal media. n = 3 (LxA<sub>4</sub>) and n = 4 (RvD<sub>2</sub>) independent experiments.