



Contents lists available at ScienceDirect

## Paediatric Respiratory Reviews



## Review

## The modifiable biopsychosocial drivers of psychological distress for adolescents with asthma: Implications for Clinical Care

Kelsey J. Sharrad<sup>a,\*</sup>, Zoe A. Kopsaftis<sup>a,b,c</sup>, Kristin V. Carson-Chahhoud<sup>a,b,d</sup>, Helen M. Stallman<sup>e</sup><sup>a</sup> Australian Centre for Precision Health, University of South Australia, Adelaide, Australia<sup>b</sup> Adelaide Medical School, University of Adelaide, Adelaide, Australia<sup>c</sup> Respiratory Medicine Unit, The Queen Elizabeth Hospital, Central Adelaide Local Health Network, Adelaide, Australia<sup>d</sup> South Australian Health and Medical Research Institute, Adelaide, Australia<sup>e</sup> Thompson Institute, University of the Sunshine Coast, Birtinya, Queensland, Australia

## Educational Aims

The reader will be able:

- To review the evidence of biopsychosocial drivers of psychological distress and asthma in adolescents.
- To examine the clinical implications of causes of psychological distress in asthma care in adolescents.

## ARTICLE INFO

## Keywords:

Asthma  
 Psychological distress  
 Adolescent  
 Child  
 Mental health  
 Psychological adaptation  
 Health behaviour

## ABSTRACT

**Purpose:** Overwhelming distress exceeds the capacity of healthy coping strategies to feel better using healthy coping strategies alone, resulting in the use of unhealthy coping strategies. Unhealthy coping strategies may exacerbate asthma symptoms and asthma can contribute to overwhelming distress. This study aimed to review the modifiable drivers of overwhelming distress in adolescents with asthma. **Methods:** The biopsychosocial drivers of psychological distress for adolescents with asthma were explored within the domains of the modifiable biopsychosocial model of health and wellbeing. **Results:** Asthma in adolescents is associated with problems in the domains of environment, developmental outcomes, sense of belonging, health behaviours, coping, and treatment of illness. **Conclusions:** The relationship between asthma and psychological distress highlights the need for holistic treatment of asthma. Further research is needed to establish causation between variables and to investigate whether interventions that address either asthma symptoms or biopsychosocial drivers of distress can improve both factors.

© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## INTRODUCTION

Asthma is a significant health problem that contributes to the global burden of disease [2]. The prevalence of asthma is higher in children and adolescents than adults [1–4], and the rate seems to increase over childhood (e.g., 10.2% in 6–7-year olds; 11.3% in 13–14 year olds) [3]. Adolescents with asthma have higher rates

of psychological distress than adolescents who do not have asthma [5,6]. One Australian study, for example, found more than 50% of adolescents with asthma had high levels of psychological distress [6]. Many of the causes of psychological distress are common across adolescents with and without asthma, for example, self-consciousness and body image problems [6]. However, some stressors are illness-related [6]. Psychological distress can impair cognitive functions needed for educational outcomes, including attention [7], behavioural inhibition [8] memory [9], planning [10] and flexible decision making [11]. Understanding the asthma-specific determinants of psychological distress is therefore critical for optimal outcomes for adolescents.

**Abbreviations:** ED, Emergency Department; EIB, Exercise-Induced Bronchoconstriction; IgE, Immunoglobulin E; PA, Physical Activity; PROMIS, Patient-Reported Outcomes Measurement Information System; QoL, Quality of Life.

\* Corresponding author. GPO Box 2471, Adelaide SA 5001, Australia.

E-mail address: [kelsey.sharrad@unisa.edu.au](mailto:kelsey.sharrad@unisa.edu.au) (K.J. Sharrad).

<https://doi.org/10.1016/j.prrv.2021.07.005>

1526-0542/© 2021 The Author(s). Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## BIOPSYCHOSOCIAL MODEL OF HEALTH AND WELLBEING

While there are genetic, biological, social and psychological contributors to psychological distress, optimal clinical care requires an understanding of which factors are modifiable [14,15]. The Biopsychosocial Model of Health and Wellbeing provides an overview of the biological, social, and psychological domains and components that are associated with health and wellbeing healthy environments, developmental competencies, sense of belonging, health behaviours, coping, resilience, and treatment of illness [12,13]. Problems in one or more of these areas is associated with overwhelming psychological distress. Overwhelming distress is unpleasant emotions that exceed the capacity of healthy coping strategies (i.e. self-soothing, relaxation/distraction, social support and professional support) to reduce the distress [14]. When distress exceeds the capacity of healthy coping strategies, people use unhealthy coping strategies to feel better (i.e. negative self-talk, harmful activities, social withdrawal, and suicidality) [15]. Chronic psychological distress during childhood and adolescence is associated with lifelong relationship problems [16], employment problems [17,18] and psychiatric illness in adulthood [19,20]. Chronic use of unhealthy coping strategies to reduce overwhelming distress can result in harm, including disorders developed from unhealthy coping harmful activities (e.g. eating disorders, substance use disorders), criminal charges (e.g., from aggression) and death by suicide. The Biopsychosocial Model of Health and Wellbeing therefore, provides a useful framework to examine the asthma-specific drivers of psychological in adolescents to inform clinical care.

## MODIFIABLE ASTHMA-RELATED BIOPSYCHOSOCIAL DRIVERS OF PSYCHOLOGICAL DISTRESS

### *Healthy environments*

Allergens, and irritants in the physical environment can exacerbate asthma symptoms. These include allergen-specific immunoglobulin E antibodies [21], airborne irritants, foods (e.g., eggs, cow's milk, and nuts), dust mites, mould, and pollen, animal dander [22,23]. Airborne irritants that exacerbate asthma include aerosols or chemical fumes, cigarette smoke. Viral infections can also trigger asthma symptoms [23].

**Social environment factors:** A healthy environment also includes a social environment that supports the health and wellbeing of the child or adolescent. A systematic review that included 25 studies of parents and caregivers of children with and without asthma found greater psychological distress in parents and caregivers of children with asthma [25]. The reason for the increased distress in these parents is unknown. Parental psychological distress (in particular, maternal distress) adversely affects asthma severity, control, and medication adherence in children—the reasons are not well understood [24]. In contrast, emotional support by parents is associated with increased perception of treatment control, reduced stress, and a reduced risk of emotional issues in children [26]. A systematic review of 106 studies found that conflict in peer and family relationships (e.g. teasing, arguing, disagreements) is related to increased self-reported asthma symptoms [26]. Professional support also impacts asthma outcomes. Empathetic relationships between patients and physicians that include clear communication, confidentiality, and shared decision-making also important to this population, and promotes honest disclosure of information by young people with chronic disease [69].

The economic environment also affects psychological distress and asthma-care outcomes. Economic disadvantage can be particularly challenging for treating asthma as optimal treatment

requires regular preventative pharmacotherapy [27,28] and is associated with poor treatment adherence [26,29]. For parents, childhood asthma can also have indirect economic impacts such as work absences, lost productivity, and healthcare costs [30]. Prevalence of delayed or unmet needs for care due to financial barriers in the United States, for example, is estimated at 8–26.6% for adolescents [27]. These threats to a healthy environment for adolescents with asthma can increase asthma symptoms and psychological distress.

### *Developmental competencies*

Adolescence is a developmental period that includes widespread biological, physical, and psychological changes [31,32]. Developmental competencies needed to be achieved during childhood for health and wellbeing are a stable sense of identity, emotional and behavioural regulation, interpersonal skills and problem-solving skills [13]. Adolescents with asthma frequently have problems across all four developmental competencies—identity, emotional and behavioural regulation, interpersonal and problem-solving. An increasing desire for independence and rejection of parental modelling [26,35]; difficulty organising time, forgetfulness, conflict with parents and confusion over responsibility of medication-taking [35], increased perception of barriers [36]; difficulty adequately expressing their health concerns or communicating their experience of symptoms [37], perceived social stigma [26]; and lack of engagement in medication decision-making [35] all contribute to issues in these domains. The importance of social feedback for identity formation during adolescence can lead to self-perceptions of being different from their peers and sickly [33], which is related to greater morbidity [33]. A qualitative review of 5 studies found that people with asthma, like individuals with other chronic diseases, thought their condition threatened their autonomy, and in turn their sense of identity [34].

### *Sense of belonging*

Feeling included and valued is a core component of health and wellbeing. A sense of belonging through peer support is associated with medication adherence [26]. Supportive environments are associated with better self-management, asthma control, and improved quality of life [26]. However, many children and adolescents with asthma report feeling socially isolated because of their illness [26,38]. Qualitative studies report adolescents experiencing social isolation due to their diagnosis and a desire not to be seen as an “ill person” or to be treated differently [35]. One qualitative study found that adolescents struggled with the incongruity of being accepted socially, and managing their condition [39]. Strategies used by some to feel a sense of belonging include ignoring symptoms and not taking medication in public [35,39].

### *Health behaviours*

Health behaviours—sleep, nutrition and physical activity—are key components of wellbeing and asthma management [26]. Psychological distress has a bidirectional relationship with health behaviours, contributing to problems and problems causing distress. Distress and rumination can disrupt sleep onset and cause maintenance insomnia and inadequate sleep can contribute to mood dysregulation. Inadequate nutrition can disrupt mood regulation and eating can be used as a coping strategy to reduce distress. Inadequate physical activity affects mood and psychological distress can make it challenging to be active.

Inadequate sleep quality and quantity are more common in adolescents with asthma, as is distress before sleep onset and

maintenance insomnia problems [40,41]. Poor quality sleep is related to increased asthma symptoms [42] and psychological distress [43]. A recent experimental study [42] found that when adolescents had 6.5 hours of sleep per night, they reported more asthma symptoms and quality of life than those who had 9.5 hours sleep.

Nutrition plays a role in the management of asthma in adolescents [44,45]. Consumption of adequate amounts of fruits and vegetables has been associated with a reduced risk of developing asthma, as well as increased asthma control, likely due to components such as antioxidants, fibre, polyunsaturated fatty acids, and vitamin D which may affect immune pathways. In contrast, dairy consumption has been associated with an increased risk of asthma exacerbation. A recent meta-review reported that Mediterranean and vegan diets which encourage consumption of fruits, vegetables, legumes, and whole grains, and reduce emphasis on animal products (namely saturated animal fat and dairy products), may reduce the risk of asthma development and exacerbation [45].

Being overweight also increases the likelihood of having asthma [46,47], more asthma symptoms [48,49], greater asthma-related healthcare utilisation [50,51], and less responsiveness to inhaled corticosteroids [52,53]—all evidence of greater difficulty controlling asthma symptoms.

There is no evidence that adolescents with asthma are overall less physically active than adolescents without asthma [54,55]. In addition to the general health benefits, physical activity in adolescents is associated with better lung function scores, fewer asthma exacerbations, hospital admissions, school absenteeism, and asthma medication usage [55–58]. In contrast, a meta-analytic review of 11 cross-sectional studies found that low physical activity was associated with new-onset asthma and increased asthma symptoms [56]. Physical activity, however, can trigger asthma symptoms of exercise-induced bronchoconstriction leading to the avoidance of physical activity to prevent symptoms [55]. Additionally, uncontrolled asthma can make physical activity more difficult, which also results in avoidance. These unpleasant experiences are greater in adolescents with asthma who are also overweight or obese [55].

### Coping

While the Health Theory of Coping suggests that all coping strategies are useful for the purpose of reducing distress [59], strategies can be considered healthy or unhealthy depending on their risk of adverse consequences. For example, healthy strategies include relaxing, distracting or self-soothing activities, social support, and professional support, while unhealthy strategies include harmful activities such as alcohol and drug use, as well as social withdrawal. A recent observational study [14] found that participants at all levels of distress used healthy coping strategies, turning to unhealthy coping strategies when the healthy strategies failed to reduce their distress.

A systematic review of 24 studies found that as with people generally, children and adolescents use a range of healthy coping strategies, including changing position, decreasing activity, taking fluids, relaxing activities such as listening to music, distracting activities such as watching TV, and social support. Interestingly, the review found that asthma patients tended to use different coping strategies than healthy participants and patients with other chronic illnesses. [60]. Similarly, a study of 553 Dutch adolescents aged 12–16 with asthma found that the use of healthy coping strategies prevented a decline in QoL scores [61]. Strategies such as thinking more positively about their condition resulted in higher QoL scores, while strategies such as increased worrying and restricting their lifestyle resulted in lower QoL scores. Positive

QoL scores were also seen for adolescents who actively sought information.

Compared with their peers, adolescents with asthma are likely to use more unhealthy coping strategies, such as alcohol, smoking [62–64], and suicidality [65], indicative of overwhelming distress. One study found adolescents with asthma are more likely to smoke than healthy adolescents and continue to smoke to cope with distress [64]. Smoking is particularly problematic in this population as it increases the risk of asthma symptoms and exacerbations and causes a reduced response to asthma medication [62].

### Resilience

Resilience is an innate capacity to bounce back after adversity [70]. A perception of one's resilience is important for health and wellbeing. The belief that you can do something—self-efficacy—improves actions [67]. There has been no research on the self-perception of resilience in adolescents with asthma directly. However, self-efficacy and outcome expectations for asthma treatment is associated with better treatment adherence in adolescents [36].

### Treatment of illness

Early and effective treatment for biopsychosocial drivers of distress and asthma management is critical to minimise the impact of distress and symptoms on health and wellbeing. It is estimated that many adolescents with asthma have not been formally diagnosed or received treatment [66]. In addition to asthma-related morbidity, these adolescents are likely to be further disadvantaged through absenteeism from school, reduced activities and sleep disturbances [66]. Of those who have been diagnosed and received treatment, treatment adherence varies considerably; estimates range from 49% to 71% [35,68]. Nonadherence contributes to poor disease control and adverse asthma outcomes [36]. In addition to other economic and developmental competency barriers to treatment adherence, other factors associated with poor medication adherence in adolescents include treatment provision factors (lack of perceived need or effectiveness, not understanding how to use medication, forgetting to use medication, healthcare relationship), medication-related factors (e.g., difficulty using an inhaler, side-effects), and social factors (stigma) [26,29,69].

There is a bidirectional relationship between psychological distress and asthma treatment adherence. Psychological distress can impair treatment adherence [29]. Subsequently, asthma that is not adequately controlled can increase the risk of asthma exacerbations, increasing psychological distress [1,70]. Psychological distress may also negatively impact self-efficacy and symptom perception and, in turn, upon medication adherence and other health behaviours, including frequency of healthcare utilisation [71]. Treatment for psychological distress in people generally with asthma typically includes pharmacotherapy and/or psychotherapy for symptoms [72,73] rather than underlying drivers of distress. There is little research on interventions in children and adolescents [72,73].

### IMPLICATIONS FOR CLINICAL CARE

This review of the biopsychosocial drivers of overwhelming distress that particularly relate to adolescents with asthma highlights its critical role in assessment and treatment—summarised in Table 1. This distress is not necessarily cause asthma symptoms and asthma may not necessarily cause the distress. However, as adolescence is the time for the onset of most mental illnesses that can have lifelong adverse impacts across functioning, the results

**Table 1**  
Modifiable biopsychosocial domains and components of health and wellbeing and asthma-related vulnerabilities.

| Domain                       | Components   | Asthma-related vulnerabilities  |
|------------------------------|--|---|
| 1 Healthy Environment        | Physical, social, cultural economic  | Allergens and irritants<br>Parental distress<br>Direct costs of medication/treatment; Indirect financial costs of asthma<br>Effect of illness on identity formation |
| 2 Developmental competencies | Healthy identity, emotional and behavioural regulation, interpersonal skills, problem-solving skills | Behavioural regulation problems with asthma management<br>Parent-child conflict related to medication<br>Problems with responsibility for medication                |
| 3 Sense of belonging         | Feeling connected to community   | Social isolation due to asthma –perceived social isolation and physical isolation   |
| 4 Health behaviours          | Sleep, nutrition, exercise   | Increased likelihood of poor sleep, nutrition<br>Low physical activity associated with onset of asthma and increased asthma symptoms                                |
| 5 Coping                     | Healthy and unhealthy strategies   | More likely to use unhealthy coping strategies – alcohol, smoking, suicidality<br>Unhealthy coping strategies exacerbate symptoms                                   |
| 6 Resilience                 | Sense of innate resilience   |   |
| 7 Treatment of illness       | Early and effective treatments for physical and psychological problems                               | Asthma is underdiagnosed<br>Treatment adherence   |

have implications for asthma healthcare provision. Holistic care is warranted that includes:

1. A thorough biopsychosocial assessment for all children and adolescents presenting with asthma symptoms to understand holistic functioning of the adolescent and the context of the family. Ongoing monitoring is needed over time to monitor change.
2. Including children, adolescents and their parents in assessment and treatment planning to ensure the plan is understood, accepted and works for the family. Encourage and support autonomy as appropriate.
3. Coping assessment and planning with all adolescents to ensure they have a coping plan that works for them to cope when they are distressed [15].
4. Referral to and communication between medical, psychological and social service providers as needed to ensure holistic care.

## CONCLUSION

This paper provided a narrative review of the literature on modifiable biopsychosocial factors of health and wellbeing that have been shown to be associated with asthma in children and adolescents. Although primarily correlational research, the relationships between asthma with psychological distress and drivers of distress highlights the need for holistic assessment and treatment of asthma. Longitudinal research that measures the domains of well-

being is needed to better understand the interrelationships between drivers of distress and asthma outcomes. Clinical trials are also needed to evaluate the extent to which ameliorating the drivers of distress indirectly impact asthma outcomes.

## DIRECTIONS FOR FUTURE RESEARCH

Future research is needed to evaluate interventions that target the biopsychosocial drivers of psychological distress in adolescents on asthma and wellbeing outcomes. Longitudinal research that measures the domains of wellbeing is needed.

## FUNDING

The production of this manuscript was supported by Early Career Fellowship funding received from NHMRC (GNT1141521), project funding received from NHMRC (GNT1108903), and project funding received from the Channel 7 Children's Research Foundation (19/10693585) by Associate Professor Carson-Chahhoud.

## CONFLICTS OF INTEREST

The authors have no conflicts of interest to disclose.

## References

- [1] GINA. Global Strategy for Asthma Management and Prevention: Global Initiative for Asthma; 2018.
- [2] Masoli M, Fabian D, Holt S, et al. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 2004;59:469–78.
- [3] ISAAC. Worldwide variations in the prevalence of asthma symptoms: the International Study of Asthma and Allergies in Childhood (ISAAC). 1998;12:315–335.
- [4] Global Asthma Network. The Global Asthma Report 2018. New Zealand: Auckland; 2018.
- [5] AIHW. Asthma. Available at: <https://www.aihw.gov.au/reports/chronic-respiratory-conditions/asthma/contents/asthma2020>.
- [6] Blanchard M, Morris J, Birrell E, et al. National young people and asthma survey: issues and opportunities in supporting the health and wellbeing of young people living with asthma. 2014.
- [7] Purcell R, Maruff P, Kyrios M, et al. Neuropsychological function in young patients with unipolar major depression. *Psychol Med* 1997;27:1277–85.
- [8] Murphy FC, Sahakian BJ, Rubinsztein JS, et al. Emotional bias and inhibitory control processes in mania and depression. *Psychol Med* 1999;29:1307–21.
- [9] Ilsley JE, Moffitt AP, O'Carroll RE. An analysis of memory dysfunction in major depression. *J Affect Disord* 1995;35:1–9.
- [10] Beats BC, Sahakian BJ, Levy R. Cognitive performance in tests sensitive to frontal lobe dysfunction in the elderly depressed. *Psychol Med* 1996;26:591–603.
- [11] Murphy FC, Rubinsztein JS, Michael A, et al. Decision-making cognition in mania and depression. *Psychol Med* 2001;31:679–93.
- [12] Stallman HM. Coping planning: a patient-centred and strengths-focused approach to suicide prevention training. *Australasian Psychiatry* 2018;26:141–4.
- [13] Stallman HM. Suicide following hospitalisation: systemic treatment failure needs to be the focus rather than risk factors. *Lancet Psychiatry* 2020;7:303.
- [14] Stallman HM, Beauquain D, Hermens DF, et al. Modelling the relationship between healthy and unhealthy coping strategies to understand overwhelming distress: A Bayesian network approach. *J Affect Disord Rep* 2021;3:100054. <https://doi.org/10.1016/j.jadr.2020.100054>.
- [15] Stallman HM. Health Theory of Coping. *Australian Psychologist* 2020;55:295–306.
- [16] Nguyen TP, Karney BR, Bradbury TN. Childhood abuse and later marital outcomes: Do partner characteristics moderate the association? *J Fam Psychol* 2017;31:82–92.
- [17] Allwood MA, Widom CS. Child abuse and neglect, developmental role attainment, and adult arrests. *J Res Crime Delinquency* 2013;50:551–78.
- [18] Currie J, Spatz Widom C. Long-term consequences of child abuse and neglect on adult economic well-being. *Child Maltreat* 2010;15:111–20.
- [19] Weich S, Patterson J, Shaw R, et al. Family relationships in childhood and common psychiatric disorders in later life: systematic review of prospective studies. *Br J Psychiatry* 2009;194:392–8.
- [20] Norman RE, Byambaa M, De R, et al. The long-term health consequences of child physical abuse, emotional abuse, and neglect: a systematic review and meta-analysis. *PLoS Med* 2012;9:e1001349.

- [21] Comberiat P, Di Cicco ME, D'Elis S, et al. How Much Asthma Is Atopic in Children? *Front Pediatr* 2017;5:122.
- [22] Foong R-X, du Toit G, Fox AT. Asthma, Food Allergy, and How They Relate to Each Other. 2017;5.
- [23] Vernon MK, Wiklund I, Bell JA, et al. What do we know about asthma triggers? A review of the literature. *J Asthma* 2012;49:991–8.
- [24] Licari A, Ciprandi R, Marsegli G, et al. Anxiety and depression in adolescents with severe asthma and in their parents: preliminary results after 1 year of treatment. *Behav Sci* 2019;9.
- [25] Easter G, Sharpe L, J. Hunt C. Systematic review and meta-analysis of anxious and depressive symptoms in caregivers of children with asthma. *J Pediatr Psychol* 2015;40:623–32.
- [26] Vazquez-Ortiz M, Angier E, Blumchen K, et al. Understanding the challenges faced by adolescents and young adults with allergic conditions: a systematic review. *Allergy* 2020;75:1850–80.
- [27] Scal P, Davern M, Ireland M, et al. Transition to adulthood: delays and unmet needs among adolescents and young adults with asthma. *J Pediatr* 2008;152:471–475.e1.
- [28] Reddel HK, Lembke K, Zwar NJ. The cost of asthma medicines. *Aust Prescr* 2018;41:34–6.
- [29] Apter AJ. Enhancing patient adherence to asthma therapy. In: Bochner BS, ed. *UpToDate*, 17.0 edition: Wolters Kluwer, 2019.
- [30] Deloitte Access Economics, Asthma Australia, National Asthma Council Australia. *The Hidden Cost of Asthma*; 2015.
- [31] Cohen R, Franco K, Motlow F, et al. Perceptions and attitudes of adolescents with asthma. *J Asthma* 2003;40:207–11.
- [32] de Benedictis D, Bush A. The challenge of asthma in adolescence. *Pediatr Pulmonol* 2007;42:683–92.
- [33] Bruzzese J-M, Bonner S, Vincent EJ, et al. Asthma education: the adolescent experience. *Patient Educ Couns* 2004;55:396–406.
- [34] Eassey D, Reddel HK, Ryan K, et al. The impact of severe asthma on patients' autonomy: a qualitative study. *Health Expect* 2019;22:528–36.
- [35] Kaplan A, Price D. Treatment adherence in adolescents with asthma. *J Asthma Allergy* 2020;13:39–49.
- [36] Rhee H, Wicks MN, Dolgoff JS, et al. Cognitive factors predict medication adherence and asthma control in urban adolescents with asthma. *Patient Preference Adherence* 2018;12:929–37.
- [37] Szeffler SJ, Chipps B. Challenges in the treatment of asthma in children and adolescents. *Ann Allergy Asthma Immunol* 2018;120:382–8.
- [38] Stewart M, Letourneau N, Masuda JR, et al. Support needs and preferences of young adolescents with asthma and allergies: "Just No One Really Seems to Understand". *J Pediatr Nurs* 2012;27:479–90.
- [39] Jonsson M, Schuster M, Protudjer JLP, et al. Experiences of daily life among adolescents with asthma – a struggle with ambivalence. *J Pediatric Nurs* 2017;35:23–9.
- [40] Fidler A, Lawless C, LaFave E, et al. Anxiety among adolescents with asthma: Relationships with asthma control and sleep quality. *Clin Pract Pediatric Psychol* 2019;7:151–6.
- [41] Meltzer LJ, Ullrich M, Szeffler SJ. Sleep duration, sleep hygiene, and insomnia in adolescents with asthma. *J Allergy Clin Immunol Pract* 2014;2:562–9.
- [42] Meltzer LJ, Beebe DW, Jump S, et al. Impact of sleep opportunity on asthma outcomes in adolescents. *Sleep Med* 2020;65:134–41.
- [43] Willis TA, Gregory AM. Anxiety disorders and sleep in children and adolescents. *Sleep Medicine Clinics* 2015;10:125–31.
- [44] Papoutsakis C, Priftis KN, Drakouli M, et al. Childhood overweight/obesity and asthma: is there a link? A systematic review of recent epidemiologic evidence. *J Acad Nutr Diet* 2013;113:77–105.
- [45] Alwarith J, Kahleova H, Crosby L, et al. The role of nutrition in asthma prevention and treatment. *Nutr Rev* 2020;78:928–38.
- [46] Azizpour Y, Delpisheh A, Montazeri Z, et al. Effect of childhood BMI on asthma: a systematic review and meta-analysis of case-control studies. *BMC Pediatrics* 2018;18:143.
- [47] Mannino DM, Mott J, Ferdinands JM, et al. Boys with high body masses have an increased risk of developing asthma: findings from the National Longitudinal Survey of Youth (NLSY). *Int J Obesity* 2006;30:6–13.
- [48] Black MH, Smith N, Porter AH, et al. Higher prevalence of obesity among children with asthma. *Obesity (Silver Spring, Md)* 2012;20:1041–7.
- [49] Trasande L, Liu Y, Fryer G, et al. Effects of childhood obesity on hospital care and costs, 1999–2005. *Health Affairs (Project Hope)* 2009;28:w751–60.
- [50] Taylor B, Mannino D, Brown C, et al. Body mass index and asthma severity in the National Asthma Survey. *Thorax* 2008;63:14–20.
- [51] Black MH, Zhou H, Takayanagi M, et al. Increased asthma risk and asthma-related health care complications associated with childhood obesity. *Am J Epidemiol* 2013;178:1120–8.
- [52] Pradeepan S, Garrison G, Dixon AE. Obesity in asthma: approaches to treatment. *Curr Allergy Asthma Rep* 2013;13:434–42.
- [53] Forno E, Lescher R, Strunk R, et al. Decreased response to inhaled steroids in overweight and obese asthmatic children. *J Allergy Clin Immunol* 2011;127:741–9.
- [54] Cassim R, Koplin JJ, Dharmage SC, et al. The difference in amount of physical activity performed by children with and without asthma: a systematic review and meta-analysis. *J Asthma* 2016;53:882–92.
- [55] Oudjedi A, Said Aissa K. Associations between obesity, asthma and physical activity in children and adolescents. *Apunts Sports Med* 2020;55:39–48.
- [56] Lochte L, Nielsen KG, Petersen PE, et al. Childhood asthma and physical activity: a systematic review with meta-analysis and Graphic Appraisal Tool for Epidemiology assessment. *BMC Pediatrics* 2016;16:50.
- [57] Teng Y-K, Huang J-L, Yeh K-W, et al. Influential factors of insufficient physical activity among adolescents with asthma in Taiwan. *PLoS ONE* 2015;9:e116417.
- [58] Dantas FMNA, Correia MAV, Silva AR, et al. Mothers impose physical activity restrictions on their asthmatic children and adolescents: an analytical cross-sectional study. *BMC Public Health* 2014;14:287.
- [59] Stallman HMJAP. *Health theory of coping*. 2020.
- [60] Barton C, Clarke D, Sulaiman N, et al. Coping as a mediator of psychosocial impediments to optimal management and control of asthma. *Respir Med* 2003;97:747–61.
- [61] Van De Ven MOM, Engels RCME, Sawyer SM, et al. The role of coping strategies in quality of life of adolescents with asthma. *Qual Life Res* 2007;16:625–34.
- [62] Lang JE, Tang M. Smoking: it's still a big problem in children with asthma. *J Pediatr (Rio J)* 2019;95:506–8.
- [63] Dowdell EB, Posner MA, Hutchinson MK. Cigarette smoking and alcohol use among adolescents and young adults with asthma. *Nurs Res Pract* 2011;2011:503201.
- [64] Vázquez-Nava F, Vázquez-Rodríguez EM, Vázquez-Rodríguez CF, et al. Epidemiological profile of smoking and nicotine addiction among asthmatic adolescents. *Public Health* 2017;149:49–56.
- [65] Barker E, Kölves K, De Leo D. The relationship between asthma and suicidal behaviours: a systematic literature review. *Eur Respir J* 2015;46:96–106.
- [66] Yeatts K, Johnston Davis K, Peden D, et al. Health consequences associated with frequent wheezing in adolescents without asthma diagnosis. *Eur Respir J* 2003;22:781–6.
- [67] Robert W, Daniel G, Allen J. Expectations and performance: an empirical test of Bandura's self-efficacy theory. *J Sport Psychol* 1979;1:320–31.
- [68] Desager K, Vermeulen F, Bodart E. Adherence to asthma treatment in childhood and adolescence – a narrative literature review. *Acta Clin Belg* 2018;73:348–55.
- [69] Wilson E. Engaging young people with a chronic illness. *Aust Fam Physician* 2017;46:572–6.
- [70] Goodwin RD, Robinson M, Sly PD, et al. Severity and persistence of asthma and mental health: a birth cohort study. *Psychol Med* 2013;43:1313–22.
- [71] Baiardini I, Sicuro F, Balbi F, et al. Psychological aspects in asthma: do psychological factors affect asthma management? *Asthma Res Practice* 2015;1:7.
- [72] Yorke J, Fleming SL, Shuldham C. Psychological interventions for children with asthma. *Cochrane Database Syst Rev* 2005.
- [73] Sharrad KJ, Sanwo O, Carson-Chahhoud KV, et al. Psychological interventions for asthma in children and adolescents. *Cochrane Database Syst Rev* 2019.