

Protocol for lapis lazuli provenance determination: evidence for an Afghan origin of the stones used for ancient carved artefacts kept at the Egyptian Museum of Florence (Italy)

Alessandro Lo Giudice^{1,2} · Debora Angelici^{2,3} · Alessandro Re^{1,2} · Gianluca Gariani¹ · Alessandro Borghi³ · Silvia Calusi^{4,5} · Lorenzo Giuntini^{4,5} · Mirko Massi^{4,5} · Lisa Castelli⁵ · Francesco Taccetti⁵ · Thomas Calligaro^{6,7} · Claire Pacheco^{6,7} · Quentin Lemasson^{6,7} · Laurent Pichon^{6,7} · Brice Moignard^{6,7} · Giovanni Pratesi⁸ · Maria Cristina Guidotti⁹

Received: 30 May 2016 / Accepted: 27 October 2016 / Published online: 8 November 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract Despite that the Badakhshan Province (Afghanistan) remains the most plausible hypothesis for the lapis lazuli used in antiquity, alternatives proposed in literature are worth to study to confirm or disprove their historical reliability. In this work, a protocol for determining the provenance of lapis lazuli rocks used for carved artefacts is described. Markers for the univocal attribution of the raw material to a source were identified analysing 45 rocks of known provenance (among which 15 georeferenced) from 4 quarry districts. To the best of our knowledge, this reference database is the widest in provenance studies on lapis lazuli. All the markers are recognisable by means of Ion Beam Analysis (IBA) techniques, in particular micro-proton-induced x-ray emission (PIXE) and micro-ionoluminescence (IL). These techniques are non-invasive and applicable in air, allowing to analyse artworks and rocks of practically any shape and dimension without sample preparation.

The protocol was applied to determine the provenance of raw material used for carved lapis lazuli artefacts kept at the Egyptian Museum of Florence, the second most important

Egyptian museum in Italy, second only to the museum of Turin. The collection in Florence has a great historical value and includes several lapis lazuli pendants, scarabs, small statuettes and amulets ascribable mainly to the first millennium BC. Following the protocol, 11 of these artefacts were analysed by means of IBA techniques. Results ascribe the raw material to the Afghan quarry district.

Keywords Lapis lazuli · Provenance · Ancient Egypt · Archaeometry · μ -PIXE · Ionoluminescence

Introduction and Background

Historical context

Lapis lazuli is a very attractive semi-precious stone. Due to its peculiar blue colour and its rareness, lapis lazuli has been used since the Neolithic Era for the manufacturing of

✉ Alessandro Lo Giudice
alessandro.logiudice@unito.it

¹ Dipartimento di Fisica, Università di Torino, Via Giuria 1, 10125 Torino, Italy

² INFN Sezione di Torino, Via Giuria 1, 10125 Torino, Italy

³ Dipartimento di Scienze della Terra, Università di Torino, Via Valperga Caluso 35, 10125 Torino, Italy

⁴ Dipartimento di Fisica, Università di Firenze, Via Sansone 1, 50019, Sesto Fiorentino, Florence, Italy

⁵ INFN Sezione di Firenze, Via Sansone 1, 50019, Sesto Fiorentino, Florence, Italy

⁶ Centre de Recherche et de Restauration des Musées de France, Palais du Louvre, Paris, France

⁷ FR3506 New AGLAE CNRS/Ministère de la Culture et de la Communication, Paris, France

⁸ Dipartimento di Scienze della Terra and Museo di Storia Naturale, Università di Firenze, Via G. La Pira 4, 50121 Florence, Italy

⁹ Museo Egizio di Firenze, Via della Pergola 65, 50121 Florence, Italy

precious objects and jewels (beads, gems, seals, small decorative artworks, etc.). The earliest evidence of the use of lapis lazuli comes from archaeological finds in Mehrgarh (Pakistan) where ornaments in lapis lazuli have been discovered in VII millennium BC burials (Jarrige 1991). During the Chalcolithic period, the number of objects made in lapis lazuli increased as confirmed by their dissemination through the Iranian plateau and the Indus Valley. The greatest success of this stone occurred in the third millennium with a diffusion all over the Middle East and in part of Central and Southern Asia (current Turkmenistan, Afghanistan and Pakistan) (Casanova 2013). Lapis lazuli played a key role as precious material particularly in the Syro-Mesopotamian area, where the majority of archaeological findings of handworks made of this rock are concentrated. Moreover, lapis lazuli was also cited many times along with gold, in the first-written poem, the Epic of Gilgamesh. Afterwards, lapis lazuli has been continuously used until the first millennium BC, when it was gradually replaced by other synthetic materials. Nevertheless, its use in glyptic, and also as pigment from VII century AD, has continued onward.

In Ancient Egypt, lapis lazuli has been used for scarabs, pendants, beads and inlaid jewels since IV millennium BC (Sofianides and Harlow 1990); in particular, many artefacts were found in Hierakonpolis and Abydos necropolis. As well as carnelian, turquoise and malachite, lapis lazuli was considered a sacred stone; in particular, it was associated with the colour of the night sky. This is why many amulets and scarabs were manufactured with this and other sacred stones, although they were often replaced by the less expensive faïence and coloured glass paste. The use of lapis lazuli did not reach in ancient Egypt a diffusion comparable with the one achieved in the Mesopotamian area, but nonetheless remains remarkable. In fact, the Tôt treasure, discovered in 1936 in Egypt (Pierrat-Bonnefois 1995), contains a huge set of ancient lapis lazuli, second only to the corpus of objects in the royal tombs of Ur. Ascribed to Amenemhat II (around 1929–1898 BC), the treasure was found in the basement of a temple dedicated to the worship of Monthu, 30 km south of Luxor. The finding includes thousands of blocks of raw lapis lazuli, minute fragments, beads and carvings. This discovery was very important because most of the objects are not of Egyptian origin, but they come from Anatolia, Mesopotamia and Iran. According to their typology, the objects are in fact distributed on a time frame of about 700 years, from the latter half of the third millennium BC to the early second millennium BC. Among the jewels, mostly ascribable to the second half of the third millennium in Mesopotamia, there are numerous lapis lazuli amulets similar to those found in Susa and Mari (Shafer 1963). For these reasons, the treasure represents

a unique evidence of lapis lazuli trade among Ancient Egypt, Eastern Mediterranean and Near East civilisations.

Sources of lapis lazuli

Where did the lapis lazuli rocks used by Ancient Egyptians, and generically in antiquity, come from? Due to the low probability of geological conditions in which it can be formed, only few sources of lapis lazuli exist in the world (Von Rosen 1988). For this reason, the possibility to associate the raw material to man-made objects is feasible and could help historians and archaeologists to reconstruct ancient trade routes, especially for the ancient time when written testimonies are scanty or absent at all (Tosi 1974). Although the distance is not the only factor to be taken into account, it is logical to assume that the sources of lapis lazuli should be where the material was used in antiquity or in the neighbour regions. In the diffusion area of lapis lazuli in ancient time (VII–I millennium BC), shown in Fig. 1, the most important source of lapis lazuli, still currently exploited, is in Badakhshan Province (Afghanistan), in particular the quarry district of Sar-e-Sang. According to the archaeological evidences, the majority of the authors agree in considering the Badakhshan Province as the only source of lapis lazuli in antiquity. This hypothesis would imply a continuous use of these quarries for nearly 9000 years, making this area one of the oldest mining sites in the world, and a trade up to almost 4000 km already in the fourth millennium BC.

However, in addition to the Afghan quarries, other potential Asian extraction sites are known (Mindat) or hypothesised (Fig. 1). In particular, Liadjura-Dara (Tajikistan), at about 200 km northeast of Sar-e-Sang, is currently the only other well-documented geological source in the diffusion area of lapis lazuli in ancient time. Moreover, also in the Swat Valley, in the north of Pakistan near the Afghan border and less than 200 km from Sar-e-Sang, veins of lapis lazuli are reported (Khaliq 2006). Other two well-known Asian sources of lapis lazuli are near Irkutsk, in the south region of Lake Baikal area (Siberia), and near Mogok in the Mandalay Region (Myanmar). In both cases, the distance from Mesopotamia is more than 6000 km and up to almost 7000 km respectively from Egypt; nevertheless, few authors did not exclude Lake Baikal area as an occasional source of lapis lazuli in antiquity (Hermann 1968).

Also, the Chagai Hills district in Pakistan has been suggested as a potential lapis lazuli source in many works (Jarrige 1988; references in Law 2014); rock samples purported to be from this district have even been used in provenance studies (Delmas 1990; Casanova 1992; Ballirano 2006). The interest in this area arises from the fact that the Chagai Hills lie just 300 km west to the Mehrgarh site, about 400 km closer than Badakhshan Province. However, doubts have been expressed on the real existence of this source, which is neither



Fig. 1 Map of the approximately diffusion area of lapis lazuli in ancient time (IV–I millennium BC). The *blue squares* are documented quarries; the *orange circles* are hypothetical sources whose existence has not been

proven; the *yellow triangle* is the first documented site where lapis lazuli was used (Mehergarh, VII millennium BC)

documented nor supported by geological data, and it has been speculated that the lapis lazuli rock samples associated with Chagai Hills actually came from the Badakhshan quarries (Law 2008; Law 2014).

In the few previous provenance studies are also mentioned analysis on rocks from not well-specified sources in “Persia” and “Levant” (Borelli 1986; Cipriani 1988), in Ural Mountains (Casanova 1992; Zöldföldi 2006), in Iran and in Pakistan (Ballirano and Maras 2006), in Russia and in Siberia (Gambardella et al. 2016) that are not, however, well documented. This is also an issue common to many museum stone collections in which there is a lack of information about the exact provenance of the stones and only a general indication is available. In these cases, it is not possible to know whether there are actually other sources of lapis lazuli or instead these rocks could be attributed to known sources. For instance, Russian and Siberian rocks could be essentially attributed to the same source, i.e. the Lake Baikal area.

Finally, some lapis lazuli sources have been proposed on the basis of ancient written testimonies, but their existence has not yet been proven, in particular the Dizmar district on the border between Armenia and Iran, the Mazandaran and Kerman regions in Iran (Hermann 1968) and the Sinai Peninsula (Nibbi 1981). These sources could be related to small local quarries now depleted or to extraction sites exploited in some historical periods and later forgotten. Some of these hypotheses are not simple to verify, taking into account that the suggested areas are often in regions of difficult access. It should also be noted that in ancient testimonies, some confusion might have been done both between lapis lazuli and other blue rocks, such as azurite or turquoise, and also between extraction sites of lapis lazuli and transit/trade zones, often object of misinterpretations. This is probably the case of the Sinai source, supposed to be exploited by Ancient Egyptians but not supported by geological data (Herrmann & Moore 1983; Aston 2000).

As a matter of fact, a systematic and exhaustive provenance study of the raw material utilised in works of art is nowadays still lacking. Despite that the origin from Badakhshan of the lapis lazuli used in antiquity remains the most plausible hypothesis, the variety of the proposed possible alternatives is worth to be studied by means of an analytical technique approach to confirm or disprove their historical reliability.

Materials and methods

Lapis lazuli rock and previous provenance studies

Lapis lazuli is generically classified as a metamorphic rock, even if this definition could not be considered exhaustive, due to the complexity of the mechanism involved in its genesis. It is characterised by the presence of the mineral lazurite (giving the blue colour to the rock), combined with other types of minerals whose presence and relative amount varies from and within quarries. Among these, there are hauyne, sodalite, nosean, K-feldspars, Na-feldspars, anorthite, mica group minerals like phlogopite, monoclinic pyroxenes like diopside, wollastonite, nepheline, titanite, apatite group minerals, calcite, dolomite and pyrite. Considering all the existent quarries, more than 30 mineral phases in lapis lazuli rocks have been reported so far in literature.

The question about the provenance of lapis lazuli used in ancient time was investigated by means of different kinds of analysis techniques, comparing the physico-chemical properties of rocks of known provenance with archaeological finds. An example are the studies carried out on the collection of the Museum of Mineralogy of Florence (Borelli 1986; Cipriani 1988) including both 22 rocks from 4 provenances (Badakhshan, Lake Baikal, Chile and Italy) and objects of the Medici Family (XV century AD). Moreover, another work was made by

Delmas and Casanova (Delmas and Casanova 1990; Casanova 1992). Using analytical techniques, they compared 21 rocks from 4 sources (Badakhshan, Pamir Mountains, Chagai Hills and Urals Mountains) with lapis lazuli waste products mainly found in the archaeological site of Shahr-i-Sokhta (III millennium BC) in the south-eastern part of Iran. Another investigation by means of microdestructive techniques was made by Ballirano (2006) who analysed pigments used in the Sistina Chapel by Michelangelo comparing the results with those obtained on 10 rocks from 4 provenances (Badakhshan, Chagai Hills, Pakistan and Iran). Moreover, Favaro (2012) analysed 9 rocks from 3 provenances (Afghanistan, Siberia and Chile) comparing results with ultramarine blue pigments. Finally, a very interesting work was recently made by Law (2014) that compared 18 rocks (Afghanistan, Chagai Hills, Myanmar and Lake Baikal) with 7 artefacts from 6 different archaeological sites in the Indo-Iranian area.

About the use of non-invasive techniques, the studies of Zöldföldi et al. (2006) carried out on 37 rocks from 4 provenances (Badakhshan, Lake Baikal, Chile, Ural Mountains) and Gambardella et al. (2016) on 34 rock samples from 7 sources (Afghanistan, Chile, Siberia, USA, Tajikistan, Iran and Myanmar) have to be cited. Finally, Calligaro (Calligaro 2011; Calligaro 2014) is the only one that studied artefacts from Ancient Egypt, comparing 11 objects from Töd treasure with 8 rocks from 4 provenances (the Badakhshan province of Afghanistan, the Pamir Mountains in Tajikistan, the borders of Lake Baikal in Russia and the Coquimbo region in Chile).

In many of these works, when comparisons between ancient artefacts and rocks were carried out, it was found a match not only with Afghanistan but also with Chagai Hills and Pamir and Ural Mountains. Although the results of these studies are innovative and interesting, it should be noted that in many cases, either the number of analysed rocks and/or objects is not enough to be representative. In these cases, the proposed markers and the provenance results should be confirmed extending the measurements over a larger number of samples, both rocks and artefacts. Moreover, the used techniques are sometimes invasive and not applicable to precious objects. Finally, it is to remember the difficulty in the availability of rocks of referenced provenance, also in museums collections. Consequently, the results showing a similarity among lapis lazuli from different sources, for example Chagai Hills and Afghanistan (Law 2014), could be attributed not to an effective exploitation of various quarries but instead to a wrong attribution of the provenance of the reference lapis lazuli rocks. For all these reasons, until now, the

studies in this field are not conclusive and the question of the determination of lapis lazuli origin by means of an analytical approach still presents some unresolved issues.

Protocol for lapis lazuli provenance determination

To solve the issue, we started in 2008 a long-term research, involving an interdisciplinary team. Until now, 45 lapis lazuli rocks of known provenance from 4 quarry districts have been analysed, creating a database that, to the best of our knowledge, is probably the widest in provenance studies on this material. Our database is composed by 21 samples from Badakhshan in Afghanistan; 4 samples from Liadjura-Dara, Pamir Mountains in Tajikistan; 11 samples from Lake Baikal area in Siberia (4 from Malaya Bistraya quarries and 7 from Sludyanka River quarries) and 9 samples from Coquimbo region in Chile. Samples from Tajikistan and Siberia are georeferenced, i.e. GPS coordinates are known. In the case of Lake Baikal, a geological expedition was organised by the authors to collect samples from two quarries in the area (Malaya Bistraya and Sludyanka River), considering that the mines are no longer exploited and it is difficult to find rocks of recorded provenance. More details about the analysed samples can be found in our previous studies (Lo Giudice 2009; Re 2011; Re 2013; Angelici et al. 2015; Re 2015).

Although obviously not suitable as a source of lapis lazuli for Ancient Egypt, the Chilean provenance was considered because the quarries are currently exploited and lapis lazuli samples from this provenance area are easy to find. Moreover, the results could be useful in future studies because there are hypotheses about utilisation of this rock by pre-Columbian civilisations (Da Cunha 1989).

It is interesting to observe that the study of lapis lazuli from all the known quarry districts and the identification of univocal markers can be useful to help in the identification of fakes and forgeries (Craddock 2009).


Due to the heterogeneity of lapis lazuli, instead of searching markers in the whole stone, we mostly focused our attention in searching minero-chemical markers inside single mineral phases. Since the crystal dimensions can vary from fraction of a micron to hundreds of microns, we decided to use mainly microscopic techniques (optical microscopy, scanning electron microscopy coupled with microanalysis, ion beam analyses and others, as explained in the section: experimental setup and measurements), which allow us to observe and to analyse one specific crystal, avoiding the contribution from the neighbouring ones. We looked for the presence or the absence of a mineral phase, minor/trace elements inside a peculiar mineral and the different luminescence of the same mineral. In particular for a high number of crystals, chemical analyses were carried out on two mineral phases that demonstrated to be suitable as markers: diopside

(CaMgSi₂O₆) and pyrite (FeS₂). The significant statistics (more than 94 diopside crystals and 80 pyrite crystals were analysed) for these mineral phases contributed to take into account the variability in trace elements also within the same sample. Even if other techniques have been used for these studies (i.e. optical and Raman microscopy, X-ray fluorescence, scanning electron microscopy with microanalysis, cathodoluminescence), it was verified that all the markers found are simultaneously detectable by means of ion beam techniques, in particular μ -proton induced x-ray emission (PIXE) and μ -ionoluminescence (IL) (Calusi 2008; Re 2011; Lo Giudice 2012). Besides, these techniques are non-invasive

and applicable in air, allowing to analyse artworks and rocks of practically any shape and dimension without sample preparation (Re et al. 2015). We also demonstrated the possibility to use μ -X-ray fluorescence (XRF) analysis to found markers based on trace elements with outcomes similar to those obtained by μ -PIXE (Angelici et al. 2015).

In Fig. 2, the protocol used to associate a lapis lazuli rock to one of the four provenances we studied until now is summarised. In the main block diagram, only “strong markers” were taken into account, which means physical-chemical features that are systematically present or absent in all the 45 analysed samples. Moreover, for Chilean and

Fig. 2 Schematic diagram of the protocol we proposed to identify the provenance of lapis lazuli by means of non-invasive techniques (in our case μ -PIXE and ionoluminescence)



Afghan samples, also other “weaker markers” are shown, which are features very frequent for these lapis lazuli quarry districts but not observed in all the samples.

The Chilean lapis lazuli are the easiest to distinguish due to the widespread presence of wollastonite (CaSiO_3), which was observed in all the analysed rock samples. Wollastonite can be recognised acquiring in a few seconds the ionoluminescence spectrum (Calusi 2008; Lo Giudice 2009; Re 2015; Czelusniak 2016). In fact, lapis lazuli from all provenances have a strong prevalent white-yellowish ionoluminescence, visible to the naked eyes, which is due to wollastonite in Chilean samples (IL spectra have a double broad bands at 560 and 620 nm) and to diopside in Asian samples (IL spectra centred at 580 nm). In addition, Chile is the only provenance in which selenium is present in amount higher than 40 ppm in pyrite crystals (Re 2013). However, this quantity is not always reached and this marker is to be considered weaker than the presence of wollastonite. Samples from Siberia are the only ones characterised by iron oxide-hydroxide minerals pseudomorphosed after pyrite, only sometimes occurring as relict, and a strontium content higher than about 180 ppm in diopside crystals (Re 2013; Re 2015). Instead, idiomorphic pyrite crystals are well preserved in lapis lazuli from Tajikistan and from Afghanistan. The trace element analysis on pyrite crystals seems to be the best way to distinguish Tajikistan from Afghanistan provenance (Re 2013). In fact, in samples from Afghanistan, pyrite crystals contain sensible contents of nickel (more than about 330 ppm) and low quantity of copper (below 240 ppm). Moreover, only diopside crystals of the Afghan samples show higher quantity of titanium (>710 ppm) or vanadium (>210 ppm) or chromium (>220 ppm) than do Tajik samples (Re 2015) or a strong ionoluminescence band at about 770 nm. It is to be stressed that although all the rock samples analysed until now have been successfully associated to one of the four sources of lapis lazuli by means of this protocol, increasing the number of analysed samples, all the quantities in ppm proposed as lower or upper limits for a provenance may have to be modified (due to the natural heterogeneity of each quarry district).

Recently, the first part of the protocol was successfully applied on precious artefacts from the Medici Collection kept in the Museum of Natural History of Florence (Re 2015).

Analysed objects from the Egyptian Museum of Florence

The 11 lapis lazuli carved objects analysed in this work belong to the collection of the Egyptian Museum of Florence. It is the second most important Egyptian museum in Italy, second only to the museum of Turin. The very first nucleus of antiquities was already settled in Florence during the eighteenth century. During the nineteenth century, this original core of pieces was implemented by other acquisitions; the promoter was Leopold II, Grand Duke of Tuscany, in the period from 1824 to 1859.

He bought some new collections of artworks and artefacts and together with Charles X of France, he funded an archaeological expedition in Egypt, the so called Franco-Tuscan expedition (1828–1829). The expedition was directed by the French Jean François Champollion and the Italian Ippolito Rosellini, considered pioneers of Egyptology. In 1882, the Egyptian Museum was officially established in the present seat and Ernesto Schiaparelli was appointed to direct the museum. Under his direction, the museum implemented the collections thanks to two excavation campaigns and numerous acquisitions until 1894, when the Egyptologist became the director of the Egyptian Museum of Turin. The collection in Florence has a great historical value and includes also several precious carved amulets made in lapis lazuli (pendants, small statues, scarabs and amulets), which are ascribable mainly to the first millennium BC, considering stylistic features and iconography.

The artefacts analysed in this work belong to different acquisitions (Tab. 1): two amulets are part of the finding of the Franco-Tuscan expedition (1828–1829), a scarab was acquired in the Theban Necropolis during the first Schiaparelli expedition (1884–1885) and other two objects come from the original Grand-Ducal Collections. A group of four objects results from purchases and private collector donations to the museum: Valle de Paz (1868, donation), Nardi (1885, purchase) and the more recent Wilson-Barker (1948, donation). The djed amulet does not belong to a definite acquisition, but certainly it is part of the finding of the Franco-Tuscan expedition or of the Grand-Ducal Collections. Finally, one of the wdjat eyes is of unknown acquisition.

In addition to the different acquisition modalities, the analysed objects were selected also on the basis of their iconography and archaeological significance (Guidotti 2015). Among the selected artefacts, there are two precious scarabs (cat. 1275 and cat. 6524). In Ancient Egypt, scarabs were used as seals or for funerary purposes on the mummies; this is the case of the Florence Museum objects. In particular, the first scarab (cat. 1275) has a small hole in the backside probably used for hanging or to sew it on the mummy, whereas the second scarab (cat. 6524) is a hearth scarab that was placed on the mummy in correspondence of the heart of the departed to make it lighter in the judgment of Osiris that every soul had to face in order to enter the afterlife. The second scarab presents a fragmented backside, probably because it was originally part of a pectoral. Another analysed funerary object is a djed pillar (cat. 1361); it represented solidity and durability and it was placed in many parts of the mummy, mainly on the throat. Moreover, three objects are a representation of gods. The first (cat. 179) is an amulet representing the Maat Goddess (Schiaparelli 1887), the personification of truth and justice and embodying the cosmic order. Maat amulets, almost always made in lapis lazuli, were placed mainly on the mummy chest. Also in this case, there is a small hole on the back.

Table 1 Objects from the Egyptian Museum of Florence analysed in this work

	<p>Catalogue number 64 Statue head: Hathor Goddess Dimension ~3 cm × 2.3 cm Period: Late (VII–IV century BC) Acquisition: Grand-Ducal Collections</p>		<p>Catalogue number 179 Amulet: Maat Goddess Dimension ~2.5 cm × 1.1 cm Period: Late (VII–IV century BC) Acquisition: Egypt, Franco-Tuscan Expedition 1828–1829</p>
	<p>Catalogue number 1275 Amulet: scarab Dimension ~2.1 cm × 1.5 cm Period: Late (VII–IV century BC) Acquisition: Grand-Ducal Collections</p>		<p>Catalogue number 1361 Amulet: djed pillar Dimension ~2.6 cm × 1 cm Period: unknown Acquisition: Egypt, Franco-Tuscan Expedition 1828–1829 or Grand-Ducal Collections</p>
	<p>Catalogue number 1600 Amulet: wdjat eye Dimension ~4.7 cm × 5.6 cm Period: 3apix Intermediate–Late (X–IV century BC) Acquisition: Egypt, unknown</p>		<p>Catalogue number 1644 Amulet: wdjat eye Dimension ~1.5 cm × 1.8 cm Period: 3th Intermediate–Late (X–IV century BC) Acquisition: Egypt, Franco-Tuscan Expedition 1828–1829</p>
	<p>Catalogue number 3914 Rectangular plate, wdjat eye Dimension ~1.4 cm × 1.7 cm Period: Late (VII–IV century BC) Acquisition: donation, Valle de Paz 1868</p>		<p>Catalogue number 6524 Hearth scarab Dimension ~5.3 cm × 3.4 cm Period: Late (VII–IV century BC) Acquisition: Egypt, Thebes (necropolis), first Schiaparelli Expedition 1884–1885</p>
	<p>Catalogue number 7298 Amulet: wdjat eye Dimension ~1.5 cm × 1.7 cm Period: New Kingdom–Late Period (XVI–IV century BC) Acquisition: purchase, Nardi 1885</p>		<p>Catalogue number 10284 Rectangular plate, wdjat eye Dimension ~2.3 cm × 2.4 cm Period: Late (VII–IV century BC) Acquisition: donation Wilson-Barker 1948</p>

Table 1 (continued)

Catalogue number 10316
 Amulet: Bes God
 Dimension ~2.4cm×1.2cm
 Period: New Kingdom–Late Period
 (XVI–IV century BC)
 Acquisition: donation Wilson-Barker
 1948

For each object, catalogue number, short description, dimensions, historical period and acquisition are indicated

The second (cat. 64) is a head of the Goddess Hathor that was originally part of a fairly rare small statue, both in terms of iconography and for the material quality. In fact, the lapis lazuli used appears different in terms of colour from that used for other artefacts. The third artefact (cat. 10316) is a stylized representation of Bes, God protector of households, pregnant women and infants. Also in this case, lapis lazuli appears to be different because it is poor in lazurite and considerable of lower quality. Finally, five wdjat eye amulets (cat. 1644, cat. 3914, cat. 7298, cat. 10284 and cat. 1600) were analysed. This kind of amulets was very common and widespread in Ancient Egypt and represents the eye of the God Horus, symbol of protection and integrity (Guidotti 2013).

Experimental setup and measurements


As previously observed, all the markers are detectable by means of Ion Beam Analysis, in particular μ -PIXE and μ -IL, non-invasive techniques applicable in air without sample preparation and that allow analysing artworks and rocks of almost any shape and dimension.

Ion Beam Analyses were performed at two external scanning microbeam facilities: the INFN LABEC Laboratory in

Florence (Giuntini 2015a, b; Giuntini 2007; Calusi 2011) and the AGLAE facility at the C2RMF in Paris (Calligaro 2000; Pichon 2014). In both cases, μ -PIXE and μ -IL analyses on Ancient Egyptian objects were performed in air, without any preparation and using 3 MeV proton beam (Fig. 3). At INFN LABEC laboratory, the beam was about 20 μ m wide and a 150 pA current was used on selected areas of 1×1 mm². At AGLAE, the beam was about 30 μ m wide and a current of about 500 pA was used on selected areas of 1.5×1.5 mm².

As required by the described protocol for lapis lazuli attribution, elemental quantitative analyses by means of μ -PIXE were carried out on pyrite and diopside crystals. To find these minerals and considering the impossibility to perform preliminary invasive analysis on artefacts, the first step was a survey of all the samples by means of optical imaging and microscopy. This approach was sufficient to identify candidate pyrite crystals and white areas characteristic of diopside. Concerning pyrite crystals, μ -PIXE mapping was performed to verify the presence of iron and sulphur as main elements and to select the more homogeneous part of the crystals for quantitative analysis, as shown in Fig. 4. Instead, diopside crystals were found not only by means of μ -PIXE mapping but also by exploiting the strong luminescence properties of this mineral that exhibit

Fig. 3 Measurement setup at the AGLAE (C2RMF in Paris) and the LABEC (INFN in Florence) laboratories during analysis of the hearth scarab cat. 6524 and the wdjat eye cat. 1600 respectively



a strong emission at about 580 nm. In particular, the broad-beam IL microscopy apparatus was used at INFN LABEC (Lo Giudice 2012; Re 2015) and μ -IL mapping at AGLAE (Pichon 2015a). Also in this case, μ -PIXE maps were used to select the more homogeneous part of the crystals for quantitative analysis. Moreover, to avoid possible systematic errors and to ensure the reliability of the results, during all measurement sessions, a number of previously analysed pyrite and diopside crystals in lapis lazuli from Afghanistan and Tajikistan rocks were measured and used as references.

The μ -PIXE spectra obtained at INFN LABEC were quantitatively analysed by means of GUPIXWIN (version 2.2.3) (Campbell 2010) using a set of reference standards: a soda-lime container from NIST and thick pure elements from Goodfellow Cambridge Ltd. Instead, at AGLAE, the elemental quantification was subsequently obtained using the TRAUPIXE program (Pichon 2015b) with the pivot procedure normalisation between major and trace concentrations. This program is based upon the same GUPIXWIN version with processing parameters adjusted using the Dr-N (Diorite) geostandard.


Results and discussion

In accordance with the protocol summarised in Fig. 2, the first step for all the samples was the identification of wollastonite or diopside as one of the main mineral phases. As expected, all Egyptian objects show a prevalent ionoluminescence centred at 580 nm, a feature characteristic of diopside that is present in all the studied rocks of Asian provenance (Calusi 2008; Lo Giudice 2009; Re 2015). Figure 5 shows some of the IL

spectra collected in correspondence of strong luminescent crystals, which were confirmed to be diopside by means of μ -PIXE elemental analysis. The absence of wollastonite excludes the Chilean provenance for all samples, a result that was expected (due to the object typology and their historical period) but nonetheless important, as it is a further proof of the reliability of the method to recognise Chilean samples.

The second step of the protocol was the measurement of strontium contents in the diopside crystals and the composition of iron-based inclusions. In fact, for all of the Siberian samples, i.e. from the Lake Baikal region both from Sludyanka River and Malaya Bistraya River districts, pyrite crystals present an altered form. In particular, they are characterised by the presence of iron oxide-hydroxide minerals (probable alteration products of pyrite), presenting an intensive compositional zoning in which sulphur-free zones prevail (Re 2013; Angelici 2015). Only in rare cases, very small pyrite crystals (below about 30 μ m) show unaltered aspect with neither oxidised margins nor core zones. For the Egyptian objects studied in this work, the majority of the mapped pyrite crystals (more than 20) are well preserved and stoichiometrically corresponding to FeS_2 , a feature characteristic of Tajikistan and Afghanistan rocks; in many cases, as in Fig. 4, crystals are very large, up to 0.5 mm^2 , and are easily visible to the naked eyes. In Siberian rocks, such features were never observed. It is interesting to observe that various pyrite crystals in Egyptian artefacts have a certain degree of chemical weathering: this is an oxidation process of pyrite where the first reaction products going in solution are ferrous iron and sulphate (Blowes 2003). The process is a superficial one, in accordance with the analysed objects, probably subject to burial, percolating waters and humidity. In

Fig. 4 Procedure for μ -PIXE analysis of pyrite crystals (hearth scarab Cat. 6524): **a**, **b** identification of a candidate pyrite crystal by means of optical imaging and microscopy, **c** μ -PIXE maps of some elements in the selected area (about 1 mm \times 1 mm). μ -PIXE spectra for quantitative analysis were collected in the more homogeneous part of the pyrite (white box)






Fig. 5 Some of the IL spectra of diopside crystals acquired from the analysed samples. A representative spectrum is shown for each object. The main feature is the characteristic peak centred at 580 nm, common to all the spectra

Fig. 4 are visible small areas and veins inside the pyrite crystal that show a loss of sulphur due to this kind of degradation. The level of degradation is heterogeneous: in fact, in the same object are present both unaltered pyrite crystals and others almost completely altered. A microscopy image of a strong chemical weathered pyrite crystal in an Egyptian artefact (sample cat. 10284) is shown in Fig. 6. It is very similar to what was observed in a previous work on Töd treasure (Calligaro 2014), where oxidation turned pyrite in brown/red-stained grains, involving adjacent areas in the process. In the case of Töd treasure, this phenomenon was mainly observed on the surface of the artefacts. A limited or absent chemical weathering was instead observed in the inner part of the cut samples. On the contrary, finding altered pyrite crystals is a feature of all the Siberian rocks, observed even in fresh cut samples, and it is related to hydrothermal processes during the petrogenesis (Re 2013; Angelici 2015).

Also, the second strong marker excluding the Siberian provenance confirms the initial attribution. Actually, the strontium content in diopside crystals of all Siberian rocks




Fig. 6 Optical image of a strong chemical weathered pyrite crystal in an Ancient Egyptian artefact (cat. 10284)

Table 2 Trace elements (expressed in ppm) in diopside crystals in lapis lazuli Egyptian artefacts; each line is one point of analysis (i.e. one diopside crystal) and when an element is under the limit of detection (LOD), it is indicated with <LOD

Cat	Object	Ti	V	Cr	Mn	Sr	Fe
64	Statue head: Hathor Goddess	60 ± 20 60 ± 20 30 ± 10	90 ± 10 110 ± 10 90 ± 10	< 15 < 13 < 12	280 ± 10 320 ± 10 290 ± 10	40 ± 20 18 ± 10 30 ± 10	380 ± 10 410 ± 20 180 ± 10
179	Amulet: Maat Goddess	1200 ± 60	330 ± 40	140 ± 20	1140 ± 40	80 ± 50	540 ± 30
1275	Amulet: scarab	2290 ± 60 2430 ± 80	190 ± 30 210 ± 30	100 ± 20 70 ± 10	920 ± 30 960 ± 40	200 ± 50 90 ± 30	870 ± 30 380 ± 20
1644	Amulet: wdjat eye	2680 ± 130 2600 ± 80 2580 ± 80	220 ± 40 210 ± 30 210 ± 30	110 ± 20 90 ± 10 110 ± 20	840 ± 50 810 ± 30 750 ± 30	190 ± 70 60 ± 20 110 ± 40	320 ± 30 340 ± 20 3210 ± 80
3914	Amulet: wdjat eye	1360 ± 40 2900 ± 70	180 ± 20 170 ± 30	120 ± 10 110 ± 20	960 ± 30 780 ± 30	160 ± 40 100 ± 30	470 ± 20 2670 ± 40
7298	Amulet: wdjat eye	2340 ± 50 2350 ± 70	250 ± 30 200 ± 20	80 ± 10	780 ± 20 610 ± 20	110 ± 20	780 ± 40 270 ± 10
10,284	Amulet: wdjat eye	1900 ± 60 1090 ± 40	200 ± 20 200 ± 20	80 ± 10 50 ± 10	640 ± 20 1790 ± 50	70 ± 20 90 ± 20	330 ± 20 810 ± 30
10,316	Amulet: Bes God	1000 ± 40 300 ± 20 340 ± 20	240 ± 20 120 ± 10 90 ± 10	90 ± 10 <13 < 11	1810 ± 40 230 ± 10 290 ± 10	90 ± 20 120 ± 20 40 ± 10	1440 ± 40 610 ± 10 820 ± 20

in the database were always higher than 160 ± 40 ppm, with a maximum of 870 ± 30 ppm and a median of the distribution at about 420 ppm (Re 2015), while the maximum strontium content measured in diopside crystals was 90 ± 40 and 140 ± 20 ppm in Afghan and Tajik rocks respectively. Diopside crystals in Egyptian objects suitable for trace element analysis were identified by means of ionoluminescence maps and on the basis of stoichiometry in order to avoid spurious signal from other phases, as we limited the analyses only to the crystals for which the stoichiometry was coincident within experimental errors to what was expected for this mineral phase. In particular, 18 points of interest were selected on 8 out of 11 Egyptian objects; the measured strontium and other trace element concentrations useful for provenances study are reported in Table 2. When possible, for every object, more than one diopside crystal was analysed to take into account the variability in trace elements within the same sample. In Fig. 7 are also shown the box charts of the same values of Table 2 in comparison with rocks from our database (Re 2015). The box charts represent the dispersion of the

experimental points: each point is an individual μ -PIXE measurement, the boxes include 50% of analysed points (from 25 to 75%) and are separated in two parts by the median line, while the bars join the maximum and the minimum detected values. In this representation (called box and whisker plot), error bars are not reported to make the graphics more readable. The square symbols in correspondence to the 0 value represent the numbers of measurements below the limit of detection (LOD). As shown in Table 2 and Figs. 7, 5 out of the 8 analysed Egyptian objects show a Sr content in diopside in the same range of Afghan and Tajikistan rocks, all other objects being also compatible within 2σ . Such results seem to exclude a Siberian provenance of the Egyptian artefacts, although it should be observed that the data distribution is slightly different from that expected for Afghan and Tajik samples. In particular, there are 4 analysed diopside crystals (i.e. 20% of analysed points) that have Sr contents between 160 and 200 ppm, slightly higher than those observed for these two provenances. Discrepancies could be due to the limited number of Afghan reference rocks in our database.




Fig. 7 Trace element contents in diopside crystals (in ppm): comparison between Egyptian artefacts and rocks from our database (Re 2015). The square symbols in correspondence to 0 value represent the numbers of measurements below the limit of detection (LOD)

Table 3 Trace elements (expressed in ppm) in pyrite crystals in lapis lazuli Egyptian artefacts; each line is one point of analysis (i.e. one pyrite crystal) and when an element is under the limit of detection (LOD) it is indicated with <LOD


Cat.	Object	Ni	Cu
1275	Amulet: scarab	790 ± 50	110 ± 30
		740 ± 50	130 ± 30
1361	Amulet: djed pillar	600 ± 30	120 ± 20
1600	Amulet: wdjat eye	1000 ± 200	<35
		970 ± 110	80 ± 20
1644	Amulet: wdjat eye	660 ± 50	140 ± 30
		600 ± 160	100 ± 30
3914	Amulet: wdjat eye	690 ± 40	100 ± 20
		650 ± 60	150 ± 40
6524	Hearth scarab	620 ± 20	100 ± 10
		680 ± 30	150 ± 10
		590 ± 120	150 ± 20
		560 ± 110	< 30

Although 21 samples are a sufficient starting point to attribute raw materials to a provenance, in the future, the analysis of further rocks could broaden the boundary values found so far because of the natural heterogeneity of the extraction sites. Moreover, it should be noted that all the samples used for our database were collected in the last century and although the proposed markers are characteristic for all the rocks of a district, small differences for quarries in the same area were observed. These differences are not discussed here but could be useful in future studies to distinguish among extraction sites in the same district and perhaps to identify the location of ancient quarries. About the other trace elements interesting for provenance study in diopside crystals (Re 2015), it must be observed that almost all the 90 values in Table 2 are in the range expected for Afghan and Tajik rocks and only 2 data related to manganese are outside (which means about 2.2% of the total). Moreover, it is interesting to observe that although the head of the Hathor Goddess (cat. 64) and the Bes God amulet (cat. 10316) are compatible with an Afghan or Tajik

provenance, they seem to be slightly different from other objects due to the very low contents of all trace elements in diopside crystals. As in the case of strontium, this could be explained with a difference in the extraction site inside the same district. As already stated, the lapis lazuli used for these artefacts appears different in terms of colour from those used for the other artefacts; in particular, the Bes God amulet (cat. 10316) is poor in lazurite and with a considerable low quality.

The third step of the protocol was the analysis of trace elements in pyrite crystals, in particular nickel and copper, which appear the most effective fingerprint to distinguish Afghan samples from Tajik samples. In Table 3, a summary of the results on 13 pyrite crystals selected on 6 Egyptian objects is reported. When possible, for each object, more than one pyrite crystal was analysed to take into account the variability of trace elements within the same sample. In the other 5 objects, the analysed pyrite crystals were not stoichiometric, due to chemical weathering; therefore, they were not suitable for a reliable quantitative analysis. In Fig. 8, the box charts of

Fig. 8 Ni and Cu contents in pyrite crystals (in ppm): comparison between Egyptian artefacts and rocks from our database (Re 2013, Angelici 2015). The square symbols in correspondence to 0 value represent the numbers of measurements below the limit of detection (LOD), which is around 100 ppm for Ni and 50 ppm for Cu






Fig. 9 Cu versus Ni contents in pyrite crystals from μ -PIXE measurements. The individual errors associated with each measurement are plotted, whereas the values below the limit of detection (LOD around 100 ppm for Ni and 50 ppm for Cu) are represented on the axes

Cu and Ni show a good match of Egyptian artefacts and Afghan rocks from our database (Re 2013; Angelici 2015). To better observe the differences between Afghan and Tajik rocks and the results on Egyptian artefacts, Cu versus Ni contents in pyrite crystals are shown in Fig. 9 (values below the LOD, which is around 100 ppm for Ni and 50 ppm for Cu, are represented on the axes). Moreover, as expected, in all the pyrite crystals, selenium was below LOD, which is a further proof of the validity of the method to recognise Chilean samples.

Finally in 6 Egyptian artefacts, also the marker related to titanium contents in diopside is compatible with an Afghan provenance of the lapis lazuli, i.e. greater than 710 ppm. It must be observed that almost all the 90 values in Table 2 are in the range expected for Afghan rocks and only 4 data related to manganese and iron are outside (which means about <5% of the total).

Summarising, on the basis of the results, the raw material used to carve 9 of the 11 lapis lazuli artefacts analysed in this work can be ascribed to an Afghan provenance. For the other 2 objects (cat. 10316 and cat. 64), a Siberian provenance is excluded. They are compatible with an Afghan or Tajik provenance, but it is not possible to distinguish among them because the analysis on the pyrite crystals were not carried out and the measurements of trace element contents in diopside crystals was not sufficient alone.

Conclusions

In this work, a protocol for determining the provenance of raw material used to carve lapis lazuli artefacts is described. Markers were obtained analysing 45 rocks of known provenance from 4 quarry districts: Badakhshan in Afghanistan,

Liadjura-Dara in Tajikistan, Lake Baikal area in Siberia and Coquimbo region in Chile. To the best of our knowledge, for every provenance, the number of studied rocks is equal or higher than in any previous provenance study on lapis lazuli. The created database constitutes a sufficient starting point to attribute raw materials to a provenance, although in the future, its representativeness must be further augmented to increase the marker reliability or to revise them. Moreover, the database must be extended to rocks from other quarry districts. An important point is that all the markers are recognisable by means of Ion Beam Analysis. These techniques are non-invasive and applicable in air, allowing to analyse precious artworks and rocks of almost any shape and dimension without sample preparation.

The protocol was applied to determine the provenance of raw material used for 11 carved lapis lazuli artefacts kept at the Egyptian Museum of Florence. Analysed objects are scarabs, amulets and small statues, mainly ascribable to the first millennium BC, considering stylistic features and iconography. For 9 objects, an Afghan provenance was deduced. The other 2 artefacts are compatible both with Afghan and Tajik provenances, and it is not possible to distinguish between them, because the analysis on pyrite crystals was not yet carried out. It is interesting to observe some differences in distribution of trace element contents in diopside crystals in comparison to referenced rocks. This information could be related to the natural heterogeneity of the extraction sites, and it could be useful in future studies to distinguish among different quarries in the same district.

Although the results of this work must be strengthened by increasing the number and the archaeological representativeness of the analysed artefacts, the scientific proof of the presence of Afghan lapis lazuli in Ancient Egypt is a further confirmation of trading of this stone up to 4000 km from its extraction site in the first millennium BC, excluding the possibility of other local sources exploited in ancient time, as it has long been supposed by the majority of archaeologists.

Acknowledgements Financial support by the Transnational Access to Research Infrastructures activity in the Seventh Framework Program of the EU (CHARISMA Grant Agreement n. 228330) is gratefully acknowledged. The authors warmly thank all the people participating and contributing to this provenance study during the last years.

The authors wish to warmly thank CHNet, the INFN network of laboratories working in the Cultural Heritage field, for supporting this research, all in terms of instrumentation, competencies and grants. Moreover, the INFN experiment CHNet-imaging is acknowledged with gratitude.

References

- Angelici D, Borghi A, Chiarelli F, Cossio R, Gariani G, Lo Giudice A, Re A, Pratesi G, Vaggelli G (2015) μ -XRF analysis of trace elements in lapis lazuli-forming minerals for a provenance study. *Microsc Microanal* 21:526–533

- Aston B, Harrell JA, Shaw I (2000) Lapis lazuli. In: Nicholson PT, Shaw I (eds) *Ancient Egyptian materials and technology*. Cambridge University Press, Cambridge, pp. 39–40
- Ballirano P, Maras A (2006) Mineralogical characterization of the blue pigment of Michelangelo's fresco 'the last judgment'. *Am Mineral* 91:997–1005
- Blowes D.W., Ptacek C.J., Jambor J.L., Weisener C.G. (2003) The geochemistry of acid mine drainage. *Treatise on Geochemistry* 9–9, 149–204, Ed. Holland H.D., University of Harvard, USA
- Borelli A, Cipriani C, Innocenti C, Trosti R (1986) Caratterizzazione del lapislazzuli. *La Gemmologia* XI:24–27
- Calligaro T, Dran JC, Ioannidou E, Moignard B, Pichon L, Salomon J (2000) Development of an external beam nuclear microprobe on the AGLAE facility of the Louvre museum. *Nucl Instrum Meth B* 161–163:328–333
- Calligaro T, Coquinot Y, Pichon L, Moignard B (2011) Advances in elemental imaging of rocks using the AGLAE external microbeam. *Nucl Instrum Meth B* 269(20):2364–2372
- Calligaro T, Coquinot Y, Pichon L, Pierrat-Bonnefois G, de Campos P, Re A, Angelici D (2014) Characterization of the lapis lazuli from the Egyptian treasure of Tod and its alteration using external μ -PIXE and μ -IBIL. *Nucl Instrum Meth B* 318:139–144
- Calusi S (2011) The external ion microbeam of the LABEC laboratory in Florence: some applications to cultural heritage. *Microsc Microanal* 17(5):661–666
- Calusi S, Colombo E, Giuntini L, Lo Giudice A, Manfredotti C, Massi M, Pratesi G, Vittono E (2008) The new ionoluminescence apparatus at the LABEC external microbeam facility. *Nucl Instrum Meth B* 266:2306–2310
- Campbell JL, Boyd NI, Grassi N, Bonnicksen P, Maxwell JA (2010) The Guelph PIXE software package IV. *Nucl Instrum Meth B* 268:3356–3363
- Casanova M (1992) The sources of the lapis lazuli found in Iran. In: *South Asian archaeology, monographs in world archaeology*. Prehistory Press 14:49–56
- Casanova M (2013) *Le lapis-lazuli dans l'Orient Ancien: production et circulation du Néolithique au IIe millénaire av. J.-C.* Editions du Comité des travaux historiques et scientifiques, Paris
- Cipriani C, Innocenti C, Trosti-Ferroni R (1988) Le collezioni del Museo di Mineralogia di Firenze. (VI) I Lapislazzuli *Museologia Scientifica* V(1–2):17–30
- Craddock PT (2009). *Scientific investigation of copies, fakes and forgeries*. Butterworth-Heinemann, Oxford, UK, pp 412
- Czelusniak C, Palla L, Massi M, Carraresi L, Giuntini L, Re A, Lo Giudice A, Pratesi G, Mazzinghi A, Ruberto C, Castelli L, Fedi ME, Liccioli L, Gueli A, Mandò PA, Taccetti F (2016) Preliminary results on time-resolved ion beam induced luminescence applied to the provenance study of lapis lazuli. *Nucl Instrum Meth B* 371:336–339
- Da Cunha C (1989). *Le lapis lazuli: son histoire, ses gisements, ses imitations*. Editions du Rocher, Paris
- Delmas AB, Casanova M (1990) The lapis lazuli sources in the ancient east. In: Maurizio Taddei (ed) *South Asian archaeology*. Istituto Italiano per il Medio ed Estremo Oriente, Roma, Serie Orientale Roma 66(1):493–505
- Favaro M, Guastoni A, Marini F, Bianchin S, Gambirasi A (2012) Characterization of lapis lazuli and corresponding purified pigments for a provenance study of ultramarine pigments used in works of art. *Anal Bioanal Chem* 402:2195–2208
- Gambardella AA, Schmidt Patterson CM, Webb SM, Walton MS (2016) Sulfur K-edge XANES of lazurite: toward determining the provenance of lapis lazuli. *Microchem J* 125:299–307
- Giuntini L, Massi M, Calusi S (2007) The external scanning proton microprobe of Firenze: a comprehensive description. *Nucl Instrum Meth A* 576(2–3):266–273
- Giuntini L, Massi M, Calusi S, Castelli L, Carraresi L, Fedi ME, Gelli N, Liccioli L, Mandò PA, Mazzinghi A, Palla L, Romano FP, Ruberto C, Taccetti F (2015a) Wide area scanning system and carbon microbeams at the external microbeam facility of the INFN LABEC laboratory in Florence. *Nucl Instrum Meth B* 348:14–17
- Giuntini L, Massi M, Calusi S, Gelli N, Castelli L, Carraresi L, Czelusniak C, Fedi ME, Gueli AM, Liccioli L, Mandò PA, Mazzinghi A, Palla L, Ruberto C, Taccetti F (2015b) The set-up for forward scattered particle detection at the external microbeam facility of the INFN-LABEC laboratory in Florence. *Nucl Instrum Meth B* 348:8–13
- Guidotti MC (2013) *La vita oltre la morte. Il corredo funerario nell'antico Egitto*, Pisa, p.17, n.11
- Guidotti MC (2015) Schede. In: *Lapislazzuli, magia del blu*. Sillabe, Livorno, pp. 302–306
- Hermann G (1968) Lapis lazuli: the early phases of its trade. *Iraq* 30(1):21–57
- Herrmann G, Moorey PRS (1983) Lapis lazuli. *Reallexikon der Assyriologie und Vorderasiatischen Archäologie* 6:489–492
- Jarrige JF (1988) *Les cités oubliées de l'Indus: archéologie du Pakistan*. Association Française d'Action Artistique, Paris, pp 28
- Jarrige JF (1991) Mehrgarh: its place in the development of ancient cultures in Pakistan. In: Janssen M, Mulloy M and Urban G (eds) *Forgotten cities on the Indus*. Verlag Philipp von Zabern, Mainz am Rhein, pp 41
- Khalik A, Ahmad J, Ahmad A (2006) Reporting of gold anomaly and showings of lapis-lazuli from Chakdara granite gneiss and its environs, district Dir, northern Pakistan. *Journal of Himalayan Earth Sciences* 39:85–87
- Law RW (2008) Inter-regional interaction and urbanism in the ancient Indus valley: a geologic provenance study of Harappa's rock and mineral assemblage. University of Wisconsin, Madison
- Law R (2014) Evaluating potential lapis lazuli sources for ancient South Asia using sulfur isotope analysis. In: Lamberger-Karlovsky CC, Genito B (eds) "My life is like the summer rose" Maurizio Tosi e l'archeologia come modo di vivere. Archaeopress, Oxford, pp. 419–429
- Lo Giudice A, Re A, Calusi S, Giuntini L, Massi M, Olivero P, Pratesi G, Albonico M, Conz E (2009) Multitechnique characterization of lapis lazuli for provenance study. *Anal Bioanal Chem* 395(7):2211–2217
- Lo Giudice A, Re A, Angelici D, Calusi S, Gelli N, Giuntini L, Massi M, Pratesi M (2012) In-air broad beam ionoluminescence microscopy as a tool for rocks and stone artworks characterisation. *Anal Bioanal Chem* 404(1):277–281
- Mindat Web Site: <http://www.mindat.org/min-2330.html>
- Nibbi A (1981) *Ancient Egypt and some eastern neighbours*, Noyes Publication, Park Ridge NJ, pp 33–55
- Pichon L, Moignard B, Lemasson Q, Pacheco C, Walter P (2014) Development of a multi-detector and a systematic imaging system on the AGLAE external beam. *Nucl Instrum Meth B* 318(A):27–31
- Pichon L, Calligaro T, Gonzalez V, Lemasson Q, Moignard B, Pacheco C (2015a) Implementation of ionoluminescence in the AGLAE scanning external microprobe. *Nucl Instrum Meth B* 348:68–72
- Pichon L, Calligaro T, Lemasson Q, Moignard B, Pacheco C (2015b) Programs for visualization, handling and quantification of PIXE maps at the AGLAE facility. *Nucl Instrum Meth B* 363:48–54
- Pierrat-Bonnefois G (1995) *Le lapis lazuli du trésor de Tôd*. In: *Les pierres précieuses de l'Orient ancien*, Dossiers du musée du Louvre. Éditions de la Réunion des musées nationaux, Paris
- Re A, Lo Giudice A, Angelici D, Calusi S, Giuntini L, Massi M, Pratesi G (2011) Lapis lazuli provenance study by means of micro-PIXE. *Nucl Instr Meth B* 269(20):2373–2377
- Re A, Angelici D, Lo Giudice A, Maupas E, Giuntini L, Calusi S, Gelli N, Massi M, Borghi A, Gallo LM, Pratesi G, Mandò PA (2013) New markers to identify the provenance of lapis lazuli: trace elements in pyrite by means of micro-PIXE. *Appl Phys A-Mater* 111:69–74
- Re A, Angelici D, Lo Giudice A, Corsi J, Allegretti S, Biondi AF, Gariani G, Calusi S, Gelli N, Giuntini L, Massi M, Taccetti F, La Torre L, Rigato V, Pratesi G (2015) Ion beam analysis for the provenance attribution of lapis lazuli used in glyptic art: the case of the "Collezione Medicea". *Nucl Instrum Meth B* 348:278–284

- Schiaparelli E (1887) Museo Archeologico di Firenze—Antichità egizie. Tipografia della R. Accademia dei Lincei, Roma, pp 104, n.842
- Shafer EH (1963) The golden peaches of Samarkand: a study of T'Ang exotics. University of California Press, Los Angeles, pp 230–234
- Sofianides AS, Harlow GE (1990) Gems and crystals from the American Museum of Natural History. Simon and Schuster, New York
- Tosi M (1974) The lapis lazuli trade across the Iranian plateau in the 3rd millennium BC. In: Gururājamañjarikā. Studi in onore di G. Tucci. Istituto Universitario Orientale, Napoli, pp. 3–22
- Von Rosen L (1988) Lapis lazuli in geological contexts and in ancient written source. *Studies in Mediterranean Archaeology and Literature*, 65; ed. by Paul Astroms forlag, Partille, Sweden
- Zöldföldi J, Richter S, Kasztovszky Z, Mihály J (2006) Where does lapis lazuli come from? Non-destructive provenance analysis by PGAA. In: *Proceeding of the 34th international symposium on Archaeometry*. Zaragoza, Spain, pp. 353–360