

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Variance components using genomic information for 2 functional traits in Italian Simmental cattle: Calving interval and lactation persistency

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1744139> since 2020-07-16T12:44:19Z

Published version:

DOI:10.3168/jds.2019-17421

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

*Alberto Cesarani, **Giustino Gaspa**, Yutaka Masuda, Lorenzo Degano, Daniele Vicario, Daniela A.L. Lourenco, Nicolò P.P. Macciotta*

Variance components using genomic information for two functional traits in Italian Simmental cattle: calving interval and lactation persistency

Journal of Dairy Science

Volume 103, Issue 6, June 2020, Pages 5227-5233

<https://doi.org/10.3168/jds.2019-17421>

The publisher's version is available at:

<https://www.sciencedirect.com/science/article/pii/S0022030220302551?via%3Dihub>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/ID:431290>

1 **Variance components using genomic information for two functional traits in Italian Simmental**
2 **cattle: calving interval and lactation persistency.** *by Cesarani et al.* Modern breeding programs
3 aim to improve functional traits, such as fertility and/or lactation persistency. Calving interval is an
4 **easily accessible** indicator of female fertility, but is lowly heritable and has moderate, unfavorable
5 correlations with yield traits. Lactation persistency is also lowly heritable. **We investigated the genetic**
6 **background of calving interval and lactation persistency and estimated the genetic parameters that**
7 **are needed to include both traits and milk production for the Italian Simmental breeding program.**
8 **There is a potential to implement a single step-genomic prediction model by using both genomic and**
9 **pedigree information.**

11 VARIANCE COMPONENTS OF FUNCTIONAL TRAITS

13 **Variance components using genomic information for two functional traits in Italian**
14 **Simmental cattle: calving interval and lactation persistency**

16 Alberto Cesarani^{14*}, Giustino Gaspa³, Yutaka Masuda⁴, Lorenzo Degano², Daniele Vicario²,
17 Daniela A.L. Lourenco⁴, Nicolò P.P. Macciotta¹

19 ¹ Department of Agricultural Sciences, University of Sassari, Sassari, Italy

20 ² Associazione Nazionale Allevatori Pezzata Rossa Italiana (ANAPRI), Udine

21 ³ Department of Agricultural, Forestry and Alimentary Sciences, University of Torino, Grugliasco

22 ⁴ Department of Animal and Dairy Science, University of Georgia, Athens, Georgia

24 * Corresponding author: acesarani@uniss.it

27

28 **ABSTRACT**

29 Functional traits, such as fertility and lactation persistency, are becoming relevant breeding
30 goals for dairy cattle. Fertility is a key element for herd profitability and animal welfare: in particular,
31 calving interval (CI) is an indicator of female fertility that can be easily recorded. Lactation
32 persistency (LPE), i.e. the ability of a cow to maintain a high milk yield after the lactation peak, is
33 economically important and it is related to several other traits, such as feed efficiency, health, and
34 reproduction. The selection of these functional traits is constrained by their low heritability. In this
35 study, variance components for CI and LPE in the Italian Simmental cattle breed were estimated
36 using genomic and pedigree information under the single-step genomic framework. A data set of
37 594,257 calving interval records (from 275,399 cows), 285,213 lactation persistency records (from
38 1563,389 cows) was considered. Phenotypes were limited up to the third parity. The pedigree
39 contained about 2 million animals, and 7,246 genotypes were available. LPE was estimated using
40 principal component analysis on test day records, with higher PC2 values associated to lower LPE
41 and lower PC2 values to higher LPE, respectively. Heritability of CI and LPE were estimated using
42 single-trait repeatability models. A multiple trait analysis using CI and production traits (milk, fat,
43 and protein yields) was performed to estimate genetic correlations among these traits. Heritability for
44 CI in the single-trait model was low (0.06 ± 0.002). Unfavorable genetic correlations were found
45 between CI and production traits. A measure of LPE was derived using principal component analysis
46 on test-day records. The heritability and repeatability of LPE were 0.11 ± 0.004 and 0.20 ± 0.02 ,
47 respectively. Genetic correlation between CI and LPE was weak but had a favorable direction.
48 Despite the low heritability estimates, results of the present work suggest the possibility of including
49 these traits in the Italian Simmental breeding program. The use of single-step approach can potentially
50 provide better results for young genotyped animals without own phenotypes.

51

52 **Keywords:** fertility, persistency, genomic selection

53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78

INTRODUCTION

In the last century, the combined action of genetic, management, and feeding advancements has led to a substantial improvement of dairy cow performances. Such results have been, however, accompanied by a general deterioration in fitness traits (Sun et al., 2019). The progressive reduction of cow fertility is one of the most relevant examples of the unfavorable consequences of selection for production traits (Ma et al., 2019). Relationships between fertility and genetic improvement for milk production have been investigated (Castillo-Juarez et al., 2000; Lucy, 2001; Hayes et al., 2009; Walsh et al., 2011). Reproduction traits have been, therefore, included in breeding programs (Philipsson and Lindhé, 2003; López-Gatius, 2003; Miglior et al., 2005) even though a clear definition of phenotypes is still an issue. For example, fertility traits can be defined in many ways: time lengths (i.e. days open), or frequencies (i.e. number of inseminations needed to conceive). Generally, these traits are difficult to record routinely and have low heritability. Among fertility traits, calving interval (CI) is one of the easiest to record (Dal Zotto et al., 2007); however, it can be biased for not being available for cows that do not re-calve because of poor fertility. Other traits like conception or pregnancy rates could be better fertility traits, but they need additional resources to be collected. A longer CI is usually a consequence of more inseminations needed for the cow to conceive, with increased costs for the herd, and therefore, is undesirable. Moreover, cows with longer CI will have fewer calvings in their life. Selection for shorter CI would increase herd profitability: more calvings per cow, more offspring that can be sold or used as replacement, decreased feed costs, and less reproductive problems (Esslemont et al., 2001; Groenendaal et al., 2004; Atashi et al., 2013).

Reproductive performance is also strongly related to lactation persistency (LPE), i.e. the ability of the cow to maintain high levels of production after the lactation peak (Dekkers et al., 1998). LPE is a production trait connected to health, reproduction, and feed costs (Koloi et al., 2018). The relationship between CI and LPE in cattle has been investigated (Atashi et al., 2013; Němečková et al., 2015). Muir et al. (2004) reported a moderate, positive genetic correlation between CI and LPE

79 suggesting an unfavorable and antagonistic relationship between these two traits. This result was
80 confirmed by Atashi et al. (2013), who found that cows with shorter CI were less persistent in milk
81 production. Although there is a general consensus on the basic concept of LPE, generally defined as
82 the ratio between the milk measured in a certain period and the total milk yield, several measurements
83 have been proposed for this trait (Gengler, 1996; Sölkner and Fuchs, 1987; Cole and VanRaden,
84 2006; Strabel and Jamrozik, 2006; Togashi and Lin, 2006). Macciotta et al. (2006) have defined the
85 LPE in Italian Simmental using a principal component (PC) approach. The advantage of this approach
86 is that the measure of LPE is independent of milk production.

87 The Italian Simmental is a dual-purpose cattle breed farmed mostly in Northeastern Italy. Its
88 breeding program aims to improve both dairy and beef traits, and an economic selection index has
89 been developed for this purpose. Recently, the introduction of new functional traits such as CI and
90 LPE has been suggested; however, variance components for these traits were not available for this
91 breed in Italy. Thus, to investigate the genetic background of these two traits, the present study
92 focused on the estimation of genetic parameters for CI and its relationship with production traits and
93 LPE in Italian Simmental cattle using a genomic approach.

94

95 MATERIALS AND METHODS

96 *Calving interval*

97 CI phenotypes recorded from 275,399 cows in the period 1983 to 2017 were used. The number
98 of parities per cow ranged from 1 to 3, and the first record was mandatory in order to include a cow
99 in the analysis. A total of 594,257 CI records that were greater than 300 days and lower than 700 days
100 were retained for analysis.

101 Heritability and repeatability for CI were estimated using repeatability single-trait model:

$$102 \quad y = \mathbf{hy} + \mathbf{par} + \mathbf{a} + \mathbf{pe} + \mathbf{e} \quad [1]$$

103 where **y** was the CI record; **hy** was the fixed effect of herd-year combination (103,467 levels); **par**
104 was the fixed effect of parity (three levels: 1-2, 2-3, 3-4); **a** was the random additive genetic effect

105 (465,633 animals in the relationship matrix); **pe** was the random effect of permanent environment
106 (465,633 levels); **e** was the random residual.

107 Genetic correlations between fertility and production traits were estimated using a multiple
108 trait animal model with the same structure of [1] considering CI, milk (MY), fat (FY) and protein
109 (PY) 305-d yield (kg). All available records for these 4 traits were included in the analysis (713,376
110 records from 274,759 cows). Average values of 5,687±1,676 kg, 221±81 kg, and 194±70 kg were
111 observed for MY, FY, and PY, respectively.

112

113 *Lactation persistency*

114 Genetic parameters for LPE were investigated using a data set with 285,213 lactation records
115 of 156,389 cows (parities ranging from 1 to 3) farmed in 5,344 herds. Each lactation (from 5 to 305
116 days in milk) was divided into seven intervals and one record per interval was kept. When more than
117 one test day per interval was available, the average value was used. Seven intervals were chosen
118 because the majority of cows have this number of controls available in the routine evaluation system
119 of Italian Simmental. The availability of at least one record before the 45th day of lactation and after
120 the 245th day of lactation was mandatory to include a cow in the analysis. Since milk, fat and protein
121 daily yields showed a similar decreasing trend along lactation, we decided to consider only milk yield
122 to define LPE. Thus, data were arranged in a multivariate framework and a principal component
123 analysis using SAS PROC PRINCOMP procedure was performed to extract eigenvalues and
124 eigenvectors of the phenotypic correlation matrix of test-day records. The second extracted principal
125 component (PC2) was used as an indicator of persistency (Macciotta et al., 2006). The PC2 scores
126 were analyzed with the following single-trait repeatability animal model:

$$127 \quad y = hy + par + a + pe + e \quad [2]$$

128 where **y** was the value of PC2 scores (i.e. LPE); **hy** was the combination between herd and year
129 (49,638 levels); **par** was the effect of parity (three levels: 1, 2 and 3); **a** was the random additive

130 genetic effect (333,003 levels); **pe** was the random effect of permanent environment (333,003 levels);
131 **e** was the random residual.

132 Finally, a two-trait model was used to estimate genetic correlations between CI and LPE of the
133 corresponding lactation: CI records between two consecutive parities (i.e. 1-2, 2-3 and 3-4) and the
134 LPE estimated for the corresponding lactations (i.e. first, second, and third lactations) were used (total
135 of 340,573 records).

136

137 ***Genetic parameter estimation***

138 Variance components and heritability (h^2) estimation for all the described models was
139 performed using a single-step genomic REML (ssGREML) approach with a combined relationship
140 matrix (**H**) built as described in Aguilar *et al.* (2010). Analyses were also done with pedigree
141 relationship matrix (**A**) with very similar results (not shown). Before the analyses, pedigree was traced
142 back for 5 generations. The full pedigree contained 1,981,728 animals of which 7,246 were genotyped
143 for 40,200 markers. Among all the genotyped animals, 3,358 were females with phenotypes and 2,045
144 were sires. The females with both phenotypes and genotypes came from 250 different herds and could
145 be dams of other phenotyped females. The other genotyped animals without phenotypes were half-
146 sib or relatives of phenotyped females. Table 1 shows the distribution of genotyped animals by year
147 of birth. Variance components were estimated using the average information REML (AIREML)
148 algorithm implemented in the blupf90 family programs (Misztal *et al.*, 2014).

149

150

RESULTS AND DISCUSSION

151 ***Calving interval in a single-trait, repeatability model***

152 The phenotypic average of CI for Italian Simmental (397.50 ± 68.32) is consistent with
153 previous reports on other cattle breeds. Values between 387 and 398 were reported for UK Holstein
154 (Wall *et al.*, 2003) and Australian Holstein (Haile-Mariam *et al.*, 2003), respectively. CI of 400 or
155 more days were found in Spanish dairy cattle (González-Recio and Alenda, 2005) and in Mexican

156 Holstein (Montaldo et al., 2010). Large values of CI were also reported for US Holstein (Tiezzi et al.,
157 2017).

158 Heritability (0.06 ± 0.002) and repeatability (0.11 ± 0.002) for CI in the present study, using a
159 single-trait model, were quite low. However, they were slightly higher than previous literature
160 reports. Values of 0.02 were estimated in Mexican Holstein (Montaldo et al., 2010) and in Xinjiang
161 Brown (Fu et al., 2017). Estimates ranging from 0.03 to 0.05 were reported for Australian dairy
162 (Haile-Mariam et al., 2003; 2008), UK Holstein (Wall et al., 2003), Irish Holstein (Olori et al., 2002),
163 Spanish dairy (González-Recio and Alenda, 2005), and Italian Brown Swiss (Dal Zotto et al., 2007).
164 It should be pointed out that heritability for fertility traits is generally low, as reported in a recent
165 review (Ma et al., 2019).

166

167 *Lactation persistency*

168 About 90% of the total phenotypic variance of test-day records was explained by the first two
169 principal components (PC). PC1 was related to the average level of milk yield whereas PC2 was
170 associated with the shape of the lactation curve. They explained about 78% and 12% of test-day
171 phenotypic variance, respectively. PC1 scores ranged from -7.60 to 14.54, with an average value of
172 0.14 ± 2.37 , and it showed a correlation of 0.97 with 305-d MY. PC2 scores ranged from -4.53 to 4.71.
173 Animals were grouped according to their PC2 scores into four classes: Class1 = from -4.53 to -1.04;
174 Class2 = from -1.03 to -0.57, Class3 = from -0.56 to -0.15; and Class4 = from -0.14 to 4.71. Average
175 milk yield for all the seven test day records were calculated separately for each PC2 class. These
176 average lactation curves for different PC2 class are shown in Figure 1. It can be clearly seen that the
177 average lactation curve for PC2 class4 exhibited the steepest negative slope. LPE tends to increase
178 for higher PC2 classes reaching the maximum in Class1. These results confirm the meaning of PC2
179 score as an indicator of the shape of the lactation curve.

180 PC2 (i.e. LPE) and MY showed a phenotypic correlation of -0.03, which was expected
181 because PC2 was defined to be independent of milk production in the calculations. However, the

182 genetic correlation between PC2 and MY was unfavorable (0.35 ± 0.03) because higher values of PC2,
183 i.e., lower LPE, are associated to higher MY. In fact, animals belonging to the Class1 (those with the
184 highest LPE) showed a slightly lower 305-d MY compared to the animals of Class4 (those with the
185 lowest LPE). LPE in this study was not defined by combining production levels at different lactation
186 stages, but using the PC2 values that just capture the shape of lactation curve without considering
187 production levels (PC are orthogonal). Thus, the genetic correlation observed between LPE and MY
188 could probably be a spurious result mediated by other variables. In the literature, estimates of genetic
189 correlation between LPE and MY show different magnitudes and signs according to the LPE
190 definition (Haile Mariam et al., 2003; Muir et al., 2004). Cole and Null (2009) found that genetic
191 correlation between LPE and MY changed magnitudes and signs according to the considered breed.
192 Some studies reported unfavorable genetic correlation between LPE and MY (Cobuci and Costa,
193 2012; Khorshidie et al., 2012; Pereira et al., 2012), whereas other studies reported favorable genetic
194 correlation between these two traits (Muir et al., 2004; Yamazaki et al., 2013).

195 Heritability (0.11 ± 0.004) and repeatability (0.20 ± 0.003) of LPE in a single-trait model were
196 rather low. Macciotta et al. (2006) reported lower heritability and repeatability for LPE using a
197 smaller, older dataset that comprised only phenotypic and pedigree information from the same Italian
198 Simmental population. In fact, heritability estimates for LPE in the literature exhibit a considerable
199 variation, with values ranging from 0.01 (Otwindowska-Mindur and Ptak, 2015) to 0.50 (Koloji et al.,
200 2018). Such relevant differences can be ascribed to statistical model, breed, and trait definition. There
201 is no consensus in literature for measuring LPE. Grayaa et al. (2019) defined LPE as the difference
202 between milk production at 280 days in milk and at the lactation peak, and estimated heritability using
203 different multi-trait models: the estimates ranged from 0.05 when milk fat percentage was considered
204 to 0.21 when milk yield was included among the response variables. Strabel and Jamrozik (2006)
205 reported heritability estimates for LPE that ranged from 0.07 to 0.18 using the eigenvectors of the
206 (co)variance matrix of RRM coefficients. Higher heritability (0.18 ± 0.02) was reported for LPE for
207 first-lactation Canadian Holstein (Muir et al., 2004). Additionally, Cole and Null (2009), observed a

208 large heterogeneity in the h^2 of LPE in several dairy cattle breeds: from 0.09 to 0.26 and 0.18 to 0.28
209 for Milking Shorthorn and Guernsey, respectively.

210

211 ***Calving interval, production traits, and lactation persistency***

212 Heritability for CI obtained with the multiple trait model was slightly higher compared to the
213 single-trait analysis (Table 2). Heritabilities for the production traits were close to the current
214 estimates for the Italian Simmental breed. Unfavorable, moderate genetic correlations were found
215 between CI and production traits (Table 2), as generally reported in literature. The magnitude of the
216 estimates obtained in the present study is not far from previous reports of about 0.5-0.6 (Pryce et al.,
217 2000; Dal Zotto et al., 2007; Fu et al., 2017). Deb et al. (2008) reported a genetic correlation of 0.4
218 between CI and MY in a native breed from Bangladesh. Antagonistic genetic correlation between
219 MY and CI was also observed by Haile-Mariam et al. (2003) and by Strapáková et al. (2016). The
220 latest reported a genetic correlation of 0.51 ± 0.11 , with higher MY observed for cows with longer
221 calving interval. Other similar reproductive traits, such as days open and days from calving to the
222 first service, show undesirable genetic correlations with milk production traits (Abe et al., 2009).

223 Heritability estimates for CI and LPE using the univariate or bivariate models were similar. A
224 weak, positive genetic correlation was observed between these two traits, whereas a near zero
225 phenotypic correlation was observed (Table 3). The positive genetic correlation reflected a favorable
226 association between CI and LPE because high values of CI are associated with high values of PC2,
227 which means lower LPE. On the contrary, lower values of CI (desirable) are related to lower values
228 of PC2 and, therefore, to higher LPE (desirable). However, reports about the genetic association
229 between CI and LPE are not always consistent. An undesirable association between CI and LPE was
230 reported by Atashi et al. (2013), who found that cows with short CI had a lower LPE. Unfavorable
231 genetic correlation between CI and LPE was reported also by Muir et al. (2004). Němečková et al.
232 (2015) reported no significant association between these two traits and Andersen et al. (2011) found
233 no significant differences on peak yield and peak day (i.e. traits associated with LPE) in different

234 calving interval groups. Haile-Mariam et al. (2003) concluded that the genetic correlation between
235 LPE and CI was almost zero. Apart from sampling effect, it should be pointed out that the different
236 definition of LPE used in the various studies may strongly affect the results.

237 An antagonistic relationship between productive and reproductive performances in cattle has
238 been observed: animals need energy to produce milk and to conceive at the same time and, therefore,
239 the energy balance during lactation is of great interest. However, fertility and production traits are
240 associated in a complex causal pattern because this relationship strongly depends on the considered
241 period of the lactation. The attention can be mainly focused on the energy balance during the lactation:
242 milk yield affects the energy balance mostly in the first part of the lactation when cows are
243 inseminated; in this period, body reserve mobilization and negative energy balance frequently occur
244 and the cow has not enough energy to conceive (Andersen et al., 2011; Beran et al., 2012; Duchacek
245 et al., 2014). Thus, a high level of milk production in early lactation (before or around the peak) could
246 reduce fertility with a subsequent delay of pregnancy. In the late part of the lactation, a switch in the
247 causal link between productive and reproductive performance occurs: in this period, energy
248 requirements for the fetal development are higher compared to the first part of lactation and, therefore,
249 pregnancy prioritizes the energy needs with an effect on milk production (that in the advanced stages
250 of lactation can be identified as LPE). More persistent cows with lower peak yield usually suffer less
251 strong energy imbalance during the lactation (Haile-Mariam et al., 2003). Results of the present study
252 highlighted a quite strong unfavorable genetic correlation between CI and MY, confirming the
253 negative relationship between reproductive and productive performance. Slightly favorable genetic
254 correlation between LPE and CI was also observed. However, such relationship between LPE and CI
255 could be influenced by MY. This is because selection for CI is unlikely to cause decline in MY, but
256 the selection for MY ignoring CI would have more negative impact on fertility due to the high
257 unfavorable genetic correlation between these two traits. Nevertheless, LPE has some positive
258 consequences for dairy cows because persistent cows could have less health and reproduction
259 problems, they are easier to manage, and there is a lower feeding cost (Sölkner and Fuchs, 1987;

260 Atashi et al., 2013). Thus, findings of the present work suggest the possibility to limit the fertility
261 deterioration caused by the selection for MY by including CI and LPE as breeding goals.

262 In spite of the low h^2 for CI and LPE that has been confirmed also in the present study, it
263 should be pointed out that genomic selection offers interesting perspectives for improving these
264 functional traits, giving more phenotypes and genotypes are collected. García-Ruiz et al. (2016)
265 showed that the genetic gain per year achieved in US Holsteins has been markedly larger for low
266 heritability traits because of the considerable amount of data (e.g. somatic cell score, productive life,
267 and daughter pregnancy rate). For estimating variance components, the main benefit of using genomic
268 information in a single-step approach is the availability of more data that is reflected in the smaller
269 standard errors (Forni et al., 2011; Veerkamp et al., 2011). The use of combined pedigree and genomic
270 information using the single-step approach could have potential benefits for young candidates that
271 have genotypes but no phenotypic records. However, in our case using A instead of H gave very close
272 estimates (data not shown) because of the small number of genotyped animals. The similar results
273 found using BLUP or ssGBLUP, even if with a small number of genotyped animals, showed the
274 robustness of the latter methodology and the possibility to have better results by increasing the
275 amount of genomic information.

276

277

CONCLUSIONS

278 Although of small magnitude, the heritabilities for CI and LPE show that these traits can be
279 improved by genomic selection. The use of multi-trait models allowed to better understand the genetic
280 connection between CI and LPE, showing that both traits should be included as breeding objective in
281 order to prevent a deterioration of fertility.

282

283

ACKNOWLEDGEMENTS

284 This work was supported by the Italian Government through the project “DUAL BREEDING
285 - *Le razze bovine a duplice attitudine: un modello alternativo di Zootecnia eco-sostenibile*” (grant
286 no. J21J18000010005).

287

- 289 Abe, H., Masuda, Y., and M. Suzuki. 2009. Relationships between reproductive traits of heifers and
290 cows and yield traits for Holsteins in Japan. *J. Dairy Sci.* 92:4055-4062.
- 291 Aguilar, I., I. Misztal, D. L. Johnson, A. Legarra, S. Tsuruta, and T. J. Lawlor. 2010. Hot topic: A
292 unified approach to utilize phenotypic, full pedigree, and genomic information for genetic
293 evaluation of Holstein final score. *J. Dairy Sci.* 93:743–752.
- 294 Andersen, F., Østerås, O., Reksen, O., Toft, N., and Y. T. Gröhn. 2011. Associations between the
295 time of conception and the shape of the lactation curve in early lactation in Norwegian dairy
296 cattle. *Acta Vet. Scand.* 53:5.
- 297 Atashi, H., Zamiri, M. J., Akhlaghi, A., Dadpasand, M., Sayyadnejad, M. B., and A. R.
298 Abdolmohammadi. 2013. Association between the lactation curve shape and calving interval
299 in Holstein dairy cows of Iran. *Iran. J. Vet. Res.* 14:88-93.
- 300 Beran, J., Stadnik, L., and J. Duchacek. 2012.: Effect of body condition score at calving on
301 reproduction indicators of dairy cows. *Reprod. Domest. Anim.* 47:69-69.
- 302 Castillo-Juarez, H., Oltenacu, P. A., Blake, R. W., McCulloch, C. E., and E. G. Cienfuegos-Rivas.
303 2000. Effect of herd environment on the genetic and phenotypic relationships among milk yield,
304 conception rate, and somatic cell score in Holstein cattle. *J. Dairy Sci.* 83:807-814.
- 305 Cobuci, J. A. and C. N. Costa. 2012. Persistency of lactation using random regression models and
306 different fixed regression modeling approaches. *Rev. Bras. Zootec.* 41:1996-2004.
- 307 Cole, J. B., and D. J. Null. 2009. Genetic evaluation of lactation persistency for five breeds of dairy
308 cattle. *J Dairy Sci.* 92: 2248-2258.
- 309 Cole, J. B., and P. M. VanRaden. 2006. Genetic evaluation and best prediction of lactation
310 persistency. *J. Dairy Sci.* 89:2722-2728.
- 311 Dal Zotto, R., De Marchi, M., Dalvit, C., Cassandro, M., Gallo, L., Carnier, P., and G. Bittante. 2007.
312 Heritabilities and genetic correlations of body condition score and calving interval with yield,
313 somatic cell score, and linear type traits in Brown Swiss cattle. *J. Dairy Sci.* 90:5737-5743.

314 Deb, G. K., Mufti, M. M., Mostari, M. P., and K. S. Huque. 2008. Genetic evaluation of Bangladesh
315 livestock research institute cattle breed-1: heritability and genetic correlation. *Bangladesh J.*
316 *Anim. Sci.* 37:25-33.

317 Dekkers, J. C. M., Ten Hag, J. H., and A. Weersink. 1998. Economic aspects of persistency of
318 lactation in dairy cattle. *Livest. Prod. Sci.* 53:237-252.

319 Duchacek, J., Beran, J., Ptacek, M., and L. Stadnik. 2014: Influence of breed, parity and body
320 condition change on reproductive performance of dairy cows. *Reprod. Domest. Anim.* 49:66-
321 66.

322 **Esslemont, R. J., Kossaibati, M. A., and J. Allcock, J. 2001. Economics of fertility in dairy cows.**
323 **BSAP Occasional Publication, 26:19-29.**

324 Forni, S., Aguilar, I., and I. Misztal. 2011. Different genomic relationship matrices for single-step
325 analysis using phenotypic, pedigree and genomic information. *Genet Sel Evol.* 43:1.

326 Fu, X., Lu, L., Huang, X., Wang, Y., Tian, K., Xu, X., Fang, J., Cheng, L., Guo, Z., and Y. Tian.
327 2017. Estimation of genetic parameters for 305 days milk yields and calving interval in Xinjiang
328 Brown cattle. *Agric Sci.* 8:46-55.

329 García-Ruiz, A., Cole, J. B., VanRaden, P. M., Wiggans, G. R., Ruiz-López, F. J., and C. P., Van
330 Tassell. 2016. Changes in genetic selection differentials and generation intervals in US Holstein
331 dairy cattle as a result of genomic selection. *Proc Natl Acad Sci.* 113(28), E3995-E4004.

332 Gengler, N. 1996. Persistency of lactation yields: a review. *Interbull Bulletin.* 12:87-96.

333 González-Recio, O., and R. Alenda. 2005. Genetic parameters for female fertility traits and a fertility
334 index in Spanish dairy cattle. *J. Dairy Sci.* 88:3282-3289.

335 Grayaa, M., Vanderick, S., Rekik, B., Ben Gara, A., Hanzen, C., Grayaa, S., Reis Mota, R.,
336 Hammami, H., and N. Gengler. 2019. Linking first lactation survival to milk yield and
337 components and lactation persistency in Tunisian Holstein cows. *Arch. Anim. Breed.* 62:153-
338 160.

- 339 Groenendaal, H., Galligan, D. T., and H. A. Mulder. 2004. An economic spreadsheet model to
340 determine optimal breeding and replacement decisions for dairy cattle. *J Dairy Sci.* 87:2146-
341 2157.
- 342 Haile-Mariam, M., Bowman, P. J., and M. E. Goddard. 2003. Genetic and environmental relationship
343 among calving interval, survival, persistency of milk yield and somatic cell count in dairy cattle.
344 *Livest. Prod. Sci.* 80:189-200.
- 345 Haile-Mariam, M., Carrick, M. J., and M. E. Goddard. 2008. Genotype by environment interaction
346 for fertility, survival, and milk production traits in Australian dairy cattle. *J. Dairy Sci.* 91:4840-
347 4853.
- 348 Hayes, B. J., Bowman, P. J., Chamberlain, A. J., and M. E. Goddard. 2009. Invited review: Genomic
349 selection in dairy cattle: Progress and challenges. *J. Dairy Sci.* 92:433-443.
- 350 Khorshidie, R., A. A. Shadparvar, N. Ghavi Hossein-Zadeh, and S. Joezy Shakalgurabi. 2012.
351 Genetic trends for 305-day milk yield and persistency in Iranian Holsteins. *Livestock Science,*
352 *144:211-217.*
- 353 Koloji, S., Pathak, K., Karunakaran, M., and A. Mandal. 2018. Lactation Persistency and its Genetic
354 Evaluation in Cattle—A Review. *Research & Reviews: Journal of Dairy Science and*
355 *Technology.* 7:1-8.
- 356 López-Gatius, F. 2003. Is fertility declining in dairy cattle?: a retrospective study in northeastern
357 Spain. *Theriogenology.* 60:89-99.
- 358 Lucy, M.C. 2001. Reproductive loss in high-producing dairy cattle: where will it end? *J. Dairy Sci.*
359 84:1277–1293.
- 360 Ma, L., Cole, J. B., Da, Y., and P. M. VanRaden. 2019. Symposium review: genetics, genome-wide
361 association study, and genetic improvement of dairy fertility traits. *J. Dairy Sci.* 102:3735-3743.
- 362 Macciotta, N. P. P., Vicario, D., and A. Cappio-Borlino. 2006. Use of multivariate analysis to extract
363 latent variables related to level of production and lactation persistency in dairy cattle. *J. Dairy*
364 *Sci.* 89:3188-3194.

365 Miglior, F., Muir, B.L., and B. J. Van Doormaal. 2005. Selection indices in Holstein cattle of various
366 countries. *J. Dairy Sci.* 88:1255–1263.

367 Misztal, I., Tsuruta, S., Lourenco, D., Aguilar, I., Legarra, A., and Z. Vitezica. 2014. Manual for
368 BLUPF90 family of programs. Accessed January 2019.
369 http://nce.ads.uga.edu/wiki/lib/exe/fetch.php?media=blupf90_all7.pdf

370 Montaldo, H. H., Castillo-Juárez, H., Valencia-Posadas, M., Cienfuegos-Rivas, E. G., and F. J. Ruiz-
371 López. 2010. Genetic and environmental parameters for milk production, udder health, and
372 fertility traits in Mexican Holstein cows. *J. Dairy Sci.* 93:2168-2175.

373 Muir, B. L., Fatehi, J., and L. R. Schaeffer. 2004. Genetic relationships between persistency and
374 reproductive performance in first-lactation Canadian Holsteins. *J. Dairy Sci.* 87:3029-3037.

375 Němečková, D., Stádník, L., and J. Čítek. 2015. Associations between milk production level, calving
376 interval length, lactation curve parameters and economic results in Holstein cows. *Mljekarstvo:*
377 *časopis za unaprjeđenje proizvodnje i prerade mlijeka*, 65(4), 243-250.

378 Olori, V. E., Meuwissen, T. H. E., and R. F. Veerkamp. 2002. Calving interval and survival breeding
379 values as measure of cow fertility in a pasture-based production system with seasonal calving.
380 *J. Dairy Sci.* 85:689-696.

381 Otwinowska-Mindur, A., and E. Ptak. 2015. Genetic analysis of lactation persistency in the Polish
382 Holstein-Friesian cows. *Anim. Sci. Pap. Rep.* 33:2.

383 Pereira, R. J., R. S. Verneque, P. S. Lopes, M. L. Santana Júnior, M. R. Lagrotta, R. A. Torres, A.
384 E. Vercesi Filho, and M. A. Machado. 2012. Milk yield persistency in Brazilian Gyr cattle based
385 on a random regression model. *Genet. Mol. Res.* 11:1599-1609.

386 Philipsson, J., and B. Lindhé. 2003. Experiences of including reproduction and health traits in
387 Scandinavian dairy cattle breeding programmes. *Livest. Prod. Sci.* 83:99-112.

388 Pryce, J. E., Coffey, M. P., and S. Brotherstone. 2000. The genetic relationship between calving
389 interval, body condition score and linear type and management traits in registered Holsteins. *J.*
390 *Dairy Sci.* 83:2664-2671.

- 391 Sölkner, J., and W. Fuchs. 1987. A comparison of different measures of persistency with special
392 respect to variation of test-day milk yields. *Livest. Prod. Sci.* 16:305-319.
- 393 Strabel, T., and J. Jamrozik. 2006. Alternative measures of lactation persistency from random
394 regression models with Legendre polynomials. In *Proceedings of the 8th World Congress on*
395 *Genetics Applied to Livestock Production, Belo Horizonte, Minas Gerais, Brazil, 13-18*
396 *August, 2006 (pp. 01-33). Instituto Prociência.*
- 397 Strapáková, E., Candrák, J., and P. Strapák. 2016. Genetic relationship of lactation persistency with
398 milk yield, somatic cell score, reproductive traits, and longevity in Slovak Holstein cattle.
399 *Archives Animal Breeding*, 59(3), 329-335.
- 400 Sun, H. Z., Plastow, G., and L. L. Guan. 2019. Invited review: Advances and challenges in application
401 of feedomics to improve dairy cow production and health. *J. Dairy Sci.* 102
402 <https://doi.org/10.3168/jds>
- 403 Tiezzi, F., de Los Campos, G., Gaddis, K. P., and C. Maltecca. 2017. Genotype by environment
404 (climate) interaction improves genomic prediction for production traits in US Holstein cattle.
405 *J. Dairy Sci.* 100:2042-2056.
- 406 Togashi, K., and C. Y. Lin. 2006. Selection for milk production and persistency using eigenvectors
407 of the random regression coefficient matrix. *J. Dairy Sci.* 89:4866-4873.
- 408 Veerkamp, R. F., Mulder, H. A., Thompson, R., and M. P. L. Calus. 2011. Genomic and pedigree-
409 based genetic parameters for scarcely recorded traits when some animals are genotyped. *J Dairy*
410 *Sci.* 94:4189–4197.
- 411 Wall, E., Brotherstone, S., Woolliams, J. A., Banos, G., and M. P. Coffey. 2003. Genetic evaluation
412 of fertility using direct and correlated traits. *J. Dairy Sci.* 86:4093-4102.
- 413 Walsh, S. W., Williams, E. J., and A. C. O. Evans. 2011. A review of the causes of poor fertility in
414 high milk producing dairy cows. *Anim. Reprod. Sci.* 123:127-138.
- 415 Yamazaki, T., Hagiya, K., Takeda, H., Sasaki, O., Yamaguchi, S., Sogabe, M., Saito, Y., Nakagawa,
416 S., Togashi, K., Suzuki, K., and Y. Nagamine. 2013. Genetic correlations between milk

417 production traits and somatic cell scores on test day within and across first and second lactations
418 in Holstein cows. *Livest Sci.* 152:120-126.
419

420 **Table 1.** Distribution of genotyped animals by birth year.

| Year of birth | Animals (n) |
|---------------|-------------|
| 1963 - 1980 | 38 |
| 1981-1990 | 245 |
| 1991-2000 | 816 |
| 2001 | 118 |
| 2002 | 126 |
| 2003 | 148 |
| 2004 | 196 |
| 2005 | 179 |
| 2006 | 169 |
| 2007 | 196 |
| 2008 | 238 |
| 2009 | 269 |
| 2010 | 318 |
| 2011 | 336 |
| 2012 | 427 |
| 2013 | 859 |
| 2014 | 1048 |
| 2015 | 790 |
| 2016 | 389 |
| 2017 | 265 |
| 2018 | 76 |

421

422

423 **Table 2.** Heritability (diagonal), phenotypic (above diagonal) and genetic (below diagonal)
424 correlation for calving interval and production traits using a 4-trait model.

| | Calving interval | Milk yield | Fat yield | Protein yield |
|------------------|------------------|------------|-----------|---------------|
| Calving interval | 0.09±0.02 | 0.17 | 0.15 | 0.16 |
| Milk yield | 0.64±0.02 | 0.26±0.02 | 0.88 | 0.96 |
| Fat yield | 0.63±0.02 | 0.86±0.01 | 0.25±0.02 | 0.89 |
| Protein yield | 0.56±0.02 | 0.95±0.01 | 0.90±0.01 | 0.22±0.02 |

425

426

427 **Table 3.** Heritability, repeatability, and correlations (phenotypic and genetic) for calving interval and
428 lactation persistency using the two-trait model.

| | Heritability | Repeatability | Correlation | |
|-----------------------|--------------|---------------|-------------|-----------|
| | | | Phenotypic | Genetic |
| Calving interval | 0.05±0.01 | 0.11±0.01 | -0.05 | 0.25±0.03 |
| Lactation persistency | 0.11±0.01 | 0.20±0.01 | | |

429


430

FIGURE CAPTION

431 **Figure 1.** Average shape for the lactation curve using quartiles of the second principal component
432 (PC2) that is associated with lactation persistency. Each curve is made using the average value of
433 milk production (kg/day) of all animals belonging to the first (Class1), second (Class2), third
434 (Class3), and fourth (Class4) quartile of PC2.

435

436 Cesarani. Figure 1.



437

438