

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas

This is a pre print version of the following article:

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1781336> since 2021-03-22T12:16:24Z

Published version:

DOI:10.1016/j.envres.2021.110999

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Environmental Research

Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas

--Manuscript Draft--

Manuscript Number:	
Article Type:	Research paper
Section/Category:	Environmental Health & Risk Assessment
Keywords:	Urban Vegetation; Remote Sensing; Public Health; Isoprostane
Corresponding Author:	Samuele De Petris University of Torino ITALY
First Author:	Samuele De Petris
Order of Authors:	Samuele De Petris Giulia Squillacioti, Dr. Roberto Bono, Prof. Enrico Borgogno-Mondino, Prof.
Abstract:	<p>Green spaces may benefit human health mainly by mitigating noise and air pollution, promoting physical or social activities and improving mental health. Based on the influence that green space exposure seems to exert on Public Health and using a multidisciplinary approach, we investigated, the association between oxidative stress (OS) and green exposure in children. Overall, 207 subjects (10-13 yrs.) living in Torino (NW- Italy) were involved in this study. Each participant provided a urinary sample, used to quantify a reliable OS biomarker (15-F2t-IsoP), and their residence addresses, used for geocoding. Green exposure was characterised by calculating i) the Soil Adjusted Vegetation Index (SAVI) within fixed buffers around each participant's home, using remotely-sensed data; ii) Tree Map accounting for evergreen/broadleaf species; iii) The percentage of green cover (PGC). Significant negative correlation (Pearson's $r = -0.758$, $p < 0.001$) between PGC and 15-F2t-IsoP was found. Greater SAVI was associated with lower OS (Pearson's $r = -0.717$, $p < 0.001$). Noticeably, evergreens seemed to determine a significant OS reduction compared to broadleaves (slope = -0.12 and -0.02, respectively; Warton-test $F = 12.48$, $p = 0.0011$). Finally, a spatial distribution of 15-F2t-IsoP estimates map, overlying with 2011 Census Data on same-aged dwellers of Torino, was generated. Predictive models accounting for green spaces influence on OS can be useful tool derived from geomatic employ in Public Health field. Future developments of such a multidisciplinary approach should be considered in urban planning and policy-makers decisions to better define priority zones to requalify in urban settings.</p>
Suggested Reviewers:	Andrea Lessio andrea.lessio@ithaca.polito.it Alessio Calantropio alessio.calantropio@polito.it Piero Boccoardo piero.boccoardo@polito.it Eija Parmes eija.parmes@vtt.fi Carmen de Keijzer carmen.dekeijzer@isglobal.org

**UNIVERSITY OF TURIN, ITALY**

Department of Agricultural, Forest and Food Sciences
Largo P. Braccini 2, 10095, Grugliasco (TO), Italy

October 14th, 2020Dear Editor of *Environmental Research*,

Please find attached the manuscript “*Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas*”. In my view, the relevance of the results can be summarised as follows. Firstly, the evidence that urban green spaces are able to influence oxidative stress in youths living in urban areas. Secondly, evergreens show a greater impact on this association compared to broadleaf species. Thirdly, the analysis on spatial distribution of oxidative stress can be a useful tool derived from GIS and remote sensing employ in Public Health, serving urban planning and policy-makers decisions to define potential priority zones to requalify in urban settings. Further key information about the presented manuscript:

- the manuscript is an original work, has not been previously published, and is not under consideration for publication elsewhere;
- the participation of all human subjects did not occur prior their informed consent was obtained;
- all authors have disclosed any potential competing interest regarding the submitted article;
- all authors have read the manuscript, agreeing on the submission to *Environmental Research*, and accepting the responsibilities for the manuscript’s contents;
- all authors have read and approved the paper and it has not been published previously nor is it being considered by any other peer-reviewed journal;

The multidisciplinary approach of this manuscript involves: biosphere, anthroposphere, atmosphere. In my point of view, this manuscript may deserve Your considerations because it fits some of subject areas included in the journal:

- Environmental risks assessment and management
- Air pollution quality and human health
- Risks and public health
- Environmental management and policy
- Environmental risks assessment and management

Please find below some suggestions for potential reviewers, as requested.

1. Andrea Lessio - andrea.lessio@ithaca.polito.it
2. Alessio Calantropio - alessio.calantropio@polito.it
3. Piero Boccardo - piero.boccardo@polito.it
4. Eija Parmes - eija.parmes@vtt.fi.
5. Carmen de Keijzer - carmen.dekeijzer@isglobal.org

Hoping that the manuscript may fulfil the scientific standards of *Environmental Research*.

Best Regards,
Samuele De Petris

Contacts:

e-mail: samuele.depetris@unito.it

1. Oxidative stress and green exposure association was investigated in children
2. Green exposure was analyzed by geomatic techniques
3. Spectral vegetation index and tree census data were used to map oxidative stress
4. Evergreens trees determine a significant oxidative stress reduction
5. Geomatics can support urban planning to improve public health

Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban Areas

DE PETRIS SAMUELE^{A*}, SQUILLACIOTI GIULIA^{B*}, BONO ROBERTO^B, BORGOGNO-MONDINO ENRICO^A

^ADepartment of Agriculture, Forest and Food sciences, University of Torino, 10095 Grugliasco (TO), Italy; samuele.depétris@unito.it (S.D.P.); enrico.borgogno@unito.it (E.B.M.)

^BDepartment of Public Health and Pediatrics, University of Torino, 10121 Torino, Italy; giulia.squillacioti@unito.it (G.S.); roberto.bono@unito.it (R.B.)

*these two authors equally contributed to the conceptualization and execution of the manuscript

Corresponding author: Samuele De Petris, samuele.depétris@unito.it

Declarations of interest: none.

Geomatics and Epidemiology: Associating Oxidative Stress and Greenness in Urban

Areas

Abstract: Green spaces may benefit human health mainly by mitigating noise and air pollution, promoting physical or social activities and improving mental health. Based on the influence that green space exposure seems to exert on Public Health and using a multidisciplinary approach, we investigated, the association between oxidative stress (OS) and green exposure in children. Overall, 207 subjects (10-13 yrs.) living in Torino (NW- Italy) were involved in this study. Each participant provided a urinary sample, used to quantify a reliable OS biomarker (15-F2t-IsoP), and their residence addresses, used for geocoding. Green exposure was characterised by calculating i) the Soil Adjusted Vegetation Index (SAVI) within fixed buffers around each participant's home, using remotely-sensed data; ii) Tree Map accounting for evergreen/broadleaf species; iii) The percentage of green cover (PGC). Significant negative correlation (*Pearson's r* = -0.758, *p* < 0.001) between PGC and 15-F2t-IsoP was found. Greater SAVI was associated with lower OS (*Pearson's r* = -0.717, *p* < 0.001). Noticeably, evergreens seemed to determine a significant OS reduction compared to broadleaves (slope = -0.12 and -0.02, respectively; *Warton-test F* = 12.48, *p* = 0.0011). Finally, a spatial distribution of 15-F2t-IsoP estimates map, overlying with 2011 Census Data on same-aged dwellers of Torino, was generated. Predictive models accounting for green spaces influence on OS can be useful tool derived from geomatic employ in Public Health field. Future developments of such a multidisciplinary approach should be considered in urban planning and policy-makers decisions to better define priority zones to requalify in urban settings.

Keywords: Urban Vegetation, Remote Sensing, Public Health, Isoprostane

1. Introduction

Green spaces are thought to benefit human health mainly through i) mitigation of noise and air pollution exposure ii) promotion of outdoor exercise and social activities iii) improvement of mental health. Participation in physical activity is significantly higher for people living close to parks, or green areas, determining a reduction of overweight, obesity and, in general, a lower body mass index (Bell et al., 2008). Physical activity is also showing a positive impact on health by reducing risk of diabetes (Dalton et al., 2016) and cardiovascular diseases (Fong et al., 2018; Pereira et al., 2012). Green space exposure proves to positively affect mental health and social engagement by reducing stress (Gong et al., 2016; Markevych et al., 2017), depressive symptoms (Garipey et al., 2015), aggressive behaviour (Younan et al. 2016), and contributing in decelerating the cognitive decline (de Keijzer et al., 2018). Moreover, maternal exposure to green space is positively associated with greater new-born's birthweight; this is mainly due to environmental pollution mitigation, opportunity of socialisation, and to exercise in contact with nature.

Studies addressing greenness and respiratory health possible association are still inconsistent. Some authors referred about a positive effect of higher exposures to greenness finding risk reduction of asthma and respiratory symptoms (Lovasi et al., 2008; Sbihi et al., 2015; Squillacioti et al., 2019b). On the other hand, some authors found adverse effect of green space exposure on asthma, rhinitis and respiratory health in general (Fuertes et al., 2014; Parmes et al., 2020).

Based on the influence that green space exposure seems to exert on Public Health, we speculated that greenness might also be involved in oxidative stress (OS) induction.

OS is a pre-pathological condition characterised by an imbalance between pro-oxidant and anti-oxidant species, in favour of pro-oxidants. OS does not represent a pathology

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

itself (Sies, 2015), but a risky condition related to the pathophysiological mechanisms behind several health impairments, such as Cardio Vascular Diseases (CVDs), diabetes and respiratory diseases. Furthermore, it is susceptible to those environmental risk factors able to act as pro-oxidants (e.g. air pollutants and tobacco smoke)(Milne et al., 2015).

At the meantime, OS may be influenced by individual characteristics of subjects like obesity, health *status*, tobacco smoking exposure and exercise training (Nikolaidis et al., 2011; van 't Erve et al., 2017). Within this context, we focused on the potential association between OS and urban green spaces, being this type of knowledge still lacking (Woo et al., 2009; Yeager et al., 2018). We operated by a multidisciplinary approach aiming at investigating strength of this potential association, specifically evaluating usefulness of geomatic tools in epidemiological studies. Many epidemiological studies investigating the relationship between green space and health benefits are already based on remotely-sensed data. Specifically, the Normalised Difference Vegetation Index (NDVI) is widely used in epidemiological research as spectral metric (Fong et al., 2018). NDVI does not provide any information about species characterisation (Fong et al., 2018). Few works, alternatively, have tried to characterise exposure with reference to land cover maps (Egorov et al., 2017; Parmes et al., 2020; Tsai et al., 2019) or tree *census* data, with the aim of taking into account type and shape of green spaces, or vegetation fraction cover (Browning and Rigolon, 2018; Lovasi et al., 2013; Pilat et al., 2012).

2. Materials

2.1. Study Area

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

The study area corresponds to the municipality of Torino (NW Italy, E: 396027 N: 4991913 reference frame WGS84 UTM32N), sizing about 130 Km² (Fig.1). In 2019 green areas covered 43% of the whole city, addressing Torino as one of the most relevant greenest cities (Baycan-Levent and Nijkamp, 2009; Li et al., 2015). With reference to green patches, the following metrics were computed: minimum and maximum patch size are 0.015 and 108 ha, respectively; patch size mode is 0.63 ha; 25th percentile is 0.38 ha; 75th percentile is 3.76 ha.




Figure 1 – Torino area localization (Reference frame: WGS84 UTM32N).

2.2. Multispectral Data

Multispectral imagery are widely used to detect and characterise vegetation in urban contexts, permitting to locally map vegetated areas (Mudele and Gamba, 2019; Rosina and Kopecká, 2016). We assumed that the maximum of vegetation vigour, in the area, occurs in the summer period (June-July, Zhou et al. 2016; De Petris S. et al. 2019;


1 Borgogno-Mondino, Sarvia, and Gomarasca 2019). Consequently, a Sentinel-2 Level
2 2A image (S2), acquired by the Copernicus Sentinel-2 MSI (Multi Spectral Instrument)
3 sensor on 14th June 2016, was obtained from the Theia CNES geoportal
4 (<https://www.Theia-land.fr/en/product/sentinel-2-surface-reflectance/>, Baetens,
5 Desjardins, and Hagolle 2019; Revel et al. 2019)). In summer time, vegetation-related
6 biomass expresses its maximum in Torino. This assumption also relies on the observed
7 inter-annual phenological behaviour of vegetation in urban context. Since
8 anthropogenic green areas show no expansion dynamics, if compared with natural ones,
9 and vegetation density does not change significantly, due to the human-management
10 (i.e. pruning, mowing), biomass shows inter-annual flat behaviour (Li et al., 2017). S2
11 Level 2A data are supplied already calibrated in at-the-ground-reflectance. Nominal
12 radiometric accuracy is about 0.01 reflectance units (European Space Agency/Centre
13 National d’Eudes Spatiales (CNES), 2019). Technical features S2 image are reported in
14 Table 1. It is worth to remind that the minimum mapping unit from S2 imagery is 100
15 m²; this size was retained appropriate if compared with the average dimension of green
16 areas in Torino.

17 [Here Table 1]

18 2.3.Epidemiological Data

19 Available epidemiologic sample refers to an on-going cohort whose subjects were
20 recruited in 2010 as part of a research project funded by the Piedmont Regional Council
21 focusing on the effects of environmental factors. This cross-sectional study involved
22 207 healthy children (10–13 years old) from secondary schools located in Torino. All
23 subjects gave the assent to participate along with their parents, who signed an informed
24 consent. Since the study involved human subjects, the study protocol required the Ethics

1 Committee approval (protocol number 826/13/08). Parents filled out a questionnaire to
2 provide general information and their home address; participants provided a sample of
3 urine that was used for quantification of a reliable biomarker of OS, namely 15-F2t-
4 urine that was used for quantification of a reliable biomarker of OS, namely 15-F2t-
5 isoprostane (15-F2t-IsoP). At this concern, 15-F2t-IsoP was quantified to measure OS
6 by a specific enzyme-linked immunosorbent assay (ELISA) kit (Oxford, MI, USA),
7 according to the manufacturer's instructions. To achieve better accuracy each sample
8 was diluted 1:4, as reported in previous works (Romanazzi et al., 2013; Squillacioti et
9 al., 2019a). Urinary creatinine (crea) was quantified in each urinary sample in order to
10 normalise urinary excretion rate and dilution. 15-F2t-IsoP was finally referred as ng/mg
11 crea. Collected home addresses (Fig. 2) were used to geocode each subject by means of
12 MMQGis QGIS plugin; positioning accuracy achieved by MMQGis is declared lower
13 than 25m (Cetl et al., 2018).



54 *Figure 2 – Position (accuracy < 25 m) of sampled subjects (207) children aged between 10 and 13) within the study area.*

56 *Reference frame is WGS84 UTM 32N.*

2.4. Auxiliary data

Auxiliary data, provided by institutional geoportals, were supplied in vector format.

A Tree Map (TM), updated at 2019 with a nominal scale 1:1000 was obtained as point layer from the Torino Geoportal. TM contains more than 160,000 individuals divided in evergreen and broadleaf species (Fig. 3). Main evergreen genera are: *Abies spp.*, *Picea spp.*, *Pinus spp.*, *Thuja spp.*, *Cedrus spp.*, *Cupressus spp.*, *Ilex spp.*, *Magnolia spp.* representing the 5% of the trees; while the broadleaf genera are: *Platanus spp.*, *Tilia spp.*, *Aesculus spp.*, *Celtis spp.*, *Acer spp.*, *Ulmus spp.*, *Carpinus spp.*, representing the 95%.

Census data (*CD*, 2011) were obtained from the ISTAT (Italian Statistics Institute) geoportal; *CD* has a nominal scale of 1:5000 and contains census sections (polygons) and the correspondent features of surveyed people, included the number of children aged between 10 and 13 years. *CD* was used to describe the spatial distribution of potential targets that was retained a driving factor to address future urban policies eventually aimed at mitigating OS effects.




Figure 3 - Tree Map (TM) providing position of trees within the study area. It was obtained from Torino Geoportal. Nominal scale is 1:1000 and reference frame WGS84 UTM 32N.

3. Methods and data processing

Spatial analyses were operated by *SAGA GIS* vs.7.1 and *QGIS* vs. 3.4.12; statistical analyses were performed using *Past* vs. 3.6.6 (Hammer et al., 2001).

3.1. Detection and characterization of urban vegetation

In urban contexts soil background makes remotely sensed pixels not pure, i.e. mixed (Small and Lu, 2006; Song, 2005). In fact, background exerts considerable influence on the average pixel spectrum, thus influencing the accuracy of detection of green areas. To minimize such an effect while mapping vegetation within urban environments (Huete, 1988), the Soil Adjusted Vegetation Index (SAVI) was calculated by eq. 1:

$$SAVI = \frac{1.5 (\rho_{NIR} - \rho_{RED})}{\rho_{NIR} + \rho_{RED} + 0.5} \quad (1)$$

where ρ_{NIR} and ρ_{RED} are S2 band 4 (665 nm) and band 8 (840 nm), respectively. Not-vegetated areas (buildings, streets etc.) were masked out by SAVI thresholding: pixel showing SAVI values < 0.45 were labelled as not-vegetated (Borgogno-Mondino et al., 2016; Burgan, 1993; Gao, 1996; Ormsby et al., 1987; Zhang et al., 2003). A map showing only vegetated areas was, therefore, generated (Fig.4), making possible to obtain preliminary information about spatial distribution of biomass in the area. Biomass is well-known to be related to spectrally derived vegetation indices (SAVI included).




Figure 4 – SAVI map obtained by S2 image. Not-vegetated pixels were masked out by SAVI thresholding (SAVI < 0.45). Reference frame is WGS84 UTM32N.

3.2. Green Exposure Assessment

1
2 Many studies attested that green spaces may positively contribute to children's health
3
4 (McMorris et al., 2015; Thiering et al., 2016). Greenness positive effects are assumed to
5
6 decay with distance and increase with exposure time. Consequently, a first reasonable
7
8 approach to quantify greenness exposure level is the characterisation of vegetated areas
9
10 falling in a buffer zone surrounding subject' houses (James et al., 2015). Considering that
11
12 children aged between 10 and 13, they can be supposed having an independent mobility
13
14 pattern that ranges between 500 and 1000 m from their houses (Fagerholm and Broberg,
15
16 2011; Mavoia et al., 2011; Tillberg Mattsson, 2002). Consequently, buffers having a fixed
17
18 radius of 500 m (B_{500}) were mapped around geocoded addresses. Different metrics, useful
19
20 to measure and characterise green space falling in each buffer, were then calculated.
21
22

23
24
25
26 With reference to the masked SAVI map pixels were counted falling in B_{500} and the
27
28 related SAVI mean value (mSAVI) computed. The percentage of green cover (PGC) was
29
30 also calculated comparing the vegetation fraction with the whole area of the buffer.
31
32 mSAVI was then multiplied by PGC, assuming the latter as a sort of weight. A new
33
34 metric, hereinafter called *Unitary SAVI* (uSAVI) was finally obtained to somehow
35
36 measure biomass density. Using TM, two more variables accounting for the number of
37
38 evergreen and broadleaf tree falling in B_{500} were calculated, namely NET (Number of
39
40 Evergreen Trees) and NBT (Number of Broadleaf Trees).
41
42
43
44
45
46
47

3.3. Statistical Analysis and Data Processing

48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Outliers detection in the 15-F2t-IsoP dataset was performed looking for values greater than $Q3 + 1.5 * (Q3 - Q1)$, where Q1 is the 25th percentile and Q3 is the 75th percentile of 15-F2t-IsoP statistical distribution. Preliminarily, the relationship between OS and uSAVI

1 was tested using the Pearson's linear correlation (r) and the correspondent linear
2 regression was calibrated by Ordinary Least Squares (OLS). Correlation was also tested
3
4 by r for the following relationship: 15-F2t-IsoP vs PGC, IsoP vs NET, and 15-F2t-IsoP
5
6 vs NBT. Obtained values showed, for all the tested relationships, a weak correlation (see
7
8 Results and Discussion section). Nevertheless, an envelope function (Ev) of these
9
10 distributions was considered bounding observations clouds. Ev can be interpreted as the
11
12 upper limit beyond which none 15-F2t-IsoP values exist referred to actual vegetation
13
14 parameters values. It was computed by dividing the 15-F2t-IsoP range into 20
15
16 equiprobable classes corresponding to a class width of about 1.4 points of 15-F2t-IsoP
17
18 value, starting from 3.56 ng/mg crea. With reference to the defined classes the
19
20 correspondent 15-F2t-IsoP and uSAVI, PGC, NET, NBT maximum values were
21
22 calculated and compared by scatterplots. Linear associations relating 15-F2t-IsoP- Ev with
23
24 the other variables, were tested using Pearson's coefficient and correspondent linear
25
26 regression modelled. Namely, NET- Ev and NBT- Ev slope values were compared
27
28 according to the Warton method (Warton et al., 2006) in order to evaluate if evergreen
29
30 trees determined a significant difference in OS reduction rate (i.e. slope value of bivariate
31
32 model) with respect to broadleaf trees. Given these relationships, in order to spatialise the
33
34 information generating a map of estimate of 15-F2t-IsoP- Ev , authors only focused on
35
36 $uSAVI-Ev$ since, among the tested metrics, from an operational point of view, it can be
37
38 easily and globally computed from free satellite data. Consequently, the same procedure
39
40 could be applied everywhere in spite of the existence of more specific databases
41
42 containing more accurate information about local vegetation. Nevertheless, other useful
43
44 information is derived from the interpretation of relationship between 15-F2t-IsoP- Ev and
45
46 PGC- Ev , NET- Ev and NB- Ev for investigation purposes, solely. With these premises
47
48 the linear regression model relating 15-F2t-IsoP- Ev to uSAVI- Ev was adopted to
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 spatialise 15-F2t-IsoP-Ev estimates in the study area. For this purpose a 500x500 m
2 squared graticule (G) was generated assuming this size consistent with a walking
3 mobility of 10–15 minutes (Wolch et al., 2014). For each cell of G, the uSAVI maximum
4 value was computed from the previously generated SAVI masked map. G was then
5 rasterised to generate uSAVI^G. A raster map of estimates of 15-F2t-IsoP-Ev (hereinafter
6 called 15-F2t-IsoP-Ev^G) was finally computed by grid calculation tools, implementing
7 the previously calibrated linear regression relating uSAVI to 15-F2t-IsoP-Ev. An
8 accuracy assessment of 15-F2t-IsoP-Ev^G was operated by a “leave-one-out” procedure
9 (Brovelli et al., 2008) to compute the Mean Absolute Error (MAE) (Willmott and
10 Matsuura, 2005).

26 **4. Results and Discussions**

27 Concerning outliers detection in the 15-F2t-IsoP dataset 4 individuals were found having a 15-
28 F2t-IsoP value > 40 ng/mg crea and removed. With reference to the test concerning relationship
29 between OS and uSAVI (Fig. 5) a Pearson’s linear correlation coefficient was found equal to -
30 0.045, $p > 0.05$. A linear regression was calibrated by OLS estimation finding a slope value of
31 -2.214, $C.Is$ 95% (upper bound =9.538 and lower bound=4.68, $p > 0.05$).
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65





Figure 5– Linear regression model (red line, dark-grey are 95% Cis) of 15-F2t-IsoP (dependent variable) and uSAVI (independent variable). Box-plot represent respectively: minimum, Q1, median, Q3, maximum of both uSAVI (up box-plot) and 15-F2t-IsoP (right) value distributions without outliers.

It is worth to stress that some of the OLS regression assumptions were violated: residuals were not normally distributed (Shapiro-Wilk test) showing autocorrelation (Durbin-Watson test) and, more in general, both r and the OLS regression model showed that OS and uSAVI were poorly correlated. Such poorly correlation values were possibly due to the fact that OS is also related with other factors like air quality, personal behaviour (i.e. diet, smoking), individual pathology. Therefore, all these relationships influence the covariance and, as well, correlation coefficient. Concerning the other tested relationships (15-F2t-IsoP vs PGC, IsoP vs NET, and 15-F2t-IsoP vs NBT) the following values were found (Tab.2).

[Here Table 2]

It is worth to remind that OS level differences are due to several risk factors, both individual and environmental (Fig. 6). Therefore, a modelling of 15-F2t-IsoP based on uSAVI does not have the aim of producing reliable absolute 15-F2t-IsoP estimates; conversely, it is expected to produce a general overview, enhancing relative differences between different areas.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 6 – OS is influenced by many individual (left ellipse) and environmental factors (right ellipse). This generates high variability in OS values if ordinary regression-based approaches are used. (Green ellipse)

Interaction between OS and greenness. OS level can be influenced by green areas increasing physical activity or mitigating air pollution (Nowak, Crane, and Stevens 2006; McMorris et al. 2015)

Figure 7 graphically shows scatterplots relating 15-F2t-IsoP to uSAVI, PGC, NET and NB. Red lines define the upper bound (*envelope*) of 15-F2t-IsoP estimates, which class *maxima*, used to generate the new metrics 15-F2t-IsoP-Ev, uSAVI-Ev, PGC-Ev, NET_Ev and NB-E, have to somehow represent. Classes used for maximum value computation are reported in Tab. 3.




Figure 7 – Scatterplots relating 15-F2t-IsoP and vegetation parameters. A linear envelope function (Red line) can be defined for each cloud defining the upper boundary of the estimates.

[Here Table 3]

Figures 8 (a-b) and 9, and table 4 report results of the bivariate models that relate 15-F2t-IsoP-Ev to PGC-Ev, 15-F2t-IsoP-Ev and uSAVI-Ev, respectively.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 8 – (A) bivariate model between 15-F2t-IsoP-Ev and uSAVI-Ev; (B) Bivariate model between 15-F2t-IsoP-Ev and PGC-Ev. Pearson correlation coefficient (r) was reported with related significance level. Dark-grey limits are 95% CIs.

A significant negative correlation was found between 15-F2t-IsoP-Ev and PGC-Ev ($r = -0.758$, $p < 0.001$) and between 15-F2t-IsoP-Ev and uSAVI-Ev ($r = -0.717$, $p < 0.001$). Results prove that higher vegetation cover seems to significantly reduce OS. In particular, if vegetation cover is composed by trees (i.e. high uSAVI), OS tends to decrease. With respect to evergreen and broadleaf tree *genera*, the correspondent regressions were tested separately (Figure 9).





Figure 9 – (Triangle) bivariate model between 15-F2t-IsoP-Ev and NET-Ev; (Circle) Bivariate model between 15-F2t-IsoP-Ev and NBT-Ev. Pearson correlation coefficient (r) was reported with related significance level. Dark-grey limits are 95% CIs. In Bold are highlighted IsoP reduction rate (slope values) of NET and NBT. NET slope is steeper negative than NBT one and the two slopes are statistically different ($p < 0.01$).

1 To test significance of slope of NET and NBT, the Warton test (Warton et al., 2006) was applied
2 finding a F value = 12.48 , $p = 0.0011$. It proved that NET and NBT bivariate model gain values
3 were significantly different. In particular, NET determines a negative steeper 15-F2t-IsoP-Ev
4 reduction rate six-time greater in respect of NBT, suggesting that evergreen trees determine a
5 stronger positive effect on OS. Statistic parameters of tested relationships are reported in Table
6
7
8
9
10
11
12 4.

13
14
15 [Here Table 4]
16
17

18 Concerning accuracy of estimates, MAE of 15-F2t-IsoP-Ev^G was computed by a leave-one-out
19 approach. It resulted equal to 5.26 15-F2t-IsoP ng/mg crea. Results proved that green areas
20 where evergreen trees dominate OS is lower. Consequently, urban planners could exploit this
21 information while designing new restoration/requalification actions, possibly promoting these
22 interventions especially in those parts of the city having a higher density of potential targets
23 (i.e. children). In order to exemplify spatial implications of the above-mentioned relationships
24 a 15-F2t-IsoP-Ev^G map was generated with a grid size of 10 m and coupled, by overlaying, with
25 CD (Fig. 10). This map is intended to represent an operational tool to describe OS risk of 10-
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 areas could be added, or old ones requalified, providing estimates of associated 15-F2t-IsoP-
3 Ev^G resulting from the adopted choices.
4
5



37
38 *Figure 10 –Map shows spatial distribution of 15-F2t-IsoP-Ev^G estimates coupled by overlaying with 2011*

39 *Census Data (children). It could be useful to urban planners to define priority zones where creating/*
40 *requalifying green areas. Dotted rectangles are critical zones where high children density and high OS values*
41 *are located. Reference frame is WGS84 UTM32N.*
42
43
44
45

46 **5. Conclusions**

47
48 This study aimed at investigating and formalising the relationship between OS and
49 greenness with reference to children living in Torino (NW Italy). Significant negative
50 correlations (Pearson's $r = -0.758$, $p < 0.001$) between PGC-Ev and 15-F2t-IsoP-Ev were
51 found indicating that a higher degree of green areas around children's houses determines a
52 lower OS levels in children. One of the potential mechanisms underlying this association
53
54
55
56
57
58
59
60
61
62
63
64
65

1 could be related to the presence of accessible green areas potentially used to perform
2 physical activity.
3

4 Moreover, our results emphasised that local biomass plays a key role in OS reduction. In
5 fact, higher values of uSAVI correspond to lower levels of OS (Pearson's $r = -0.717$, $p <$
6 0.001). High biomass can be generally related to a strong presence of trees: green areas
7 where trees density was higher proved to be significantly related to lower OS values. Trees
8 are thought to provide positive effects to health by reducing exposure to particular matter
9 (PM_{10} or $PM_{2.5}$) or ozone (Fausto Manes et al., 2012; F. Manes et al., 2012; Nowak et al.,
10 2006). In particular, evergreen trees seem to determine a significant 15-F2t-IsoP-Ev
11 reduction rate in respect to broadleaves (NET slope = -0.12 , NBT slope = -0.02 , Warton-
12 test $F = 12.48$, $p = 0.0011$). With respect to evergreen trees, this can be related to: (a) a
13 higher content of cuticular resins, which have an important role in pollution absorption and
14 mitigation (Marando et al., 2016; Sawidis et al., 2012); (b) a total leaf area greater than
15 broadleaf that, considering the annual phenology of urban trees, provides larger absorption
16 surfaces for pollutants. Furthermore, in northern Italy, especially in Po river watershed, air
17 quality dramatically decreases in winter (Bigi and Ghermandi, 2014; Finardi and Pellegrini,
18 2004) when only the evergreen species can provide the mitigation effect on air pollution.
19 uSAVI and OS bivariate model (15-F2t-IsoP-Ev^G map) can be used for future developments
20 related to urban planning strategies and policy makers decisions, making possible to define
21 priority zones where requalification or new greenspaces should be considered. These plans
22 might take into account the here presented results according to three main strategies: (a)
23 green cover improvement; (b) trees or shrubs (i.e. high biomass density cover) planting in
24 place of grass; (c) evergreen trees selection rather than broadleaves.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 It is worth to remind that calibrated uSAVI-OS model describes the Ev function containing
56 OS estimates, meaning that one can infer only about the upper limits of expected 15-F2t-
57
58
59
60
61
62
63
64
65

1 IsoP, with no possibility of knowing its exact local value. This is mainly due to the high
2 intrinsic variability in OS-induced factors (individual and environmental). Since this study
3
4 mainly refers to children, further works are expected to test the association between
5
6 greenness and OS in differently aged subjects and different geographic urban areas where
7
8 the sole intensity of urbanization can produce different level of OS (Squillacioti et al.,
9
10 2020). Expectation is that this approach could promote the highest public health standards
11
12
13 even with the best urban green design *criteria*.
14
15
16
17
18
19
20

21 **References**

- 22
23
24 Baetens, L., Desjardins, C., Hagolle, O., 2019. Validation of Copernicus Sentinel-2 Cloud Masks
25 Obtained from MAJA, Sen2Cor, and FMask Processors Using Reference Cloud Masks
26 Generated with a Supervised Active Learning Procedure. *Remote Sensing* 11, 433.
- 27 Baycan-Levent, T., Nijkamp, P., 2009. Planning and management of urban green spaces in Europe:
28 Comparative analysis. *Journal of Urban Planning and Development* 135, 1–12.
- 29 Bell, J.F., Wilson, J.S., Liu, G.C., 2008. Neighborhood greenness and 2-year changes in body mass
30 index of children and youth. *American journal of preventive medicine* 35, 547–553.
- 31 Bigi, A., Ghermandi, G., 2014. Long-term trend and variability of atmospheric PM10 concentration in
32 the Po Valley.
- 33
34 Borgogno-Mondino, E., Lessio, A., Gomasasca, M.A., 2016. A fast operative method for NDVI
35 uncertainty estimation and its role in vegetation analysis. *European Journal of Remote Sensing*
36 49, 137–156.
- 37 Borgogno-Mondino, E., Sarvia, F., Gomasasca, M.A., 2019. Supporting Insurance Strategies in
38 Agriculture by Remote Sensing: A Possible Approach at Regional Level, in: *International*
39 *Conference on Computational Science and Its Applications*. Springer, pp. 186–199.
- 40 Brovelli, M.A., Crespi, M., Fratarcangeli, F., Giannone, F., Realini, E., 2008. Accuracy assessment of
41 high resolution satellite imagery orientation by leave-one-out method. *ISPRS Journal of*
42 *Photogrammetry and Remote Sensing* 63, 427–440.
- 43
44 Browning, M., Rigolon, A., 2018. Do Income, Race and Ethnicity, and Sprawl Influence the
45 Greenspace-Human Health Link in City-Level Analyses? Findings from 496 Cities in the
46 United States. *International Journal of Environmental Research and Public Health* 15, 1541.
47 <https://doi.org/10.3390/ijerph15071541>
- 48 Burgan, R.E., 1993. Monitoring vegetation greenness with satellite data. US Department of Agriculture,
49 Forest Service, Intermountain Research Station.
- 50
51 Cetl, V., Kliment, T., Jogun, T., 2018. A comparison of address geocoding techniques—case study of the
52 city of Zagreb, Croatia. *Survey Review* 50, 97–106.
- 53 Dalton, A.M., Jones, A.P., Sharp, S.J., Cooper, A.J.M., Griffin, S., Wareham, N.J., 2016. Residential
54 neighbourhood greenspace is associated with reduced risk of incident diabetes in older people:
55 a prospective cohort study. *BMC Public Health* 16, 1171. [https://doi.org/10.1186/s12889-016-](https://doi.org/10.1186/s12889-016-3833-z)
56 [3833-z](https://doi.org/10.1186/s12889-016-3833-z)
- 57 de Keijzer, C., Tonne, C., Basagaña, X., Valentín, A., Singh-Manoux, A., Alonso, J., Antó, J.M.,
58 Nieuwenhuijsen, M.J., Sunyer, J., Dadvand, P., 2018. Residential surrounding greenness and
59
60
61
62
63
64
65

- cognitive decline: a 10-year follow-up of the Whitehall II cohort. *Environmental health perspectives* 126, 077003.
- 1
2 De Petris S., Berretti R., Sarvia F., Borgogno-Mondino E., 2019. Precision arboriculture: a new
3 approach to tree risk management based on geomatics tools, in: *SPIE Remote Sensing*, 2019.
4 Presented at the Remote Sensing for Agriculture, Ecosystems, and Hydrology XXI, SPIE.
5 <https://doi.org/10.1117/12.2532778>
- 6
7 Egorov, A.I., Griffin, S.M., Converse, R.R., Styles, J.N., Sams, E.A., Wilson, A., Jackson, L.E., Wade,
8 T.J., 2017. Vegetated land cover near residence is associated with reduced allostatic load and
9 improved biomarkers of neuroendocrine, metabolic and immune functions. *Environmental*
10 *Research* 158, 508–521. <https://doi.org/10.1016/j.envres.2017.07.009>
- 11 European Space Agency/Centre National d’Eudes Spatiales (CNES), 2019. 3rd Sentinel-2 Validation
12 Team Meeting - 12-14 March 2019 - Toulouse, France . Toulouse, France, p. 30.
- 13 Fagerholm, N.C., Broberg, A., 2011. Mapping and characterising children’s daily mobility in urban
14 residential areas in Turku, Finland. *Fennia-International Journal of Geography* 189, 31–46.
- 15 Finardi, S., Pellegrini, U., 2004. 6.13 SYSTEMATIC ANALYSIS OF METEOROLOGICAL
16 CONDITIONS CAUSING SEVERE URBAN AIR POLLUTION EPISODES IN THE
17 CENTRAL PO VALLEY.
- 18 Fong, K.C., Hart, J.E., James, P., 2018. A Review of Epidemiologic Studies on Greenness and Health:
19 Updated Literature Through 2017, Current environmental health reports. NLM (Medline).
20 <https://doi.org/10.1007/s40572-018-0179-y>
- 21
22 Fuertes, E., Markevych, I., Berg, V., 2014. Greenness and allergies: evidence of differential associations
23 in two areas in Germany. *J Epidemiol Community Health*. <https://doi.org/10.1136/jech>
- 24 Gao, B.-C., 1996. NDWI—A normalized difference water index for remote sensing of vegetation liquid
25 water from space. *Remote sensing of environment* 58, 257–266.
- 26 Garipey, G., Kaufman, J.S., Blair, A., Kestens, Y., Schmitz, N., 2015. Place and health in diabetes: the
27 neighbourhood environment and risk of depression in adults with Type 2 diabetes. *Diabetic*
28 *Medicine* 32, 944–950. <https://doi.org/10.1111/dme.12650>
- 29
30 Gong, Y., Palmer, S., Gallacher, J., Marsden, T., Fone, D., 2016. A systematic review of the relationship
31 between objective measurements of the urban environment and psychological distress.
32 *Environment International* 96, 48–57. <https://doi.org/10.1016/j.envint.2016.08.019>
- 33 Hammer, Ø., Harper, D.A., Ryan, P.D., 2001. PAST: paleontological statistics software package for
34 education and data analysis. *Palaeontologia electronica* 4, 9.
- 35 Huete, A., 1988. Huete, AR A soil-adjusted vegetation index (SAVI). *Remote Sensing of Environment*.
36 *Remote sensing of environment* 25, 295–309.
- 37 James, P., Banay, R.F., Hart, J.E., Laden, F., 2015. A review of the health benefits of greenness. *Current*
38 *epidemiology reports* 2, 131–142.
- 39
40 Li, F., Song, G., Liujun, Z., Yanan, Z., Di, L., 2017. Urban vegetation phenology analysis using high
41 spatio-temporal NDVI time series. *Urban Forestry & Urban Greening* 25, 43–57.
- 42 Li, X., Zhang, C., Li, W., Ricard, R., Meng, Q., Zhang, W., 2015. Assessing street-level urban greenery
43 using Google Street View and a modified green view index. *Urban Forestry & Urban Greening*
44 14, 675–685.
- 45 Lovasi, G.S., O’Neil-Dunne, J.P.M., Lu, J.W.T., Sheehan, D., Perzanowski, M.S., MacFaden, S.W.,
46 King, K.L., Matte, T., Miller, R.L., Hoepner, L.A., Perera, F.P., Rundle, A., 2013. Urban Tree
47 Canopy and Asthma, Wheeze, Rhinitis, and Allergic Sensitization to Tree Pollen in a New York
48 City Birth Cohort. *Environmental Health Perspectives* 121, 494–500.
49 <https://doi.org/10.1289/ehp.1205513>
- 50
51 Lovasi, G.S., Quinn, J.W., Neckerman, K.M., Perzanowski, M.S., Rundle, A., 2008. Children living in
52 areas with more street trees have lower prevalence of asthma. *Journal of Epidemiology &*
53 *Community Health* 62, 647–649. <https://doi.org/10.1136/jech.2007.071894>
- 54 Manes, F., Blasi, C., Salvatori, E., Capotorti, G., Galante, G., Feoli, E., Incerti, G., 2012. Natural
55 vegetation and ecosystem services related to air quality improvement: tropospheric ozone
56 removal by evergreen and deciduous forests in Latium (Italy). *Annali di Botanica* 2, 79–86.
- 57 Manes, Fausto, Incerti, G., Salvatori, E., Vitale, M., Ricotta, C., Costanza, R., 2012. Urban ecosystem
58 services: tree diversity and stability of tropospheric ozone removal. *Ecological Applications* 22,
59 349–360.
60
61
62
63
64
65

- 1 Marando, F., Salvatori, E., Fusaro, L., Manes, F., 2016. Removal of PM10 by forests as a nature-based
2 solution for air quality improvement in the Metropolitan city of Rome. *Forests* 7, 150.
- 3 Markevych, I., Schoierer, J., Hartig, T., Chudnovsky, A., Hystad, P., Dzhambov, A.M., de Vries, S.,
4 Triguero-Mas, M., Brauer, M., Nieuwenhuijsen, M.J., Lupp, G., Richardson, E.A., Astell-Burt,
5 T., Dimitrova, D., Feng, X., Sadeh, M., Standl, M., Heinrich, J., Fuertes, E., 2017. Exploring
6 pathways linking greenspace to health: Theoretical and methodological guidance.
7 *Environmental Research* 158, 301–317. <https://doi.org/10.1016/j.envres.2017.06.028>
- 8 Mavoia, S., Oliver, M., Witten, K., Badland, H.M., 2011. Linking GPS and travel diary data using
9 sequence alignment in a study of children's independent mobility. *International Journal of*
10 *Health Geographics* 10, 64.
- 11 McMorris, O., Villeneuve, P.J., Su, J., Jerrett, M., 2015. Urban greenness and physical activity in a
12 national survey of Canadians. *Environmental research* 137, 94–100.
- 13 Milne, G.L., Dai, Q., Roberts, L.J., 2015. The isoprostanes - 25 years later. *Biochimica et Biophysica*
14 *Acta - Molecular and Cell Biology of Lipids* 1851, 433–445.
15 <https://doi.org/10.1016/j.bbali.2014.10.007>
- 16 Mudele, O., Gamba, P., 2019. Mapping vegetation in urban areas using Sentinel-2, in: 2019 Joint Urban
17 Remote Sensing Event (JURSE). IEEE, pp. 1–4.
- 18 Nikolaidis, M.G., Kyparos, A., Vrabas, I.S., 2011. F2-isoprostane formation, measurement and
19 interpretation: The role of exercise. *Progress in Lipid Research* 50, 89–103.
20 <https://doi.org/10.1016/J.PLIPRES.2010.10.002>
- 21 Nowak, D.J., Crane, D.E., Stevens, J.C., 2006. Air pollution removal by urban trees and shrubs in the
22 United States. *Urban forestry & urban greening* 4, 115–123.
- 23 Ormsby, J.P., Choudhury, B.J., Owe, M., 1987. Vegetation spatial variability and its effect on vegetation
24 indices. *International Journal of Remote Sensing* 8, 1301–1306.
- 25 Parmes, E., Pesce, G., Sabel, C.E., Baldacci, S., Bono, R., Brescianini, S., D'Ippolito, C., Hanke, W.,
26 Horvat, M., Lieder, H., Maio, S., Marchetti, P., Marcon, A., Medda, E., Molinier, M., Panunzi,
27 S., Pärkkä, J., Polańska, K., Prud'homme, J., Ricci, P., Snoj Tratnik, J., Squillacioti, G., Stazi,
28 M.A., Maesano, C.N., Annesi-Maesano, I., 2020. Influence of residential land cover on
29 childhood allergic and respiratory symptoms and diseases: Evidence from 9 European cohorts.
30 *Environmental Research* 183, 108953. <https://doi.org/10.1016/j.envres.2019.108953>
- 31 Pereira, G., Foster, S., Martin, K., Christian, H., Boruff, B.J., Knuiman, M., Giles-Corti, B., 2012. The
32 association between neighborhood greenness and cardiovascular disease: an observational
33 study. *BMC Public Health* 12, 466. <https://doi.org/10.1186/1471-2458-12-466>
- 34 Pilat, M.A., McFarland, A., Snelgrove, A., Collins, K., Waliczek, T.M., Zajicek, J., 2012. The effect of
35 tree cover and vegetation on incidence of childhood asthma in metropolitan statistical areas of
36 Texas. *HortTechnology* 22, 631–637.
- 37 Revel, C., Lonjou, V., Marcq, S., Desjardins, C., Fougny, B., Coppolani-Delle Luche, C., Guillemot,
38 N., Lacamp, A.-S., Lourme, E., Miquel, C., 2019. Sentinel-2A and 2B absolute calibration
39 monitoring. *European Journal of Remote Sensing* 52, 122–137.
- 40 Romanazzi, V., Pirro, V., Bellisario, V., Mengozzi, G., Peluso, M., Pazzi, M., Bugiani, M., Verlatto, G.,
41 Bono, R., 2013. 15-F2t isoprostane as biomarker of oxidative stress induced by tobacco smoke
42 and occupational exposure to formaldehyde in workers of plastic laminates. *Science of The*
43 *Total Environment* 442, 20–25. <https://doi.org/10.1016/j.scitotenv.2012.10.057>
- 44 Rosina, K., Kopecká, M., 2016. Mapping of urban green spaces using Sentinel-2A data: Methodical
45 aspects, in: 6th International Conference on Cartography and GIS, Albena. Bulgarian
46 Cartographic Association (in Print). pp. 562–568.
- 47 Sawidis, T., Krystallidis, P., Veros, D., Chettri, M., 2012. A study of air pollution with heavy metals in
48 Athens city and Attica basin using evergreen trees as biological indicators. *Biological trace*
49 *element research* 148, 396–408.
- 50 Sbihi, H., Tamburic, L., Koehoorn, M., Brauer, M., 2015. Greenness and Incident Childhood Asthma:
51 A 10-Year Follow-up in a Population-based Birth Cohort. *American journal of respiratory and*
52 *critical care medicine* 192, 1131–3. <https://doi.org/10.1164/rccm.201504-0707LE>
- 53 Sies, H., 2015. Oxidative stress: a concept in redox biology and medicine. *Redox biology* 4, 180–3.
54 <https://doi.org/10.1016/j.redox.2015.01.002>

- 1 Small, C., Lu, J.W., 2006. Estimation and vicarious validation of urban vegetation abundance by spectral
2 mixture analysis. *Remote Sensing of Environment* 100, 441–456.
- 3 Song, C., 2005. Spectral mixture analysis for subpixel vegetation fractions in the urban environment:
4 How to incorporate endmember variability? *Remote Sensing of Environment* 95, 248–263.
- 5 Squillaciotti, G., Bellisario, V., Grignani, E., Mengozzi, G., Bardaglio, G., Dalmaso, P., Bono, R.,
6 2019a. The Asti Study: The Induction of Oxidative Stress in A Population of Children
7 According to Their Body Composition and Passive Tobacco Smoking Exposure. *International*
8 *Journal of Environmental Research and Public Health* 16, 490.
9 <https://doi.org/10.3390/ijerph16030490>
- 10 Squillaciotti, G., Bellisario, V., Grosso, A., Ghelli, F., Piccioni, P., Grignani, E., Corsico, A., Bono, R.,
11 2020. Formaldehyde, Oxidative Stress, and FeNO in Traffic Police Officers Working in Two
12 Cities of Northern Italy. *International Journal of Environmental Research and Public Health* 17,
13 1655.
- 14 Squillaciotti, G., Bellisario, V., Levra, S., Piccioni, P., Bono, R., 2019b. Greenness Availability and
15 Respiratory Health in a Population of Urbanised Children in North-Western Italy. *International*
16 *Journal of Environmental Research and Public Health* 17, 108.
17 <https://doi.org/10.3390/ijerph17010108>
- 18 Thiering, E., Markevych, I., Brüske, I., Fuertes, E., Kratzsch, J., Sugiri, D., Hoffmann, B., von Berg, A.,
19 Bauer, C.-P., Koletzko, S., 2016. Associations of residential long-term air pollution exposures
20 and satellite-derived greenness with insulin resistance in German adolescents. *Environmental*
21 *health perspectives* 124, 1291–1298.
- 22 Tillberg Mattsson, K., 2002. Children's (in) dependent mobility and parents' chauffeuring in the town
23 and the countryside. *Tijdschrift voor economische en sociale geografie* 93, 443–453.
- 24 Tsai, W.-L., Leung, Y.-F., McHale, M.R., Floyd, M.F., Reich, B.J., 2019. Relationships between urban
25 green land cover and human health at different spatial resolutions. *Urban Ecosystems* 22, 315–
26 324. <https://doi.org/10.1007/s11252-018-0813-3>
- 27 van 't Erve, T.J., Kadiiska, M.B., London, S.J., Mason, R.P., 2017. Classifying oxidative stress by F2-
28 isoprostane levels across human diseases: A meta-analysis. *Redox Biology* 12, 582–599.
29 <https://doi.org/10.1016/j.redox.2017.03.024>
- 30 Warton, D.I., Wright, I.J., Falster, D.S., Westoby, M., 2006. Bivariate line-fitting methods for allometry.
31 *Biological reviews* 81, 259–291.
- 32 Willmott, C.J., Matsuura, K., 2005. Advantages of the mean absolute error (MAE) over the root mean
33 square error (RMSE) in assessing average model performance. *Climate research* 30, 79–82.
- 34 Wolch, J.R., Byrne, J., Newell, J.P., 2014. Urban green space, public health, and environmental justice:
35 The challenge of making cities 'just green enough.' *Landscape and urban planning* 125, 234–
36 244.
- 37 Woo, J., Tang, N., Suen, E., Leung, J., Wong, M., 2009. Green space, psychological restoration, and
38 telomere length. *Lancet (London, England)* 373, 299–300. [https://doi.org/10.1016/S0140-6736\(09\)60094-5](https://doi.org/10.1016/S0140-6736(09)60094-5)
- 39 Yeager, R., Riggs, D.W., DeJarnett, N., Tollerud, D.J., Wilson, J., Conklin, D.J., O'Toole, T.E.,
40 McCracken, J., Lorkiewicz, P., Xie, Z., Zafar, N., Krishnasamy, S.S., Srivastava, S., Finch, J.,
41 Keith, R.J., DeFilippis, A., Rai, S.N., Liu, G., Bhatnagar, A., 2018. Association Between
42 Residential Greenness and Cardiovascular Disease Risk. *Journal of the American Heart*
43 *Association* 7, e009117. <https://doi.org/10.1161/JAHA.118.009117>
- 44 Zhang, X., Friedl, M.A., Schaaf, C.B., Strahler, A.H., Hodges, J.C., Gao, F., Reed, B.C., Huete, A.,
45 2003. Monitoring vegetation phenology using MODIS. *Remote sensing of environment* 84,
46 471–475.
- 47 Zhou, D., Zhao, S., Zhang, L., Liu, S., 2016. Remotely sensed assessment of urbanization effects on
48 vegetation phenology in China's 32 major cities. *Remote Sensing of Environment* 176, 272–
49 281.

Table 1. Technical features of the S2 images obtained from Theia CNES provider (<https://www.Theia-land.fr/en/product/sentinel-2-surface-reflectance/>)

Band ID	Central wavelength (nm)	Bandwidth (nm)	Nominal Geometric resolution by Theia CNES provider
B2	490	98	10 m
B3	560	45	10 m
B4	665	38	10 m
B5	705	19	10 m (20 m native resolution)
B6	740	18	10 m (20 m native resolution)
B7	775	28	10 m (20 m native resolution)
B8	840	145	10 m
B8A	865	33	10 m (20 m native resolution)
B11	1610	143	10 m (20 m native resolution)
B12	2200	242	10 m (20 m native resolution)

Table 2 – Coefficients of linear regression models between 15-F2t-IsoP vs PGC, IsoP vs NET, and 15-F2t-IsoP vs NBT.

All dependent variables were poorly correlated with OS.

Dependent variable	<i>r</i>	<i>p</i>-value	R²	Slope	<i>p</i>(slope)	Slope (95% C.Is)	Intercept
uSAVI	-0.045	0.520	0.002	-2.214	0.520	(-9.538, 4.685)	11.967
PGC	-0.046	0.519	0.002	-2.126	0.519	(-8.747, 4.819)	12.035
NBT	-0.028	0.691	0.001	-0.001	0.691	(-0.003, 0.002)	11.699
NET	-0.030	0.673	0.001	-0.005	0.673	(-0.025, 0.014)	11.523

Table 3 – Twenty equi-probable classes of 15-F2t-IsoP (divided by about 1.4 ng/mg crea step) and vegetation parameters. In order to calculate the envelope function of each scatterplot, maximum values of equi-probable class were extracted. For each equi-probable class maximum value and coefficient of variation (%) were reported. Slash sign denotes only one value within equi-probable class.

15-F2t-IsoP		uSAVI		NBT		NET		PGC	
Max	CV%	Max	CV%	Max	CV%	Max	CV%	Max	CV%
3.56	39.9	0.47	46.9	990	55.2	102	91.5	0.514	43.5
4.93	10.8	0.42	38.8	1874	58.9	89	84.3	0.487	34.6
6.35	7.7	0.44	37.5	1239	51.7	115	95.8	0.512	34.5
7.52	4.2	0.57	51.6	1016	37.6	113	94.9	0.586	46.5
9.16	5.5	0.63	46.2	1310	48.6	136	100.0	0.619	40.8
10.55	4.2	0.65	45.5	1262	45.8	244	99.4	0.616	39.9
11.85	3.2	0.44	40.3	1144	33.0	115	80.7	0.492	35.9
13.35	3.0	0.50	35.1	1124	30.2	165	91.4	0.497	31.2
14.56	3.2	0.42	45.8	1103	72.6	108	119.2	0.477	43.5
16.05	2.6	0.42	43.1	1144	41.8	115	79.6	0.473	41.1
17.17	1.2	0.44	60.6	923	60.0	117	167.9	0.517	57.6
18.74	2.2	0.47	46.1	1107	50.3	92	108.4	0.484	44.3
20.23	1.8	0.36	43.7	1163	55.5	79	105.8	0.426	42.6
21.18	1.4	0.42	36.7	1111	58.6	79	137.0	0.457	33.1
22.88	1.8	0.41	59.7	1118	88.9	76	95.8	0.471	60.3
24.2	2.7	0.56	20.7	892	2.2	77	51.3	0.575	13.5
26.81	/	0.41	/	995	/	101	/	0.469	/
28.98	0.7	0.34	69.5	1109	52.1	19	/	0.411	66.9
34.34	/	0.11	/	206	/	0	/	0.139	/
39.65	/	0.16	/	170	/	6	/	0.190	/

Table 4 – Bivariate relationships statistics of 15-F2t-IsoP-Ev and vegetation parameters. Significance levels: * p

<0.05 , ** $p <0.01$, *** $p < 0.001$.

Proxy	r	R^2	Slope	Slope (95% C.I.s)	Intercept
PGC	-0.758***	0.57	-62.75***	(-83.671 -34.792)	34.26
uSAVI	-0.717***	0.51	-0.01***	(-0.016, -0.005)	0.59
NET	-0.696***	0.48	-0.12***	(-0.192, -0.023)	30.02
NBT	-0.743***	0.55	-0.02***	(-0.031, -0.010)	39.17

S.D.P.: Conceptualization, Methodology, Software, Formal analysis, Writing - Original Draft; **G.S.:** Conceptualization, Methodology, Software, Formal analysis, Writing - Original Draft, Data curation; **R.B.:** Supervision, Project administration, Writing - Review & Editing; **E.B.M.:** Supervision, Project administration, Writing - Review & Editing

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.