

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**New phases for analytical scale extraction from plants: Current and future trends**

**This is a pre print version of the following article:**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1798336> since 2021-08-28T15:40:29Z

*Published version:*

DOI:10.1016/j.trac.2021.116288

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# 1 New phases for micro-scale extractions from plants: 2 current and future trends

3 Giulia Mastellone<sup>§</sup>, Arianna Marengo<sup>§</sup>, Barbara Sgorbini, Patrizia Rubiolo,  
4 Cecilia Cagliero\*

5 *<sup>1</sup>Dipartimento di Scienza e Tecnologia del Farmaco, Università degli Studi di Torino, I-  
6 10125, Turin, Italy*

## 8 Abstract

9 Analyses in the plant field have to cover a wide range of topics, starting from  
10 metabolomics to the quali-quantitative determination of a single or a class of metabolites,  
11 often dealing with very complex matrices. The development of an appropriate sample  
12 preparation method is fundamental to obtain complete information, because downstream  
13 analyses will detect only the metabolites previously extracted. The choice of the proper  
14 extraction phase is therefore of utmost importance and the most critical parameter to be  
15 considered. Although most of the studies on plants still mainly adopt traditional extraction  
16 techniques, a wide number of new phases with improved features in terms of tunability,  
17 selectivity and, most important, sustainability have been developed in the past decades  
18 and applied to the study of natural products. This review provides an overview of the new  
19 extraction phases developed and adopted for the study of plant endogenous and  
20 exogenous metabolites.

---

22 <sup>§</sup>Both authors contributed equally to this work

23 \*Corresponding author: cecilia.cagliero@unito.it, Via Pietro Giuria, 9, I-10125, Torino, Italy,  
24 Tel: +39-011-6707133

26 **Keywords:** extraction phases, natural products, ionic liquids, deep eutectic solvents, natural  
27 sorbents, molecular imprinted polymers, metal organic frameworks, carbon-based materials,  
28 hybrid materials, green analytical chemistry

30 **Abbreviations:** **ABS:** aqueous biphasic system, **Bio-IL:** bio-renewable IL, **CD:** cyclodextrin,  
31 **ChCl:** choline chloride, **CNT:** carbon nanotube, **COF:** covalent-organic framework, **DES:**  
32 deep eutectic solvent, **DI:** direct immersion, **DLLME:** dispersive liquid-liquid microextraction,

33 **FT-IR**: Fourier transform infrared spectroscopy, **GO**: graphene oxide, **rGO**: reduced graphene  
34 oxide, **HBA**: hydrogen bond acceptor, **HBD**: hydrogen bond donor, **IL**: ionic liquid, **HS**:  
35 headspace, **IL-HIM**: ionic liquid–hybrid molecularly imprinted material, **ILMHDE**: IL-  
36 mediated microwave-assisted hydro distillation liquid-liquid extraction, **ILUMASDE**: ionic  
37 liquid-based ultrasound/microwave-assisted simultaneous distillation extraction, **MAE**:  
38 microwave assisted extraction, **MCE**: mechanochemical extraction, **MNP**: magnetic  
39 nanoparticle, **MIL**: magnetic ionic liquid, **MIP**: molecular imprinted polymer, **MOF**: metal  
40 organic framework, **MSPD**: matrix solid phase dispersion **MWCNT**: multiwalled carbon  
41 nanotube, **NADES**: natural deep eutectic solvent, **PANI**: polyaniline, **PCB**:  
42 polychlorobiphenyl, **PDMS**: polydimethylsiloxane, **PIL**: polymeric ionic liquid, **PPy**:  
43 polypyrrole, **PSA**: primary-secondary amine, **PVC**: polyvinyl chloride, **QuEChERS**: Quick,  
44 Easy, Cheap, Effective, Rugged, and Safe, **UAE**: ultrasound assisted extraction, **SLE**: solid-  
45 liquid extraction, **SPME**: solid phase microextraction, **SPE**: solid phase extraction, **dSPE**:  
46 dispersive SPE , **μ-dSPE**: miniaturized version of dispersive SPE , **m-SPE**: magnetic solid  
47 phase extraction, **ToF**: time of flight , **TRIL**: temperature-responsive hydrophobic ionic liquid,  
48 **UCTS**: upper critical solution temperature, **UHPLC**: ultra-high performance liquid  
49 chromatography.

50

51

---

52 **1. Introduction**

53 Natural product analyses must cover a wide range of topics, starting from the study  
54 of plant metabolomes to the quality control and determination of endogenous and  
55 exogenous compounds. In both cases, sample preparation plays a fundamental role. In  
56 fact, all plant metabolites should be extracted in metabolomics studies. However, a  
57 universal extraction phase or technique is practically impossible to find because of the  
58 complexity of plant systems and the chemical diversity of metabolites [1]. On the other  
59 hand, the determination of single or multiple metabolites, which are often present at trace  
60 levels in plant materials, is required in quality control analyses. The analytical challenge  
61 here is the development of methods that can detect target compound(s) at low limits, as  
62 often required by regulations, meaning that proper enrichment is required [2].

63 The development of appropriate sample preparation methods in the plant field faces  
64 particular challenges as the complexity of plant biological systems can affect the reliable  
65 determination of target analytes. In fact, the metabolome of a plant comprises a high  
66 variety of compounds, ranging from different classes of primary metabolites (sugars,  
67 proteins, fats and nucleic acids) to a very wide range of specialized metabolites (such as  
68 terpenoids, alkaloids, phenolic compounds and others). Another issue in plant analysis is  
69 the presence of chlorophylls and other pigments; these molecules are present in high  
70 amounts in the photosynthetic parts of the plant, but poorly compatible with downstream  
71 analyses, and with HPLC systems in particular, because of their strong interaction with  
72 commonly used reverse stationary phases. At the same time, it is important to remember  
73 that plant cells are protected by thick and robust lignocellulosic walls that should be  
74 disrupted if the efficient extraction of intracellular metabolites is to be achieved.

75 A number of mechanical means, including ultrasound and microwaves, can be  
76 adopted to increase the efficiency of metabolite extraction. However, the choice of the  
77 appropriate extraction phase is of utmost importance, and is the most critical parameter

78 to be considered. The extraction phase should either cover the widest possible array of  
79 metabolites in metabolomics studies, or increase the selectivity and the enrichment of  
80 target analytes in studies dealing with the determination of specific compounds.

81 In this sense, most studies on plants still principally adopt traditional extraction  
82 techniques, including liquid-liquid extraction and Soxhlet extraction, that entail the high  
83 consumption of toxic organic solvents. However, the use of miniaturizing techniques and,  
84 in particular, of new classes of more sustainable extraction phases is gaining ground as  
85 they meet the criteria of Green Analytical Chemistry [3]. Furthermore, these techniques  
86 can be tailored to increase extraction selectivity towards specific compounds [4,5].

87 This review aims to provide an overview of the extraction phases that have been  
88 developed and adopted, over the last three years (from 2017 to 2020), for the study of  
89 endogenous and exogenous plant metabolites. A critical evaluation of the main features  
90 of each phase will be provided as well as some ideas for new applications in the plant  
91 field (see also figure 1 and table 1).

92 The first part of this review deals with the evaluation of new green extraction  
93 solvents (ionic liquids and deep eutectic solvents) in the liquid-phase extraction of natural  
94 products, while the second part is devoted to the evaluation of innovative solid sorbents  
95 for the clean-up and pre-concentration of plant extracts. The potential, limits and future  
96 trends of each extraction phase will be highlighted in both cases.

97

## 98 **2. Liquid extraction phases**

99 The choice of extraction solvent is of utmost importance when dealing with the  
100 extraction of plant metabolites, and the search for new green solvents has become one of  
101 the most significant subjects in the sample preparation field. In this respect, ionic liquids  
102 (ILs) have received a great deal of attention as they may be able to replace organic  
103 solvents in the extraction of natural products. Moreover, deep eutectic solvents (DESS)

104 are also gaining position in the field of plant sample preparation. In fact, these “green”  
105 solvents can be combined with a range of extraction techniques, such as ultrasound  
106 assisted extraction (UAE) and microwave-assisted extraction (MAE), and can be  
107 exploited in liquid-phase microextractions.

108 In this section, the potential of using ILs and DESs as liquid extraction phases in  
109 the plant field will be presented and critically discussed together with their possible  
110 weaknesses. Figure 1b provides a representation of the structure of the liquid extraction  
111 phases that have been investigated, while table 2 reports a number of representative  
112 applications.

113

## 114 **2.1 Ionic Liquids**

115 Ionic liquids (ILs) are an interesting class of molten salts whose physicochemical  
116 properties have inspired the rapid development of new applications in analytical  
117 chemistry and, in particular, in the sample-preparation field [6]. They are organic salts  
118 that possess melting points at or below 100 °C. In most cases, they are composed of an  
119 organic cation and an organic or inorganic anion, and are characterized by high thermal  
120 stability, low vapor pressure and varying viscosity, conductivity and miscibility in  
121 different solvents.

122 One of the main features of ILs, compared to traditional organic solvents, is their  
123 ability to incorporate functional groups, within the IL structure, that selectively interact  
124 with specific analytes or a class of analytes. The chemical structure of ILs (type of  
125 functional groups and/or anion/cation combination) can be tailored to provide various  
126 solvation interactions (such as hydrogen bonding, electrostatic, hydrophobic and  $\pi$ - $\pi$   
127 interactions) and capabilities with both polar and apolar compounds, thus giving ILs with  
128 very different properties in terms of water solubility and solution behavior [6,7]. In

129 addition, since they present negligible vapor pressure at room temperature, they prevent  
130 solvent loss to the atmosphere and decrease the environmental footprint, thus making  
131 them a sustainable alternative to organic solvents.

132 Many studies have also demonstrated the ability of ionic liquids to dissolve  
133 cellulose and other cell-wall components (such as hemicellulose and lignin), via the  
134 decrystallization of the cellulose portion of lignocellulosic biomass, and simultaneously  
135 disruption of the lignin and hemicellulose network [8,9]. In fact, the anions of ionic  
136 liquids can form strong hydrogen bonds with the hydroxyl groups of cellulose, and thus  
137 eliminate their intramolecular hydrogen bonds and disrupt cell walls, causing cell lysis.  
138 This ability can be exploited for the pre-treatment of plant material before its extraction,  
139 for instance before or during hydrodistillation, at laboratory scale, to increase essential  
140 oil yield [10,11].

141 The above-mentioned properties have allowed ILs to find many applications as  
142 solvents in extraction and separation processes for bioactive compounds, either in simple  
143 solid-liquid extractions (SLE) or in microwave-assisted (MA) and ultrasound-assisted  
144 (UA) extractions [5,12]. In particular, because of their high polarity and high thermal  
145 stability, ILs are heated rapidly and uniformly under microwave irradiation being  
146 considered as excellent microwave absorbers. They can also be used in UAE, in particular  
147 for extracting thermolabile compounds in short times and at room temperature [5].

148 Because of their versatility, ILs can be adopted in liquid-phase extractions of natural  
149 products under different extraction modes. They can be used as surfactants above their  
150 critical micellar concentration, as reported by Mastellone *et al.* in a study that exploited  
151 1-hexadecyl-3-butyl imidazolium bromide as IL surfactant for the extraction of phenolic  
152 compounds from *Vitis vinifera* L. leaves [13], and in another work by Moučková *et al.*  
153 that evaluated a set of structurally different IL-based surfactants for the flavonoid profile

154 determination of *Mangifera* spp. and *Passiflora* spp. leaves [14]. They can also form  
155 aqueous biphasic systems (ABS) when the aqueous IL solution is mixed with inorganic  
156 salts that promote phase separation and increase analyte(s) enrichment. This approach has  
157 been adopted, for instance, for the extraction of eight ginsenosides from the flower buds  
158 of *Panax ginseng* C.A.Mey; in this study, the extraction of the analytes has been carried  
159 out by aqueous IL-based UAE and it has been followed by the separation of the IL phase  
160 with an ABS, using NaH<sub>2</sub>PO<sub>4</sub>, and by the direct analysis by HPLC-UV [15]. Finally,  
161 dispersive liquid-liquid microextraction (DLLME) can be used when dealing with liquid  
162 samples, such as herbal teas. In this case, hydrophobic ILs are dispersed in the water  
163 solution with the aid of a dispersive solvent and the IL-phase is recovered and analyzed.  
164 In the IL-DLLME procedure, ultrasound energy can be applied to accelerate the mass  
165 transfer process and to improve the extraction efficiency. In addition, the solubility of  
166 target analytes in ILs can be optimized by changing the temperature. An ultrasonic-  
167 assisted temperature-controlled IL-DLLME approach has recently been coupled to  
168 HPLC-UV and applied for the determination of pyrethroid residues in herbal teas [16].

169 ILs can also incorporate a paramagnetic element into their structure, either in the  
170 cation or in the anion giving rise to the magnetic ionic liquids (MILs), a sub-class of ILs  
171 whose application in sample preparation is rapidly increasing [6,7]. MILs have found  
172 some applications in the plant field, in particular in the extraction of sinomenine from  
173 *Sinomenium acutum* (Thunb.) Rehder & E.H.Wilson [17] and in the extraction, adopting  
174 an ABS approach, of berberine from the rhizome of *Coptis* spp [18]. They also have been  
175 used for the isolation of DNA from *Arabidopsis thaliana* (L.) Heynh. and other plants  
176 [19]. These compounds have great potential for the extraction of endogenous and  
177 exogenous metabolites from vegetable matrices because they combine all of the features  
178 of ILs with the ability to exploit their paramagnetic characteristics that allow them to be



179 easily separated using an external magnetic field, in a very rapid process that avoids  
180 centrifugation and filtration steps. MILs therefore deserve to be taken into more serious  
181 consideration when the extraction phase for the sample preparation of plant matrices is  
182 selected.

183 ILs can also be incorporated in solid phase platforms, either alone (as polymeric  
184 ionic liquids, PILs) or in combination with other sorbents to form composites. PILs are  
185 polyelectrolytes obtained through the polymerization of IL monomers. PILs not only  
186 retain the main features of ILs, but also present the inherent properties of polymers with  
187 enhanced mechanical stability and conductivity properties. Thus, they are the best option  
188 to exploit the outstanding properties of ILs in sorbent-based microextraction strategies.  
189 For instance, they can be used in solid phase microextraction (SPME) as sorbent coatings  
190 directly bonded and crosslinked to rugged supports. PIL-based SPME fibers can be  
191 applied in both direct immersion (DI) and headspace (HS) extraction modes and can be  
192 coupled to both HPLC and GC analysis. For instance, PIL coatings have found use in the  
193 determination of acrylamide in brewed coffee and coffee powder by DI-SPME [20,21]  
194 and in the profiling of the volatile compounds from *Echinacea* spp. flowers by HS-SPME  
195 [22], in both cases coupled to GC-MS analysis.

196 In addition, a number of IL-based composites have been developed to exploit the  
197 unique properties of ILs in sorbent-based strategies, such as in solid-phase extraction  
198 (SPE) [23]. In the plant field, for instance, a bionic multi-tentacled ionic liquid-modified  
199 silica gel has been synthesized and used as a selective adsorbent in the separation of tea  
200 polyphenols from green tea [24]. Ionic liquids and  $\beta$ -cyclodextrin functionalized  
201 magnetic graphene oxide materials have been synthesized for the extraction of plant  
202 growth regulators in vegetable samples that were pre-extracted with 1% acetic acid in  
203 acetonitrile [25]. ILs have also been widely employed in combination with natural

204 sorbents, molecular imprinted polymers (MIPs) and carbon-based materials; these  
205 composites will be better described in section 3.

206         Thanks to the tunability of their structure and, consequently, of their properties, ILs  
207 can be applied to both metabolomics studies and the selective extraction of specific  
208 endogenous and exogenous compounds. When dealing with metabolomics studies, ILs  
209 can be exploited for the simultaneous determination of volatile and non-volatile  
210 components [26-28]. Different approaches for the determination of these classes of  
211 compounds in plant matrices have recently been developed. In the first case, an efficient  
212 IL-mediated microwave-assisted hydrodistillation has been concatenated with liquid-  
213 liquid extraction (ILMHDE) for the distillation of essential oil and the simultaneous  
214 extraction of a set of non-volatile flavonoids (astragalin, quercetin, luteolin, kaempferol  
215 and apigenin) from the inflorescences of *Helichrysum arenarium* (L.) Moench. An IL  
216 solution is added to the plant material and submitted to microwave-assisted  
217 hydrodistillation, and the hydrosol is then collected in two separation columns, the first  
218 of which recovers the volatile components, while the second collects the non-volatile  
219 compounds after the addition of an organic solvent. [27]. ILs can also be used in  
220 combination with enzyme digestion, as clearly shown in a study in which a solution of 1-  
221 butyl-3-methylimidazolium bromide and a mixture of cellulase and pectinase have been  
222 used as pretreatment before the simultaneous extraction, through a modified  
223 hydrodistillation concatenated Soxhlet extraction apparatus, of procyanidins and essential  
224 oil from the pinecones of *Pinus koraiensis* Siebold & Zucc. [28]. Finally, an ionic liquid-  
225 based ultrasound/microwave-assisted simultaneous distillation extraction (ILUMASDE)  
226 approach can be used. Again, an IL-water solution is added to the plant material and  
227 submitted to UAE and/or MAE but, in this case, the non-volatile components are directly  
228 determined in the solvent solution, while the volatile components are collected after the

229 condensation of the vapor phase. This approach has been successfully applied to the  
230 determination of the flavonoids and volatile components in sea buckthorn (*Elaeagnus*  
231 *rhamnoides* (L) A. Nelson) leaves, and has provided a drastic reduction in extraction  
232 times, higher analyte recovery and similar volatile composition, compared to  
233 conventional hydrodistillation. [26]. The ability to extract hydrophilic and hydrophobic  
234 non-volatile compounds at the same time is also very interesting. For instance, an ionic-  
235 liquid-based ultrasonic-assisted extraction has been used to simultaneously extract  
236 gingerols and polysaccharides from ginger (*Zingiber officinale* Roscoe) with higher  
237 yields and reduced extraction times for both classes of compound [29]. The use of  
238 temperature-responsive hydrophobic ionic liquids (TRILs) for the simultaneous  
239 extraction of hydrophilic and hydrophobic compounds, followed by their separation and  
240 pre-concentration at lower temperatures, is another remarkable alternative. In fact, TRILs  
241 are soluble in water above a certain temperature (the upper critical solution temperature,  
242 UCST) and immiscible below the UCST. As illustrated in figure 2b, the extraction process  
243 is carried out at high temperature, exploiting the different polarities of the two solvents,  
244 and then the *in-situ* separation and pre-concentration of the target compounds is achieved  
245 by reducing the temperature below the UCST, meaning that the extraction solvent will  
246 switch from being single-phase to two-phase. This approach has been developed  
247 specifically for the extraction of bioactive compounds from medicinal plants, and has  
248 been applied to *Chrysanthemum* spp., as a case study [30].

249 On the other hand, the structure of ILs can be tuned for the selective recovery of  
250 plant metabolites belonging to specific chemical classes As previously mentioned, a  
251 cholinium-based MIL has been exploited for the selective extraction of berberine from  
252 the rhizome of *Coptis* spp [18] but numerous other examples can be found in literature,  
253 as reported in a recent review by Xiao et al [31]. Furthermore, ILs can be exploited for

254 the selective extraction and pre-concentration of toxic exogenous compounds. As already  
255 reported, an IL-based SPME and an IL-based *in-situ* DLLME method have been  
256 developed for the extraction of acrylamide from coffee [20,21,32] and an IL-based  
257 DLLME approach has been adopted for the selective extraction and determination of  
258 pyrethroid residues in herbal teas [16].

259 Despite the undoubted advantages of IL-based extractions, these molecules also  
260 have some limits. One drawback is the difficulty in recovering extracted compounds  
261 because of the negligible vapor pressure of ILs. This limit mostly affects their use in the  
262 isolation of natural compounds, and less their analytical application. However, the poor  
263 volatility of ILs also means that they are not directly compatible with LC-MS analyses or  
264 the direct injection into a GC system; as a result, a back-extraction of the analytes from  
265 the IL phase is required before their analysis, when these techniques are adopted. The  
266 difficulty in recycling and recovering ILs on an industrial scale is also a critical issue.

267 In addition, although they were originally considered to be green chemicals, some  
268 concerns about the toxicity and potential environmental impact of ILs has started to arise  
269 in recent years. In fact, most of them are obtained from fossil sources and show poor  
270 degradability. In this respect, the current trend is to design ILs to be more degradable and  
271 less toxic. In particular, ILs with fluorine-free anions and with more biodegradable  
272 cations (e.g. pyridinium or guanidinium) have been developed, in part thanks to the fact  
273 that shorter IL side chains in any of the moieties give less toxic ILs. More eco-friendly  
274 guanidinium-based ILs have been applied, for instance, in the above-mentioned  
275 extraction of flavonoids from *Mangifera* spp. and *Passiflora* spp. leaves [14]. These  
276 derivatives, although more environmentally sustainable, still require a large amount of  
277 non-renewable solvents in their synthesis, meaning that the use of anionic or cationic  
278 counterparts that are fully or partly derived from natural compounds would therefore be

279 an ideal and sustainable approach to the development of a new generation of bio-  
280 renewable ILs. (Bio-ILs) [33]. For instance, matrinium-based ionic liquids, derived from  
281 the alkaloid matrine, isolated from *Sophora flavescens* Aiton, have optimal  
282 physicochemical properties and low cytotoxicity [34]. Unfortunately, these Bio-ILs have  
283 rarely been applied in sample preparation and no applications can be found in the plant  
284 field. Future advancements in the development of bio-renewable ionic liquids are  
285 therefore expected and their application in the extraction of plant matrices should be  
286 explored.

287

## 288 **2.2. Deep eutectic solvents (DES)**

289 Deep eutectic solvents are systems that are formed by a hydrogen-bond acceptor  
290 (HBA) and a hydrogen-bond donor (HBD). As the term “eutectic” suggests, they are  
291 characterized by a lower melting point than the individual components. They are usually  
292 associated with ILs because they have similar physicochemical properties, including low  
293 vapor pressure, low flammability and thermal stability. However, DES can be prepared  
294 by incorporating ionic or neutral species, unlike ILs (see paragraph 2.1). Moreover, DESs  
295 overcome some of the limits of ILs due to their easy preparation, the low cost of their  
296 synthetic raw materials, as well as their lower toxicity and higher biodegradability [35–  
297 38]. Since their introduction by Abbot and colleagues in 2003 [39], they have gained  
298 popularity in the extraction field thanks to the extraction capacities they present, which  
299 are related to H-bonding interactions, polarity and viscosity. The large number of HBAs  
300 and HBDs that are available for the preparation of DESs allows many combinations of  
301 eutectic mixtures to be easily obtained with different types of extraction performance and  
302 characteristics that are suitable for a range of matrices.

303 In this sense, DESs have been widely explored as valid alternatives to organic  
304 solvents for the extraction of bioactive compounds from different plant materials, with a  
305 focus on non-volatile endogenous compounds [40]. Of the various components that can  
306 be used to obtain a DES, HBAs and HBDs of natural origin, such as amines, sugars,  
307 carboxylic acids and polyalcohols, are an attractive source for the preparation of  
308 biodegradable and biocompatible DES. These natural deep eutectic solvents (NADES)  
309 have found promising applications in the plant field, according to current trends towards  
310 the use of green solvents [36,38,41–45]. The solubility of most natural compounds in  
311 NADES, which is due to multiple types of interactions including hydrogen bonding,  
312 dispersion forces and electrostatic interactions [35], can be explained by a hypothesis  
313 developed by Choi *et al.* [46] who proposed that NADES are naturally present in plant  
314 cells and act as a third liquid phase in living organisms. In fact, they observed that some  
315 primary metabolites (sugars, organic acids and bases, amino acids) naturally occur in  
316 similarly high molar ratios in the plant cells, thus probably explaining the multistep  
317 biosynthesis of non-water-soluble compounds. In addition, several studies have  
318 demonstrated the ability of DESs to cleave lignin-carbohydrate complexes by disrupting  
319 covalent linkages and hydrogen bonding, thus affecting the structure of the cell wall and  
320 accelerating the release of the compounds [35].

321 The heating and stirring method is the most common of the procedures employed  
322 to prepare DES for the analysis of plants. It normally requires the mixing of two or three  
323 components under magnetic stirring, and keeping the temperature at 40-90°C until a  
324 homogeneous liquid is produced. [36,38,41–44,47]. When preparing a DES, it is  
325 important that hydrogen bonds are formed between the two components (HBA and HBD),  
326 and an FT-IR spectrum of the mixture can therefore be a useful tool to evaluate their  
327 correct formation [42,47]. Moreover, the DES instability over time can lead to the  
328 formation of crystals in the mixture, and a visual inspection is a simple way to evaluate

329 whether the correct procedure and HBA/HBD ratio were used to obtain a stable  
330 preparation. For instance, Kaltsa *et al.* [44] have examined several lactic-acid-based DESs  
331 for the presence of crystals, at regular intervals over 6 weeks. A range of different  
332 physicochemical properties were observed according to the HBA and HBD chosen, and  
333 the ratio used. For example, the widely used HBA choline chloride (ChCl) can be  
334 combined with polyols (e.g. ethylene glycol, glycerol...), carboxylic acids, amides,  
335 alcohols, sugars and amino acids to give mixtures with different polarities and viscosities  
336 [36,38,42,43]. As a result, the higher viscosity of sugar-based DES (e.g Choline chloride-  
337 D-glucose) led to a lower extraction capacity for flavonoid glycosides and aglycones from  
338 the Chinese herbal medicine Platycladi Cacumen (*Platycladus orientalis* (L.) Franco),  
339 compared to other DESs [43]. Viscosity is a key point when dealing with DES because it  
340 can affect the mass transfer of analytes during extraction. The main ways of reducing  
341 viscosity include higher extraction temperatures, which may result in the degradation of  
342 the target compounds if not properly controlled, and the addition of water [43]. The  
343 presence of water clearly affects the polarity of the solvent. Consequently, it is not  
344 surprising that glycosides are more influenced by water content than aglycones, even  
345 though the extraction of the more hydrophilic glycosides does not improve with higher  
346 percentages of water. In fact, the correlation between analyte solubility and water content  
347 is not linear, and the careful optimization of this parameter is fundamental to achieve  
348 good performance, also considering that the presence of water may also interfere with the  
349 formation of the hydrogen bonding network [35,41].

350       Ultrasounds [28,30,33,35,36,38,39] or microwaves [36] are often employed during  
351 an extraction to enhance the transfer of analytes from the vegetal matrix to the eutectic  
352 solvent, resulting in a greener approach due to the decrease of extraction time and energy  
353 consumption [5]. However, prolonged or excessive heating are generally not  
354 recommended as they can lead to the degradation of thermally unstable analytes, and also

355 of some HBDs and HBAs [35]. Yao *et al.* have compared the performance of heat-stirring,  
356 microwave and ultrasound extraction in the analysis of phenol compounds from *Pyrola*  
357 *asarifolia* subsp. *incarnata* (DC.) Haber & Hir. Takah, and microwave-assisted extraction  
358 led to better results thanks to its fast and focused heating [36].

359 A further aspect in the use of DES, that meets the Green Chemistry principles, is  
360 the absence of harmful organic solvents, both in the synthesis of DES and during the  
361 extraction process, unlike other sorbent-assisted extractions (e.g MWCNTs, MOFs...).

362 This is a key point in the plant field. As previously mentioned, the complex composition  
363 of vegetal materials, including the presence of pigments (e.g. chlorophyll), usually means  
364 that a pre-treatment process is needed to eliminate these interferences, and large quantities  
365 of harmful organic solvents, such as MeOH, EtOH, acetone and hexane, are consumed.

366 On the contrary, the common procedure with DES involves a solid-liquid extraction of  
367 the dried plant at 25-70°C in hydrophilic media. The percentage of water added to the  
368 eutectic mixture prevent the transfer of chlorophyll and other apolar interferences from  
369 the plant material to the extract [36,43,44,47,49]. However, it is important to highlight  
370 that a certain volume of solvent is normally used to dilute the sample before the analysis,  
371 this step is necessary to decrease the viscosity of the final extract in order to prevent  
372 damage to the chromatographic system [38,41,43,47,49]. Nevertheless, as discussed for  
373 ILs, the direct injection of the DESs in the chromatographic system when LC is coupled  
374 to MS is not possible because of their low volatility. The recovery of the target analytes  
375 from the eutectic mixture is therefore mandatory for LC-MS analyses. This latter involves  
376 the use of macroporous resins [41,43,50] or antisolvents that disrupt the hydrogen  
377 bonding between the two components of the DES [35,40]. In this sense, Fu *et al.* [42]  
378 performed a purification step by solid phase extraction (SPE) using a DES-modified  
379 adsorbent and a ChCl-based DES for the elution of polyphenols from palm bark samples  
380 (*Trachycarpus fortunei* (Hook.) H.Wendl), obtaining an enrichment of the analytes



381 without using organic solvents and expensive SPE phases. Moreover, the same volume  
382 of DES and water was eluted through the SPE column, achieving a proper dilution of the  
383 final extract.

384 As previously mentioned, DESs have found many applications in the plant field,  
385 mainly because of their ability to extract a very wide range of compounds. In particular,  
386 they have been extensively applied for the extraction of phenolic compounds that have  
387 been shown to dissolve better in DESs than in lipids or water [51]. Aside from phenolic  
388 compounds, other plant specialized metabolites have been extracted using DES. In  
389 particular, Duan *et al.* [38] have thoroughly investigated the behavior of several DES  
390 species in the extraction of various bioactive compounds, including alkaloids, saponins  
391 and anthraquinones. For the alkaloids, carboxylic acid-based DES (e.g. betaine laevulinic  
392 acid) resulted to be a more efficient solvent, than MeOH, less suitable for extracting  
393 partially ionized compounds. On the other hand, acid-based DESs are not suitable for the  
394 extraction of saponins, and better results were obtained when using dimethylurea or  
395 methylurea as HBD. Finally, anthraquinones were poorly extracted by the majority of the  
396 hydrophilic DES as a result of the low polarity of these compounds. In general, the study  
397 demonstrated that sugar- and alcohol-based DESs had a lower extraction capacity than  
398 MeOH towards low-polarity analytes. In this respect, hydrophobic DESs can be used to  
399 extend the application of these solvents to the analysis of poorly polar compounds as the  
400 properties of DESs can be tailored as a function of HBA and HBD structure. For instance,  
401 a HBA consisting of methyl trioctyl ammonium chloride combined with 1-propanol in a  
402 1:4 ratio has been successfully applied for the extraction of artemisin, which is an apolar  
403 terpenoid used in malaria treatment, from *Artemisia annua* L. [52]. A new generation of  
404 hydrophobic NADES has also recently been developed by mixing menthol with different  
405 HBDs [53]. The combination of menthol and 1-propanol has been used for the isolation  
406 of taxanes from *Taxus wallichiana* var. *chinensis* (Pilg.) Florin (syn. *Taxus chinensis*

407 (Pilg.) Rehder) needles, and was found to be more efficient than conventional solvents,  
408 such as MeOH or EtOH [47]. Despite the apolar nature of taxanes, the DES with 80%  
409 water content showed better extraction efficiency than a common hydroalcoholic solvent.  
410 It could also boast of less solvent consumption, confirming that the presence of water  
411 plays a critical role when using DES. The same HBA has been combined with different  
412 carboxylic acids in an efficient extraction of four cannabinoids from *Cannabis sativa* L.  
413 inflorescences. [48]. It is interesting to highlight that the extraction yield was not only  
414 affected by the viscosity of the mixture, but also by the different alkyl chain lengths of  
415 the HBD used, which may have improved interactions with more lipophilic compounds.  
416 NADESs (together with ILs) have also been investigated in the extraction of ecdysteroids,  
417 and, in particular, 20-hydroxyecdysone, from spinach (*Spinacia oleracea* L.) and several  
418 advantages were found, including higher selectivity, simplicity, efficiency, and rapidity,  
419 as well as the recovery and much lower consumption of organic solvents, in comparison  
420 with conventional methods [54]. In addition, it has been shown that DESs are also able  
421 to extract volatile compounds. In fact, the headspace SPME of the DES extracts of  
422 peppermint leaves and tobacco powder shows a richer volatile profile than that of  
423 conventional methods [55,56].

424 Very interestingly, DESs are also able to preserve the bioactivity of the compounds.  
425 Several studies have demonstrated the ability of DESs to stabilize the antioxidant capacity  
426 of natural products [35], as shown, for instance, for the choline chloride-based extract of  
427 phenolic compounds from *Rosmarinus officinalis* L., which provided higher antioxidant  
428 capacity and lower degradation kinetics than the corresponding ethanolic extract [57].  
429 This feature opens the perspective of the direct use of NADES extracts in the health field.  
430 In fact, NADESs composed by HBAs and HBDs that are commonly introduced in our  
431 daily diet (such as sugars, amino acids or citric acids) can likely be directly used for food  
432 and pharmaceutical purposes [41]. However, current studies rarely investigate the

433 bioactivity, biotoxicity and metabolism of these derivatives. In this sense, Zhao *et al* [58]  
434 have tested the biocompatibility and biodegradability of a series of DESs that were used  
435 for the extraction of rutin from *Styphnolobium japonicum* (L.) Schott. Bacterial growth  
436 inhibition was higher for organic acid-based DES than for alcohol- or sugar-based DES,  
437 although the antimicrobial effects of all of the tested DESs were lower than those of some  
438 conventional ILs. In addition, the biodegradability level of the organic acid-based DES  
439 was the lowest, although its value was still within acceptable limits.

440         The outstanding features of this innovative class of solvents have led to increasing  
441 interest in a number of applications, in particular in the plant field. The psychochemical  
442 characteristics and environmentally friendly profile of DES perfectly match the principles  
443 of Green Analytical Chemistry, including the replacement of harmful organic solvents,  
444 and the natural origin of the majority of the HBAs and HBDs available. At the same time,  
445 however, it is important to highlight that there is a need to overcome some drawbacks in  
446 their use. These disadvantages are mainly related to their viscosity, which can negatively  
447 affect extraction time (up to 2 hours), extraction capacity and compatibility with the main  
448 analytical platforms, if not properly controlled. These future challenges may provide the  
449 basis for new perspectives in the use of DESs.

450

### 451 **3. Solid sorbents**

452         A wide range of new sorbents has recently been developed for sample preparation.  
453 Several of them have been applied in the plant field, in particular for the clean-up and  
454 pre-concentration of specific analytes from water and organic extracts. In general, they  
455 show good potential in terms of efficiency and sustainability. However, it should be noted  
456 that plant matrices are commonly solids, as are the sorbents, meaning that previous  
457 sample extraction has to be carried out to allow a solid phase extraction with the sorbents.  
458 In most cases, it is performed with organic solvents, and this decreases the sustainability

459 of the whole process. However, greener alternatives are available in the form of matrix  
460 solid-phase dispersion extraction (MSPD), and via pre-extraction with the green solvents  
461 described above.

462 In this section, the potential of the main classes of solid sorbents that have been  
463 exploited in the extraction of natural products will be discussed, together with their main  
464 limits and future trends. Figure 1b reports an illustration of the structures of the  
465 investigated sorbents and table 2 reports some representative applications.

466

### 467 **3.1. Natural sorbents**

468 Natural sorbents are certainly the greenest alternative to conventional materials in  
469 extraction procedures, and they comply with the increasing concern about environmental  
470 sustainability and safety. Apart from their low ecological impact, these sorbents are  
471 cheap, compared with other extraction phases, and easily available from natural sources.  
472 They include sorbents with a range of characteristics, but the common principle is the  
473 exploitation of natural products as raw materials for the preparation of sustainable and  
474 efficient extraction phases. Over the years, many natural sorbents have been demonstrated  
475 to possess good extraction capacities together with suitable characteristics for extraction,  
476 such as mechanical strength, gelling properties and tunability [59–61].

477 According to literature, two main classes have been explored as sorbent phases in  
478 plant extraction: i) biopolymers; and ii) food by-products. The first are isolated from  
479 living organisms and consist of monomeric units (e.g. nucleotides, monosaccharides,  
480 amino acids) that are covalently bonded to form a polymeric chain. Polysaccharides have  
481 undergone the most widespread explorations in the extraction of endogenous and  
482 exogenous compounds from plants. Chitosan and cellulose have been used in different  
483 formats, including as dispersants [62–64], SPE cartridges [65,66], nanoparticles [67,68],  
484 and fibers [69], thanks to their versatility. Unlike cellulose, which is abundant in the walls

485 of vegetal cells, chitosan is mainly derived from the skeletons of marine animals, as a  
486 deacylated form of chitin. Both materials present surfaces that are rich in hydrophilic  
487 groups, which can easily form hydrogen bonding and electrostatic interactions with target  
488 analytes [59]. Sporopollenin, a major component of the external walls of pollen grains, is  
489 another interesting biopolymer. It has gained attention as a sorbent material for its  
490 impressive resistance to high temperatures and chemical treatments. However, it has been  
491 less thoroughly explored than other [61]. Its adsorption capacity comes from the presence  
492 of several different functional structures on its surface, including hydroxylic groups and  
493 saturated and unsaturated aliphatic chains with aromatics. In this regard, Qian Lu and  
494 colleagues have tested pollen grains as SPE sorbents for the determination of plant growth  
495 regulators and *trans* resveratrol in different vegetal samples, and have reported that this  
496 biopolymer presents excellent separation and purification properties [70,71].

497 The use of food by-products as sorbent phases in the extraction of vegetable  
498 matrices has been documented in the literature, and some interesting applications have  
499 been proposed [72–74]. The valorization of biomass that is normally discharged as food  
500 waste is an added value for this type of sorbent in view of the circular economy paradigm.  
501 For example, Jiang *et al.* have used crab shells from a local restaurant as the starting  
502 materials for a chitosan-rich powder that has been used for the isolation of anthraquinones  
503 from *Senna* spp. (syn. *Cassia*) [72]. Food by-products can also be used to obtain carbon-  
504 based materials, and thus avoid the use of fossil-fuel based precursors. In two different  
505 works, sorbents that were obtained from fruit peels have been able to extract pesticides  
506 from vegetal complex samples. Hierarchical porous carbon from banana peel presents a  
507 large surface that is rich in graphitized pores, and these pores are capable of selective  
508 interactions with the aromatic groups of carbamate pesticides [73]. In a second work,  
509 magnetic nanoparticles with a carbon structure have been prepared from the hydrothermal

510 carbonization of pomelo peel, and the FT-IR spectra showed that the surface was rich in  
511 oxygen-containing groups [74].

512 A further point that must be taken into account when attempting to develop a  
513 greener extraction approach, is to consider all the step of the procedure, and not only one  
514 aspect, such as the sorbent. Thus, the decrease in consumption of toxic solvents and  
515 energy, the production of waste and the reusability of the sorbent should all be considered.  
516 In this sense, the elimination of use of toxic solvents, which are usually necessary to  
517 desorb the analytes, is a true challenge when the extraction phase is a solid sorbent.  
518 However, as mentioned before, microextraction techniques can help to reduce solvent  
519 consumption or green solvents (e.g. ILs) can be employed as adequate substitutes [72].  
520 An evaluation of the reusability of a sorbent is also an important factor in reducing  
521 analysis costs and waste production. In general, many of these natural sorbents can be re-  
522 used several times [69,70,74]. However, Li *et al.* [68] have observed that their cellulose-  
523 MIP microspheres showed reduced adsorption capacity, by more than 10%, after five  
524 cycles of extraction. This may have been due to a loss in adsorption cavitation during the  
525 adsorption-desorption process. On the other hand, the performance of graphene oxide  
526 functionalized cotton fibers was found to be almost unaltered after 50 extraction  
527 procedures, and the regeneration process only entailed sonication in a MeOH/ACN  
528 solution for 5 minutes [69].

529 The use of natural sorbent as such should be preferable in order to develop green  
530 methods, simplify the process and reduce the number of tedious synthetic steps that are  
531 normally required when numerous phases are involved. In many examples, natural  
532 materials exhibit excellent extraction capacities thanks to their specific functional groups  
533 and porous structures, which allow proper interactions with the target analytes without  
534 further functionalization [62–64,66,70,73]. On the other hand, the functionalization  
535 and/or combination of different sorption mechanisms can enhance extraction

536 performance, and a number of works have developed hybrid phases starting from natural  
537 sorbents. For example, the incorporation of magnetic particles into a sorbent material is  
538 a well-known method with which to simplify extraction procedures. In this way, the  
539 sorbent can be easily collected from the plant sample with a magnet, and thus further  
540 separation steps are avoided [67,68,74,75]. Some interesting applications have combined  
541 solid-phase extraction with IL-based extraction for a range of different purposes. The  
542 liquid phase can act as a green solvent for the elution of analytes from a natural sorbent  
543 [72] or in a pre-extraction step, followed by extraction with the sorbent [67]. In this latter  
544 case, the comparison with an organic solvent extraction and the only ionic-liquid  
545 extraction without sorbent enrichment, clearly showed that the IL-chitosan combination  
546 had a higher capacity for the extraction of natural pigments from the rhizomes of  
547 *Curcuma longa L.*

548 Various extraction techniques have been developed for the isolation of compounds  
549 from plant samples, and the versatility of these natural materials can grant significant  
550 advantages. Several SPE cartridges have been prepared with chitosan [65], pollen [70]  
551 cotton [69] and treated food by-products [73], with these being simply packed into a  
552 cartridge and the compounds of interest eluted with a suitable organic solvent. Matrix  
553 solid-phase dispersion (MSPD) is an extraction method in which a sample is directly  
554 ground and mixed with a solid-phase sorbent and the analytes are subsequently isolated,  
555 via adsorption onto a SPE cartridge, and eluted with a certain volume of solvent. One of  
556 the main advantages of MSPD is its ability to directly extract solid materials, which is  
557 particularly important for plant samples that normally require pre-treatment before  
558 extraction. As shown in figure 2e, cellulose and chitosan can act as optimal dispersants  
559 and sorbents in MSPD, and have been successfully employed for the extraction of phenols  
560 [63] triterpenoid acids [62], anthraquinones [72] and caffeoylquinic acid derivatives [64]  
561 from various vegetal samples. Er *et al.* [75] and Xu *et al.* [67] have coupled natural

562 sorbents with a miniaturized version of magnetic dispersive SPE (m- $\mu$ -dSPE) to develop  
563 a truly sustainable procedure. The method only required a small amount of chitosan  
564 nanoparticles, which were dispersed into the liquid sample, and a few microliters of  
565 organic solvent to elute the compounds from the sorbent.

566 These natural materials have been designed to improve the environmental  
567 friendliness of extraction approaches, but also to enhance the performance of analytical  
568 procedures. Comparisons with common approaches have shown similar or better results  
569 in terms of sample and solvent amount, extraction time, LOD and RSD in many examples  
570 [62,66,67,72–74].

571

### 572 **3.2. Molecular imprinted polymers (MIPs)**

573 Molecular imprinted polymers (MIPs) are a peculiar group of extraction materials  
574 with unique “mimic molecular recognition” properties. Interestingly, an interaction site  
575 with high affinity for a chosen template is formed during the synthesis of this polymeric  
576 matrix. The high selectivity of MIPs is therefore based on the geometry of this cavity and  
577 on the presence of specific functional groups that are complementary to the target analyte  
578 [76,77].

579 Regarding extraction in the plant field, MIPs are arising as a valid approach when  
580 the goal is to isolate target compounds from the rich phyto-complexes that are typical in  
581 plants. Indeed, most extraction phases lack high selectivity for a specific analyte or class  
582 of compounds, meaning that pretreatment and/or purification steps for analyte enrichment  
583 are usually necessary before extraction. The isolation of exogenous compounds (e.g.  
584 antibiotics, pollutants and toxins) from food or environmental samples is one of the most  
585 highly developed applications of MIPs in analytical chemistry [77–79]. The use of MIPs  
586 in plant analyses follows a different trend, that is the extraction of endogenous bioactive



587 compounds, including molecules with widely differing chemical structures, thanks to the  
588 possibility of obtaining tailor-made MIPs [80–86].

589         When a new material is prepared, an accurate characterization of its morphology  
590 (e.g. using scanning electron microscopy, IR spectra) is necessary to verify that the  
591 synthesis has occurred correctly. This is especially true with MIPs, since the synthetic  
592 protocol involves several reagents and steps. In particular, a monomer interacts with the  
593 template (i.e. molecule with the same or similar chemical features as the target analytes)  
594 to form the specific cavity that characterizes every MIP. An initiator induces the  
595 polymerase chain reaction, while the porogen assures the formation of pores that facilitate  
596 interaction with the target analyte during extraction. Finally, a cross-linker ensures the  
597 formation of a rigid structure, which must be preserved after the removal of the template,  
598 at the end of the polymerization process. Normally, MIPs with different composition/ratio  
599 for each component are prepared to find the optimal procedure. The choice of the reagents  
600 to use (especially the monomer) clearly depends on the template, and therefore on the  
601 target compound to be analyzed. For example, coumarins were easily able to form  
602 hydrogen-bonding and ionic interactions with carboxylic groups when methacrylic acid  
603 was tested as a functional monomer [80]. Also, the correct selection of the porogen  
604 solvent seems to play a key role in the recognition ability of MIPs, and can directly  
605 influence their morphology. In this respect, chloroform, acetonitrile and toluene have  
606 been documented to be the most effective [80,82,83]. Moreover, many polymerization  
607 techniques have been developed over the years, including emulsion, precipitation and  
608 bulk polymerization, which is the most commonly used [76,78,79]. According to *Gao et*  
609 *al.*, the precipitation method is the most suitable for the synthesis of spherical  
610 nano/microparticles and has been successfully applied in the preparation of nanosized-  
611 MIPs, for the extraction of tiliroside from *Edgeworthia gardneri* (Wall.) Meisn [82]. Ionic  
612 liquids can be used as porogenic solvents when preparing MIPs as they form discrete

613 polar and non-polar microenvironments that lead to an enhanced interaction between the  
614 template and functional monomer, and limit nonspecific binding. In addition, ILs have  
615 been shown to accelerate polymerization and improve the effect of imprinting. This  
616 strategy has been applied for the isolation of corilagin, a bioactive ellagitannin, from a  
617 *Phyllanthus urinaria* L. water extract: corilagin-MIP monoliths were prepared using 4-  
618 vinylpyridine as a functional monomer, and ethylene glycol dimethacrylate as the cross-  
619 linking monomer, using a mixture of 1-butyl-3-methylimidazoliumtetrafluoroborate  
620 (ionic liquid) and N,N-dimethylformamide-dimethyl sulfoxide as a porogen [87]. Ionic  
621 liquids can also be used in combination with MIPs to form hybrid materials that combine  
622 the selectivity of MIPs with the ion-exchanging and electrostatic-attraction ability of ionic  
623 liquids. An IL-hybrid molecularly imprinted material (IL-HIM) has, in fact, been used as  
624 a sorbent for the solid-phase extraction of two plant-growth regulators, namely 6-  
625 benzyladenine (6-BA) and 4-chlorophenoxyacetic acid (4-CPA) from bean sprouts,  
626 coupled to HPLC analysis. The hybrid material is suitable for the simultaneous  
627 determination of the two target analytes in bean sprouts, but accurate selectivity and  
628 competitive experiments showed that the adsorption of 6-BA onto IL-HIM is based on  
629 selective imprinted recognition, whereas the adsorption of 4-CPA is mainly dependent on  
630 ion-exchange interactions [88].

631 While MIPs have mostly been associated with SPE, the primary goal in their use is  
632 currently the incorporation of miniaturized extraction techniques to improve the accuracy  
633 and precision of the method and, at the same time, decrease sample and sorbent  
634 consumption [89]. However, the combination of MIPs and microextraction methods in  
635 the plant field has yet to be thoroughly investigated. Some SPME coatings have recently  
636 been developed and successfully used for the enrichment of low concentration bioactive  
637 compounds, such as auxins [83] and for the rapid detection of the potentially toxic  
638 pyrrolizidine alkaloids [86]. In the first example, a pre-treated silica fiber was inserted

639 into a prepolymer solution of the MIP, and the glass tube was transferred to an oven at  
640 60°C for 8 hours for the polymerization step. It was reported that the porogen affected the  
641 thickness of the coating, and therefore the adsorption capacity. In particular, the authors  
642 observed that the repetition of the polymerization cycle on the same fiber for up to five  
643 times assured the maximum extraction rate by increasing the binding sites on the fiber.  
644 Moreover, the MIP coating had good chemical stability, which was demonstrated by  
645 immersing the fiber in different organic solvents for a couple of hours. A different  
646 approach was employed for the preparation of the pyrrolizidine alkaloid-MIP, where the  
647 quartz fiber was first treated with the template and the monomer, while the crosslinker  
648 was added subsequently. In this case, monocrotaline was chosen as the dummy template  
649 to simulate the alkaloid structure as the standard materials for these latter structures are  
650 quite expensive. Most importantly, if the template is not properly eliminated it can be  
651 released during extraction, causing false positive results in the sample. For this reason,  
652 the use of the target analyte as template should be carefully handled. Both of the SPME-  
653 MIP applications showed high selectivity for the selected analytes and better sensitivity,  
654 in terms of LOD, than conventional extraction techniques.

655 A further application for MIPs as electrochemical sensors, which find some  
656 applications in plant analysis, is worth mentioning. Again, several procedures have been  
657 used to prepare the MIP-functionalized electrodes. Alipour *et al.* have prepared a sensor  
658 that is based on magnetic MIPs for the determination of rosmarinic acid from four plant  
659 extracts (*Salvia officinalis* L., *Zataria multiflora* Boiss., *Mentha longifolia* (L.) L., and  
660 *Rosmarinus officinalis* L.). The magnetic MIP has been mixed with graphene powder to  
661 obtain a carbon electrode. The magnetic component improves the kinetic rate of electron  
662 transfer, while the MIP increase the selectivity of the system for the target analyte. The  
663 plant material has been treated with different organic solvents before the analysis and the  
664 voltammograms has been recorded at 0.0-0.35 V [84]. Yang and colleagues used the same

665 principle to prepare a portable and cheap sensor for the rapid determination of gallic acid  
666 in edible plants. In this case, MIP-based sensor has been placed in a PVC tube and insert  
667 in an aqueous solution of the sample, as represented in figure 2d [85]. pH dependency  
668 was investigated in both examples as it has an influence on the electrochemical behavior  
669 of the molecules. Indeed, very acidic pH values reduced the carboxylic ionization of gallic  
670 acid, while the upper pH range led to the degradation of the compound.

671         Apart from selectivity, versatility is one of the main advantages of MIPs. Their  
672 characteristics, including durability over a wide range of pH values and temperatures,  
673 make them suitable for innovative applications as extraction phases. A recent review by  
674 Arabi *et al.* gives a comprehensive overview of the new trends in MIP-based SPE,  
675 including nano-scale MIPs, composite MIPs, hollow porous MIPs and many others [78].  
676 Again, the application of these novelties in plant field have been less frequently explored  
677 compared to other application fields. However, progress on this is being made. The  
678 preparation of surface-imprinting MIPs that are based on boron nitride and deep eutectic  
679 solvents for the isolation of flavonoids, and that give a high specific surface area of  
680 extraction is an interesting example [90].

681         Although MIPs show outstanding features, there are still some critical points.  
682 Current and future effort will undoubtedly be directed towards the improvement of these  
683 aspects. In particular, the synthesis of MIPs requires many steps and reagents, including  
684 high amounts of toxic solvents to remove the template. The incorporation of green  
685 materials/solvents should be considered as a means to avoid the large-scale consumption  
686 of hazardous substances. The synthesis of water-compatible MIPs is another hot topic,  
687 because the presence of water can negatively affect non-covalent binding with the analyte,  
688 decreasing interactions with it [77]. However, some water-compatible MIPs, that can be  
689 directly used in aqueous media, have been developed. These phases are particularly  
690 important in the plant field as vegetal matrices are normally pre-treated with polar

691 solvents to give hydrophilic extracts. Finally, the high selectivity of MIPs can be  
692 unfavorable when the goal is to isolate a class of compounds and not a single analyte.  
693 Again, this is particularly true for plant extraction, in which specialized metabolites are  
694 numerous and heterogeneous. Moreover, compounds from the same chemical class can  
695 present quite different functional groups in their structure, and molecules with low  
696 similarity to the template may have some difficulties interacting with MIP cavities. To  
697 solve these problems, Sun *et al.* have developed an off-line two-dimensional MIP-SPE  
698 for the extraction of ellagitannins from pomegranate [81]. In particular, ellagic acid and  
699 punicalagin MIP SPE columns were prepared and used as the first and second dimensions,  
700 respectively. Different fractions were collected after the elution to study the distribution  
701 of the analytes as a function of the purification procedure. Careful optimization of each  
702 elution step and elution solvent is needed to obtain good enrichment for each ellagitannin.  
703 Moreover, the use of the ellagic tannin MIP as the first dimension is important to improve  
704 the capacity and the recovery of the analytes, while using the punicalagin MIP column as  
705 the second step allows the non-specific adsorption of the ellagic acid step to be overcome,  
706 and good selectivity to be achieved.

707         Although the “molecular key and lock” theory that is behind the functioning of MIP  
708 materials is somewhat dated [91], these extraction phases can be considered valuable and  
709 promising tools in the extraction of natural products, especially when coupled with  
710 innovative approaches such as miniaturization, green chemistry and automation.

711

712

### 713 **3.3. Magnetic nanoparticles (MNPs)**

714         Metallic materials have also found applications in microextraction in the plant field.  
715 They have been used because of their high thermal, mechanical and chemical stability,  
716 and feasible functionalization. Of the various nanoparticles to have been developed, those

717 made from transition metals (Fe, Co and Ni), and from their oxides, and, in particular,  
718 magnetite ( $\text{Fe}_3\text{O}_4$ ), are the most commonly used [92]. Given their superparamagnetic  
719 properties, which permit their easy and quick isolation when applying an external  
720 magnetic field, they have emerged as useful tools for the development of magnetic solid-  
721 phase extraction (m-SPE) protocols that avoid centrifugation and separation steps, as  
722 previously mentioned.

723 Moreover, these magnetic nanoparticles (MNPs) are often modified to give  
724 composite materials with a magnetic core and a functional shell that can provide more  
725 active sites for adsorption, while protecting the magnetic core. These composite materials  
726 are most commonly used for the extraction of plant samples. For instance, cellulose MNPs  
727 have been developed for the extraction and determination of polychlorobiphenyls (PCB)  
728 in juice samples [93] and magnetic  $\beta$ -cyclodextrin (CD) modified graphene oxide  
729 nanoparticles ( $\text{Fe}_3\text{O}_4@\text{SiO}_2/\text{GO}/\beta\text{-CD}$ ) have been applied as a sorbent for m-SPE of plant  
730 growth regulators from vegetable methanolic extracts [94,95]. MNPs that were  
731 functionalized with tetraethyl orthosilicate, 3-(trimethoxysilyl)propyl methacrylate,  
732 polymethacrylate and combinations of the three have very recently been successfully  
733 adopted for the determination of brominated flame retardants in red fruit samples  
734 (strawberries, blueberries, and raspberries) using GC-MS and QuEChERS extraction. The  
735 MNPs were used for the clean-up of the extracts, and proved to have excellent adsorption  
736 ability for pigment matrices and other co-extracts, and to be very easy to recover simply  
737 using an external magnet [96].

738 Very interestingly, MNPs that were coated with an amino-terminated  
739 supramolecular cucurbit-[6]-uril pseudorotaxane motif have been developed for the  
740 extraction of polar compounds and, in particular, salvianolic acids. The investigated  
741 MNPs showed improved adsorption capacity towards the target analytes, compared to the  
742 non-functionalized NPs, while the combination of electrostatic interactions, van der

743 Waals interactions, hydrophilic interactions and hydrogen bonding was thought to be the  
744 driving force behind the interaction between the analytes and sorbent. In addition, the  
745 developed MNPs provided good recyclability and stability. These qualities were  
746 exploited when the sorbent material was successfully adopted for the determination of  
747 four salvianolic acids from *Salvia miltiorrhiza* Bunge root water extracts, showing  
748 excellent recovery and quantification performance [97].

749 Most of the magnetic composite nanoparticles applied in the plant field are  
750 composed by magnetic nanoparticles coated with metal organic frameworks (MOFs) that  
751 are used because of their large surface areas, high porosity, uniform pore sizes and  
752 structural diversity. These composite sorbents will be thoroughly described in the  
753 following section.

754

### 755 **3.4 Reticular materials**

756 Reticular materials are another important group of novel solid sorbents. They are  
757 large crystalline frameworks made up of building blocks linked by either covalent or  
758 coordination bonds, depending on the type of reticular material. These materials are  
759 characterized by impressive porosity, good chemical and thermal stability, and by the  
760 possibility of post-synthetic modification, which makes them suitable sorbents for  
761 microextractions. The large variety in building blocks means that it is possible to prepare  
762 innumerable reticular materials with different selectivities. The most common types of  
763 reticular materials are metal-organic frameworks (MOFs) and covalent-organic  
764 frameworks (COFs), which are mainly differentiated by the organic or inorganic nature  
765 of their building blocks. COFs only include organic molecules with non-metallic light  
766 elements (C, H, O, N, B and Si), which are then linked by covalent bonds that are ordered  
767 in two- or three-dimensional networks, while MOFs are made up of metal ions or metallic  
768 clusters and organic linkers as building blocks, connected by strong coordination bonds

769 [98]. Of the two classes of reticular materials, MOFs have been more widely applied in  
770 the plant field.

771       Mainly used for the extraction of contaminants from plant extracts [99,100], MOFs  
772 have also shown great potential as sorbents in combination with microextraction methods  
773 to extract active compounds from plants and plant products. However, their applications  
774 in this field are still quite limited as only a few studies on this possible use can be found.  
775 For example, a copper-based MOF has been adopted for the selective isolation of  
776 chamazulene, a compound with antiphlogistic effects, from a chamomile hexane extract.  
777 The driving force behind this selectivity is probably the chemical structure of the target  
778 compound, which is different from the other components in the extract, as chamazulene  
779 is the only aromatic compound [101]. A zeolitic imidazolate framework has also been  
780 investigated for use as a model MOF for the selective extraction of 3,4-dihydroxy-8,9-  
781 methylenepterocarpan (an antioxidant flavonoid derivative) from an *Caragana jubata*  
782 (Pall.) Poir ethanolic extract. In this case, an adsorption mechanism study has shown that  
783 the target compound is adsorbed onto the MOF surface, and that the *o*-phenolic hydroxyl  
784 group plays a significant role in the selective adsorption of the investigated flavonoid  
785 [102].

786       As can be seen in the above-cited examples, one of the main advantages of the use  
787 of MOFs in solid-phase extractions is the ability to selectively extract specific analytes  
788 and increase their enrichment. However, it should be noted that the raw plant material has  
789 to be submitted to analyte pre-extraction. This step is usually carried out with organic  
790 solvents, thus decreasing the sustainability of the extraction method. In addition, a back-  
791 extraction with another solvent (usually in low amounts) is necessary to release the target  
792 analytes from the MOF for downstream analytical determinations. In fact, attention must  
793 be paid to the fact that analytes are strongly adsorbed by MOFs, meaning that their  
794 desorption must be carefully optimized.



795 MOFs can be used in MSPD extractions as a very interesting alternative that can  
796 avoid analyte pre-extraction. As previously mentioned, MSPD has the significant  
797 advantages of lower organic-solvent consumption and cost, and of showing, at the same  
798 time, high extraction efficiency and selectivity. Furthermore, the subsequent purification  
799 procedure is very simple and short. MSPD can be used for the extraction and clean-up of  
800 solid, semi-solid and highly viscous samples, thus making it suitable for extractions from  
801 plant matrices. It is followed by the elution of the extracted analytes with a small amount  
802 of a suitable solvent. For instance, a zirconium-based MOF (MOF-808) has been applied  
803 for the MSPD extraction of ginsenosides from *Panax ginseng* C.A. Meyer. In detail, the  
804 ginseng leaves were directly ground with MOF-808 and the mixture was packed between  
805 two Teflon frits in a glass cartridge. Only two hundred microliters of methanol were used  
806 for the elution and the collected filtrate was analyzed using UHPLC-ToF-MS. This  
807 protocol gave higher extraction efficiency, simpler operation, a cleaner extract and lower  
808 organic-reagent consumption than a traditional extraction [103].

809 Besides the use of MOFs alone, the current trend is their combination with other  
810 materials to obtain (nano)composites. As previously mentioned, MOFs are very  
811 commonly combined with magnetic nanoparticles to increase the magnetic susceptibility.  
812 The combination of magnetic Fe<sub>2</sub>O<sub>4</sub> nanoparticles with a zinc-based metal organic  
813 framework (MOF-5) has been successfully applied for the extraction of colchicine, a  
814 bioactive alkaloid, from autumn and spring *Colchicum* spp. root ethanolic extracts.  
815 Extraction was combined with ultrasound sonication that provided an incremental mass  
816 transfer and the sorbent was easily separated from the sample matrix by an external  
817 magnetic field. Colchicine sorption occurs by combination of various mechanisms such  
818 as ion exchange, hydrogen bonding, electrostatic,  $\pi$ - $\pi$ , and dipole-ion interaction. In  
819 particular, ionic bonding between the negatively charged functional groups of colchicine  
820 and the positively charged centers of both Zn<sup>2+</sup> in MOF and Fe<sup>3+</sup> in Fe<sub>2</sub>O<sub>4</sub> may be the

821 main reason behind the fast and high colchicine adsorption on this hybrid phase [104].  
822 SiO<sub>2</sub>@Fe<sub>3</sub>O<sub>4</sub> magnetic nanoparticles have been anchored onto an aluminum-based MOF  
823 (MIL53) and used for the extraction of aflatoxin B1 from several herbal teas via magnetic  
824 SPE coupled with spectrofluorimetric determination (see figure 2c) [105]. Furthermore,  
825 SiO<sub>2</sub>@Fe<sub>3</sub>O<sub>4</sub> magnetic nanoparticles have also been grafted onto a titanium-based MOF  
826 for the extraction of caffeic acid from methanolic/water extracts and coupled to HPLC-  
827 UV analysis [106]. Magnetic metal-organic-framework nanoparticles can also be adopted  
828 to speed-up the mechanochemical extraction (MCE) of analytes from plant samples. MCE  
829 is a room-temperature process that disrupts biomolecular lipid layers and cell walls using  
830 a multi-directional, mechanical in-liquid collision between the sample and specialized  
831 ceramic beads to achieve the complete release of target compounds, while preserving  
832 their chemical integrity, into the surrounding solvents. This technique can be combined  
833 with m-SPE with Fe<sub>3</sub>O<sub>4</sub>@MOF, meaning that the extraction of the analytes and their  
834 enrichment can be performed in one single step with the very fast removal of the  
835 extraction phase using a simple magnet. This approach has been successfully applied in  
836 the determination of organochlorine pesticides in tea leaves, but also shows great  
837 potential for other applications in the plant field [107].

838 Combining MOF with silica nanoparticles is another interesting possibility. As an  
839 example, a chromium-based MOF (MIL-101), combined with polyaniline (PANI), which  
840 was used as a conducting polymer, has been doped with silica nanoparticles and used for  
841 the efficient solid-phase extraction of thymol and carvacrol from *Mentha longifolia* (L.)  
842 L. plant extracts and a honey sample [108].

843 In addition, MOFs can be coated onto different carrier substrates. For instance, a  
844 zirconium-based MOF has been coated onto cotton fibers thanks to modifications to the  
845 cellulosic surface of the cotton substrate with carboxyl groups, which serve as active sites  
846 for the growing Zr-MOF. The sorbent, packed into a pipette tip, has been successfully

847 used for the extraction of phenoxy herbicides from cucumber-water extracts [109]. MOFs  
848 have also been coated onto polymeric monolithic columns and adopted for the on-line  
849 SPE and determination of ursolic acid from several ethanolic plant extracts [110,111] and  
850 of aristolochic acid from methanolic plant extracts [112].

851 MOFs have also recently been grafted onto SPME fibers [113], as a means to couple  
852 the simplicity and performance of the technique with the previously mentioned  
853 outstanding properties of these materials. However, their use in the plant field has not yet  
854 been widely explored, but future applications are expected.

855

### 856 **3.5. Carbon-based materials**

857 Carbon-based materials also present exceptional physical and chemical properties,  
858 such as versatility, large surface areas with delocalized  $\pi$ -electrons, and easily modifiable  
859 surfaces. They constitute a group of different hexagonal structures of  $sp^2$  hybridized  
860 carbon atoms. A single layer of these materials is called graphene. This carbon allotrope,  
861 together with graphene oxide (GO), and reduced graphene oxide (rGO), are the most  
862 representative two-dimensional forms. The difference between them lies in the presence  
863 of oxygen functional groups on their surface and their water solubility, with graphene and  
864 rGO that are non-polar and hydrophobic materials with high affinity towards carbon ring  
865 structures, while GO is hydrophilic. The three-dimensional forms of graphene are single  
866 or multiple wall carbon nanotubes (CNTs). CNTs share the same exceptional properties  
867 as graphene and its derivatives, but with a few differences as the inner walls of these  
868 materials are blocked by steric hindrance [114].

869 Both two and three-dimensional carbon-based materials have been applied in the  
870 plant field although graphene and GO are rarely used alone because they are prone to  
871 form aggregates. Ionic liquids can be adopted to prevent the aggregation of graphene,  
872 increasing, at the same time, the ability of  $\pi$ - $\pi$  interactions, ionic exchange, electrostatic

873 interactions and hydrogen bonding between the sorbent and the analytes. An IL-  
874 functionalized graphene adsorbent has, in fact, been developed for the pipette-tip solid-  
875 phase extraction of auxins from soybean sprouts [115].

876 However, graphene and, even more, GO are mainly applied to modify other sorbent  
877 materials to form composites [69,116,117]. For instance, GO has been added to a  
878 composite made of sporopollenin (see paragraph 3.1) and Fe<sub>3</sub>O<sub>4</sub> MNPs to increase the  
879 adsorption capacity and selectivity towards polar compound of the sorbent material. The  
880 composite has been then adopted to the determination of polar organophosphorus  
881 pesticides in several vegetables that had been pre-treated with water and methanol [116].

882 Due to their high hydrophobicity, MWCNTs can be successfully applied for the  
883 clean-up of plant extract. In fact, they can provide very high chlorophyll removal and a  
884 decrease in the matrix effect, therefore making them very useful in HPLC analyses, in  
885 particular when coupled to mass spectrometry [118]. For instance, MWCNTs have been  
886 successfully applied to remove interferences from vegetables submitted to the  
887 QuEChERS procedure for the determination of exogenous compounds as paraben  
888 residues. MWCNTs provided up to a 94% reduction in chlorophyll and lower matrix  
889 effects compared to the commonly used primary-secondary amine (PSA) sorbent thus  
890 making the extracts fully compatible with HPLC–MS/MS analyses [119].

891 MWCNTs can also be combined with other adsorbent materials to form MWCNT  
892 composites that have become promising sorbent materials for SPE in recent years. In  
893 particular, combinations with magnetic nanoparticles increase the number of nanotube  
894 active sites, and provide excellent adsorption capacity. Fe<sub>3</sub>O<sub>4</sub>/MWCNT composites have  
895 been used as SPE sorbents for simultaneous determination by UHPLC-MS/MS of  
896 zearalenone and other mycotoxins in the acetonitrile/water extracts of *Salvia miltiorrhiza*  
897 Bunge roots and ryzhome [120] and for the extraction of a set of 11 chiral pesticides from  
898 several plant extracts [121]. MWCNTs can also be grafted with β-cyclodextrins. In this

899 case, the cyclodextrins increase the absorption capacity of the composite thanks to their  
900 ability to form host–guest inclusion complexes with the target compounds in the inner  
901 cavities of their cone-shaped structures. For instance, multiwalled carbon nanotube-β-  
902 cyclodextrin composites have been incorporated into a poly(butyl methacrylate-ethylene  
903 dimethacrylate) monolith for the online extraction of psoralen and isopsoralen from a  
904 methanolic *Cullen corylifolium* (L.) Medik. (syn. *Psoralea corylifolia* L.) fruit extract  
905 [122].

906 MWCNTs composites incorporated in SPME fibers show a great potential also for  
907 *in-vivo* analyses. A very interesting application has been reported in a study in which a  
908 multiwalled carbon nanotubes/polyaniline-polypyrrole@polydimethylsiloxane  
909 (MWCNTs/PANI-PPy@PDMS) fiber has been prepared by electrodeposition strategy  
910 followed by polymer surface modification. The fiber was then used for the *in-vivo*  
911 extraction of pesticides from garlic by direct contact SPME, as represented in figure 2a  
912 [123].

913

#### 914 **4. Concluding remarks**

915 Although new extraction phases are rarely originally developed for the extraction  
916 of plants and plant products, these samples can represent a good assessment for the  
917 efficiency of the new materials because of the complexity of natural matrices and the  
918 chemical variety of natural compounds.

919 Most of the studies that deal with the analytical characterization of natural products,  
920 in particular for metabolomics purposes, still adopt traditional extraction techniques and  
921 solvents. However, it is possible to note, from a literature survey, that there is a wide  
922 range of new phases that can cover all the range of topics dealing with plant analysis and  
923 increasing, at the same time, the sustainability and the selectivity of the extraction  
924 process.

925 New liquid phases, such as ionic liquids and deep eutectic solvents, can be directly  
926 used for the extraction of analytes from solid plant materials. These green solvents are  
927 significantly more sustainable than organic solvents at a microscale level and, at the same  
928 time, have the tremendous advantage that they can be tailored to the extraction of the  
929 classes and compounds of interest, thus decreasing the number of interfering components.  
930 In this respect, deep eutectic solvents and, in particular, natural deep eutectic solvents  
931 have found a huge number of applications in the extraction of natural samples and have  
932 the potential to replace traditional solvents, at least partly.

933 Furthermore, new solid sorbents also show great potential in the plant field. They  
934 can be used for the clean-up of complex samples as well as to increase enrichment and  
935 selectivity towards specific compounds. In this respect, the use of natural sorbents is a  
936 more sustainable approach for natural-product extraction, and it is expected that new  
937 biocompatible and efficient sorbents will be developed in the near future. The use of  
938 molecular imprinted polymers is still the most common approach to increase the  
939 extraction selectivity. Although this method cannot be considered “new”, the current  
940 trend is to combine these compounds with other extraction phases to produce composites  
941 that improve extraction efficiency and selectivity. In general, the merging of different  
942 classes of sorbents to produce hybrid materials is the current trend in analytical sample  
943 preparation, and the development of new hybrid phases has just started. These hybrid  
944 phases merit thorough investigation in the future, in particular for plant applications.

945

#### 946 **Acknowledgments**

947 The work was financially supported by the ‘Ricerca Locale’ (Ex60%2019) project of the  
948 University of Turin, Turin (Italy). This article is based upon work from the Sample Preparation  
949 Study Group and Network supported by the Division of Analytical Chemistry of the European  
950 Chemical Society.

951

952 **References**

- 953 [1] M.Y. Mushtaq, Y.H. Choi, R. Verpoorte, E.G. Wilson, Extraction for metabolomics:  
954 Access to the metabolome, *Phytochem. Anal.* 25 (2014) 291–306.  
955 <https://doi.org/10.1002/pca.2505>.
- 956 [2] E.V.S. Maciel, A.L. de Toffoli, E.S. Neto, C.E.D. Nazario, F.M. Lanças, New materials  
957 in sample preparation: Recent advances and future trends, *TrAC - Trends Anal. Chem.*  
958 119 (2019). <https://doi.org/10.1016/j.trac.2019.115633>.
- 959 [3] A. Gałuszka, Z. Migaszewski, J. Namieśnik, The 12 principles of green analytical  
960 chemistry and the SIGNIFICANCE mnemonic of green analytical practices, *TrAC -*  
961 *Trends Anal. Chem.* 50 (2013) 78–84. <https://doi.org/10.1016/j.trac.2013.04.010>.
- 962 [4] M. Ramos, A. Jiménez, M.C. Garrigós, Il-based advanced techniques for the extraction  
963 of value-added compounds from natural sources and food by-products, *TrAC - Trends*  
964 *Anal. Chem.* 119 (2019). <https://doi.org/10.1016/j.trac.2019.07.027>.
- 965 [5] F. Chemat, M. Abert-Vian, A.S. Fabiano-Tixier, J. Strube, L. Uhlenbrock, V. Gunjevic,  
966 G. Cravotto, Green extraction of natural products. Origins, current status, and future  
967 challenges, *TrAC Trends Anal. Chem.* 118 (2019) 248–263.  
968 <https://doi.org/10.1016/j.trac.2019.05.037>.
- 969 [6] K.D. Clark, M.N. Emaus, M. Varona, A.N. Bowers, J.L. Anderson, Ionic liquids:  
970 solvents and sorbents in sample preparation, *J. Sep. Sci.* 41 (2018) 209–235.  
971 <https://doi.org/10.1002/jssc.201700864>.
- 972 [7] M.J. Trujillo-Rodríguez, H. Nan, M. Varona, M.N. Emaus, I.D. Souza, J.L. Anderson,  
973 *Advances of Ionic Liquids in Analytical Chemistry*, *Anal. Chem.* 91 (2019) 505–531.  
974 <https://doi.org/10.1021/acs.analchem.8b04710>.
- 975 [8] R.P. Swatloski, S.K. Spear, J.D. Holbrey, R.D. Rogers, Dissolution of cellulose with ionic  
976 liquids, *J. Am. Chem. Soc.* 124 (2002) 4974–4975. <https://doi.org/10.1021/ja025790m>.

- 977 [9] A. Brandt, J. Gräsvik, J.P. Hallett, T. Welton, Deconstruction of lignocellulosic biomass  
978 with ionic liquids, *Green Chem.* 15 (2013) 550–583. <https://doi.org/10.1039/c2gc36364j>.
- 979 [10] G. Flamini, B. Melai, L. Pistelli, C. Chiappe, How to make a green product greener: Use  
980 of ionic liquids as additives during essential oil hydrodistillation, *RSC Adv.* 5 (2015)  
981 69894–69898. <https://doi.org/10.1039/c5ra12649e>.
- 982 [11] D. Lanari, M.C. Marcotullio, A. Neri, A design of experiment approach for ionic liquid-  
983 based extraction of toxic components-minimized essential oil from *Myristica fragrans*  
984 *houutt*, *Molecules.* 23 (2018). <https://doi.org/10.3390/molecules23112817>.
- 985 [12] S.P.M. Ventura, F.A. E Silva, M. V. Quental, D. Mondal, M.G. Freire, J.A.P. Coutinho,  
986 Ionic-Liquid-Mediated Extraction and Separation Processes for Bioactive Compounds:  
987 Past, Present, and Future Trends, *Chem. Rev.* 117 (2017) 6984–7052.  
988 <https://doi.org/10.1021/acs.chemrev.6b00550>.
- 989 [13] G. Mastellone, I. Pacheco-Fernández, P. Rubiolo, V. Pino, C. Cagliero, Sustainable  
990 Micro-Scale Extraction of Bioactive Phenolic Compounds from *Vitis vinifera* Leaves  
991 with Ionic Liquid-Based Surfactants, *Molecules.* 25 (2020).  
992 <https://doi.org/10.3390/molecules25133072>.
- 993 [14] K. Moučková, I. Pacheco-Fernández, J.H. Ayala, P. Bajeroová, V. Pino, Evaluation of  
994 structurally different ionic liquid-based surfactants in a green microwave-assisted  
995 extraction for the flavonoids profile determination of *mangifera* sp. And *passiflora* sp.  
996 leaves from canary Islands, *Molecules.* 25 (2020) 1–17.  
997 <https://doi.org/10.3390/molecules25204734>.
- 998 [15] Q. Liang, J. Zhang, X. Su, Q. Meng, J. Dou, Extraction and separation of eight  
999 ginsenosides from flower buds of *Panax ginseng* using aqueous ionic liquid-based  
1000 ultrasonic-assisted extraction coupled with an aqueous biphasic system, *Molecules.* 24  
1001 (2019) 1–12. <https://doi.org/10.3390/molecules24040778>.
- 1002 [16] R. Zhang, Z. Tan, J. Zhao, Y. Wen, S. Fan, C. Liu, Determination of pyrethroid residues



- 1003 in herbal tea using temperature-controlled ionic liquid dispersive liquid-liquid  
1004 microextraction by high performance liquid chromatography, *Sci. Rep.* 10 (2020) 1–8.  
1005 <https://doi.org/10.1038/s41598-020-61755-z>.
- 1006 [17] Q. Li, S. Wu, C. Wang, Y. Yi, W. Zhou, H. Wang, F. Li, Z. Tan, Ultrasonic-assisted  
1007 extraction of sinomenine from *Sinomenium acutum* using magnetic ionic liquids coupled  
1008 with further purification by reversed micellar extraction, *Process Biochem.* 58 (2017)  
1009 282–288. <https://doi.org/10.1016/j.procbio.2017.04.030>.
- 1010 [18] L.R. Nie, H. Song, A. Yohannes, S. Liang, S. Yao, Extraction in cholinium-based  
1011 magnetic ionic liquid aqueous two-phase system for the determination of berberine  
1012 hydrochloride in: *Rhizoma coptidis*, *RSC Adv.* 8 (2018) 25201–25209.  
1013 <https://doi.org/10.1039/c8ra01745j>.
- 1014 [19] A. Marengo, C. Cagliero, B. Sgorbini, J.L. Anderson, M.N. Emaus, C. Bicchi, C.M.  
1015 Berteà, P. Rubiolo, Development of an innovative and sustainable one-step method for  
1016 rapid plant DNA isolation for targeted PCR using magnetic ionic liquids, *Plant Methods.*  
1017 15 (2019) 1–11. <https://doi.org/10.1186/s13007-019-0408-x>.
- 1018 [20] C. Cagliero, T.D. Ho, C. Zhang, C. Bicchi, J.L. Anderson, Determination of acrylamide  
1019 in brewed coffee and coffee powder using polymeric ionic liquid-based sorbent coatings  
1020 in solid-phase microextraction coupled to gas chromatography-mass spectrometry, *J.*  
1021 *Chromatogr. A.* 1449 (2016) 2–7. <https://doi.org/10.1016/j.chroma.2016.04.034>.
- 1022 [21] C. Cagliero, H. Nan, C. Bicchi, J.L. Anderson, Matrix-compatible sorbent coatings based  
1023 on structurally-tuned polymeric ionic liquids for the determination of acrylamide in  
1024 brewed coffee and coffee powder using solid-phase microextraction, *J. Chromatogr. A.*  
1025 1459 (2016) 17–23. <https://doi.org/10.1016/j.chroma.2016.06.075>.
- 1026 [22] M. Kaya, M. Merdivan, P. Tashakkori, P. Erdem, J.L. Anderson, Analysis of Echinacea  
1027 flower volatile constituents by HS-SPME-GC/MS using laboratory-prepared and  
1028 commercial SPME fibers, *J. Essent. Oil Res.* 31 (2019) 91–98.

- 1029 <https://doi.org/10.1080/10412905.2018.1545707>.
- 1030 [23] A. Gutiérrez-Serpa, P.I. Napolitano-Tabares, J. Šulc, I. Pacheco-Fernández, V. Pino,  
1031 Role of ionic liquids in composites in analytical sample preparation, *Separations*. 7  
1032 (2020) 1–32. <https://doi.org/10.3390/separations7030037>.
- 1033 [24] W. Zhang, X. Feng, Y. Alula, S. Yao, Bionic multi-tentacled ionic liquid-modified silica  
1034 gel for adsorption and separation of polyphenols from green tea (*Camellia sinensis*)  
1035 leaves, *Food Chem.* 230 (2017) 637–648.  
1036 <https://doi.org/10.1016/j.foodchem.2017.03.054>.
- 1037 [25] S. Cao, J. Chen, G. Lai, C. Xi, X. Li, L. Zhang, G. Wang, Z. Chen, A high efficient  
1038 adsorbent for plant growth regulators based on ionic liquid and  $\beta$ -cyclodextrin  
1039 functionalized magnetic graphene oxide, *Talanta*. 194 (2019) 14–25.  
1040 <https://doi.org/10.1016/j.talanta.2018.10.013>.
- 1041 [26] C. Li, J. Zhang, C. Zhao, L. Yang, W. Zhao, H. Jiang, X. Ren, W. Su, Y. Li, J. Guan,  
1042 Separation of the main flavonoids and essential oil from seabuckthorn leaves by  
1043 ultrasonic/microwave-assisted simultaneous distillation extraction, *R. Soc. Open Sci.* 5  
1044 (2018). <https://doi.org/10.1098/rsos.180133>.
- 1045 [27] X. Liu, X. Jing, G. Li, A process to acquire essential oil by distillation concatenated  
1046 liquid-liquid extraction and flavonoids by solid-liquid extraction simultaneously from  
1047 *Helichrysum arenarium* (L.) Moench inflorescences under ionic liquid-microwave  
1048 mediated, *Sep. Purif. Technol.* 209 (2019) 164–174.  
1049 <https://doi.org/10.1016/j.seppur.2018.07.028>.
- 1050 [28] K. Hou, M. Bao, L. Wang, H. Zhang, L. Yang, H. Zhao, Z. Wang, Aqueous enzymatic  
1051 pretreatment ionic liquid–lithium salt based microwave–assisted extraction of essential  
1052 oil and procyanidins from pinecones of *Pinus koraiensis*, *J. Clean. Prod.* 236 (2019).  
1053 <https://doi.org/10.1016/j.jclepro.2019.07.056>.
- 1054 [29] X. Kou, Y. Ke, X. Wang, M.R.T. Rahman, Y. Xie, S. Chen, H. Wang, Simultaneous

- 1055 extraction of hydrophobic and hydrophilic bioactive compounds from ginger (*Zingiber*  
1056 *officinale* Roscoe), *Food Chem.* 257 (2018) 223–229.  
1057 <https://doi.org/10.1016/j.foodchem.2018.02.125>.
- 1058 [30] T. Wang, Q. Wang, P. Li, H. Yang, Temperature-responsive ionic liquids to set up a  
1059 method for the simultaneous extraction and: In situ preconcentration of hydrophilic and  
1060 lipophilic compounds from medicinal plant matrices, *Green Chem.* 21 (2019) 4133–  
1061 4142. <https://doi.org/10.1039/c9gc00995g>.
- 1062 [31] J. Xiao, G. Chen, N. Li, Ionic liquid solutions as a green tool for the extraction and  
1063 isolation of natural products, *Molecules.* 23 (2018).  
1064 <https://doi.org/10.3390/molecules23071765>.
- 1065 [32] C. Zhang, C. Cagliero, S.A. Pierson, J.L. Anderson, Rapid and sensitive analysis of  
1066 polychlorinated biphenyls and acrylamide in food samples using ionic liquid-based in  
1067 situ dispersive liquid-liquid microextraction coupled to headspace gas chromatography,  
1068 *J. Chromatogr. A.* 1481 (2017) 1–11. <https://doi.org/10.1016/j.chroma.2016.12.013>.
- 1069 [33] J. Hulsbosch, D.E. De Vos, K. Binnemans, R. Ameloot, Biobased Ionic Liquids:  
1070 Solvents for a Green Processing Industry?, *ACS Sustain. Chem. Eng.* 4 (2016) 2917–  
1071 2931. <https://doi.org/10.1021/acssuschemeng.6b00553>.
- 1072 [34] Z. Wang, J. Zhang, B. Lu, Y. Li, Y. Liang, J. Yuan, M. Zhao, B. Wang, C. Mai, J. Zhang,  
1073 Novel bio-renewable matrinium-based ionic liquids derived from Chinese herb  
1074 medicine: Synthesis, physicochemical properties and biological activity, *J. Mol. Liq.* 296  
1075 (2019). <https://doi.org/10.1016/j.molliq.2019.111822>.
- 1076 [35] M.Q. Farooq, N.M. Abbasi, J.L. Anderson, Deep eutectic solvents in separations:  
1077 Methods of preparation, polarity, and applications in extractions and capillary  
1078 electrochromatography, *J. Chromatogr. A.* 1633 (2020).  
1079 <https://doi.org/10.1016/j.chroma.2020.461613>.
- 1080 [36] X.H. Yao, D.Y. Zhang, M.H. Duan, Q. Cui, W.J. Xu, M. Luo, C.Y. Li, Y.G. Zu, Y.J. Fu,

- 1081 Preparation and determination of phenolic compounds from *Pyrola incarnata* Fisch. with  
1082 a green polyols based-deep eutectic solvent, *Sep. Purif. Technol.* 149 (2015) 116–123.  
1083 <https://doi.org/10.1016/j.seppur.2015.03.037>.
- 1084 [37] E.L. Smith, A.P. Abbott, K.S. Ryder, Deep Eutectic Solvents (DESs) and Their  
1085 Applications, *Chem. Rev.* 114 (2014) 11060–11082. <https://doi.org/10.1021/cr300162p>.
- 1086 [38] L. Duan, L.L. Dou, L. Guo, P. Li, E.H. Liu, Comprehensive Evaluation of Deep Eutectic  
1087 Solvents in Extraction of Bioactive Natural Products, *ACS Sustain. Chem. Eng.* 4 (2016)  
1088 2405–2411. <https://doi.org/10.1021/acssuschemeng.6b00091>.
- 1089 [39] A.P. Abbott, G. Capper, D.L. Davies, R.K. Rasheed, V. Tambyrajah, Novel solvent  
1090 properties of choline chloride/urea mixtures, *Chem. Commun.* (2003) 70–71.  
1091 <https://doi.org/10.1039/b210714g>.
- 1092 [40] J. Huang, X. Guo, T. Xu, L. Fan, X. Zhou, S. Wu, Ionic deep eutectic solvents for the  
1093 extraction and separation of natural products, *J. Chromatogr. A.* 1598 (2019) 1–19.  
1094 <https://doi.org/10.1016/j.chroma.2019.03.046>.
- 1095 [41] W.W. Oomen, P. Begines, N.R. Mustafa, E.G. Wilson, R. Verpoorte, Y.H. Choi, Natural  
1096 Deep Eutectic Solvent Extraction of Flavonoids of *Scutellaria baicalensis* as a  
1097 Replacement for Conventional Organic Solvents, *Molecules.* 25 (2020) 1–11.  
1098 <https://doi.org/10.3390/molecules25030617>.
- 1099 [42] N. Fu, R. Lv, Z. Guo, Y. Guo, X. You, B. Tang, D. Han, H. Yan, K.H. Row,  
1100 Environmentally friendly and non-polluting solvent pretreatment of palm samples for  
1101 polyphenol analysis using choline chloride deep eutectic solvents, *J. Chromatogr. A.*  
1102 1492 (2017) 1–11. <https://doi.org/10.1016/j.chroma.2017.02.036>.
- 1103 [43] B. Zhuang, L.L. Dou, P. Li, E.H. Liu, Deep eutectic solvents as green media for  
1104 extraction of flavonoid glycosides and aglycones from *Platycladi Cacumen*, *J. Pharm.*  
1105 *Biomed. Anal.* 134 (2017) 214–219. <https://doi.org/10.1016/j.jpba.2016.11.049>.
- 1106 [44] O. Kaltsa, A. Lakka, S. Grigorakis, I. Karageorgou, G. Batra, E. Bozinou, S. Lalas, D.P.

- 1107 Makris, A green extraction process for polyphenols from elderberry (*Sambucus nigra*)  
1108 flowers using deep eutectic solvent and ultrasound-assisted pretreatment, *Molecules*. 25  
1109 (2020) 1–17. <https://doi.org/10.3390/molecules25040921>.
- 1110 [45] W. Ma, K.H. Row, Solid-phase extraction of catechins from green tea with deep eutectic  
1111 solvent immobilized magnetic molybdenum disulfide molecularly imprinted polymer,  
1112 *Molecules*. 25 (2020). <https://doi.org/10.3390/molecules25020280>.
- 1113 [46] Y.H. Choi, J. van Spronsen, Y. Dai, M. Verberne, F. Hollmann, I.W.C.E. Arends, G.J.  
1114 Witkamp, R. Verpoorte, Are natural deep eutectic solvents the missing link in  
1115 understanding cellular metabolism and physiology?, *Plant Physiol*. 156 (2011) 1701–  
1116 1705. <https://doi.org/10.1104/pp.111.178426>.
- 1117 [47] X.H. Fan, Y.H. Chang, L.T. Wang, Y.W. Zhu, M.Z. Dong, M.J. Lv, J.Y. An, Q. Yang,  
1118 J. Jiao, D. Meng, Y.J. Fu, A simple and efficient sample preparation for taxanes in *Taxus*  
1119 *chinensis* needles with natural menthol-based aqueous deep eutectic solvent, *J. Sep. Sci.*  
1120 43 (2020) 1339–1347. <https://doi.org/10.1002/jssc.201900754>.
- 1121 [48] T. Křížek, M. Bursová, R. Horsley, M. Kuchař, P. Tůma, R. Čabala, T. Hložek, Menthol-  
1122 based hydrophobic deep eutectic solvents: Towards greener and efficient extraction of  
1123 phytocannabinoids, *J. Clean. Prod.* 193 (2018) 391–396.  
1124 <https://doi.org/10.1016/j.jclepro.2018.05.080>.
- 1125 [49] C. Cai, W. Yu, C. Wang, L. Liu, F. Li, Z. Tan, Green extraction of cannabidiol from  
1126 industrial hemp (*Cannabis sativa* L.) using deep eutectic solvents coupled with further  
1127 enrichment and recovery by macroporous resin, *J. Mol. Liq.* 287 (2019).  
1128 <https://doi.org/10.1016/j.molliq.2019.110957>.
- 1129 [50] J. Sang, B. Li, Y.Y. Huang, Q. Ma, K. Liu, C.Q. Li, Deep eutectic solvent-based  
1130 extraction coupled with green two-dimensional HPLC-DAD-ESI-MS/MS for the  
1131 determination of anthocyanins from: *Lycium ruthenicum* Murr. fruit, *Anal. Methods*. 10  
1132 (2018) 1247–1257. <https://doi.org/10.1039/c8ay00101d>.

- 1133 [51] M. Ruesgas-Ramón, M.C. Figueroa-Espinoza, E. Durand, Application of Deep Eutectic  
1134 Solvents (DES) for Phenolic Compounds Extraction: Overview, Challenges, and  
1135 Opportunities, *J. Agric. Food Chem.* 65 (2017) 3591–3601.  
1136 <https://doi.org/10.1021/acs.jafc.7b01054>.
- 1137 [52] J. Cao, M. Yang, F. Cao, J. Wang, E. Su, Well-Designed Hydrophobic Deep Eutectic  
1138 Solvents As Green and Efficient Media for the Extraction of Artemisinin from *Artemisia*  
1139 *annua* Leaves, *ACS Sustain. Chem. Eng.* 5 (2017) 3270–3278.  
1140 <https://doi.org/10.1021/acssuschemeng.6b03092>.
- 1141 [53] B.D. Ribeiro, C. Florindo, L.C. Iff, M.A.Z. Coelho, I.M. Marrucho, Menthol-based  
1142 eutectic mixtures: Hydrophobic low viscosity solvents, *ACS Sustain. Chem. Eng.* 3  
1143 (2015) 2469–2477. <https://doi.org/10.1021/acssuschemeng.5b00532>.
- 1144 [54] S. Bajkacz, K. Rusin, A. Wolny, J. Adamek, K. Erfurt, A. Chrobok, Highly efficient  
1145 extraction procedures based on natural deep eutectic solvents or ionic liquids for  
1146 determination of 20-hydroxyecdysone in spinach, *Molecules.* 25 (2020) 1–14.  
1147 <https://doi.org/10.3390/molecules25204736>.
- 1148 [55] K.M. Jeong, Y. Jin, D.E. Yoo, S.Y. Han, E.M. Kim, J. Lee, One-step sample preparation  
1149 for convenient examination of volatile monoterpenes and phenolic compounds in  
1150 peppermint leaves using deep eutectic solvents, *Food Chem.* 251 (2018) 69–76.  
1151 <https://doi.org/10.1016/j.foodchem.2018.01.079>.
- 1152 [56] J. Nie, G. Yu, Z. Song, X. Wang, Z. Li, Y. She, M. Lee, Microwave-assisted deep  
1153 eutectic solvent extraction coupled with headspace solid-phase microextraction followed  
1154 by GC-MS for the analysis of volatile compounds from tobacco, *Anal. Methods.* 9 (2017)  
1155 856–863. <https://doi.org/10.1039/c6ay03076a>.
- 1156 [57] J.B. Barbieri, C. Goltz, F.B. Cavalheiro, A.T. Toci, L. Igarashi-Mafra, M.R. Mafra,  
1157 Corrigendum to “Deep eutectic solvents applied in the extraction and stabilization of  
1158 rosemary (*Rosmarinus officinalis* L.) phenolic compounds” [*Ind. Crops Prod.* 144 (2020)

- 1159 112049] (*Industrial Crops & Products* (2020) 144, (S0926669019310593), (10.1016/j.in,  
1160 *Ind. Crops Prod.* 154 (2020). <https://doi.org/10.1016/j.indcrop.2020.112722>.
- 1161 [58] B.Y. Zhao, P. Xu, F.X. Yang, H. Wu, M.H. Zong, W.Y. Lou, Biocompatible Deep  
1162 Eutectic Solvents Based on Choline Chloride: Characterization and Application to the  
1163 Extraction of Rutin from *Sophora japonica*, *ACS Sustain. Chem. Eng.* 3 (2015) 2746–  
1164 2755. <https://doi.org/10.1021/acssuschemeng.5b00619>.
- 1165 [59] I. Pacheco-Fernández, D.W. Allgaier-Díaz, G. Mastellone, C. Cagliero, D.D. Díaz, V.  
1166 Pino, Biopolymers in sorbent-based microextraction methods, *TrAC Trends Anal.*  
1167 *Chem.* 125 (2020) 115839. <https://doi.org/10.1016/j.trac.2020.115839>.
- 1168 [60] N.H. Godage, E. Gionfriddo, Use of natural sorbents as alternative and green extractive  
1169 materials: A critical review, *Anal. Chim. Acta.* 1125 (2020) 187–200.  
1170 <https://doi.org/10.1016/j.aca.2020.05.045>.
- 1171 [61] G. Mafra, M.T. García-Valverde, J. Millán-Santiago, E. Carasek, R. Lucena, S.  
1172 Cárdenas, Returning to nature for the design of sorptive phases in solid-phase  
1173 microextraction, *Separations.* 7 (2020) 1–22.  
1174 <https://doi.org/10.3390/separations7010002>.
- 1175 [62] J. Cao, L.Q. Peng, J.J. Xu, Microcrystalline cellulose based matrix solid phase dispersion  
1176 microextraction for isomeric triterpenoid acids in loquat leaves by ultrahigh-performance  
1177 liquid chromatography and quadrupole time-of-flight mass spectrometry, *J. Chromatogr.*  
1178 *A.* 1472 (2016) 16–26. <https://doi.org/10.1016/j.chroma.2016.10.034>.
- 1179 [63] L.Q. Peng, Q. Li, Y. xu Chang, M. An, R. Yang, Z. Tan, J. Hao, J. Cao, J.J. Xu, S.S. Hu,  
1180 Determination of natural phenols in olive fruits by chitosan assisted matrix solid-phase  
1181 dispersion microextraction and ultrahigh performance liquid chromatography with  
1182 quadrupole time-of-flight tandem mass spectrometry, *J. Chromatogr. A.* 1456 (2016) 68–  
1183 76. <https://doi.org/10.1016/j.chroma.2016.06.011>.
- 1184 [64] W. Wu, W. Qian, H. Hao, Y. kang, Y. Wang, Y. Deng, C. Ni, J. Huang, W. Weng,

- 1185 Determination of caffeoylquinic acid derivatives in *Azolla imbricata* by chitosan-based  
1186 matrix solid-phase dispersion coupled with HPLC–PDA, *J. Pharm. Biomed. Anal.* 163  
1187 (2019) 197–203. <https://doi.org/10.1016/j.jpba.2018.10.008>.
- 1188 [65] H. Shu, Y.H. Ge, X.Y. Xu, P.Q. Guo, Z.M. Luo, W. Du, C. Chang, R.L. Liu, Q. Fu,  
1189 Hybrid-type carbon microcoil-chitosan composite for selective extraction of aristolochic  
1190 acid I from Aristolochiaceae medicinal plants, *J. Chromatogr. A.* 1561 (2018) 13–19.  
1191 <https://doi.org/10.1016/j.chroma.2018.05.037>.
- 1192 [66] E.M.C. de Matos, L.C. Ribeiro, O.D. Prestes, J.A.G. da Silva, B.S. de Farias, L.A. d. A.  
1193 Pinto, R. Zanella, Multiclass Method for the Determination of Pesticide Residues in Oat  
1194 Using Modified QuEChERS with Alternative Sorbent and Liquid Chromatography with  
1195 Tandem Mass Spectrometry, *Food Anal. Methods.* 12 (2019) 2835–2844.  
1196 <https://doi.org/10.1007/s12161-019-01641-1>.
- 1197 [67] J.J. Xu, Q. Li, J. Cao, E. Warner, M. An, Z. Tan, S.L. Wang, L.Q. Peng, X.G. Liu,  
1198 Extraction and enrichment of natural pigments from solid samples using ionic liquids  
1199 and chitosan nanoparticles, *J. Chromatogr. A.* 1463 (2016) 32–41.  
1200 <https://doi.org/10.1016/j.chroma.2016.08.012>.
- 1201 [68] Z. Li, S. Lu, J. Jin, T. Wang, Preparation of a new cellulose magnetic molecularly  
1202 imprinted polymer micro-spheres to extract and analyze the indole-3-acetic acid in plant  
1203 tissues, *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 1092 (2018) 343–349.  
1204 <https://doi.org/10.1016/j.jchromb.2018.06.023>.
- 1205 [69] Q. Zeng, Y. Ruan, L. Sun, F. Du, L. Guo, Z. Cheng, G. Ruan, J. Li, Development of  
1206 Graphene Oxide Functionalized Cotton Fiber Based Solid Phase Extraction Combined  
1207 with Liquid Chromatography-Fluorescence Detection for Determination of Trace  
1208 Auxins in Plant Samples, *Chromatographia.* 81 (2018) 861–869.  
1209 <https://doi.org/10.1007/s10337-018-3518-0>.
- 1210 [70] Q. Lu, Q. Zhao, Q.W. Yu, Y.Q. Feng, Use of Pollen Solid-Phase Extraction for the



- 1211 Determination of trans-Resveratrol in Peanut Oils, *J. Agric. Food Chem.* 63 (2015)  
1212 4771–4776. <https://doi.org/10.1021/jf505938w>.
- 1213 [71] Q. Lu, J.H. Wu, Q.W. Yu, Y.Q. Feng, Using pollen grains as novel hydrophilic solid-  
1214 phase extraction sorbents for the simultaneous determination of 16 plant growth  
1215 regulators, *J. Chromatogr. A.* 1367 (2014) 39–47.  
1216 <https://doi.org/10.1016/j.chroma.2014.09.071>.
- 1217 [72] L. Jiang, J. Wang, H. Zhang, C. Liu, Y. Tang, C. Chu, New vortex-synchronized matrix  
1218 solid-phase dispersion method for simultaneous determination of four anthraquinones in  
1219 *Cassiae semen*, *Molecules.* 24 (2019) 6–13. <https://doi.org/10.3390/molecules24071312>.
- 1220 [73] M. Li, C. Jiao, X. Yang, C. Wang, Q. Wu, Z. Wang, Solid phase extraction of carbamate  
1221 pesticides with banana peel derived hierarchical porous carbon prior to high performance  
1222 liquid chromatography, *Anal. Methods.* 9 (2017) 593–599.  
1223 <https://doi.org/10.1039/c6ay02678h>.
- 1224 [74] K. Ren, W. Zhang, S. Cao, G. Wang, Z. Zhou, Carbon-based fe<sub>3</sub>o<sub>4</sub> nanocomposites  
1225 derived from waste pomelo peels for magnetic solid-phase extraction of 11 triazole  
1226 fungicides in fruit samples, *Nanomaterials.* 8 (2018).  
1227 <https://doi.org/10.3390/nano8050302>.
- 1228 [75] E. Öztürk Er, E. Maltepe, S. Bakirdere, A novel analytical method for the determination  
1229 of cadmium in sorrel and rocket plants at ultratrace levels: Magnetic chitosan hydrogels  
1230 based solid phase microextraction-slotted quartz tube-flame atomic absorption  
1231 spectrophotometry, *Microchem. J.* 143 (2018) 393–399.  
1232 <https://doi.org/10.1016/j.microc.2018.08.019>.
- 1233 [76] M. Singh, S. Singh, S.P. Singh, S.S. Patel, Recent advancement of carbon nanomaterials  
1234 engrained molecular imprinted polymer for environmental matrix, *Trends Environ. Anal.*  
1235 *Chem.* 27 (2020) e00092. <https://doi.org/10.1016/j.teac.2020.e00092>.
- 1236 [77] N. Tarannum, S. Khatoon, B.B. Dzantiev, Perspective and application of molecular

1237 imprinting approach for antibiotic detection in food and environmental samples: A  
1238 critical review, *Food Control*. 118 (2020) 107381.  
1239 <https://doi.org/10.1016/j.foodcont.2020.107381>.

1240 [78] M. Arabi, A. Ostovan, A.R. Bagheri, X. Guo, L. Wang, J. Li, X. Wang, B. Li, L. Chen,  
1241 Strategies of molecular imprinting-based solid-phase extraction prior to  
1242 chromatographic analysis, *TrAC - Trends Anal. Chem.* 128 (2020) 115923.  
1243 <https://doi.org/10.1016/j.trac.2020.115923>.

1244 [79] M.I. Malik, H. Shaikh, G. Mustafa, M.I. Bhangar, Recent Applications of Molecularly  
1245 Imprinted Polymers in Analytical Chemistry, *Sep. Purif. Rev.* 48 (2019) 179–219.  
1246 <https://doi.org/10.1080/15422119.2018.1457541>.

1247 [80] A. Machyňáková, K. Hroboňová, Synthesis and Evaluation of Molecularly Imprinted  
1248 Polymers as Sorbents for Selective Extraction of Coumarins, *Chromatographia*. 80  
1249 (2017) 1015–1024. <https://doi.org/10.1007/s10337-017-3325-z>.

1250 [81] G. Sun, Y. Liu, H. Ahat, A. Shen, X. Liang, X. Xue, Y. Luo, J. Yang, Z.S. Liu, H.A.  
1251 Aisa, “Two-dimensional” molecularly imprinted solid-phase extraction coupled with  
1252 crystallization and high performance liquid chromatography for fast semi-preparative  
1253 purification of tannins from pomegranate husk extract, *J. Chromatogr. A*. 1505 (2017)  
1254 35–42. <https://doi.org/10.1016/j.chroma.2017.05.033>.

1255 [82] D. Gao, Q.F. Fu, L.J. Wang, D.D. Wang, K.L. Zhang, F.Q. Yang, Z.N. Xia, Molecularly  
1256 imprinted polymers for the selective extraction of tiliroside from the flowers of  
1257 *Edgeworthia gardneri* (wall.) Meisn, *J. Sep. Sci.* 40 (2017) 2629–2637.  
1258 <https://doi.org/10.1002/jssc.201700240>.

1259 [83] Q. Zhang, L. Fan, Q. Lu, X. Yu, M. Liang, J. Nie, N. Li, L. Zhang, Preparation and  
1260 application of molecularly imprinted polymer solid-phase microextraction fiber for the  
1261 selective analysis of auxins in tobacco, *J. Sep. Sci.* 42 (2019) 2687–2695.  
1262 <https://doi.org/10.1002/jssc.201900265>.

- 1263 [84] S. Alipour, P.A. Azar, S.W. Husain, H.R. Rajabi, Determination of Rosmarinic acid in  
1264 plant extracts using a modified sensor based on magnetic imprinted polymeric  
1265 nanostructures, *Sensors Actuators, B Chem.* 323 (2020) 128668.  
1266 <https://doi.org/10.1016/j.snb.2020.128668>.
- 1267 [85] T. Yang, Q. Zhang, T. Chen, W. Wu, X. Tang, G. Wang, J. Feng, W. Zhang, Facile  
1268 potentiometric sensing of gallic acid in edible plants based on molecularly imprinted  
1269 polymer, *J. Food Sci.* 85 (2020) 2622–2628. <https://doi.org/10.1111/1750-3841.15346>.
- 1270 [86] Z. Luo, G. Chen, X. Li, L. Wang, H. Shu, X. Cui, C. Chang, A. Zeng, Q. Fu, Molecularly  
1271 imprinted polymer solid-phase microextraction coupled with ultra high performance  
1272 liquid chromatography and tandem mass spectrometry for rapid analysis of pyrrolizidine  
1273 alkaloids in herbal medicine, *J. Sep. Sci.* 42 (2019) 3352–3362.  
1274 <https://doi.org/10.1002/jssc.201900665>.
- 1275 [87] C. Wang, X.J. Li, J. Yang, Y.X. Zhao, Z.S. Liu, H.A. Aisa, Preparation of ionic liquid-  
1276 mediated imprinted monolith for selective capture and purification of corilagin, *J.*  
1277 *Chromatogr. B Anal. Technol. Biomed. Life Sci.* 1041–1042 (2017) 98–103.  
1278 <https://doi.org/10.1016/j.jchromb.2016.12.023>.
- 1279 [88] Y. Han, C. Yang, Y. Zhou, D. Han, H. Yan, Ionic Liquid-Hybrid Molecularly Imprinted  
1280 Material-Filter Solid-Phase Extraction Coupled with HPLC for Determination of 6-  
1281 Benzyladenine and 4-Chlorophenoxyacetic Acid in Bean Sprouts, *J. Agric. Food Chem.*  
1282 65 (2017) 1750–1757. <https://doi.org/10.1021/acs.jafc.6b03922>.
- 1283 [89] E. Turiel, A. Martín-Esteban, Molecularly imprinted polymers-based microextraction  
1284 techniques, *TrAC - Trends Anal. Chem.* 118 (2019) 574–586.  
1285 <https://doi.org/10.1016/j.trac.2019.06.016>.
- 1286 [90] X. Li, Y. Dai, K.H. Row, Preparation of two-dimensional magnetic molecularly  
1287 imprinted polymers based on boron nitride and a deep eutectic solvent for the selective  
1288 recognition of flavonoids, *Analyst.* 144 (2019) 1777–1788.

- 1289 <https://doi.org/10.1039/c8an02258e>.
- 1290 [91] E. Fischer, Einfluss der Configuration auf die Wirkung der Enzyme, Chem. Ber. 27  
1291 (1894) 2985-299, <https://doi.org/10.1002/cber.18940270364>
- 1292 [92] D. Xiao, T. Lu, R. Zeng, Y. Bi, Preparation and highlighted applications of magnetic  
1293 microparticles and nanoparticles : a review on recent advances, Microchim. Acta (2016)  
1294 2655–2675. <https://doi.org/10.1007/s00604-016-1928-y>.
- 1295 [93] F. Abujaber, F.J.G. Bernardo, R.C.R. Martín-doimeadios, Talanta Magnetic cellulose  
1296 nanoparticles as sorbents for stir bar-sorptive dispersive microextraction of  
1297 polychlorinated biphenyls in juice samples, Talanta 201 (2019) 266–270.  
1298 <https://doi.org/10.1016/j.talanta.2019.04.005>
- 1299 [94] J. Chen, S. Cao, M. Zhu, C. Xi, L. Zhang, X. Li, G. Wang, Y. Zhou, Z. Chen, Fabrication  
1300 of a high selectivity magnetic solid phase extraction adsorbent based on  $\beta$ -cyclodextrin  
1301 and application for recognition of plant growth regulators, J. Chromatogr. A. 1547  
1302 (2018) 1–13. <https://doi.org/10.1016/j.chroma.2018.03.004>.
- 1303 [95] J.Y. Chen, S.R. Cao, C.X. Xi, Y. Chen, X.L. Li, L. Zhang, G.M. Wang, Y.L. Chen, Z.Q.  
1304 Chen, A novel magnetic  $\beta$ -cyclodextrin modified graphene oxide adsorbent with high  
1305 recognition capability for 5 plant growth regulators, Food Chem. 239 (2018) 911–919.  
1306 <https://doi.org/10.1016/j.foodchem.2017.07.013>.
- 1307 [96] V. Cruz Fernandes, M. Freitas, J.G. Pacheco, V. Fernandes Domingues, C. Delerue-  
1308 Matos, Evaluation of the QuEChERS and magnetic micro dispersive solid-phase  
1309 extraction of brominated flame retardants in red fruits with determination by GC/MS,  
1310 Food Chem. 309 (2020) 125572. <https://doi.org/10.1016/j.foodchem.2019.125572>.
- 1311 [97] Q. Zhang, D.D. Zhou, J.W. Zhang, D. Gao, F.Q. Yang, H. Chen, Z.N. Xia, Amino-  
1312 terminated supramolecular cucurbit [6] uril pseudorotaxane complexes immobilized on  
1313 magnetite@silica nanoparticles: A highly efficient sorbent for salvianolic acids, Talanta.  
1314 195 (2019) 354–365. <https://doi.org/10.1016/j.talanta.2018.11.086>.

- 1315 [98] A. Gutiérrez-Serpa, I. Pacheco-Fernández, J. Pasán, V. Pino, Metal–organic frameworks  
1316 as key materials for solid-phase microextraction devices—a review, *Separations*. 6  
1317 (2019). <https://doi.org/10.3390/separations6040047>.
- 1318 [99] H. Duo, Y. Wang, L. Wang X. Lu, X. Liang, Zirconium(IV)-based metal–organic  
1319 frameworks (UiO-67) as solid-phase extraction adsorbent for the extraction of  
1320 phenoxyacetic acid herbicides from vegetables, *J. Sep. Sci.* (2018).  
1321 <https://doi.org/10.1002/jssc.201800784>
- 1322 [100] T. Si, L.Liu, X. Liang, H. Duo, L. Wang, S. Wang, Solid-phase extraction of  
1323 phenoxyacetic acid herbicides in complex samples with a zirconium(IV)-based metal–  
1324 organic framework, *J. Sep. Sci.* (2019).<https://doi.org/10.1002/jssc.201900243>
- 1325 [101] R.M. Abdelhameed, H. Abdel-Gawad, M. Taha, B. Hegazi, Separation of bioactive  
1326 chamazulene from chamomile extract using metal-organic framework, *J. Pharm.*  
1327 *Biomed. Anal.* 146 (2017) 126–134. <https://doi.org/10.1016/j.jpba.2017.08.020>.
- 1328 [102] Y.Y. Cui, C.X. Yang, X.D. Yang, X.P. Yan, Zeolitic imidazolate framework-8 for  
1329 selective extraction of a highly active anti-oxidant flavonoid from *Caragana Jubata*, *J.*  
1330 *Chromatogr. A.* 1544 (2018) 8–15. <https://doi.org/10.1016/j.chroma.2018.02.046>.
- 1331 [103] R. Zhang, N. Xu, Y. Wang, X. Liu, S. Wang, J. Cao, Metal–organic framework assisted  
1332 matrix solid-phase dispersion microextraction of saponins using response surface  
1333 methodology, *Electrophoresis*. 41 (2020) 1354–1363.  
1334 <https://doi.org/10.1002/elps.202000042>.
- 1335 [104] S. Bahrani, M. Ghaedi, K. Dashtian, A. Ostovan, M.J.K. Mansoorkhani, A. Salehi, MOF-  
1336 5(Zn)-Fe<sub>2</sub>O<sub>4</sub> nanocomposite based magnetic solid-phase microextraction followed by  
1337 HPLC-UV for efficient enrichment of colchicine in root of *colchicum* extracts and  
1338 plasma samples, *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 1067 (2017) 45–52.  
1339 <https://doi.org/10.1016/j.jchromb.2017.09.044>.
- 1340 [105] Z. Durmus, B. Zengin Kurt, I. Gazioğlu, E. Sevgi, C. Kizilarlan Hancer,

- 1341 Spectrofluorimetric Determination of Aflatoxin B1 in Winter Herbal Teas via Magnetic  
1342 Solid Phase Extraction Method by using Metal–Organic Framework (MOF) Hybrid  
1343 Structures Anchored with Magnetic Nanoparticles, *Appl. Organomet. Chem.* 34 (2020)  
1344 2–11. <https://doi.org/10.1002/aoc.5375>.
- 1345 [106] Z. Moradi, E. Alipanahpour Dil, A. Asfaram, Dispersive micro-solid phase extraction  
1346 based on Fe<sub>3</sub>O<sub>4</sub>@SiO<sub>2</sub>@Ti-MOF as a magnetic nanocomposite sorbent for the trace  
1347 analysis of caffeic acid in the medical extracts of plants and water samples prior to  
1348 HPLC-UV analysis, *Analyst.* 144 (2019) 4351–4361.  
1349 <https://doi.org/10.1039/c9an00120d>.
- 1350 [107] Y. Zhou, J. Zhu, J. Yang, Y. Lv, Y. Zhu, W. Bi, X. Yang, D.D.Y. Chen, Magnetic  
1351 nanoparticles speed up mechanochemical solid phase extraction with enhanced  
1352 enrichment capability for organochlorines in plants, *Anal. Chim. Acta.* 1066 (2019) 49–  
1353 57. <https://doi.org/10.1016/j.aca.2019.03.049>.
- 1354 [108] A. Ghiasvand, S. Koonani, F. Yazdankhah, S. Farhadi, A comparison study on a  
1355 sulfonated graphene-polyaniline nanocomposite coated fiber for analysis of nicotine in  
1356 solid samples through the traditional and vacuum-assisted HS-SPME, *J. Pharm. Biomed.*  
1357 *Anal.* 149 (2018) 271–277. <https://doi.org/10.1016/j.jpba.2017.11.020>.
- 1358 [109] Y. Su, S. Wang, N. Zhang, P. Cui, Y. Gao, T. Bao, Zr-MOF modified cotton fiber for  
1359 pipette tip solid-phase extraction of four phenoxy herbicides in complex samples,  
1360 *Ecotoxicol. Environ. Saf.* 201 (2020) 110764.  
1361 <https://doi.org/10.1016/j.ecoenv.2020.110764>.
- 1362 [110] X. Pang, H. Liu, H. Yu, M. Zhang, L. Bai, H. Yan, Monolithic Column Prepared with  
1363 UiO-66-2COOH MOF as Monomer for Enrichment and Purification of Ursolic Acid in  
1364 Plants by Online Solid-Phase Extraction, *Chromatographia.* 83 (2020) 1121–1131.  
1365 <https://doi.org/10.1007/s10337-020-03931-x>.
- 1366 [111] X. Pang, H. Liu, H. Yu, M. Zhang, L. Bai, H. Yan, A metal organic framework polymer

- 1367 monolithic column as a novel adsorbent for on-line solid phase extraction and  
1368 determination of ursolic acid in Chinese herbal medicine, *J. Chromatogr. B Anal.*  
1369 *Technol. Biomed. Life Sci.* 1125 (2019).  
1370 <https://doi.org/10.1016/j.jchromb.2019.121715>.
- 1371 [112] M. Zhang, H. Liu, Y. Han, L. Bai, H. Yan, On-line enrichment and determination of  
1372 aristolochic acid in medicinal plants using a MOF-based composite monolith as  
1373 adsorbent, *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 1159 (2020).  
1374 <https://doi.org/10.1016/j.jchromb.2020.122343>.
- 1375 [113] I. P. Fernández, M. Rentero, J.H. Ayala, J. Pasàn, V. Pino, Green solid-phase  
1376 microextraction fiber coating based on the metal-organic framework CIM-80 ( Al ):  
1377 Analytical performance evaluation in direct immersion and headspace using gas  
1378 chromatography and mass spectrometry for the analysis of water, urine, brewed coffee,  
1379 *Anal. Chim. Acta* (2020). <https://doi.org/10.1016/j.aca.2020.08.009>.
- 1380 [114] Y. Zhu, S. Murali, W. Cai, X. Li, J.W. Suk, J.R. Potts, R.S. Ruoff, Graphene and  
1381 graphene oxide: Synthesis, properties, and applications, *Adv. Mater.* 22 (2010) 3906–  
1382 3924. <https://doi.org/10.1002/adma.201001068>.
- 1383 [115] H. Zhang, X. Wu, Y. Yuan, D. Han, F. Qiao, H. Yan, An ionic liquid functionalized  
1384 graphene adsorbent with multiple adsorption mechanisms for pipette-tip solid-phase  
1385 extraction of auxins in soybean sprouts, *Food Chem.* 265 (2018) 290–297.  
1386 <https://doi.org/10.1016/j.foodchem.2018.05.090>.
- 1387 [116] A. Markus, A.O. Gbadamosi, A.S. Yusuff, A. Agi, J. Oseh, Magnetite-  
1388 sporopollenin/graphene oxide as new preconcentration adsorbent for removal of polar  
1389 organophosphorus pesticides in vegetables, *Environ. Sci. Pollut. Res.* 25 (2018) 35130–  
1390 35142. <https://doi.org/10.1007/s11356-018-3402-3>.
- 1391 [117] L.A. Akamine, D.A. Vargas Medina, F.M. Lanças, Magnetic solid-phase extraction of  
1392 gingerols in ginger containing products, *Talanta.* 222 (2021) 121683.

- 1393 <https://doi.org/10.1016/j.talanta.2020.121683>.
- 1394 [118] D. Jiang, D. Wei, L. Wang, S. Ma, Y. Du, M. Wang, Multiwalled carbon nanotube for  
1395 one-step cleanup of 21 mycotoxins in corn and wheat prior to ultraperformance liquid  
1396 chromatography–tandem mass spectrometry analysis, *Toxins (Basel)*. 10 (2018) 1–17.  
1397 <https://doi.org/10.3390/toxins10100409>.
- 1398 [119] S. Song, Z. Zhang, N. Zou, R. Chen, L. Han, C. Pan, Y. Sapozhnikova, Determination  
1399 of Six Paraben Residues in Fresh-cut Vegetables Using QuEChERS with Multi-walled  
1400 Carbon Nanotubes and High-Performance Liquid Chromatography–Tandem Mass  
1401 Spectrometry, *Food Anal. Methods*. 10 (2017) 3972–3979.  
1402 <https://doi.org/10.1007/s12161-017-0970-7>.
- 1403 [120] K. Jiang, P. Huang, L. Luan, K. Fan, W. Guo, Z. Zhao, Y. Wu, Z. Han, Iron (II, III)  
1404 oxide/multi-walled carbon nanotube composite as solid-phase extraction sorbent  
1405 followed by ultra-high performance liquid chromatography tandem mass spectrometry  
1406 for simultaneous determination of zearalenone and type A trichothecenes in *Salv*, *J.*  
1407 *Chromatogr. A*. 1482 (2017) 1–10. <https://doi.org/10.1016/j.chroma.2016.12.058>.
- 1408 [121] H. Liu, T. Lin, Q. Li, A magnetic multi-walled carbon nanotube preparative method for  
1409 analyzing asymmetric carbon, phosphorus and sulfur atoms of chiral pesticide residues  
1410 in Chinese herbals by chiral liquid chromatography-quadrupole/linear ion trap mass  
1411 spectrometry determinat, *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 1148  
1412 (2020) 122152. <https://doi.org/10.1016/j.jchromb.2020.122152>.
- 1413 [122] X. Ling, L. Zou, Z. Chen, Polymeric monolith column composited with multiwalled  
1414 carbon nanotubes- $\beta$ -cyclodextrin for the selective extraction of psoralen and isopsoralen,  
1415 *J. Sep. Sci.* 40 (2017) 3718–3724. <https://doi.org/10.1002/jssc.201700562>.
- 1416 [123] T. Chen, H. Xu, In vivo investigation of pesticide residues in garlic using solid phase  
1417 microextraction-gas chromatography-mass spectrometry, *Anal. Chim. Acta.* 1090  
1418 (2019) 72–81. <https://doi.org/10.1016/j.aca.2019.09.011>.





## Artwork

**Figure 1.** Overview of the main extraction phases employed as sorbent in liquid and solid phase extraction in plant field (A) and scheme of the structural characteristics of the main extraction phases employed in plant field (B).

**Figure 2.** Representative examples of extraction procedures using innovative sorbents in plant field: A) *in-vivo* SPME using a MWCNTs-based SPME fiber, adapted from Ref. [123], with permission from Elsevier; B) switchable temperature-responsive ionic liquid-based method for the simultaneous extraction and *in-situ* separation of lipophilic and hydrophilic compounds, adapted from Ref. [30], with permission from The Royal Society of Chemistry; C) m-SPE using a MOF-NPs hybrid material, adapted from Ref. [105], with permission from Wiley D) potentiometric analysis using MIP-based electrochemical sensor, adapted from Ref. [85], with permission from Institute of Food Technologists; and E) MSPD microextraction using microcrystalline cellulose as sorbent, adapted from Ref. [62], with permission from Elsevier.

## Tables

**Table 1.** Main features of innovative extraction phases in plant field.

<b>Feature</b>	<b>ILs</b>	<b>DESs</b>	<b>Natural sorbents</b>	<b>MIPs</b>	<b>MNPs</b>	<b>Reticular materials</b>	<b>Carbon materials</b>
Pre-treatment of the sample	+/-	-	+/- (MSPD <sup>a</sup> )	+	+	+/- (MSPD <sup>a</sup> )	+
Extraction time (min)	10-30	30-120	1-5	+/- <sup>b</sup>	+/- <sup>b</sup>	+/- <sup>b</sup>	+/- <sup>b</sup>
Solvent consumption	-	-	+	+	+	+	+
Hybrid phases	+ [25]	+ [45]	+ [75]	+ [87]	+ [117]	+ [107]	+ [116]
Example of automatation	-	-	-	+ [83]	-	-	+ [122]
Microextraction	+ [32, 18]	- <sup>c</sup>	+ [67]	+ [86]	+ [96]	+ [104]	+ [123]
Re-use	+ [18]	+ [42]	+ [69]	+ [83]	+ [97]	+ [102]	+ [117]
Commercially available	+	+	+/-	+	+	+	+

+/- depend on the application/technique

<sup>a</sup> matrix solid-phase dispersion

<sup>b</sup> from few seconds to more than 30 minutes

<sup>c</sup> possible, but no applications found in plant field

**Table 2.** Representative applications of innovative extraction phases in plant field.

Sample	Analytes	Phase	Additional materials	Extraction method <sup>a</sup>	Analytical platform	Ref.
IL-based extraction						
<i>Camellia sinensis</i> (L.) Kuntze	polyphenols	multi-tentacled PVAlm <sup>+</sup> Pro <sup>-</sup> @SiO <sub>2</sub> <sup>b</sup>	silica	dSPE	HPLC-MS/MS	[24]
<i>Elaeagnus rhamnoides</i> (L.) A.Nelson	essential oil, non-volatile compounds	[C <sub>4</sub> mIm] <sup>+</sup> [Br] <sup>-c</sup>	/	UA/MA distillation extraction	HPLC-UV/GC-MS	[26]
<i>Coptis</i> spp.	berberine hydrochloride	MIL two phase system [N <sub>115</sub> 2OH] <sup>+</sup> [Br] <sup>-</sup> and [TEMPO-OSO <sub>3</sub> ] <sup>-</sup> [Na] <sup>+d</sup>	/	magnetic ABS	HPLC-UV	[18]
<i>Chrysanthemum</i> spp.	lipophilic and hydrophilic endogenous compounds	Temperature-responsive [HMIM] <sup>+</sup> [BF <sub>4</sub> ] <sup>-c</sup>	/	<i>in-situ</i> solid-liquid extraction	UHPLC-UV-QTOF-MS	[30]
DES-based extraction						
<i>Trachycarpus fortunei</i> (Hook.) H.Wendl.	polyphenols	ChCl/Ph <sup>f</sup> sorbent ChCl DES eluent	/	SPE	HPLC-MS	[42]
<i>Mentha × piperita</i> L.	volatile monoterpenes, phenols	ChCl/D(+)-glucose	/	solid-liquid UAE	UHPLC-QTOF-MS/GC-MS	[55]
<i>Rosmarinus officinalis</i> L.	phenols	ChCl/1,2-Propanediol	/	solid-liquid UAE	HPLC-UV	[57]
<i>Taxus wallichiana</i> var. <i>chinensis</i> (Pilg.) Florin	taxanes	menthol/1-propanol	/	solid-liquid UAE	HPLC-UV	[47]
Natural sorbent-based extraction						
<i>Eriobotrya japonica</i> (Thunb.) Lindl.	triterpenoid acid	MCC <sup>g</sup>	/	MSPD	UHPLC-QTOF-MS	[62]
Fruits	triazole fungicides	carbon (from pomelo peel) NPs	Fe <sub>3</sub> O <sub>4</sub>	m-SPE	GC-MS	[74]
<i>Azolla imbricata</i> (Roxb. ex Griff.) Nakai	caffeoylquinic acid derivatives	chitosan	/	MSPD	HPLC-UV	[64]
MIP-based extraction						

<i>Phyllanthus urinaria</i> L.	corilagin	Corilagin-MIP	[BMIM] <sup>+</sup> [BF <sub>4</sub> ] <sup>-h</sup>	SPE	HPLC-UV	[87]
<i>Tussilago farfara</i> L.	pyrrolizidine alkaloids	PA <sup>i</sup> -MIP	/	SPME	UHPLC	[86]
Edible plant	gallic acid	GA <sup>l</sup> -MIP	/	/	IES <sup>e</sup> potentiometric analysis	[85]
MNP-based extraction						
<i>Salvia miltiorrhiza</i> Bunge	salvianolic acids	magnetite@silica NPs	curcubit pseudorotaxane complex	m-dSPE	HPLC-MS	[97]
Reticular material-based extraction						
<i>Panax C.A.Mey</i>	<i>ginseng</i> saponins	MOF-808	/	μ-MSPD	UHPLC-QTOF-MS	[103]
<i>Camellia (L.) Kuntze</i>	<i>sinensis</i> organochlorine pesticides	MIL-100	Fe <sub>3</sub> O <sub>4</sub>	m-SPE	UHPLC-UV/GC-MS	[107]
Carbon material-based extraction						
Vegetables	parabens	MWCNTs	/	QuEChERS (SPE)	HPLC-MS/MS	[119]
<i>Allium sativum</i> L.	pesticides	MWCNTs	PANI-PPy@PDMS <sup>m</sup>	DI/ <i>in-vivo</i> / <i>in-vitro</i> SPME	GC-MS	[123]

<sup>a</sup> for the acronyms of the techniques, see abbreviation list

<sup>b</sup> silica particles with polyvinyl alcohol chains modified by N-methyl imidazole proline salt

<sup>c</sup> 1-butyl-3-methyl imidazole bromine salt

<sup>d</sup> alkyl-(2-hydroxyethyl)-dimethylammonium bromides, sodium 4-sulfonatoxy-2,2,6,6-tetramethyl piperidine-1-yl oxyl

<sup>e</sup> 1-hexyl-3-methylimidazolium tetrafluoroborate

<sup>f</sup> phenol

<sup>g</sup> microcrystalline cellulose

<sup>h</sup> 1-butyl-3-methylimidazolium tetrafluoroborate

<sup>i</sup> pyrrolizidine alkaloid


<sup>l</sup> gallic acid

<sup>m</sup> polyaniline-polypyrrole@polydimethylsiloxane

A)


### Liquid phase extraction

- Ionic liquids
- Deep eutectic solvents





### Solid phase extraction

- Natural sorbents
- MIPs
- Magnetic NPs
- Reticular and carbon-based materials




### Hybrid phases

Association of different phases




B)


ILs




Biopolymers




MIPs




Magnetic NPs



DESs



Reticular materials



Carbon-based materials

