

SCIENTIFIC REPORTS

OPEN

Zolpidem is a potent stoichiometry-selective modulator of $\alpha 1\beta 3$ GABA_A receptors: evidence of a novel benzodiazepine site in the $\alpha 1$ - $\alpha 1$ interface

Received: 03 March 2016

Accepted: 06 June 2016

Published: 27 June 2016

Ahmad Tarmizi Che Has¹, Nathan Absalom¹, Petra S. van Nieuwenhuijzen¹, Andrew N. Clarkson^{1,2}, Philip K. Ahring¹ & Mary Chebib¹

Zolpidem is not a typical GABA_A receptor hypnotic. Unlike benzodiazepines, zolpidem modulates tonic GABA currents in the rat dorsal motor nucleus of the vagus, exhibits residual effects in mice lacking the benzodiazepine binding site, and improves speech, cognitive and motor function in human patients with severe brain injury. The receptor by which zolpidem mediates these effects is not known. In this study we evaluated binary $\alpha 1\beta 3$ GABA_A receptors in either the $3\alpha 1:2\beta 3$ or $2\alpha 1:3\beta 3$ subunit stoichiometry, which differ by the existence of either an $\alpha 1$ - $\alpha 1$ interface, or a $\beta 3$ - $\beta 3$ interface, respectively. Both receptor stoichiometries are readily expressed in *Xenopus* oocytes, distinguished from each other by using GABA, zolpidem, diazepam and Zn²⁺. At the $3\alpha 1:2\beta 3$ receptor, clinically relevant concentrations of zolpidem enhanced GABA in a flumazenil-sensitive manner. The efficacy of diazepam was significantly lower compared to zolpidem. No modulation by either zolpidem or diazepam was detected at the $2\alpha 1:3\beta 3$ receptor, indicating that the binding site for zolpidem is at the $\alpha 1$ - $\alpha 1$ interface, a site mimicking the classical $\alpha 1$ - $\gamma 2$ benzodiazepine site. Activating $\alpha 1\beta 3$ ($3\alpha 1:2\beta 3$) receptors may, in part, mediate the physiological effects of zolpidem observed under distinct physiological and clinical conditions, constituting a potentially attractive drug target.

γ -aminobutyric acid receptors of type A (GABA_A) are members of the Cys-loop family of ligand-gated ion channels that mediate most of the inhibitory neurotransmission in the central nervous system (CNS). These receptors are pentameric assemblies of individual subunits including $\alpha 1$ -6, $\beta 1$ -3, $\gamma 1$ -3, δ , ϵ , π and θ . The majority of receptors are composed of α , β , and γ or δ subunits¹⁻⁴. Depending on the subunit composition, the receptors are located either at the synapse ($\alpha 1/2/3\beta 2$ receptors) where they are exposed to brief (ms) bursts of high concentrations of GABA when the neuron fires (termed phasic inhibition) or at extrasynaptic regions ($\alpha 1/4/6\beta$, $\alpha 5\beta 2$ or $\alpha 4/6\beta 3$ receptors) where receptors typically experience long lasting (min to hours) exposure to relatively low but consistent concentrations of GABA (termed tonic inhibition)²⁻⁴. The most abundant receptors are $\alpha 1\beta 2$ receptors, and these are activated and modulated by a variety of pharmacologically and clinically unrelated agents including benzodiazepines, barbiturates, anaesthetics and neurosteroids, all of which bind at distinct binding sites located within the receptor complex¹⁻⁴.

Zolpidem is an imidazopyridine, a non-benzodiazepine, that binds with high-affinity to $\alpha 1$ containing GABA_A receptors ($\alpha 1\beta 2$)^{5,6} and with lower affinity to $\alpha 2$ or $\alpha 3$ containing receptors ($\alpha 2\beta 2$ and $\alpha 3\beta 2$)^{7,8}. In contrast, zolpidem is significantly more efficacious at $\alpha 2$ or $\alpha 3$ than $\alpha 1$ containing receptors⁹. The binding site for zolpidem at these receptors is located at the interface between the principle side (+) of the α and the complementary side (-) of $\gamma 2$ subunit^{4,10,11}. When zolpidem binds to this site, there is a structural change to the protein

¹Faculty of Pharmacy, The University of Sydney, Sydney, New South Wales, 2006, Australia. ²Department of Anatomy, Brain Health Research Centre and Brain Research New Zealand, School of Medical Sciences, University of Otago, Dunedin, 9054, New Zealand. Correspondence and requests for materials should be addressed to M.C. (email: mary.collins@sydney.edu.au)

complex resulting in an increase in GABA potency⁴. The effects of zolpidem can be competitively blocked by the neutral modulator flumazenil¹².

Like many benzodiazepines, zolpidem exhibits sedative and hypnotic effects via $\alpha 1$ containing receptors, but unlike benzodiazepines, zolpidem reverses cognitive and motor deficits in neuropathological states, improves motor function in patients with Parkinson's disease, progressive supranuclear palsy and stroke^{13–17}. Studies also show that there are residual effects with zolpidem in the $\gamma 2^{F771}$ mouse where binding by zolpidem to $\alpha(+)-\gamma 2(-)$ is attenuated^{18,19}. Further, zolpidem can modulate tonic GABA currents in the rat dorsal motor nucleus of the vagus²⁰ and primary motor cortex¹⁵, implicating extrasynaptic receptors containing $\alpha 5$ and δ subunits. However $\alpha 5$ and δ containing receptors do not significantly respond to zolpidem^{7,8} and neither do $\gamma 1/3$ containing receptors^{21,22}. Thus we sought to determine whether binary $\alpha\beta$ receptors are potential targets that mediate zolpidem's atypical effects.

Binary $\alpha\beta$ receptors lack the benzodiazepine $\alpha-\gamma 2$ interface and co-exist with classical GABA_ARs^{23–25}. The physiological role of these receptors remains elusive^{25–27}, due to a lack of pharmacological tools that differentiate these receptors from their ternary counterparts. Interestingly, there are reports that certain benzodiazepines are able to enhance GABA actions at these receptors by binding to sites other than the classical $\alpha-\gamma 2$ interface^{28–30}.

In pentameric $\alpha\beta$ receptors, the third subunit is replaced with either an $\alpha 1$ or a $\beta 3$ subunit leading to two distinct receptors that differ in subunit stoichiometry, $2\alpha:3\beta^{31,32}$ or $3\alpha:2\beta^{33}$. The consequence of this is that $3\alpha:2\beta$ receptors contain an $\alpha-\alpha$ interface whereas $2\alpha:3\beta$ receptors contain a $\beta-\beta$ interface, that are clearly distinct. Drugs that selectively bind to either the $\alpha-\alpha$ or $\beta-\beta$ interface will have stoichiometric selective effects as demonstrated by the *in vivo* and *in vitro* effects of drugs at the related nicotinic acetylcholine $\alpha 4\beta 2$ receptors^{34,35}.

In this study, we evaluated the effects of GABA, Zn^{2+} , zolpidem and diazepam at binary $\alpha 1\beta 3$ GABA_A receptors expressed as $3\alpha 1:2\beta 3$ and $2\alpha 1:3\beta 3$ subunit stoichiometries in *Xenopus* oocytes using two electrode voltage clamp electrophysiology. Both receptors expressed readily, and were distinguished from each other by their pharmacology. The $2\alpha 1:3\beta 3$ receptor was highly sensitive to GABA and Zn^{2+} , while the $3\alpha 1:2\beta 3$ receptor was less sensitive to GABA and Zn^{2+} . Interestingly zolpidem and diazepam were potent allosteric modulators of $3\alpha 1:2\beta 3$ receptors and the effects were attenuated by flumazenil. While zolpidem displayed high efficacy at the $3\alpha 1:2\beta 3$ stoichiometry, the efficacy of diazepam was significantly lower. No modulation by either zolpidem or diazepam was detected at $2\alpha 1:3\beta 3$ receptors. These results demonstrate that the $\alpha 1-\alpha 1$ interface contains a binding site for zolpidem mimicking the classical $\alpha 1-\gamma 2$ benzodiazepine site. This novel target may explain why zolpidem enhances tonic GABA currents as binary $\alpha\beta$ receptors are reported to be extrasynaptic, and may be a target for its motor effects after severe brain damage.

Results

Different $\alpha 1\beta 3$ receptors are expressed by injecting *Xenopus* oocytes with different cRNA ratios. The assembly of Cys-loop receptors in *Xenopus laevis* oocytes into specific stoichiometries can be directed by injecting cRNA with variant subunit ratios^{34,36}. To determine whether varying the injection ratios led to the expression of different receptors, we injected five mixtures of $\alpha 1$ and $\beta 3$ cRNAs into oocytes ($\alpha 1 + \beta 3$ in 1:1, 5:1, 10:1, 20:1 and 30:1 ratios). To allow for comparisons between injection ratios, all the cRNA was derived from a single stock. The total amount of cRNA injected ranged from 2.5 to 4.0 ng/oocyte and the functional properties of receptors including holding currents, GABA sensitivity and maximum peak current amplitudes were compared.

Initially, receptors expressed by the most extreme injection ratios 1:1 and 30:1, were compared. The holding currents of oocytes injected with $\alpha 1 + \beta 3$ (1:1) cRNA were -150 ± 93 nA, $n = 6$ when voltage-clamped at -60 mV, suggesting constitutive receptor activity (Fig. 1A). In contrast, oocytes injected with $\alpha 1 + \beta 3$ (30:1) had negligible holding current levels that averaged -13 ± 10 nA, $n = 9$ (Fig. 1A; $p < 0.05$).

GABA activated receptors expressed from both injection ratios in a concentration-dependent manner, with maximum similar peak-current amplitudes ranging from 1200 to 4200 nA ($p > 0.05$). Fitting peak-current amplitudes as a function of the GABA concentration to the Hill equation revealed higher GABA sensitivity with receptors from the 1:1 cRNA ratio compared with those formed from the 30:1 ratio. The derived EC_{50} values were significantly different ($p < 0.0001$) with a 10-fold difference between receptors at the 1:1 and 30:1 ratios (2.8 μ M and 26 μ M, respectively) (Fig. 1B; Table 1). The GABA sensitivity from the 30:1 injection ratio was similar to that of the ubiquitous $\alpha 1\beta 3\gamma 2$ receptor (Fig. 1B; Table 1).

In contrast, receptors formed from 5:1, 10:1, and 20:1 cRNA injection ratios resulted in less uniform receptor populations. This resulted in intermediate GABA sensitivities, concentration-response curves with shallow Hill slopes and comparatively high standard errors for individual data points (data not shown). Thus, receptors formed from different injection ratios had different functional properties that are likely the result of the expression of distinct receptor populations with altered subunit stoichiometries.

Expressing stoichiometry specific $\alpha 1\beta 3$ receptors using concatenated constructs. To determine whether data obtained by varying cRNA ratios are from uniform receptor populations that consist of $2\alpha:3\beta$ and $3\alpha:2\beta$ stoichiometries, the functional properties of receptors expressed with a concatenated construct linking the $\beta 3$ and $\alpha 1$ subunits were compared to receptors expressed with the 1:1 or 30:1 cRNA injection ratios. For this experiment, the N-terminal of the $\alpha 1$ subunit was linked to the C-terminal of the $\beta 3$ subunit to generate the $\beta 3-\alpha 1$ concatenated construct as previously described³⁷. When cRNA transcribed from the $\beta 3-\alpha 1$ construct was injected into oocytes (2.5 ng/oocyte) no GABA-elicited currents were observed ($n = 10$). Therefore, to express receptors in a $2\alpha:3\beta$ and $3\alpha:2\beta$ stoichiometry, $\beta 3-\alpha 1$ was co-injected with either the $\beta 3$ or $\alpha 1$ subunit, respectively.

First, the properties of $2\alpha:3\beta$ receptors were compared to receptors formed by the 1:1 injection ratio. Both holding current levels and the GABA sensitivities of oocytes expressing receptors from the concatenated $\beta 3-\alpha 1 + \beta 3$ (1:2) construct were similar to oocytes injected with free $\alpha 1 + \beta 3$ (1:1) subunits, with holding currents




Figure 1. GABA-evoked responses at $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ GABA_A receptors. *Xenopus laevis* oocytes were injected with cRNA and subjected to two-electrode voltage-clamp electrophysiology as described in the methods. For experimentation, oocytes were clamped at -60 mV and full GABA concentration response relationships were obtained on each oocyte. (A) Representative GABA-evoked traces from oocytes injected with the denoted cRNA mixtures. Bars above each trace indicate application periods and GABA concentrations and “/” a wash period. Dotted lines indicate a 0 nA baseline and holding currents were -150 ± 93 nA, $n = 6$ for $\alpha 1 + \beta 3$ (1:1) and -13 ± 10 nA, $n = 9$ for $\alpha 1 + \beta 3$ (30:1). (B,C) Baseline subtracted peak current amplitudes for full GABA concentration-response curves at oocytes injected with the indicated cRNA mixtures using free subunits (B) or a concatenated $\beta 3$ - $\alpha 1$ construct (C) were fitted to the Hill equation using non-linear regression (fixed bottom of 0 and slope of 1) and normalized to the maximal fitted value ($I_{GABA_max_fit}$). Averaged normalized data points are depicted as means \pm S.E. as a function of the GABA concentration, fitted to the Hill equation and regression results are presented in Table 1. Each data point represents experiments from $n = 5$ –9 oocytes from ≥ 2 batches. (D) $\alpha 1\beta 3$ GABA_A receptors can express in two stoichiometries of $2\alpha 1:3\beta 3$ (left) and $3\alpha 1:2\beta 3$ (right). The two binding sites for GABA at the $\beta 3(+)$ - $\alpha 1(-)$ subunit interface are indicated by red arrowheads.

of -153 ± 65 nA, $n = 6$ compared to -150 ± 93 nA, $n = 6$ when voltage-clamped at -60 mV. The constitutive current observed with both $\alpha 1 + \beta 3$ (1:1) and $\beta 3$ - $\alpha 1 + \beta 3$ (1:2) could be the result of homomeric $\beta 3$ receptors being expressed. To determine whether or not the observed holding currents are due to homomeric $\beta 3$ receptors, we evaluated oocytes expressing $\alpha 1 + \beta 3$ (1:1) and $\beta 3$ - $\alpha 1 + \beta 3$ (1:2) using histamine (1 mM), an agonist that activates only homomeric $\beta 3$ receptors^{38,39}. We found that all cells that exhibited a significant holding current injected with either $\alpha 1 + \beta 3$ (1:1) (4 out of 4 cells) or $\beta 3$ - $\alpha 1 + \beta 3$ (1:2) (1 out of 4 cells) had some response to histamine (data not shown), indicating that homomeric $\beta 3$ receptors are the major contributor to the holding current. Irrespectively, GABA activated expressed receptors in a concentration-dependent manner, with maximum peak-current amplitudes ranging from 1000 to 2000 nA. Indistinguishable EC_{50} values of $1.4 \mu\text{M}$ and $2.8 \mu\text{M}$ were obtained from oocytes injected with $\beta 3$ - $\alpha 1 + \beta 3$ (1:2) and $\alpha 1 + \beta 3$ (1:1), respectively ($p > 0.05$, Fig. 1B,C, Table 1).

Next, properties of $3\alpha 1:2\beta 3$ stoichiometry receptors were compared to receptors formed by the 30:1 injection ratio. Holding currents of oocytes injected with either $\beta 3$ - $\alpha 1 + \alpha 1$ (1:2) or free $\alpha 1 + \beta 3$ (30:1) subunits were negligible with values of -22 ± 8 nA, $n = 7$ and -13 ± 10 nA, $n = 9$, respectively. GABA activated expressed receptors in a concentration-dependent manner, with maximum peak-current amplitudes ranging from 1400 to 2900 nA. The GABA EC_{50} values of $41 \mu\text{M}$ and $26 \mu\text{M}$ from $\beta 3$ - $\alpha 1 + \alpha 1$ (1:2) and $\alpha 1 + \beta 3$ (30:1) injections were similar ($p > 0.05$, Fig. 1B,C, Table 1).

Receptor	cRNA ratio	EC ₅₀ (μM)	E _{max} (%)	n
α1 + β3	1:1	2.8 (2.4–3.3)	99 (96–102)	6
α1 + β3	30:1	26 (21–32)	96 (91–100)	9
α1 + β3 + γ2	1:1:5	53 (45–61)	99 (96–102)	5
α1 + β3 (1 μM Zolpidem)	30:1	5.6 (4.5–7.0)	97 (93–100)	5
α1 + β3 + γ2 (10 μM Zolpidem)	1:1:5	13 (12–15)	98 (96–100)	5
β3-α1 + β3	1:2	1.4 (1.2–1.8)	98 (94–101)	6
β3-α1 + α1	1:2	41 (35–48)	100 (97–103)	7
β3-α1 + α1 (1 μM Zolpidem)	1:2	7.1 (5.7–9.0)	91 (87–95)	7

Table 1. GABA concentration response relationships at various GABA_A receptors. *Xenopus laevis* oocytes were injected with cRNA mixtures containing the indicated GABA_A receptor subunits. Background subtracted peak current amplitudes for full GABA concentration-response curves in presence or absence of zolpidem were fitted to the Hill equation (fixed bottom of 0 and slope of 1) using non-linear regression and normalized to the maximal fitted value. Averaged normalized data points were next fitted to the Hill equation and resultant EC₅₀ values and maximal efficacies (E_{max}) are presented as mean with 95% confidence intervals for n experiments.

Thus, there were no observable differences in the functional properties of α1β3 receptors in either the 2α:3β or 3α:2β stoichiometries between receptors expressed by different subunit ratios or a concatenated construct (Fig. 1D). Although the chosen cRNA ratios for the individual subunits appeared sufficient to ensure predominantly uniform receptor populations, the use of concatenated receptors may potentially yield more uniform populations. However, this methodology inherently carries a risk that the physical linkage of subunits could affect receptor pharmacology. Hence, for the remainder of the manuscript, conclusions will be drawn from data generated using both methodologies.

α1β3 receptors show stoichiometry-specific sensitivity to Zn²⁺ ions. Zn²⁺ ions are used to differentiate α1β3 from α1β3γ2 GABA_A receptors²⁵. Both α1-β3 and β3-β3 interfaces contribute to the binding pockets for Zn²⁺⁴⁰. Since 3α1:2β3 receptors do not contain a β3-β3 interface, sensitivity to Zn²⁺ will likely be altered as a result of differing subunit stoichiometries. Therefore, Zn²⁺ inhibition of α1β3 receptor currents elicited by GABA (at EC₅₀ concentrations) was measured in oocytes.

Zn²⁺ (10 μM) inhibited 87 ± 10% (n = 5) of the GABA-induced current at 2α1:3β3 receptors formed by injection of α1 + β3 (1:1) cRNA (Fig. 2A). In contrast, only 13 ± 10% (n = 6) inhibition was observed at receptors formed by injecting α1 + β3 (30:1). Next, inhibitory concentration-response relationships were performed to ascertain the potency of Zn²⁺ at each stoichiometry. At the 2α1:3β3 stoichiometry, Zn²⁺ inhibited receptors from the 1:1 ratio with an IC₅₀ value of 0.84 μM (Fig. 2B). A similar IC₅₀ value of 1.6 μM for Zn²⁺ inhibition at concatenated receptors was observed with β3-α1 + β3 (1:2) injection (Fig. 2C). These two IC₅₀ values were not significantly different from each other (p > 0.05). At the highest concentration of Zn²⁺, both receptors were fully inhibited.

At the 3α1:2β3 receptors, the maximal tested concentration of Zn²⁺ (100 μM) only displayed partial inhibition (26 ± 17%, n = 6) at receptors from the 30:1 ratio (Fig. 2B). Similar partial inhibition was observed using the concatenated β3-α1 + α1 (1:2) construct (Fig. 2C). In both cases, the level of inhibition was too low to allow for meaningful fitting to the Hill equation. The partial inhibition of Zn²⁺ at 3α1:2β3 receptors mimicked observations at α1β3γ2 receptors (Fig. 2B,C).

These data demonstrate that receptors with a β3-β3 interface have high sensitivity to inhibition by Zn²⁺ (Fig. 2D). While it was previously suggested that residues located in the α1-β3 interface also contribute to Zn²⁺ sensitivity⁴⁰ this appears to require substantially higher concentrations. In addition, α1β3γ2 receptors that also lack the β3-β3 interface, were likewise relatively insensitive to Zn²⁺ inhibition at concentrations below 100 μM.

Zolpidem is a positive modulator of α1β3 with a 3α1:2β3 subunit stoichiometry. The α1-α1 interface of 3α1:2β3 receptors is homologous to the α1-γ2 interface that binds benzodiazepines and non-benzodiazepines. Zolpidem has *in vivo* effects that are not related to binding in the classical α1-γ2 benzodiazepine site, and we wanted to evaluate whether this α1 preferring modulator displayed any efficacy at α1β3 possessing an α1-α1 interface. Zolpidem (1 μM) was co-applied with a low GABA concentration (~EC₅₋₁₀) to evaluate potential modulation of control currents at 3α1:2β3 and compared with 2α1:3β3 and α1β3γ2 receptors.

As expected, zolpidem (1 μM) had no effect at 2α:3β receptors expressed by a 1:1 ratio of free subunits (Fig. 3A). In contrast, 3α1:2β3 receptors expressed from the 30:1 ratio were positively modulated by zolpidem. This resulted in an increase in current amplitudes of more than 100% compared with the current elicited by the GABA alone. This enhancement was inhibited by co-application of flumazenil (1 μM), a benzodiazepine site neutral antagonist (Fig. 3B). As expected, zolpidem (1 μM) also enhanced GABA-elicited currents at α1β3γ2 receptors, an enhancement that could be inhibited by flumazenil (Fig. 3C).

To determine the potency of zolpidem potentiation at 3α1:2β3 receptors, full concentration-response relationships were obtained and compared with those from the α1β3γ2 receptor. When injecting free subunits using α1 + β3 (30:1) and α1 + β3 + γ2 (1:1:5) cRNA, zolpidem had a significantly lower EC₅₀ value of 0.10 μM at α1β3 receptors compared to an EC₅₀ value of 0.48 μM at α1β3γ2 (p < 0.01) (Fig. 3D; Table 2). However, when injecting cRNA containing the concatenated construct using β3-α1 + α1 (1:2) and β3-α1 + γ2 (1:2), the EC₅₀ value




Figure 2. Zn^{2+} inhibition of GABA-evoked currents from $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ GABA_A receptors. *Xenopus laevis* oocytes were injected with cRNA and subjected to two-electrode voltage clamp electrophysiology as described in the methods. Control currents (I_{control}) were evoked using a GABA concentration corresponding to $\sim EC_{50}$ and inhibition by Zn^{2+} was evaluated by co-applications with GABA_{control}. (A) Representative GABA-evoked current traces from oocytes injected with the denoted cRNA mixtures. Bars above each trace indicate GABA and Zn^{2+} application periods. Dotted lines indicate the peak current amplitude by GABA_{control} and “/” a wash period. (B,C) Concentration–response relationships of Zn^{2+} inhibition of GABA_{control}-evoked currents at $\alpha 1\beta 3$ or $\alpha 1\beta 3\gamma 2$ receptors stemming from injecting the indicated cRNA mixtures using free subunits (B) or a concatenated $\beta 3\text{-}\alpha 1$ construct (C). Averaged Zn^{2+} inhibition values were depicted as means \pm S.E.M as a function of the Zn^{2+} concentration and fitted to the Hill equation by non-linear regression. Regression results for $\alpha 1 + \beta 3$ (1:1) were $IC_{50} = 0.84$ (95% CI: 0.53–1.3), $nH = -0.5 \pm 0.32$ and for $\beta 3\text{-}\alpha 1 + \beta 3$ (1:2) were $IC_{50} = 1.6$ (95% CI: 1.1–2.4), $nH = -0.6 \pm 0.06$. For the remaining cRNA mixtures, Zn^{2+} inhibition at the maximal tested concentration was too low to allow for meaningful fitting. Each data point represent experiments from $n = 5\text{--}8$ oocytes of ≥ 2 batches. (D) The depiction of $\alpha 1\beta 3$ GABA_A receptor stoichiometries from Fig. 1D was modified to indicate Zn^{2+} binding in the $\beta 3(+)\text{-}\beta 3(-)$ subunit interface (red/orange arrowhead).

of $0.030 \mu\text{M}$ at $\alpha 1\beta 3$ receptors was not significantly different to the EC_{50} value of $0.050 \mu\text{M}$ at $\alpha 1\beta 3\gamma 2$ receptors ($p > 0.05$) (Fig. 3E; Table 2). Efficacy levels at $\alpha 1\beta 3$ receptors were 120% and 110% for the 30:1 ratio and concatenated receptors, respectively. Higher values were observed at $\alpha 1\beta 3\gamma 2$ receptors with 340% and 350% at $\alpha 1 + \beta 3 + \gamma 2$ (1:1:5) vs. $\beta 3\text{-}\alpha 1 + \gamma 2$ (1:2), respectively.

Hence zolpidem modulated GABA-evoked currents at $3\alpha 1:2\beta 3$ receptors. Importantly, the potency of zolpidem at $\alpha 1\beta 3$ ($3\alpha 1:2\beta 3$) receptors was comparable with $\alpha 1\beta 3\gamma 2$ receptors regardless of whether free subunits or concatenated subunits were injected. While the potency of zolpidem was similar at both receptors, these experiments suggest that zolpidem enhanced $\alpha 1\beta 3\gamma 2$ receptors with greater efficacy. However, in this type of assay efficacy of a modulator is highly dependent on the utilized GABA concentration and should be treated with caution.

Modulatory mechanism of action of zolpidem at $\alpha 1\beta 3$ receptors possessing a $3\alpha 1:2\beta 3$ subunit stoichiometry. The hallmark feature of allosteric modulation via the $\alpha 1\text{-}\gamma 2$ benzodiazepine site by e.g. diazepam and zolpidem is an increase in the apparent potency of GABA. This causes a shift in the GABA concentration–response curve to the left in presence of zolpidem with little accompanying change in the maximal current amplitudes. In order to assess how zolpidem affects $3\alpha 1:2\beta 3$ stoichiometry, GABA concentration–response relationships were measured in the presence of zolpidem and compared to that in its absence.




Figure 3. Zolpidem modulation of GABA-evoked currents from $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ GABA_A receptors. *Xenopus laevis* oocytes were injected with cRNA and subjected to two-electrode voltage clamp electrophysiology as described in the methods. Control currents (I_{control}) were evoked using a GABA concentration corresponding to $\sim EC_{5-10}$ and modulation by zolpidem was evaluated by co-applications with GABA_{control}. (A–C) Representative GABA-evoked current traces from oocytes injected with the denoted cRNA mixtures. Bars above each trace indicate GABA, zolpidem (Zolp) and flumazenil (Flu) concentrations and application periods. Dotted lines indicate the peak current amplitude by GABA_{control} and “/” a wash period. For the specific traces, zolpidem had no robust effects at receptors from $\alpha 1 + \beta 3$ (1:1) injection (A), but showed 130% modulation at receptors from $\alpha 1 + \beta 3$ (30:1) injection which was inhibited 95% by co-application of flumazenil (B). Zolpidem likewise modulated receptors from injection of $\alpha 1 + \beta 3 + \gamma 2$ (1:1:5) by 160% which could be inhibited 85% by flumazenil (C). (D,E) Concentration-response relationships of zolpidem modulation of GABA_{control}-evoked currents at $\alpha 1\beta 3$ or $\alpha 1\beta 3\gamma 2$ receptors stemming from injecting the indicated cRNA mixtures using free subunits (D) or a concatenated $\beta 3 - \alpha 1$ construct (E). Average modulatory values were depicted as means \pm S.E.M as a function of the zolpidem concentration and fitted to the Hill equation by non-linear regression. Each data point represents experiments from $n = 5-7$ oocytes from ≥ 2 batches and regression results are presented in Table 1.

Receptor	cRNA ratio	EC ₅₀ (μM)	E _{max} (%)	n
$\alpha 1 + \beta 3$	30:1	0.10 (0.06–0.16)	120 (110–130)	5
$\alpha 1 + \beta 3 + \gamma 2$	1:1:5	0.48 (0.31–0.78)	340 (300–380)	6
$\beta 3 - \alpha 1 + \alpha 1$	1:2	0.032 (0.015–0.070)	111 (96–126)	9
$\beta 3 - \alpha 1 + \gamma 2$	1:2	0.045 (0.035–0.057)	348 (333–363)	5

Table 2. Zolpidem modulation of GABA-evoked currents at various GABA_A receptors. *Xenopus laevis* oocytes were injected with cRNA mixtures containing the indicated GABA_A receptor subunits. Zolpidem was co-applied with a control concentration of GABA corresponding to an EC₅₋₁₀ value. Modulatory efficacies of zolpidem were calculated as percentage change from the GABA_{control}-evoked currents. Averaged efficacies were next fitted to the Hill equation (fixed bottom of 0 and slope of 1) and resultant EC₅₀ values and maximal efficacies (E_{max}) are presented as mean with 95% confidence intervals for n experiments.

At $3\alpha 1:2\beta 3$ receptors expressed by injecting $\alpha 1 + \beta 3$ (30:1) cRNA, zolpidem (1 μM) left-shifted the GABA concentration-response curve by decreasing the EC₅₀ value from 26 μM to 5.6 μM (Fig. 4A, Table 1). Hence, zolpidem modulation caused a significant 5-fold change ($p < 0.0001$) of the GABA potency with no observed change in the maximum GABA-evoked peak current amplitudes. At receptors obtained by the concatenated $\beta 3 - \alpha 1 + \alpha 1$ (1:2) construct, zolpidem (1 μM) likewise caused a significant ($p < 0.0001$) 6-fold change in the potency of GABA, with an increase in the EC₅₀ value from 41 μM to 7.1 μM (Fig. 4B, Table 1). At $\alpha 1\beta 3\gamma 2$ receptors, zolpidem (10 μM)




Figure 4. Mechanism of zolpidem modulatory actions at $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ GABA_A receptors. *Xenopus laevis* oocytes were injected with cRNA and subjected to two-electrode voltage clamp electrophysiology as described in the methods. (A–C) Full GABA concentration–response relationships were obtained in presence of the indicated concentrations of zolpidem (Zolp) at receptors stemming from injection of cRNA of free subunits (A,C) or a concatenated construct (B). Baseline subtracted GABA + zolpidem peak current amplitudes were normalized to a maximal GABA control response (3 mM) in the same oocytes. Averaged normalized data points are depicted as means \pm S.E.M. as a function of the GABA concentration and fitted to the Hill equation with regression results are presented in Table 1. Each data point represents experiments from $n = 5–7$ oocytes from ≥ 2 batches. GABA concentration response relationships in absence of zolpidem (from Fig. 1) are included for comparison. (D) The depiction of $\alpha 1\beta 3$ GABA_A receptor stoichiometries from Fig. 2D was modified to indicate zolpidem binding in the $\alpha 1(+)-\alpha 1(-)$ subunit interface (red/green arrowhead).

significantly decreased the EC_{50} value from 53 μ M to 13 μ M (Fig. 4C; Table 1; $p < 0.0001$) with no change in the maximum peak current amplitudes.

These data demonstrate that the mechanism of modulatory action by zolpidem is increase of the GABA potency at both $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ receptors. This increase ranged from 4 to 6 fold, which is in agreement with previous observations at $\alpha 1\beta 3\gamma 2$ receptors⁸. The modulatory effect was not accompanied by any change in maximal GABA-evoked peak current amplitudes and the similar magnitude of the changes in EC_{50} values suggest that zolpidem has a similar efficacy at the two receptor types. Although zolpidem is not structurally a classical benzodiazepine, it binds at a similar site within the $\alpha 1-\gamma 2$ of $\alpha 1\beta 3\gamma 2$ receptors as benzodiazepines. Taken together, zolpidem is most likely binding at the $\alpha 1-\alpha 1$ interface of $3\alpha:2\beta$ receptors to modulate receptor function analogous to the modulation of receptor function via binding at the $\alpha 1-\gamma 2$ interface (Fig. 4D).

Diazepam enhances GABA currents at $\alpha 1\beta 3$ receptors with a $3\alpha 1:2\beta 3$ subunit stoichiometry albeit with low efficacy. Finally, we determined whether $\alpha 1\beta 3$ ($3\alpha 1:2\beta 3$) receptors could be modulated by the classical benzodiazepine diazepam, which binds with equal affinity to GABA_A receptors containing $\alpha 1$, $\alpha 2$, $\alpha 3$ and $\alpha 5$ subunits^{41,42}. Like zolpidem, diazepam (1 μ M) enhanced GABA-induced currents at $\alpha 1\beta 3$ receptors obtained with $\alpha 1 + \beta 3$ (30:1) cRNA and this effect was inhibited by co-application of flumazenil (Fig. 5A). However, the enhancement by diazepam was low in comparison with zolpidem (Fig. 5B). A full concentration–response relationship revealed an EC_{50} value for diazepam of 0.040 μ M (Fig. 5C). A similar EC_{50} value of 0.020 μ M was estimated for diazepam modulation of concatenated receptors and the observed potencies




Figure 5. Diazepam modulation of GABA-evoked currents from $\alpha 1\beta 3$ GABA_A receptors. *Xenopus laevis* oocytes were injected with cRNA and subjected to two-electrode voltage clamp electrophysiology as described in the methods. Control currents (I_{control}) were evoked using a GABA concentration corresponding to $\sim EC_{5-10}$ and modulation by diazepam was evaluated by co-applications with GABA_{control}. (A,B) Representative GABA-evoked current traces from oocytes injected with the denoted cRNA mixtures. Bars above each trace indicate GABA, diazepam (Diaz) and flumazenil (Flu) concentrations and application periods. Dotted lines indicate the peak current amplitude by GABA_{control} and “/” a wash period. (C) Concentration-response relationships of diazepam modulation of GABA_{control}-evoked currents at $\alpha 1\beta 3$ receptors stemming from injecting the indicated cRNA mixtures using free subunits or a concatenated $\beta 3 - \alpha 1$ construct. Averaged modulatory values were depicted as means \pm S.E as a function of the diazepam concentration and fitted to the Hill equation by non-linear regression. Each data point represents experiments from $n = 6$ oocytes from ≥ 2 batches. Regression results with 95% confidence intervals for $\alpha 1 + \beta 3$ (30:1) were: EC_{50} value of $0.040 \mu\text{M}$ (0.010–0.12) and E_{max} value of 40% (33–48) whereas results for $\beta 3 - \alpha 1 + \alpha 1$ (1:2) were: EC_{50} value of $0.020 \mu\text{M}$ (0.010–0.04) and E_{max} value of 51% (46–56).

are similar to that reported for $\alpha 1\beta 3\gamma 2$ receptors⁸. From the concentration-response relationships, it is evident that the enhancement by diazepam displayed lower efficacy compared with zolpidem. When using free subunit cRNA diazepam enhanced GABA-elicited currents by a maximum of 40% whereas use of concatenated subunits resulted in 51% modulation and these were not significantly different from each other (Fig. 5C; $p > 0.05$). This represents half or less of the modulation observed with zolpidem (Table 2). Analogous to zolpidem, diazepam ($1 \mu\text{M}$) did not change the maximal GABA-evoked current amplitudes at $2\alpha 1:3\beta 3$ receptors obtained by injecting $\alpha 1 + \beta 3$ (1:1) (data not shown).

Discussion

Binary GABA_A receptors co-exist with their ternary counterparts^{23–25} and are gaining attention for their sensitivity to certain benzodiazepines^{25–27}. When evaluating the pharmacology of binary receptors the concept of stoichiometry-specific binding sites becomes important, because each receptor stoichiometry will have a specific binding interface that will contribute to a distinct pharmacological profile for that receptor^{34–36}. For pentameric $\alpha 1\beta 3$ receptors, the third subunit position, normally occupied by a $\gamma 2$ subunit in $\alpha 1\beta 3\gamma 2$ receptors, is substituted with either an $\alpha 1$ or a $\beta 3$ subunit leading to two distinct receptors, that differ by the presence of either an $\alpha 1 - \alpha 1$ or a $\beta 3 - \beta 3$ interface (Fig. 1D). Pharmacologically targeting either the $\alpha 1 - \alpha 1$ or $\beta 3 - \beta 3$ interface leads to stoichiometric selective effects as demonstrated by the *in vivo* and *in vitro* effects of drugs at the related nicotinic acetylcholine $\alpha 4\beta 2$ receptors^{34,35}. In this study, we demonstrate that binary $\alpha 1\beta 3$ receptors composed of $2\alpha 1:3\beta 3$ or $3\alpha 1:2\beta 3$ subunit stoichiometry were differentiated from each other using GABA, Zn^{2+} , zolpidem and diazepam.

Two methodologies were used to express the two receptor stoichiometries ($2\alpha 1:3\beta 3$ or $3\alpha 1:2\beta 3$) in *Xenopus laevis* oocytes. In the first method, we injected cRNA mixtures with variant subunits ratios, causing relative overexpression of one subunit versus the other, and hence, increasing the likelihood of receptor assembly with more of the overexpressed subunit. While this is an efficient method, it cannot always ensure uniform receptor populations or rule out the formation of “obscure” receptor populations such as $4\alpha:1\beta$ receptors. In the second method, concatenated $\beta 3 - \alpha 1$ subunits were used, ensuring that linkers force the assembly process. While this is efficient for creating uniform receptor populations, the presence of linkers may affect receptor pharmacology. Nevertheless, we found that both methods enabled the assembly of $\alpha 1\beta 3$ receptors, which were indistinguishable in their pharmacology using either methodology and concluded the receptors to be either $2\alpha 1:3\beta 3$ or $3\alpha 1:2\beta 3$ receptors. This contrasted the original study using concatenated $\beta 2 - \alpha 1$ subunits and individual $\alpha 1$ or $\beta 2$ subunit mRNA. It is not clear why Baumann and colleagues did not see functional receptors when using $\beta 2 - \alpha 1$ and individual $\alpha 1$ subunits⁴³. However the study used a 1:1 injection ratio while we used a 2-fold excess of $\alpha 1$ subunit mRNA. Further we used the $\beta 3$ subunit and not $\beta 2$ and there may be differences in the formation of receptors depending on the type of β subunit.

Xenopus oocytes expressing $\alpha 1\beta 3$ ($2\alpha 1:3\beta 3$) receptors consistently displayed voltage-clamp holding-current levels constituting approximately 10% of maximal GABA-evoked peak current amplitudes. This suggests the existence of homomeric $\beta 3$ receptors contributing to the holding current as histamine was able to activate cells that exhibited constitutive currents^{38,39}. In contrast, expression of $3\alpha 1:2\beta 3$ receptors did not result in levels significantly different from un-injected control oocytes. However, the potency of GABA differed approximately 10-fold between $2\alpha 1:3\beta 3$ and $3\alpha 1:2\beta 3$ receptors with $2\alpha 1:3\beta 3$ displaying the highest GABA sensitivity. The lower GABA sensitivity at $3\alpha 1:2\beta 3$ receptors was not significantly different to the “classical” $\alpha 1\beta 3\gamma 2$ receptor indicating “normal” receptor function.

Previous studies demonstrated that Zn^{2+} inhibits binary GABA_A $\alpha 1\beta 3$ receptor function by an allosteric mechanism and this action is critically dependent on the composition of $\alpha 1$ and $\beta 3$ subunits^{40,44}. This is consistent with our data showing that binary GABA_A receptors can be differentiated by Zn^{2+} , showing that $2\alpha 1:3\beta 3$ receptors are easily blocked by Zn^{2+} while $3\alpha 1:2\beta 3$ receptors are not. Thus, the presence of an extra $\beta 3$ subunit, and thereby a $\beta 3$ - $\beta 3$ interface, confers the resulting binary receptors with several unique characteristics such as, higher sensitivity for GABA and higher sensitivity for Zn^{2+} inhibition. The high sensitivity to Zn^{2+} inhibition is intriguing given their extrasynaptic location. Physiological brain concentrations of Zn^{2+} can reach up to $100\mu M$ ^{45–48}, *i.e.* the sweet-spot for completely inhibiting $2\alpha 1:3\beta 3$ receptors, and one could speculate that these receptors may in part be responsible for a Zn^{2+} -regulated tonic current.

In contrast, binary receptors with an $\alpha 1$ - $\alpha 1$ interface have more resemblance to ternary $\alpha 1\beta 3\gamma 2$ receptors. They exhibit a lower degree of constitutive activity, and lower sensitivity for GABA and Zn^{2+} inhibition. Furthermore, we demonstrate that zolpidem efficiently modulates GABA activity at $3\alpha 1:2\beta 3$ but not $2\alpha 1:3\beta 3$ receptors with a binding site for zolpidem most likely at the $\alpha 1$ - $\alpha 1$ interface. Several lines of evidence point to zolpidem's actions occurring via the $\alpha 1$ - $\alpha 1$ interface of the $3\alpha 1:2\beta 3$ receptor in a manner that mimics the potentiation of zolpidem via binding to the classical $\alpha 1$ - $\gamma 2$ benzodiazepine site: i) Like zolpidem, diazepam is a positive allosteric modulator of $3\alpha 1:2\beta 3$ receptors albeit with lower efficacy levels; ii) the effects of zolpidem and diazepam are inhibited by the antagonist flumazenil at a concentration normally used to inhibit benzodiazepines at the $\alpha 1$ - $\gamma 2$ benzodiazepine site; and iii) the modulatory effect of zolpidem at $3\alpha 1:2\beta 3$ receptors was one of increasing the potency of GABA without significant change to maximal current amplitudes. These data comprise signature features of modulatory actions by benzodiazepines via the classical $\alpha 1$ - $\gamma 2$ interface benzodiazepine site^{4,10,11}, however, as such a site is not available in $2\alpha 1:3\beta 3$ receptors, zolpidem's actions must occur via the $\alpha 1$ - $\alpha 1$ interface. Interestingly, the potency of zolpidem at $3\alpha 1:2\beta 3$ receptors was equal to or marginally higher than that observed at $\alpha 1\beta 3\gamma 2$ receptors suggesting that the observed effects could have clinical relevance.

In addition to phasic currents, zolpidem modulates tonic GABA currents in the rat dorsal motor nucleus of the vagus²⁰ and primary motor cortex¹⁵. This is intriguing as the subunit composition of zolpidem-sensitive extrasynaptic receptors is unlikely to be due to $\alpha 5$, $\gamma 1/3$ or δ containing GABA_A receptors, as such receptors do not significantly respond to zolpidem^{8,21}. Recently, $\alpha 1\beta 3\gamma 2$ receptors were reported to exist at extrasynaptic sites⁴⁹ but the residual effects of zolpidem in $\gamma 2F771$ knockin mice excludes these receptors as a target for zolpidem. Although non-GABA_A receptors may also contribute to zolpidem's effects, binary $\alpha 1\beta 3$ receptors represent 10% of the total number of extrasynaptic receptors in hippocampal pyramidal neurons²⁵. Thus binary $\alpha 1\beta 3$ receptors that possess an $\alpha 1$ - $\alpha 1$ subunit interface may contribute to the residual zolpidem effects reported in $\gamma 2F771$ knockin mice that lack zolpidem-sensitive α - $\gamma 2$ interfaces^{19,50,51}.

Binary $\alpha 1\beta 3$ receptors that are positively modulated by zolpidem may also have significant clinical effects in a number of neurological conditions where the ratio and location of various GABA receptor subunits are altered, for example, stroke^{52–54}, or epilepsy^{55,56}. Under these conditions, dramatic changes are occurring in the brain. After stroke the $\alpha 1$ mRNA significantly increases^{52–54}, resulting in an increase in $\alpha 1$ protein⁵². At the same time β -subunit mRNA decreases, and thus may affect the formation of GABA_A receptors in a way that the subunit composition of the receptors is altered favouring receptors that possess an $\alpha 1$ - $\alpha 1$ subunit interface. In order to ascertain that the increase in $\alpha 1$ protein after stroke results in a change in receptor function that mimics the pharmacology of binary $\alpha 1\beta 3$ receptors possessing an $\alpha 1$ - $\alpha 1$ interface, further work including patch clamp recordings are required.

In corroboration, there are increasing clinical reports documenting that zolpidem is not a typical hypnotic, improving cognitive and motor function in human patients with severe brain injury. For instance, acute administration of zolpidem in a patient suffering from brain injury resulted in a transient improvement in aphasia⁵⁷. Brefel-Courbon and colleagues (2007) reported improved motor performance and neuropsychological status in a patient who was given zolpidem⁵⁸. In addition, zolpidem has benefits on motor disorders in patients with Parkinson's disease^{13–15}, has a transient improvement of spinocerebellar ataxia¹⁶, and improve symptoms of progressive supranuclear palsy¹⁷. Whilst flumazenil inhibits the effect of zolpidem indicating a GABA_A receptor is involved with these patients⁵⁹, classical benzodiazepines are not very effective in improving speech or motor function¹⁷ indicating an unusual GABA_A receptor to be the target.

Similar positive modulation is seen by zolpidem at $\alpha 1$ - $\alpha 1$ and $\alpha 1$ - $\gamma 2$ interfaces, however the complete molecular interactions involving ligand binding and the accompanying conformational changes to the receptor complex still needs to be resolved. While the principal face (+) of the $\alpha 1$ subunit is the same in the two sites, the complementary face (–) of $\alpha 1$ and $\gamma 2$ are obviously different. Although the actions of zolpidem, diazepam and flumazenil suggest substantial overlap between the two sites, other compound structures could potentially be fully selective for the $\alpha 1$ - $\alpha 1$ interface. Future studies are required to ascertain the structure-function requirements for this interface.

In conclusion, this study demonstrates GABA_A $\alpha 1\beta 3$ receptors can express in either $2\alpha 1:3\beta 3$ and $3\alpha 1:2\beta 3$ stoichiometry possessing either a $\beta 3$ - $\beta 3$ or $\alpha 1$ - $\alpha 1$ interface, respectively. This difference leads to stoichiometry-specific binding sites, fundamentally changing receptor pharmacology. Indeed, zolpidem at clinically relevant concentrations enhanced GABA actions at $3\alpha 1:2\beta 3$ but not $2\alpha 1:3\beta 3$ receptors by binding to the

$\alpha 1$ - $\alpha 1$ interface in a manner mimicking its actions at the classical benzodiazepine site. These receptors may be the molecular target for zolpidem's described extrasynaptic and clinical effects and constitute a potentially interesting drug target for neurological conditions (e.g. stroke and epilepsy) where the $\alpha 1$ mRNA and protein levels are significantly upregulated.

Methods

Chemicals. GABA (4-aminobutanoic acid), flumazenil (Ethyl 8-fluoro-5-methyl-6-oxo-5,6-dihydro-4H-benzo[f]imidazo[1,5-a][1,4]diazepine-3-carboxylate), pyruvate, theophylline, gentamycin and zinc chloride were obtained from Sigma Aldrich, Sydney, Australia. Diazepam was obtained from Apin Chemicals Ltd, Oxon, UK, and zolpidem (*N,N*-dimethyl-2-(6-methyl-2-*p*-tolylimidazo[1,2-*a*]pyridin-3-yl)acetamide) from Chemieliva Pharmaceutical, Chongqing, China. Tricaine was purchased from Western Chemical, USA.

GABA_A receptor subunit cRNA. Human $\alpha 1$ and $\gamma 2$ cDNA subcloned in pcDM8 and $\beta 3$ in pGEMHE, were linearized with the appropriate restriction endonucleases (NotI for $\alpha 1$ and $\gamma 2$, NheI for $\beta 3$ subunit, respectively). Concatenated $\beta 3$ - $\alpha 1$ subunits were developed as previously described⁴³, subcloned in pNS3z and linearized with NotI. cRNA was produced from linearized plasmids using the 'mMessage mMachine' T7 transcript kit from Ambion (Austin, TX, USA) as previously described^{60,61}. A total of 2–4 ng of cRNA was injected per oocyte. When using free subunits of $\alpha 1$, $\beta 3$ and $\gamma 2$, cRNAs were mixed in variant ratios. To ensure incorporation of a free subunit in the pentameric complex, concatenated $\beta 3$ - $\alpha 1$ subunit cRNA was injected with either $\alpha 1$ or $\gamma 2$ cRNAs in a ratio of 1:2. However to avoid any potential formation of $\beta 3$ homomeric receptors concatenated $\beta 3$ - $\alpha 1$ subunit cRNA was injected with $\beta 3$ in a 1:1 ratio.

Xenopus oocyte extraction and preparation. All procedures using *Xenopus laevis* frogs were approved by the animal ethics committee of The University of Sydney (AEC No. 2013/5269) and are in accordance to the National Health and Medical Research Council (NHMRC) of Australia. In brief, *Xenopus laevis* were anaesthetized using 0.2% w/v solution of tricaine methanesulfonate or tricaine-S (Western Chemical, USA). A transverse incision of about 3 mm in length was made through the outer layer of her skin on the lateral ventral surface. Another incision was made through the connective tissue and muscle layer to reach the ovary wall. A section of the ovary was carefully removed onto the surface of the frog. A small section of ovary was separated and transferred to a tube containing the OR2 solution (82.5 mM NaCl, 5 mM HEPES, 2 mM MgCl₂ and 2 mM KCl; pH 7.4). Oocytes were separated manually from their follicles and digested using 40 mg collagenase A diluted in 15 ml OR2 solution at 18 °C for about 1 hour until the oocytes were fully detached from the follicles and the ovary tissue.

Stage V-VI oocytes were injected with 50.6 nl cRNA solution composed of GABA_ARs subunit cRNAs in the required ratio. The injected oocytes were incubated in ND96 solution (96 mM NaCl, 5 mM HEPES, 2 mM MgCl₂, 1 mM KCl and 1.8 mM CaCl₂; pH 7.4) supplemented with 2.5 mM Na-pyruvate, 0.5 mM theophylline, 50 mg/ml gentamycin and 50 mg/ml tetracycline for 2–5 days at 18 °C.

Two-electrode voltage-clamp electrophysiology. The electrophysiological experiments were performed by the two-electrode voltage clamp technique. The membrane potential was measured using an Oocyte Clamp OC-725C amplifier (Warner Instruments Corp, CT, USA) by the voltage electrode with current simultaneously injected by the current electrode to maintain the potential difference across the oocyte membrane at –60 mV. Data were acquired with a LabChart v. 3.5.2 analogue to digital converter and currents low-pass-filtered at 1 kHz and sampled at 3 kHz were measured offline with LabChart v. 3.5.2 software. The bath solution contained the ND96 solution and electrodes were filled with solution of 3 M KCl (0.5–2 M Ω). Solutions were bath applied using a gravity-fed perfusion system running at 5 ml/min.

Data Analysis. For Zn²⁺ inhibition studies, control currents (I_{control}) were evoked using a GABA concentration corresponding to ~EC₅₀ and inhibition by Zn²⁺ was evaluated by co-applications with GABA_{control}. Averaged Zn²⁺ inhibition in the presence of GABA EC₅₀ were depicted as means \pm S.E.M. as a function of the Zn²⁺ concentration and fitted to the Hill equation by non-linear regression. Enhancement of GABA-gated Cl[–] currents was measured by co-applying modulators with a GABA concentration that elicited 5% of the maximal current amplitude as determined at the beginning of each experiment. The enhancement of the GABA-gated Cl[–] current (I_{GABA}) was defined as $I = I_{\text{max}} / (1 + [EC_{50} / (A)]^{n_H})$; A is the agonist concentration, I is the current and I_{max} is the maximum current, EC₅₀ is the concentration of GABA that produces a response that is 50% of the maximum current, n_H is the Hill Coefficient. EC₅₀ values are expressed as mean with 95% confidence intervals and Hill coefficients (n_H) are expressed as mean \pm S.E.M.

Concentration–response curves were generated and the data fitted by non-linear regression analysis using GraphPad Prism software (version 5.0). Statistical significance was calculated using unpaired Student *t*-test with a confidence interval of $p < 0.05$ to compare parameters derived from individual experiments and data given as mean with 95% confidence intervals from at least 5 oocytes and at least 2 different batches. The logEC₅₀, derived from individual comparisons, were used for statistical comparisons.

References

- Chebib, M. & Johnston, G. A. GABA-Activated ligand gated ion channels: medicinal chemistry and molecular biology. *J Med Chem* **43**, 1427–1447 (2000).
- Farrant, M. & Nusser, Z. Variations on an inhibitory theme: phasic and tonic activation of GABA(A) receptors. *Nat Rev Neurosci* **6**, 215–229 (2005).
- Sieghart, W. Structure, pharmacology, and function of GABA_A receptor subtypes. *Adv Pharmacol* **54**, 231–263 (2006).
- Sieghart, W. Allosteric modulation of GABA_A receptors via multiple drug-binding sites. *Adv Pharmacol* **72**, 53–96 (2015).
- Sancar, F., Ericksen, S. S., Kucken, A. M., Teissere, J. A. & Czajkowski, C. Structural determinants for high-affinity zolpidem binding to GABA_A receptors. *Mol Pharmacol* **71**, 38–46 (2007).

6. Hanson, S. M., Morlock, E. V., Satyshur, K. A. & Czajkowski, C. Structural requirements for eszopiclone and zolpidem binding to the gamma-aminobutyric acid type-A (GABA_A) receptor are different. *J Med Chem* **51**, 7243–7252 (2008).
7. Baur, R. & Sigel, E. Replacement of histidine in position 105 in the alpha(5) subunit by cysteine stimulates zolpidem sensitivity of alpha(5)beta(2)gamma(2) GABA_A receptors. *J Neurochem* **103**, 2556–2564 (2007).
8. Ramerstorfer, J., Furtmuller, R., Vogel, E., Huck, S. & Sieghart, W. The point mutation gamma 2F771 changes the potency and efficacy of benzodiazepine site ligands in different GABA_A receptor subtypes. *Eur J Pharmacol* **636**, 18–27 (2010).
9. Sanna, E. *et al.* Comparison of the effects of zaleplon, zolpidem, and triazolam at various GABA_A receptor subtypes. *Eur J Pharmacol* **451**, 103–110 (2002).
10. Sigel, E. Mapping of the benzodiazepine recognition site on GABA_A receptors. *Curr Topics Med Chem* **2**, 833–839 (2002).
11. Ernst, M., Brauchart, D., Boesch, S. & Sieghart, W. Comparative modeling of GABA_A receptors: limits, insights, future developments. *Neurosci* **119**, 933–943 (2003).
12. Patat, A. *et al.* Flumazenil antagonizes the central effects of zolpidem, an imidazopyridine hypnotic. *Clin Pharmacol Ther* **56**, 430–436 (1994).
13. Chen, Y. Y., Sy, H. N. & Wu, S. L. Zolpidem improves akinesia, dystonia and dyskinesia in advanced Parkinson's disease. *J Clin Neurosci* **15**, 955–956 (2008).
14. Hall, S. D. *et al.* GABA-mediated changes in inter-hemispheric beta frequency activity in early-stage Parkinson's disease. *Neurosci* **281C**, 68–76 (2014).
15. Prokic, E. J. *et al.* Cortical oscillatory dynamics and benzodiazepine-site modulation of tonic inhibition in fast spiking interneurons. *Neuropharmacol* **95**, 192–205 (2015).
16. Clauss, R., Sathekge, M. & Nel, W. Transient improvement of spinocerebellar ataxia with zolpidem. *New Eng J Med* **351**, 511–512 (2004).
17. Chang, A. Y. & Weirich, E. Trial of Zolpidem, Eszopiclone, and Other GABA Agonists in a Patient with Progressive Supranuclear Palsy. *Case Rep Med* **2014**, 107064 (2014).
18. Cope, D. W. *et al.* Loss of zolpidem efficacy in the hippocampus of mice with the GABA_A receptor gamma2 F771 point mutation. *Eur J Neurosci* **21**, 3002–3016 (2005).
19. May, A. C., Fleischer, W., Kletke, O., Haas, H. L. & Sergeeva, O. A. Benzodiazepine-site pharmacology on GABA_A receptors in histaminergic neurons. *Br J Pharmacol* **170**, 222–232 (2013).
20. Gao, H. & Smith, B. N. Zolpidem modulation of phasic and tonic GABA currents in the rat dorsal motor nucleus of the vagus. *Neuropharmacol* **58**, 1220–1227 (2010).
21. Khom, S. *et al.* Pharmacological properties of GABA_A receptors containing gamma1 subunits. *Mol Pharmacol* **69**, 640–649 (2006).
22. Buhr, A. & Sigel, E. A point mutation in the gamma2 subunit of gamma-aminobutyric acid type A receptors results in altered benzodiazepine binding site specificity. *Proc Nat Acad Sci* **94**, 8824–8829 (1997).
23. Brickley, S. G., Cull-Candy, S. G. & Farrant, M. Single-channel properties of synaptic and extrasynaptic GABA_A receptors suggest differential targeting of receptor subtypes. *J Neurosci* **19**, 2960–2973 (1999).
24. Sieghart, W. & Sperk, G. Subunit composition, distribution and function of GABA_A receptor subtypes. *Curr Topics Med Chem* **2**, 795–816 (2002).
25. Mortensen, M. & Smart, T. G. Extrasynaptic alphabeta subunit GABA_A receptors on rat hippocampal pyramidal neurons. *J Physiol* **577**, 841–856 (2006).
26. Bencsits, E., Ebert, V., Tretter, V. & Sieghart, W. A significant part of native gamma-aminobutyric AcidA receptors containing alpha4 subunits do not contain gamma or delta subunits. *J Biol Chem* **274**, 19613–19616 (1999).
27. Sinkkonen, S. T., Luscher, B., Luddens, H. & Korpi, E. R. Autoradiographic imaging of altered synaptic alphabeta2 and extrasynaptic alphabeta GABA_A receptors in a genetic mouse model of anxiety. *Neurochem Int* **44**, 539–547 (2004).
28. Walters, R. J., Hadley, S. H., Morris, K. D. & Amin, J. Benzodiazepines act on GABA_A receptors via two distinct and separable mechanisms. *Nat. Neurosci.* **3**, 1274–1281 (2000).
29. Baur, R. *et al.* Covalent modification of GABA_A receptor isoforms by a diazepam analogue provides evidence for a novel benzodiazepine binding site that prevents modulation by these drugs. *J Neurochem* **106**, 2353–2363 (2008).
30. Ramerstorfer, J. *et al.* The GABA_A receptor alpha + beta- interface: a novel target for subtype selective drugs. *J Neurosci* **31**, 870–877 (2011).
31. Tretter, V., Ehya, N., Fuchs, K. & Sieghart, W. Stoichiometry and assembly of a recombinant GABA_A receptor subtype. *J Neurosci* **17**, 2728–2737 (1997).
32. Farrar, S. J., Whiting, P. J., Bonnert, T. P. & McKernan, R. M. Stoichiometry of a ligand-gated ion channel determined by fluorescence energy transfer. *J Biol Chem* **274**, 10100–10104 (1999).
33. Boileau, A. J., Pearce, R. A. & Czajkowski, C. Tandem subunits effectively constrain GABA_A receptor stoichiometry and recapitulate receptor kinetics but are insensitive to GABA_A receptor-associated protein. *J Neurosci* **25**, 11219–11230 (2005).
34. Harpsøe, K. *et al.* Unraveling the high- and low-sensitivity agonist responses of nicotinic acetylcholine receptors. *J Neurosci* **31**, 10759–10766 (2011).
35. Ahring, P. K. *et al.* Engineered alpha4beta2 nicotinic acetylcholine receptors as models for measuring agonist binding and effect at the orthosteric low-affinity alpha4-alpha4 interface. *Neuropharmacol* **92**, 135–145 (2015).
36. Chua, H. C., Absalom, N. L., Hanrahan, J. R., Viswas, R. & Chebib, M. The Direct Actions of GABA, 2'-Methoxy-6-Methylflavone and General Anaesthetics at beta3gamma2L GABA_A Receptors: Evidence for Receptors with Different Subunit Stoichiometries. *PLoS one* **10**, e0141359 (2015).
37. Kaur, K. H., Baur, R. & Sigel, E. Unanticipated structural and functional properties of delta-subunit-containing GABA_A receptors. *J Biol Chem* **284**, 7889–7896 (2009).
38. Saras, A. *et al.* Histamine action on vertebrate GABA_A receptors: direct channel gating and potentiation of GABA responses. *J Biol Chem* **283**, 10470–10475 (2008).
39. Hoerbelt, P. *et al.* Mutagenesis and computational docking studies support the existence of a histamine binding site at the extracellular beta3 + beta3- interface of homooligomeric beta3 GABA receptors. *Neuropharmacol* **108**, 252–263 (2016).
40. Hosie, A. M., Dunne, E. L., Harvey, R. J. & Smart, T. G. Zinc-mediated inhibition of GABA_A receptors: discrete binding sites underlie subtype specificity. *Nat. Neurosci.* **6**, 362–369 (2003).
41. Pritchett, D. B. & Seeburg, P. H. Gamma-aminobutyric acidA receptor alpha 5-subunit creates novel type II benzodiazepine receptor pharmacology. *J Neurochem* **54**, 1802–1804 (1990).
42. Wieland, H. A. & Luddens, H. Four amino acid exchanges convert a diazepam-insensitive, inverse agonist-preferring GABA_A receptor into a diazepam-preferring GABA_A receptor. *J Med Chem* **37**, 4576–4580 (1994).
43. Baumann, S. W., Baur, R. & Sigel, E. Subunit arrangement of gamma-aminobutyric acid type A receptors. *J Biol Chem* **276**, 36275–36280 (2001).
44. Smart, T. G., Moss, S. J., Xie, X. & Huganir, R. L. GABA_A receptors are differentially sensitive to zinc: dependence on subunit composition. *Br J Pharmacol* **103**, 1837–1839 (1991).
45. Frederickson, C. J. *et al.* Synaptic release of zinc from brain slices: factors governing release, imaging, and accurate calculation of concentration. *J Neurosci Methods* **154**, 19–29 (2006).
46. Frederickson, C. J. *et al.* Concentrations of extracellular free zinc (pZn)_e in the central nervous system during simple anesthetization, ischemia and reperfusion. *Experimental neurology* **198**, 285–293 (2006).

47. Qian, J. & Noebels, J. L. Visualization of transmitter release with zinc fluorescence detection at the mouse hippocampal mossy fibre synapse. *J Physiol* **566**, 747–758 (2005).
48. Vogt, K., Mellor, J., Tong, G. & Nicoll, R. The actions of synaptically released zinc at hippocampal mossy fiber synapses. *Neuron* **26**, 187–196 (2000).
49. Patel, B., Bright, D. P., Mortensen, M., Frolund, B. & Smart, T. G. Context-Dependent Modulation of GABA_AR-Mediated Tonic Currents. *J Neurosci* **36**, 607–621 (2016).
50. Cope, D. W. *et al.* Enhanced tonic GABA_A inhibition in typical absence epilepsy. *Nature medicine* **15**, 1392–1398 (2009).
51. Cope, D. W. *et al.* Abolition of zolpidem sensitivity in mice with a point mutation in the GABA_A receptor gamma2 subunit. *Neuropharmacol* **47**, 17–34 (2004).
52. Kharlamov, E. A., Downey, K. L., Jukkola, P. I., Grayson, D. R. & Kelly, K. M. Expression of GABA_A receptor alpha1 subunit mRNA and protein in rat neocortex following photothrombotic infarction. *Brain Res* **1210**, 29–38 (2008).
53. Liu, J. *et al.* Quantitative reverse transcription-polymerase chain reaction of GABA_A alpha1, beta1 and gamma2S subunits in epileptic rats following photothrombotic infarction of neocortex. *Epilepsy Res* **52**, 85–95 (2002).
54. Hiu, T. *et al.* Enhanced phasic GABA inhibition during the repair phase of stroke: a novel therapeutic target. *Brain* **139** (Pt 2), 468–80 (2016).
55. Raol, Y. H. *et al.* Enhancing GABA_A receptor alpha 1 subunit levels in hippocampal dentate gyrus inhibits epilepsy development in an animal model of temporal lobe epilepsy. *J Neurosci* **26**, 11342–11346 (2006).
56. Raol, Y. H. *et al.* Increased GABA_A-receptor alpha1-subunit expression in hippocampal dentate gyrus after early-life status epilepticus. *Epilepsia* **47**, 1665–1673 (2006).
57. Cohen, L., Chaaban, B. & Habert, M. O. Transient improvement of aphasia with zolpidem. *New Eng J Med* **350**, 949–950 (2004).
58. Brefel-Courbon, C. *et al.* Clinical and imaging evidence of zolpidem effect in hypoxic encephalopathy. *Ann Neurol* **62**, 102–105 (2007).
59. Clauss, R. P. *et al.* Cerebral blood perfusion after treatment with zolpidem and flumazenil in the baboon. *Arzneimittel-Forschung* **52**, 740–744 (2002).
60. Karim, N. *et al.* 2'-Methoxy-6-methylflavone: A novel anxiolytic and sedative with subtype selective activating and modulating actions at GABA_A receptors. *Br J Pharmacol* **165**, 880–896 (2012).
61. Karim, N. *et al.* 3-Hydroxy-2'-methoxy-6-methylflavone: A potent anxiolytic with a unique selectivity profile at GABA_A receptor subtypes. *Biochem Pharmacol* **82**, 1971–1983 (2011).

Acknowledgements

This work was funded by the National Health and Medical Research Council of Australia (NHMRC) Project Grant 1081733. A.T.C.H. was financially supported by the Ministry of Higher Education of Malaysia and Department of Neurosciences, University of Science, Malaysia (USM). P.K.A. was co-funded by Imk Almene Fond, Denmark.

Author Contributions

A.T.C.H. -Designed and performed experiments; contributed to writing of main text; N.A. -Designed experiments; contributed to writing of main text; P.S.v.N. -contributed to writing of main text; A.N.C. -Designed experiments; contributed to writing of main text; P.K.A. -Designed experiments; contributed to writing of main text; M.C. -Designed experiments; contributed writing of main text; All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Che Has, A. T. *et al.* Zolpidem is a potent stoichiometry-selective modulator of $\alpha 1\beta 3$ GABA_A receptors: evidence of a novel benzodiazepine site in the $\alpha 1$ - $\alpha 1$ interface. *Sci. Rep.* **6**, 28674; doi: 10.1038/srep28674 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>