



IO vs OI in Higher-Order Recursion Schemes

Axel Haddad

► **To cite this version:**

Axel Haddad. IO vs OI in Higher-Order Recursion Schemes. Workshop on Fixed Points in Computer Science, Mar 2012, Tallinn, Estonia. pp.23-30, 2012, <10.4204/EPTCS.77.4>. <hal-00865681>

HAL Id: hal-00865681

<https://hal.archives-ouvertes.fr/hal-00865681>

Submitted on 24 Sep 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IO vs OI in Higher-Order Recursion Schemes

Axel Haddad

LIAFA (Université Paris 7 & CNRS) & LIGM (Université Paris Est & CNRS)

We propose a study of the modes of derivation of higher-order recursion schemes, proving that value trees obtained from schemes using innermost-outermost derivations (IO) are the same as those obtained using unrestricted derivations.

Given that higher-order recursion schemes can be used as a model of functional programs, innermost-outermost derivations policy represents a theoretical view point of call by value evaluation strategy.

1 Introduction

Recursion schemes have been first considered as a model of computation, representing the syntactical aspect of a recursive program [15, 2, 3, 4]. At first, (order-1) schemes were modelling simple recursive programs whose functions only take values as input (and not functions). Since, higher-order versions of recursion schemes [11, 5, 6, 7, 8, 9] have been studied.

More recently, recursion schemes were studied as generators of infinite ranked trees and the focus was on deciding logical properties of those trees [12, 8, 10, 1, 13, 14].

As for programming languages, the question of the evaluation policy has been widely studied. Indeed, different policies results in the different evaluation [8, 9, 7]. There are two main evaluations policy for schemes: outermost-innermost derivations (OI) and inner-outermost IO derivations, respectively corresponding to call by need and call by value in programming languages.

Standardization theorem for the lambda-calculus shows that for any scheme, outermost-innermost derivations (OI) lead to the same tree as unrestricted derivation. However, this is not the case for IO derivations. In this paper we prove that the situation is different for schemes. Indeed, we establish that the trees produced using schemes with IO policy are the same as those produced using schemes with OI policy. For a given a scheme of order n , we can use a simplified continuation passing style transformation, to get a new scheme of order $n + 1$ in which IO derivations will be the same as OI derivations in the initial scheme (Section 3). Conversely, in order to turn a scheme into another one in which unrestricted derivations lead to the same tree as IO derivations in the initial scheme, we adapt Kobayashi's [13] recent results on HORS model-checking, to compute some key properties over terms (Section 4.1). Then we embed these properties into a scheme turning it into a self-correcting scheme of the same order of the initial scheme, in which OI and IO derivations produce the same tree (Section 4.2).

2 Preliminaries

Types are defined by the grammar $\tau ::= o \mid \tau \rightarrow \tau$; o is called the **ground type**. Considering that \rightarrow is associative to the right (i.e. $\tau_1 \rightarrow (\tau_2 \rightarrow \tau_3)$ can be written $\tau_1 \rightarrow \tau_2 \rightarrow \tau_3$), any type τ can be written uniquely as $\tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o$. The integer k is called the **arity** of τ . We define the **order of a type** by $\text{order}(o) = 0$ and $\text{order}(\tau_1 \rightarrow \tau_2) = \max(\text{order}(\tau_1) + 1, \text{order}(\tau_2))$. For instance $o \rightarrow o \rightarrow o \rightarrow o$ is a type

of order 1 and arity 3, $(o \rightarrow o) \rightarrow (o \rightarrow o)$, that can also be written $(o \rightarrow o) \rightarrow o \rightarrow o$ is a type of order 2. Let $\tau^\ell \rightarrow \tau'$ be a shortcut for $\underbrace{\tau \rightarrow \dots \rightarrow \tau}_{\ell \text{ times}} \rightarrow \tau'$.

Let Γ be a finite set of symbols such that to each symbol is associated a type. Let Γ^τ denote the set of symbols of type τ . For all type τ , we define the set of **terms** of type $\mathcal{T}^\tau(\Gamma)$ as the smallest set satisfying: $\Gamma^\tau \subseteq \mathcal{T}^\tau(\Gamma)$ and $\bigcup_{\tau'} \{t \ s \mid t \in \mathcal{T}^{\tau' \rightarrow \tau}(\Gamma), s \in \mathcal{T}^{\tau'}(\Gamma)\} \subseteq \mathcal{T}^\tau(\Gamma)$. If a term t is in $\mathcal{T}^\tau(\Gamma)$, we say that t has type τ . We shall write $\mathcal{T}(\Gamma)$ as the set of terms of any type, and $t : \tau$ if t has type τ . The arity of a term t , $\text{arity}(t)$, is the arity of its type. Remark that any term t can be uniquely written as $t = \alpha \ t_1 \dots t_k$ with $\alpha \in \Gamma$. We say that α is the **head** of the term t . For instance, let $\Gamma = \{F : (o \rightarrow o) \rightarrow o \rightarrow o, G : o \rightarrow o \rightarrow o, H : (o \rightarrow o), a : o\}$: $F \ H$ and $G \ a$ are terms of type $o \rightarrow o$; $F(G \ a)$ ($H(H \ a)$) is a term of type o ; $F \ a$ is not a term since F is expecting a first argument of type $o \rightarrow o$ while a has type o .

Let $t : \tau, t' : \tau'$ be two terms, $x : \tau'$ be a symbol of type τ' , then we write $t_{[x \rightarrow t']}$ the term obtained by substituting all occurrences of x by t' in the term t . A τ -**context** is a term $C[\bullet^\tau] \in \mathcal{T}(\Gamma \uplus \{\bullet^\tau : \tau\})$ containing exactly one occurrence of \bullet^τ ; it can be seen as an application turning a term into another, such that for all $t : \tau, C[t] = C[\bullet^\tau]_{[\bullet^\tau \rightarrow t]}$. In general we will only talk about ground type context where $\tau = o$ and we will omit to specify the type when it is clear. For instance, if $C[\bullet] = F \bullet (H(H \ a))$ and $t' = G \ a$ then $C[t'] = F(G \ a)(H(H \ a))$.

Let Σ be a set of symbols of order at most 1 (i.e. each symbols has type o or $o \rightarrow \dots \rightarrow o$) and $\perp : o$ be a fresh symbol. A **tree** t over $\Sigma \uplus \perp$ is a mapping $t : \text{dom}^t \rightarrow \Sigma \uplus \perp$, where dom^t is a prefix-closed subset of $\{1, \dots, m\}^*$ such that if $u \in \text{dom}^t$ and $t(u) = a$ then $\{j \mid uj \in \text{dom}^t\} = \{1, \dots, \text{arity}(a)\}$. Note that there is a direct bijection between ground terms of $\mathcal{T}^o(\Sigma \uplus \perp)$ and finite trees. Hence we will freely allow ourselves to treat ground terms over $\Sigma \uplus \perp$ as trees. We define the partial order \sqsubseteq over trees as the smallest relation satisfying $\perp \sqsubseteq t$ and $t \sqsubseteq t'$ for any tree t , and $a \ t_1 \dots t_k \sqsubseteq a \ t'_1 \dots t'_k$ iff $t_i \sqsubseteq t'_i$. Given a (possibly infinite) sequence of trees t_0, t_1, t_2, \dots such that $t_i \sqsubseteq t_{i+1}$ for all i , one can prove that the set of all t_i has a supremum that is called the **limit tree** of the sequence.

A **higher order recursion scheme (HORS)** $G = \langle \mathcal{V}, \Sigma, \mathcal{N}, \mathcal{R}, S \rangle$ is a tuple such that: \mathcal{V} is a finite set of typed symbols called **variables**; Σ is a finite set of typed symbols of order at most 1, called the **set of terminals**; \mathcal{N} is a finite set of typed symbols called **set of non-terminals**; \mathcal{R} is a set of **rewrite rules**, one per non terminal $F : \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o \in \mathcal{N}$, of the form $F \ x_1 \dots x_k \rightarrow e$ with $e : o \in \mathcal{T}(\Sigma \uplus \mathcal{N} \uplus \{x_1, \dots, x_k\})$; $S \in \mathcal{N}$ is the **initial non-terminal**.

We define the **rewriting relation** $\rightarrow_G \in \mathcal{T}(\Sigma \uplus \mathcal{N})^2$ (or just \rightarrow when G is clear) as $t \rightarrow_G t'$ iff there exists a context $C[\bullet]$, a rewrite rule $F \ x_1 \dots x_k \rightarrow e$, and a term $F \ t_1 \dots t_k : o$ such that $t = C[F \ t_1 \dots t_k]$ and $t' = C[e_{[x_1 \rightarrow t_1] \dots [x_k \rightarrow t_k]}]$. We call $F \ t_1 \dots t_k : o$ a **redex**. Finally we define \rightarrow_G^* as the reflexive and transitive closure of \rightarrow_G .

We define inductively the **\perp -transformation** $(\cdot)^\perp : \mathcal{T}^o(\mathcal{N} \uplus \Sigma) \rightarrow \mathcal{T}^o(\Sigma \uplus \{\perp : o\})$: $(F \ t_1 \dots t_k)^\perp = \perp \ \forall F \in \mathcal{N}$ and $(a \ t_1 \dots t_k)^\perp = a \ t_1^\perp \dots t_k^\perp$ for all $a \in \Sigma$. We define a **derivation**, as a possibly infinite sequence of terms linked by the rewrite relation. Let $t_0 = S \rightarrow_G t_1 \rightarrow_G t_2 \rightarrow_G \dots$ be a derivation, then one can check that $(t_0)^\perp \sqsubseteq (t_1)^\perp \sqsubseteq (t_2)^\perp \sqsubseteq \dots$, hence it admits a limit. One can prove that the set of all such limit trees has a greatest element that we denote $\|G\|$ and refer to as the **value tree** of G . Note that $\|G\|$ is the supremum of $\{t^\perp \mid S \rightarrow_G^* t\}$. Given a term $t : o$, we denote by G_t the scheme obtained by transforming G such that it starts derivations with the term t , formally, $G_t = \langle \mathcal{V}, \Sigma, \mathcal{N} \uplus \{S'\}, \mathcal{R} \uplus \{S' \rightarrow t\}, S' \rangle$. One can prove that if $t \rightarrow t'$ then $\|G_t\| = \|G_{t'}\|$.

Example. Let $G = \langle \mathcal{V}, \Sigma, \mathcal{N}, \mathcal{R}, S \rangle$ be the scheme such that: $\mathcal{V} = \{x : o, \phi : o \rightarrow o, \psi : (o \rightarrow o) \rightarrow o \rightarrow o\}$, $\Sigma = \{a : o^3 \rightarrow o, b : o \rightarrow o \rightarrow o, c : o\}$, $\mathcal{N} = \{F : ((o \rightarrow o) \rightarrow o \rightarrow o) \rightarrow (o \rightarrow o) \rightarrow o \rightarrow o, H : (o \rightarrow$


$o) \rightarrow o \rightarrow o, I, J, K : o \rightarrow o, S : o\}$, and \mathcal{R} contains the following rewrite rules:

$$\begin{array}{lll} F \psi \phi x & \rightarrow & \psi \phi x \\ J x & \rightarrow & b (J x) (J x) \end{array} \quad \begin{array}{ll} I x & \rightarrow x \\ K x & \rightarrow K (K x) \end{array} \quad \begin{array}{ll} H \phi x & \rightarrow a (J x) (K x) (\phi x) \\ S & \rightarrow F H I c \end{array}$$

Here is an example of finite derivation:

$$\begin{aligned} S &\rightarrow F H I c \rightarrow H I c \rightarrow a (J c) (K c) (I c) \\ &\rightarrow a (J c) (K (K c)) (I c) \rightarrow a (J c) (K (K (K c))) (I c) \end{aligned}$$

If one extends it by always rewriting a redex of head K , its limit is the tree $a \perp \perp \perp$, but this is not the value tree of G . The value tree $\|G\|$ is depicted below.



Evaluation Policies

We now put constraints on the derivations we allow. If there are no constraints, then we say that the derivations are unrestricted and we let $\text{Acc}^G = \{t : o \mid S \rightarrow^* t\}$ be the set of accessible terms using unrestricted derivations. Given a rewriting $t \rightarrow t'$ such that $t = C[F s_1 \dots s_k]$ and $t' = C[e_{[\forall j x_j \mapsto s_j]}]$ with $F x_1 \dots x_k \rightarrow e \in \mathcal{R}$.

- We say that $t \rightarrow t'$ is an **outermost-innermost** (OI) rewriting (written $t \rightarrow_{OI} t'$) there is no redex containing the occurrence of \bullet as a subterm of $C[\bullet]$.
- We say that $t \rightarrow t'$ is an **innermost-outermost** (IO) rewriting (written $t \rightarrow_{IO} t'$), if for all j there is no redex as a subterm of s_j .

Let $\text{Acc}_{OI}^G = \{t : o \mid S \rightarrow_{OI}^* t\}$ be the set of accessible terms using OI derivations and $\text{Acc}_{IO}^G = \{t : o \mid S \rightarrow_{IO}^* t\}$ be the set of accessible terms using IO derivations. There exists a supremum of Acc_{OI}^G (resp. Acc_{IO}^G) which is the maximum of the limit trees of OI derivations (resp. IO derivations). We write it $\|G\|_{OI}$ (resp. $\|G\|_{IO}$). For all recursive scheme G , $(\text{Acc}^G)^\perp = (\text{Acc}_{OI}^G)^\perp$, in particular $\|G\|_{OI} = \|G\|$. But $\|G\|_{IO} \sqsubseteq \|G\|$ and in general, the equality does not hold (see the example in the next section).

3 From OI to IO

Fix a recursion scheme $G = \langle \mathcal{V}, \Sigma, \mathcal{N}, \mathcal{R}, S \rangle$. Our goal is to define another scheme $\overline{G} = \langle \overline{\mathcal{V}}, \Sigma, \overline{\mathcal{N}}, \overline{\mathcal{R}}, I \rangle$ such that $\|\overline{G}\|_{IO} = \|G\|$. The idea is to add an extra argument (Δ) to each non terminal, that will be required to rewrite it (hence the types are changed). We feed this argument to the outermost non terminal, and duplicate it to subterms only if the head of the term is a terminal. Hence all derivations will be IO-derivations.

We define the $(\overline{\cdot})$ **transformation** over types by $\overline{o} = o \rightarrow o$, and $\overline{\tau_1} \rightarrow \overline{\tau_2} = \overline{\tau_1} \rightarrow \overline{\tau_2}$. In particular, if $\tau = \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o$ then $\overline{\tau} = \overline{\tau_1} \rightarrow \dots \rightarrow \overline{\tau_k} \rightarrow o \rightarrow o$. Note that for all τ , $\text{order}(\overline{\tau}) = \text{order}(\tau) + 1$.

For all $x : \tau \in \mathcal{V}$ we define $\bar{x} : \bar{\tau}$ as a fresh variable. Let ar_{max} be the maximum arity of terminals, we define $\eta_1, \dots, \eta_{ar_{max}} : o \rightarrow o$ and $\delta : o$ as fresh variables, and we let $\bar{\mathcal{V}} = \{\bar{x} : \bar{\tau} \mid x \in \mathcal{V}\} \uplus \{\eta_1, \dots, \eta_{ar_{max}}\} \uplus \{\delta : o\}$. Note that δ is the only variable of type o . For all $a : \tau \in \Sigma$ define $\bar{a} : \bar{\tau}$ as a fresh **non-terminal** and for all $F : \tau \in \mathcal{N}$ define $\bar{F} : \bar{\tau}$ as a fresh non-terminal. Let $\bar{\mathcal{N}} = \{\bar{a} : \bar{\tau} \mid a \in \Sigma\} \uplus \{\bar{F} : \bar{\tau} \mid F \in \mathcal{N}\} \uplus \{\Delta : o, I : o\}$. Note that I and Δ are the only symbols in $\bar{\mathcal{N}}$ of type o .

Let $t : \tau \in \mathcal{T}(\mathcal{V} \uplus \Sigma \uplus \mathcal{N})$, we define inductively the term $\bar{t} : \bar{\tau} \in \mathcal{T}(\bar{\mathcal{V}} \uplus \bar{\mathcal{N}})$: If $t = x \in \mathcal{V}$ (resp. $t = a \in \Sigma, t = F \in \mathcal{N}$), we let $\bar{t} = \bar{x} \in \bar{\mathcal{V}}$ (resp. $\bar{t} = \bar{a} \in \bar{\Sigma}, \bar{t} = \bar{F} \in \bar{\mathcal{N}}$), if $t = t_1 t_2 : \tau$ then $\bar{t} = \bar{t}_1 \bar{t}_2$.

Let $F x_1 \dots x_k \rightarrow e$ be a rewrite rule of \mathcal{R} . We define the (valid) rule $\bar{F} \bar{x}_1 \dots \bar{x}_k \delta \rightarrow \bar{e} \Delta$ in $\bar{\mathcal{R}}$. Let $a \in \Sigma$ of arity k , we define the rule $\bar{a} \eta_1 \dots \eta_k \delta \rightarrow a (\eta_1 \Delta) \dots (\eta_k \Delta)$ in $\bar{\mathcal{R}}$. We also add the rule $I \rightarrow \bar{S} \Delta$ to $\bar{\mathcal{R}}$. Finally let $\bar{G} = \langle \bar{\mathcal{V}}, \bar{\Sigma}, \bar{\mathcal{N}}, \bar{\mathcal{R}}, I \rangle$.

Example. Let $G = \langle \mathcal{V}, \Sigma, \mathcal{N}, \mathcal{R}, S \rangle$ be the order-1 recursion scheme with $\Sigma = \{a, c : o\}$, $\mathcal{N} = \{S : o, F : o \rightarrow o \rightarrow o, H : o \rightarrow o\}$, $\mathcal{V} = \{x, y : o\}$, and the following rewrite rules:

$$S \rightarrow F (H a) c \quad F x y \rightarrow y \quad H x \rightarrow H (H x)$$

Then we have $\|G\|_{OI} = c$ while $\|G\|_{IO} = \perp$ (indeed, the only IO derivation is the following $S \rightarrow F (Ha) c \rightarrow F (H (H a)) c \rightarrow F (H (H (H a))) c \rightarrow \dots$). The order-2 recursion scheme $\bar{G} = \langle \bar{\mathcal{V}}, \bar{\Sigma}, \bar{\mathcal{N}}, \bar{\mathcal{R}}, I \rangle$ is given by $\bar{\mathcal{N}} = \{I, \Delta : o, \bar{S}, \bar{a}, \bar{c} : o \rightarrow o, \bar{F} : (o \rightarrow o) \rightarrow (o \rightarrow o) \rightarrow o \rightarrow o, \bar{H} : (o \rightarrow o) \rightarrow o \rightarrow o\}$, $\bar{\mathcal{V}} = \{\delta : o, \bar{x}, \bar{y} : o \rightarrow o\}$ and the following rewrite rules:

$$\begin{array}{llll} I & \rightarrow & \bar{S} \Delta & \bar{S} \delta \rightarrow \bar{F} (\bar{H} \bar{a}) \bar{c} \Delta & \bar{F} \bar{x} \bar{y} \delta \rightarrow \bar{y} \Delta \\ \bar{H} \bar{x} \delta & \rightarrow & \bar{H} (\bar{H} \bar{x}) \Delta & \bar{c} \delta \rightarrow c & \bar{a} \delta \rightarrow a \end{array}$$

Note that in the term $\bar{F} (\bar{H} \bar{a}) \bar{c} \Delta$, the subterm $\bar{H} \bar{a}$ is no longer a redex since it lacks its last argument, hence it cannot be rewritten, then the only IO derivation, which is the only unrestricted derivation is $I \rightarrow \bar{S} \Delta \rightarrow \bar{F} (\bar{H} \bar{a}) \bar{c} \Delta \rightarrow \bar{c} \Delta \rightarrow c$. Therefore $\|\bar{G}\|_{IO} = \|\bar{G}\| = c = \|G\|$.

Lemma 1. Any derivation of \bar{G} is in fact an OI and an IO derivation. Hence that $\|\bar{G}\|_{IO} = \|\bar{G}\|$.

Proof (Sketch). The main idea is that the only redexes will be those that have Δ as last argument of the head non-terminal. The scheme is constructed so that Δ remains only on the outermost non-terminals, that is why any derivation is an OI derivation. Furthermore, we have that if $t = \bar{F} t_1 \dots t_k \Delta$ is a redex, then none of the t_i contains Δ , therefore they do not contain any redex, hence t is an innermost redex. \square

Note that OI derivations in \bar{G} acts like OI derivations in G , hence $\|G\| = \|\bar{G}\|$.

Theorem 2 (OI vs IO). Let G be an order- n scheme. Then one can construct an order- $(n+1)$ scheme \bar{G} such that $\|G\| = \|\bar{G}\|_{IO}$.

4 From IO to OI

The goal of this section is to transform the scheme G into a scheme G'' such that $\|G''\| = \|G\|_{IO}$. The main difference between IO and OI derivations is that some redex would lead to \perp in IO derivation while OI derivations could be more productive. For example take $F : o \rightarrow o$ such that $F x \rightarrow c$, and $H : o$ such that $H \rightarrow a H$, with $a : o \rightarrow o$ and $c : o$ being terminal symbols. The term $F H$ has a unique OI derivation, $F H \rightarrow_{OI} c$, it is finite and it leads to the value tree associated. On the other hand, the (unique) IO derivation is the following $F H \rightarrow F(a H) \rightarrow F(a(a H)) \rightarrow \dots$ which leads to the tree \perp .

The idea of the transformation is to compute a tool (based on a type system) that decides if a redex would produce \perp with IO derivations (Section 4.1); then we embed it into G and force any such redex to produce \perp even with unrestricted derivations (Section 4.2).

4.1 The Type System

Given a term $t : \tau \in \mathcal{T}(\Sigma \uplus \mathcal{N})$, we define the two following properties on t : $\mathcal{P}_\perp(t)$ = “The term t has type o and its associated IO valuation tree is \perp ”, and $\mathcal{P}_\infty(t)$ = “the term t has not necessarily ground type, it contains a redex r such that any IO derivation from r producing it’s IO valuation tree is infinite”. Note that $\mathcal{P}_\infty(t)$ is equivalent to “the term t contains a redex r such that $\|G_r\|_{IO}$ is either infinite or contains \perp ”. In this section we describe a type system, inspired from the work of Kobayashi [13], that characterises if a term verifies these properties.

Let \mathcal{Q} be the set $\{q_\perp, q_\infty\}$. Given a type τ , we define inductively the sets $(\tau)^{atom}$ and $(\tau)^\wedge$ called respectively set of atomic mappings and set of conjunctive mappings:

$$(o)^{atom} = \mathcal{Q}, \quad (o)^\wedge = \{\wedge\{\theta_1, \dots, \theta_i\} \mid \theta_1, \dots, \theta_i \in \mathcal{Q}\}, \quad (\tau_1 \rightarrow \tau_2)^{atom} = \{q_\infty\} \uplus \{(\tau_1)^\wedge \rightarrow (\tau_2)^{atom}\}$$

$$(\tau_1 \rightarrow \tau_2)^\wedge = \{\wedge\{\theta_1, \dots, \theta_i\} \mid \theta_1, \dots, \theta_i \in (\tau_1 \rightarrow \tau_2)^{atom}\}.$$

We will usually use the letter θ to represents atomic mappings, and the letter σ to represent conjunctive mappings. Given a conjunctive mapping σ (resp. an atomic mapping θ) and a type τ , we write $\sigma :: \tau$ (resp. $\theta ::_a \tau$) the relation $\sigma \in (\tau)^\wedge$ (resp. $\theta \in (\tau)^{atom}$). For the sake of simplicity, we identify the atomic mapping θ with the conjunctive mapping $\wedge\{\theta\}$.

Given a term t and a conjunctive mapping σ , we define a judgment as a tuple $\Theta \vdash t \triangleright \sigma$, pronounce “from the environment Θ , one can prove that t matches the conjunctive mapping σ ”, where the environment Θ is a partial mapping from $\mathcal{V} \uplus \mathcal{N}$ to conjunctive mapping. Given an environment Θ , $\alpha \in \mathcal{V} \uplus \mathcal{N}$ and a conjunctive mapping σ , we define the environment $\Theta' = \Theta, \alpha \triangleright \sigma$ as $Dom(\Theta') = Dom(\Theta) \cup \{\alpha\}$ and $\Theta'(\alpha) = \sigma$ if $\alpha \notin Dom(\Theta)$, $\Theta'(\alpha) = \sigma \wedge \Theta(\alpha)$ otherwise, and $\Theta'(\beta) = \Theta(\beta)$ if $\beta \neq \alpha$.

We define the following judgement rules:

$$\frac{\Theta \vdash t \triangleright \theta_1 \quad \dots \quad \Theta \vdash t \triangleright \theta_n}{\Theta \vdash t \triangleright \wedge\{\theta_1, \dots, \theta_n\}} (Set) \quad \frac{}{\Theta, \alpha \triangleright \wedge\{\theta_1, \dots, \theta_n\} \vdash \alpha \triangleright \theta_i} (At) \text{ (for all } i)$$

$$\frac{}{\Theta \vdash a \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_{i \leq \text{arity}(a)} \rightarrow q_\infty} (\Sigma) \text{ (for } a \in \Sigma \text{ and } \exists j \sigma_j = q_\infty)$$

$$\frac{\Theta \vdash t_1 \triangleright \sigma \rightarrow \theta \quad \Theta \vdash t_2 \triangleright \sigma}{\Theta \vdash t_1 t_2 \triangleright \theta} (App) \quad \frac{}{\Theta \vdash t \triangleright q_\infty \rightarrow q_\infty} (q_\infty \rightarrow q_\infty I) \text{ (if } t : \tau_1 \rightarrow \tau_2) \quad \frac{\Theta \vdash t_1 \triangleright q_\infty}{\Theta \vdash t_1 t_2 \triangleright q_\infty} (q_\infty I)$$

Remark that there is no rules that directly involves q_\perp , but it does not mean that no term matches q_\perp , since it can appear in Θ . Rules like (At) or (App) may be used to state that a term matches q_\perp .

We say that (G, t) matches the conjunctive mapping σ written $\vdash (G, t) \triangleright \sigma$ if there exists an environment Θ , called a witness environment of $\vdash (G, t) \triangleright \sigma$, such that (1) $Dom(\Theta) = \mathcal{N}$, (2) $\forall F : \tau \in \mathcal{N} \quad \Theta(F) :: \tau$, (3) if $F x_1 \dots x_k \rightarrow e \in \mathcal{R}$ and $\Theta \vdash F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_{i \leq k} \rightarrow q$ then either there exists j such that $q_\infty \in \sigma_j$, or $i = k$ and $\Theta, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e \triangleright q$, (4) $\Theta \vdash t \triangleright \sigma$.

The following two results state that this type system matches the properties \mathcal{P}_\perp and \mathcal{P}_∞ and furthermore we can construct a universal environment, Θ^* , that can correctly judge any term.

Theorem 3 (Soundness and Completeness). *Let G be an HORS, and t be term (of any type), $\vdash (G, t) \triangleright q_\infty$ (resp. $\vdash (G, t) \triangleright q_\perp$) if and only if $\mathcal{P}_\infty(t)$ (resp. $\mathcal{P}_\perp(t)$) holds.*

Proposition 4 (Universal Witness). *There exists an environment Θ^* such that for all term t , the judgment $\vdash (G, t) \triangleright \sigma$ holds if and only if $\Theta^* \vdash t \triangleright \sigma$.*

Proof (Sketch). To compute Θ^* , we start with an environment Θ_0 satisfying Properties (1) and (2) ($Dom(\Theta_0) = \mathcal{N}$ and $\forall F : \tau \in \mathcal{N} \ \Theta_0(F) :: \tau$) that is able to judge any term $t : \tau$ with any conjunctive mapping $\sigma :: \tau$.

Then let \mathcal{F} be the mapping from the set of environments to itself, such that for all $F : \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o \in \mathcal{N}$, if $F \ x_1 \dots x_k \rightarrow e \in \mathcal{R}$ then,

$$\begin{aligned} \mathcal{F}(\Theta)(F) = & \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \mid q \in Q \wedge \forall i \ \sigma_i :: \tau_i \wedge \Theta, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e : q \} \\ & \cup \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_{i \leq k} \rightarrow q_\infty \mid \wedge \forall i \ \sigma_i :: \tau_i \wedge \exists j \ q_\infty \in \sigma_j \} \\ & \cup \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q_\perp \mid \forall i \ \sigma_i :: \tau_i \wedge \exists j \ q_\infty \in \sigma_j \}. \end{aligned}$$

We iterate \mathcal{F} until we reach a fixpoint. The environment we get is Θ^* , it verifies properties (1) (2) and (3). Furthermore we can show that this is the maximum of all environment satisfying these properties, i.e. if $\vdash (G, t) \triangleright \sigma$ then $\Theta^* \vdash t \triangleright \sigma$. \square

4.2 Self-Correcting Scheme

For all term $t : \tau \in \mathcal{T}(\Sigma \uplus \mathcal{N})$, we define $\llbracket t \rrbracket \in (\tau)^\wedge$, called the semantics of t , as the conjunction of all atomic mappings θ such that $\Theta^* \vdash t \triangleright \theta$ (recall that Θ^* is the environment of Proposition 4). In particular $\mathcal{P}_\perp(t)$ (resp. $\mathcal{P}_\infty(t)$) holds if and only if $q_\perp \in \llbracket t \rrbracket$ (resp. $q_\infty \in \llbracket t \rrbracket$). Given two terms $t_1 : \tau_2 \rightarrow \tau$ and $t_2 : \tau_2$ the only rules we can apply to judge $\Theta^* \vdash t_1 \ t_2 \triangleright \theta$ are (*App*), ($q_\infty \rightarrow q_\infty I$) and ($q_\infty I$). We see that θ only depends on which atomic mappings are matched by t_1 and t_2 . In other words $\llbracket t_1 \ t_2 \rrbracket$ only depends on $\llbracket t_1 \rrbracket$ and $\llbracket t_2 \rrbracket$, we write $\llbracket t_1 \rrbracket \llbracket t_2 \rrbracket = \llbracket t_1 \ t_2 \rrbracket$.

In this section, given a scheme $G = \langle \mathcal{V}, \Sigma, \mathcal{N}, \mathcal{R}, S \rangle$, we transform it into $G' = \langle \mathcal{V}', \Sigma, \mathcal{N}', \mathcal{R}', S \rangle$ which is basically the same scheme except that while it is producing an IO derivation, it evaluates $\llbracket t' \rrbracket$ for any subterm t' of the current term and label t' with $\llbracket t' \rrbracket$. Note that if $t \rightarrow_{IO} t'$, then $\llbracket t \rrbracket = \llbracket t' \rrbracket$. Since we cannot syntactically label terms, we will label all symbols by the semantics of their arguments, e.g. if we want to label $F \ t_1 \dots t_k$, we will label F with the k -tuple $(\llbracket t_1 \rrbracket, \dots, \llbracket t_k \rrbracket)$.

A problem may appear if some of the arguments are not fully applied, for example imagine we want to label $F \ H$ with $H : o \rightarrow o$. We will label F with $\llbracket H \rrbracket$, but since H has no argument we do not know how to label it. The problem is that we cannot wait to label it because once a non-terminal is created, the derivation does not deal explicitly with it. The solution is to create one copy of H per possible semantics for its argument (here there are four of them: $\wedge\{\}, \wedge\{q_\perp\}, \wedge\{q_\infty\}, \wedge\{q_\perp, q_\infty\}$). This means that $F^{\llbracket H \rrbracket}$ would not have the same type as F : F has type $(o \rightarrow o) \rightarrow o$, but $F^{\llbracket G \rrbracket}$ will have type $(o \rightarrow o)^4 \rightarrow o$. Hence, $F \ H$ will be labelled the following way: $F^{\llbracket H \rrbracket} \ H^{\wedge\{\}} \ H^{\wedge\{q_\perp\}} \ H^{\wedge\{q_\infty\}} \ H^{\wedge\{q_\perp, q_\infty\}}$. Note that even if F has 4 arguments, it only has to be labelled with one semantics since all four arguments represent different labelling of the same term. We now formalize these notions.

Let us generalize the notion of semantics to deals with terms containing some variables. Given an environment on the variables $\Theta^\mathcal{V}$ such that $Dom(\Theta^\mathcal{V}) \subseteq \mathcal{V}$ and if $x : \tau$ then $\Theta^\mathcal{V}(x) :: \tau$, and given a term $t : \tau \in \mathcal{T}(\Sigma \uplus \mathcal{N} \uplus Dom(\Theta^\mathcal{V}))$, we define $\llbracket t \rrbracket_{\Theta^\mathcal{V}} \in (\tau)^\wedge$, as the conjunction of all atomic mappings θ such that $\Theta^*, \Theta^\mathcal{V} \vdash t \triangleright \theta$. Given two terms $t_1 : \tau_2 \rightarrow \tau$ and $t_2 : \tau_2$ we still have that $\llbracket t_1 \ t_2 \rrbracket_{\Theta^\mathcal{V}}$ only depends on $\llbracket t_1 \rrbracket_{\Theta^\mathcal{V}}$ and $\llbracket t_2 \rrbracket_{\Theta^\mathcal{V}}$.

To a type $\tau = \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o$ we associate the integer $\lceil \tau \rceil = Card(\{(\sigma_1, \dots, \sigma_k) \mid \forall i \ \sigma_i \in (\tau_i)^\wedge\})$ and a complete ordering of $\{(\sigma_1, \dots, \sigma_k) \mid \forall i \ \sigma_i \in (\tau_i)^\wedge\}$ denoted $\vec{\sigma}_1^\tau, \vec{\sigma}_2^\tau, \dots, \vec{\sigma}_{\lceil \tau \rceil}^\tau$. We define inductively the type $\tau^+ = (\tau_1^+)^{\lceil \tau_1 \rceil} \rightarrow \dots \rightarrow (\tau_k^+)^{\lceil \tau_k \rceil} \rightarrow o$.

To a non terminal $F : \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o$ (resp. a variable $x : \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o$) and a tuple $\sigma_1 :: \tau_1, \dots, \sigma_k :: \tau_k$, we associate the non-terminal $F^{\sigma_1, \dots, \sigma_k} : \tau_1^{[\tau_1]} \rightarrow \dots \rightarrow \tau_k^{[\tau_k]} \rightarrow o \in \mathcal{N}'$ (resp. a variable $x^{\sigma_1, \dots, \sigma_k} : \tau_1^{[\tau_1]} \rightarrow \dots \rightarrow \tau_k^{[\tau_k]} \rightarrow o \in \mathcal{V}'$).

Given a term $t : \tau = \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o \in \mathcal{T}(\mathcal{V} \uplus \Sigma \uplus \mathcal{N})$ and an environment on the variables $\Theta^{\mathcal{V}}$ such that $Dom(\Theta^{\mathcal{V}}) \subseteq \mathcal{V}$ contains all variables in t , we define inductively the term $t_{\Theta^{\mathcal{V}}}^{+\sigma_1, \dots, \sigma_k} : \tau^+ \in \mathcal{T}(\mathcal{V}' \uplus \Sigma' \uplus \mathcal{N}')$ for all $\sigma_1 :: \tau_1, \dots, \sigma_k :: \tau_k$. If $t = F \in \mathcal{N}$ (resp. $t = x \in \mathcal{V}$), $t_{\Theta^{\mathcal{V}}}^{+\sigma_1, \dots, \sigma_k} = F^{\sigma_1, \dots, \sigma_k}$ (resp. $t_{\Theta^{\mathcal{V}}}^{+\sigma_1, \dots, \sigma_k} = x^{\sigma_1, \dots, \sigma_k}$), if $t = a \in \Sigma$, $t_{\Theta^{\mathcal{V}}}^{+\sigma_1, \dots, \sigma_k} = a$. Finally consider the case where $t = t_1 t_2$ with $t_1 : \tau' \rightarrow \tau$ and $t_2 : \tau'$. Let $\sigma = \llbracket t_2 \rrbracket_{\Theta^{\mathcal{V}'}}$. Remark that $t_1^{+\sigma, \sigma_1, \dots, \sigma_k} : (\tau'^+)^{[\tau']} \rightarrow \tau^+$. We define $(t_1 t_2)_{\Theta^{\mathcal{V}'}}^{+\sigma_1, \dots, \sigma_k} = t_1^{+\sigma, \sigma_1, \dots, \sigma_k} t_2_{\Theta^{\mathcal{V}'}}^{+\tilde{\sigma}_1^{\tau'}}$... $t_2_{\Theta^{\mathcal{V}'}}^{+\tilde{\sigma}_1^{\tau'}}$. Note that since this transformation is only duplicating and anoting, given a term $t^{+\sigma_1, \dots, \sigma_k}$ we can uniquely find the unique term t associated to it.

Let $F : \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o \in \mathcal{N}$, $\sigma_1 :: \tau_1, \dots, \sigma_k :: \tau_k$, and $\Theta^{\mathcal{V}} = x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k$. If $F x_1 \dots x_k \rightarrow e \in \mathcal{R}$, we define in \mathcal{R}' the rule $F^{\sigma_1, \dots, \sigma_k} x_1^{+\tilde{\sigma}_1^{\tau_1}} \dots x_1^{+\tilde{\sigma}_1^{\tau_1}} \dots x_k^{+\tilde{\sigma}_1^{\tau_k}} \dots x_k^{+\tilde{\sigma}_1^{\tau_k}} \rightarrow e_{\Theta^{\mathcal{V}'}}^+$. Finally, recall that $G' = \langle \mathcal{V}', \Sigma, \mathcal{N}', \mathcal{R}', S \rangle$.

The following theorem states that G' is just a labeling version of G and that it acts the same.

Theorem 5 (Equivalence between G and G'). *Given a term $t : o$, $\|G'_{t^+}\|_{IO} = \|G_t\|_{IO}$.*

We transform G' into the scheme G'' that will directly turn into \perp a redex t such that $q_{\perp} \in \llbracket t \rrbracket$. For technical reason, instead of adding \perp we add a non terminal $Void : o$ and a rule $Void \rightarrow Void$. $G' = \langle \mathcal{V}', \Sigma, \mathcal{N}' \uplus \{Void : o\}, \mathcal{R}'', S \rangle$ such that \mathcal{R}'' contains the rule $Void \rightarrow Void$ and for all $F \in \mathcal{N}$, if $q_{\perp} \in \llbracket F \rrbracket \sigma_1 \dots \sigma_k$ then $F^{\sigma_1, \dots, \sigma_k} x_1^{+\tilde{\sigma}_1^{\tau_1}} \dots x_1^{+\tilde{\sigma}_1^{\tau_1}} \dots x_k^{+\tilde{\sigma}_1^{\tau_k}} \dots x_k^{+\tilde{\sigma}_1^{\tau_k}} \rightarrow Void$ otherwise we keep the rule of \mathcal{R}' .

The following theorem concludes Section 4.

Theorem 6 (IO vs OI). *Let G be a higher-order recursion scheme. Then one can construct a scheme G'' having the same order of G such that $\|G''\| = \|G\|_{IO}$.*

Proof (Sketch). First, given a term $t : o$, one can prove that $\|G''_{t^+}\|_{IO} = \|G'_{t^+}\|_{IO}$.

Then take a redex t such that $\|G''_t\|_{IO} = \perp$, i.e. $q_{\perp} \in \llbracket G_t \rrbracket$. There is only one OI derivation from $t : t \rightarrow Void \rightarrow Void \rightarrow \dots$, then $\|G''_t\| = \perp$. We can extend this result saying that if there is the symbol \perp at node u in $\|G''_t\|_{IO}$, then there is \perp at node u in $\|G''_t\|$. Hence, since $\|G''_t\|_{IO} \sqsubseteq \|G''_t\|$, we have $\|G''\| = \|G''\|_{IO}$. Then $\|G''\| = \|G''\|_{IO} = \|G'\|_{IO} = \|G\|_{IO}$. □

5 Conclusion

We have shown that value trees obtained from schemes using innermost-outermost derivations (IO) are the same as those obtained using unrestricted derivations. More precisely, given an order- n scheme G we create an order- $(n+1)$ scheme \bar{G} such that $\|\bar{G}\|_{IO} = \|G\|$. However, the increase of the order seems unavoidable. We also create an order- n scheme G'' such that $\|\bar{G}''\| = \|G\|_{IO}$. In this case the order does not increase, however the size of the scheme deeply increases while it remains almost the same in \bar{G} .

References

- [1] Klaus Aehlig (2006): *A Finite Semantics of Simply-Typed Lambda Terms for Infinite Runs of Automata*. In: "Proc. of Computer Science Logic, 20th Annual Conference of the EACSL", Lecture Notes in Comput. Sci. 4207, Springer-Verlag, pp. 104–118, doi:10.1007/11874683_7.
- [2] Bruno Courcelle (1978): *A Representation of Trees by Languages I*. Theoret. Comput. Sci. 6, pp. 255–279, doi:10.1016/0304-3975(78)90008-7.
- [3] Bruno Courcelle (1978): *A Representation of Trees by Languages II*. Theoret. Comput. Sci. 7, pp. 25–55, doi:10.1016/0304-3975(78)90039-7.
- [4] Bruno Courcelle & Maurice Nivat (1978): *The Algebraic Semantics of Recursive Program Schemes*. In: Proc. 7th Symposium, Mathematical Foundations of Computer Science 1978, Lecture Notes in Comput. Sci. 64, Springer-Verlag, pp. 16–30.
- [5] Werner Damm (1977): *Higher type program schemes and their tree languages*. In: Theoretical Computer Science, 3rd GI-Conference, Lecture Notes in Comput. Sci. 48, Springer-Verlag, pp. 51–72.
- [6] Werner Damm (1977): *Languages Defined by Higher Type Program Schemes*. In: Proc. 4th Colloq. on Automata, Languages, and Programming (ICALP), Lecture Notes in Comput. Sci. 52, Springer-Verlag, pp. 164–179.
- [7] Werner Damm (1982): *The IO- and OI-Hierarchies*. Theoret. Comput. Sci. 20, pp. 95–207, doi:10.1016/0304-3975(82)90009-3.
- [8] Joost Engelfriet & Erik Meineche Schmidt (1977): *IO and OI. I*. J. Comput. System Sci. 15(3), pp. 328–353, doi:10.1016/S0022-0000(77)80034-2.
- [9] Joost Engelfriet & Erik Meineche Schmidt (1978): *IO and OI. II*. J. Comput. System Sci. 16(1), pp. 67–99, doi:10.1016/0022-0000(78)90051-X.
- [10] Matthew Hague, Andrzej S. Murawski, C.-H. Luke Ong & Olivier Serre (2008): *Collapsible Pushdown Automata and Recursion Schemes*. In: Proceedings of the 23rd Annual IEEE Symposium on Logic in Computer Science (LICS), IEEE Computer Society, pp. 452–461.
- [11] Klaus Indermark (1976): *Schemes with Recursion on Higher Types*. In: Proc. 5th Symposium, Mathematical Foundations of Computer Science 1976, Lecture Notes in Comput. Sci. 45, Springer-Verlag, pp. 352–358.
- [12] Teodor Knapik, Damian Niwiński & Pawel Urzyczyn (2002): *Higher-Order Pushdown Trees Are Easy*. In: Proceedings of the 5th International Conference on Foundations of Software Science and Computation Structures (FoSSaCS), Lecture Notes in Comput. Sci. 2303, Springer-Verlag, pp. 205–222, doi:10.1007/3-540-45931-6_15.
- [13] Naoki Kobayashi (2009): *Types and higher-order recursion schemes for verification of higher-order programs*. In: Proc. 36th ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages (POPL), ACM, pp. 416–428.
- [14] Naoki Kobayashi & C.-H. Luke Ong (2009): *A Type System Equivalent to the Modal Mu-Calculus Model Checking of Higher-Order Recursion Schemes*. In: Proceedings of the 24th Annual IEEE Symposium on Logic in Computer Science (LICS), IEEE Computer Society, pp. 179–188.
- [15] M. Nivat (1972): *On the interpretation of recursive program schemes*. In: Symposia Matematica.

Appendices

A From OI to IO

Complement of Definitions

A n holes context is a term $C[\bullet_1^{\tau_1}, \dots, \bullet_n^{\tau_n}] \in \mathcal{T}(\Gamma \uplus \{\bullet_i^{\tau_i} : \tau_i \mid 1 \leq i \leq n\})$ containing exactly one occurrence of \bullet_i for all i . (We will generally omit to write the type τ_i in the notation $\bullet_i^{\tau_i}$).

For all $i_1, \dots, i_k \leq n$ we are interested in the application

$$t_1, \dots, t_k \mapsto (C[\bullet_1, \dots, \bullet_n])_{[\forall j \leq k \bullet_{i_j} \mapsto t_j]}$$

with $t_j \in \mathcal{T}_{\tau_{i_j}}(\Gamma)$ for all j . (notice that the order of the substitution is not important). One can consider $(C[\bullet_1, \dots, \bullet_n])_{[\forall j \leq k \bullet_{i_j} \mapsto t_j]}$ as a $n - k$ holes context. We may write $C[\bullet_1] \dots [\bullet_n]$ to denote the context $C[\bullet_1, \dots, \bullet_n]$ and extend this notation to $(C[\bullet_1, \dots, \bullet_n])_{[\forall j \leq k \bullet_{i_j} \mapsto t_j]}$, for example, given the context $C[\bullet_1][\bullet_2][\bullet_3]$, we define $C[t_1][\bullet_2][t_3] = (C[\bullet_1][\bullet_2][\bullet_3])_{[\bullet_1 \mapsto t_1, \bullet_3 \mapsto t_3]}$

Given a one hole context $C[\bullet]$, we define inductively the **head symbol sequence** $\text{hss}(C)$ which is a finite sequence of symbols of Γ : if $C[\bullet] = \bullet$, then $\text{hss}(C)$ is the empty sequence, if $C[\bullet] = \alpha t_1, \dots, t_{i-1} C'[\bullet] t_{i+1} \dots t_k$, then $\text{hss}(C) = \alpha, \text{hss}(C')$.

Proposition 7. *Given a n holes context $C[\bullet_1] \dots [\bullet_n]$, for all i , for all $t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_n$ and $s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_n$:*

$$\text{hss}(C[t_1] \dots [t_{i-1}][\bullet][t_{i+1}] \dots [t_n]) = \text{hss}(C[s_1] \dots [s_{i-1}][\bullet][s_{i+1}] \dots [s_n])$$

of Proposition 7. We prove this proposition by induction on the size of the context, for all n . If $C[\bullet_i] = \bullet_i$ is a 1 hole context, then the result is proven. If $C[\bullet_1] \dots [\bullet_n] = a t_1 \dots t_n$ there exists exactly one k such that t_k contains one occurrence of \bullet_i , if we look all the occurrences of some \bullet_j in t_k , we can state that t_k is a l holes context $C'[\bullet_{j_1}] \dots [\bullet_{j_l}]$ for some l . Moreover

$$\text{hss}(C[t_1] \dots [t_{i-1}][\bullet][t_{i+1}] \dots [t_n]) = a, \text{hss}(C'[t_{j_1}] \dots [\bullet_i][t_{j_l}]) \text{ and}$$

$$\text{hss}(C[s_1] \dots [s_{i-1}][\bullet][s_{i+1}] \dots [s_n]) = a, \text{hss}(C'[s_{j_1}] \dots [\bullet_i][s_{j_l}])$$

Since $\text{hss}(C'[t_{j_1}] \dots [\bullet_i][t_{j_l}]) = \text{hss}(C'[s_{j_1}] \dots [\bullet_i][s_{j_l}])$ by hypothesis of induction, we have

$$\text{hss}(C[t_1] \dots [t_{i-1}][\bullet][t_{i+1}] \dots [t_n]) = \text{hss}(C[s_1] \dots [s_{i-1}][\bullet][s_{i+1}] \dots [s_n])$$

□

Proposition 8. *Let t be a term, and let $C_1[\bullet]$ and $C_2[\bullet]$ be two contexts such that $t = C_1[t_1]$ and $t = C_2[t_2]$, then: either there exists a context C'_1 such that $C_2[\bullet] = C_1[C'_1[\bullet]]$ (1), Or there exists a context C'_2 such that $C_1[\bullet] = C_2[C'_2[\bullet]]$ (2), or there exists a two hole context $C[\bullet_1][\bullet_2]$ such that $C[t_1][\bullet] = C_2[\bullet]$ and $C[\bullet][t_2] = C_1[\bullet]$ (3).*

of Proposition 8. We proceed by induction. If $C_1[\bullet] = \bullet$, then $C_2[\bullet] = C_1[C_2[\bullet]]$, in the same way, if $C_2[\bullet] = \bullet$, $C_1[\bullet] = C_2[C_1[\bullet]]$. Else $C_1[\bullet] = a s_1 \dots s_{i-1} C'_1[\bullet] s_{i+1} \dots s_k$ and $C_2[\bullet] = a s_1 \dots s_{j-1} C'_2[\bullet] s_{j+1} \dots s_k$. If $i \neq j$ (we assume w.l.o.g. that $i < j$) we set $C[\bullet_1][\bullet_2] = a s_1 \dots s_{i-1} C'_1[\bullet_1] s_{i+1} \dots s_{j-1} C'_2[\bullet_2] s_{j+1} \dots s_k$, we have that $C[t_1][\bullet] = C_2$ and $C[\bullet][t_2] = C_1[\bullet]$. If $i = j$, then $C'_1[t_1] = C'_2[t_2]$, by induction:

- Either there exists a two hole context $C'[\bullet_1][\bullet_2]$ such that $C'[t_1][\bullet] = C_2'[\bullet]$ and $C'[\bullet][t_2] = C_1'[\bullet]$, in that case we set $C[\bullet_1][\bullet_2] = a s_1 \dots s_{i-1} C'[\bullet_1][\bullet_2] s_{i+1} \dots s_k$, and we have $C[t_1][\bullet] = C_2[\bullet]$ and $C[\bullet][t_2] = C_1[\bullet]$,
- Or there exists a context C_1'' such that $C_2'[\bullet] = C_1''[C_1'[\bullet]]$, in that case $C_2[\bullet] = C_1[C_1''[\bullet]]$.
- Or there exists a context C_2'' such that $C_1'[\bullet] = C_2''[C_2'[\bullet]]$, in that case $C_1[\bullet] = C_2[C_2''[\bullet]]$.

□

Correctness of the Transformation

We remark that \bar{t} does not contain any occurrence of Δ , I or δ . We also remark that any subterms of \bar{t} has type $\bar{\tau}$ for some type τ , hence it can not have ground type and in particular, it is not a redex. Moreover, given two terms t_1, t_2 , if the term $\overline{t_1 t_2}$ is valid, then it is equal to $\overline{t_1} \overline{t_2}$, in particular, it is an “overlined” term. It follows by induction, that given three terms t, t_1, t_2 , if the term $\bar{t}_{[\bar{t}_1 \rightarrow \bar{t}_2]}$ is well defined, then it is equal to $\overline{t_{[t_1 \rightarrow t_2]}}$ which is also well defined.

Proof of Lemma 1. First We need the following claim.

Claim 9. *For all accessible term t (with unrestricted derivations), for all context $C[\bullet]$ such that $t = C[\text{red}]$ with red being a redex, $\text{hss}(C)$ only contains terminal symbols. Furthermore, the term red doesn't contain occurrence of any terminal symbol.*

Proof of Claim 9. We prove it by induction. I satisfies Claim 9, $\bar{S} \Delta$ too. Assume that $t = C[\bar{F} t_1 \dots t_k]$ satisfies Claim 9, with $k = \text{arity}(\bar{F})$ and $F \in \mathcal{N}$. If $\bar{F} x_1 \dots x_k \rightarrow_{\bar{G}} \bar{e} \Delta \in \bar{\mathcal{R}}$, let $t' = C[\bar{e}_{[\forall i x_i \rightarrow t_i]} \Delta]$. Let $C'[\bullet]$ a context and $\text{red} = \gamma s_1 \dots s_{\text{arity}(\gamma)}$ with $\gamma \in \bar{\mathcal{N}}$ a redex such that $t' = C'[\text{red}]$.

First, we notice that since $\bar{e}_{[\forall i x_i \rightarrow t_i]} \Delta$ is a ground type term only containing non-terminal symbols, it is a redex, let $\rho r_1 \dots r_{\text{arity}(\rho)-1} \Delta = \bar{e}_{[\forall i x_i \rightarrow t_i]} \Delta$.

Using Proposition 8 we now that there are four options:

1. either $C[\bullet] = C'[\gamma s_1 \dots s_{i-1} C''[\bullet] s_{i+1} \dots s_{\text{arity}(\gamma)}]$ with C'' a context,
2. or $C[\bullet] = C[\rho r_1 \dots r_{i-1} C''[\bullet] r_{i+1} \dots r_{\text{arity}(\rho)-1} \Delta]$ with C'' a context,
3. or $C'[\bullet] = C[\bullet]$,
4. or there exists a two holes context $\mathbb{C}[\bullet_1][\bullet_2]$ such that $C[\bullet] = \mathbb{C}[\bullet][\text{red}]$ and $C'[\bullet] = \mathbb{C}[\bar{e}_{[\forall i x_i \rightarrow t_i]} \Delta][\bullet]$.

Option 1 is impossible, otherwise γ would be an element of $\text{hss}(C)$.

Option 2 would imply that $\bar{e}_{[\forall i x_i \rightarrow t_i]} = \rho r_1 \dots r_{i-1} C''[\text{red}] r_{i+1} \dots r_{\text{arity}(\rho)-1}$ then $\bar{e}_{[\forall i x_i \rightarrow t_i]}$ contains a ground typed term, which can't be true, see Remark A.

If **Option 3** is true. Then $\text{hss}(C') = \text{hss}(C)$ which by induction only contains terminal symbols. Since there is no terminal symbols in \bar{e} and in t_i for all i , there is no terminal in $\bar{e} \Delta = \text{red}$. Hence t' satisfies Claim 9.

If **Option 4** is true, then $t = \mathbb{C}[\bar{F} t_1 \dots t_n][\text{red}]$. Then by induction, $\text{hss}(\mathbb{C}[\bar{F} t_1 \dots t_n][\bullet])$ only contains terminal symbols. But, using Proposition 7, we know that

$$\text{hss}(C'[\bullet]) = \text{hss}(\mathbb{C}[\bar{e}_{[\forall i x_i \rightarrow t_i]} \Delta][\bullet]) = \text{hss}(\mathbb{C}[\bar{F} t_1 \dots t_n][\bullet]).$$

Then $\text{hss}(C'[\bullet])$ only contains terminal symbols. Furthermore, since red is a subterm of t , by induction it only contains non-terminal symbols, which proves that t' satisfies Claim 9.

Assume that $t = C[\bar{a} t_1 \dots t_k]$ satisfies Claim 9 with $a \in \Sigma$ and $k = \text{arity}(a)$, and let $t' = C[a(t_1 \Delta) \dots (t_k \Delta)]$. We can prove that t' satisfies Claim 9 in a similar way. □

Let $t = C[\text{red}]$ an accessible term with red a redex, let exp be the rewrite expression of red , and let's look at the derivation $t = C[\text{red}] \rightarrow_{\bar{G}} t' = C[\text{exp}]$.

Claim 9 tells us that $\text{hss}(C)$ only contains terminals, hence there is no redex containing an occurrence of \bullet in C , hence the derivation is OI. Assume that $\text{red} = \gamma t_1 \dots t_{i-1} C'(t) t_{i+1} \dots t_{\text{arity}(\gamma)}$ with C' a context and t a term.

Then $t = C[\gamma t_1 \dots t_{i-1} C'(t) t_{i+1} \dots t_{\text{arity}(\gamma)}]$ then $\text{hss}(C[\gamma t_1 \dots t_{i-1} C'(\bullet) t_{i+1} \dots t_{\text{arity}(\gamma)}])$ contains a non terminal symbol γ , hence Claim 9 tells that t is not a redex, so no non-trivial subterm of red is a redex, so the derivation is IO. \square

Proof of Theorem 2. Lemma 1 shows that we only have to prove that $\|G\|_{OI} = \|\bar{G}\|$. Concretely we will show that:

$$\forall t \in \text{Acc}_G, \exists t' \in \text{Acc}_{\bar{G}} : t^\perp \sqsubseteq (t')^\perp \quad (1)$$

$$\forall t' \in \text{Acc}_{\bar{G}}, \exists t \in \text{Acc}_G : (t')^\perp \sqsubseteq t^\perp \quad (2)$$

Definition 1 ($\|\cdot\|$ -transformation). We define inductively the transformation $\|\cdot\| : \mathcal{T}^o(\Sigma \uplus \mathcal{N}) \rightarrow \mathcal{T}^o(\Sigma \uplus \bar{\mathcal{N}})$:

- $\|a t_1 \dots t_{\text{arity}(a)}\| = a \|t_1\| \dots \|t_{\text{arity}(a)}\|$ for all $a \in \Sigma$,
- $\|\text{red}\| = \bar{\text{red}} \Delta$ for red a redex.

Remark 1. Notice that $t^\perp = (\|t\|)^\perp$.

Claim 10. If $t \in \mathcal{T}^o(\Sigma \uplus \mathcal{N})$ then $\bar{t} \Delta \rightarrow_{\bar{G}} \|t\|$.

Proof. The proof is done by induction. If t is a redex then $\|t\| = \bar{t} \Delta$. If $t = a t_1 \dots t_k$ with $a \in \Sigma$ and $k = \text{arity}(a)$, assume that for all i , $\bar{t}_i \Delta \rightarrow \|t_i\|$. We have $\bar{t} \Delta = \bar{a} \bar{t}_1 \dots \bar{t}_k$ so $\bar{t} \rightarrow_{\bar{G}} a (\bar{t}_1 \Delta) \dots (\bar{t}_k \Delta)$. Hence $\bar{t} \Delta \rightarrow_{\bar{G}}^* a \|t_1\| \dots \|t_k\| = \|t\|$. \square

Claim 11. For all t , if $t \in \text{Acc}_G$, then $\|t\| \in \text{Acc}_{\bar{G}}$. *This claim implies property (1).*

Proof of Claim 11. We prove this by induction. If $t = S$, $\|t\| = S \Delta$, and $I \rightarrow S \Delta$, so $\|t\| \in \text{Acc}_{\bar{G}}$.

Let $t = C[F t_1 \dots t_k] \in \text{Acc}_G$ with $k = \text{arity}(F)$ and $\text{hss}(C)$ contains only terminal symbols. Assume that $\|t\| \in \text{Acc}_{\bar{G}}$. Given that $F x_1 \dots x_k \rightarrow e \in \mathcal{R}$, let $t' = C[e_{[\forall i x_i \rightarrow t_i]}]$.

First, given a ground type context $C[\bullet^o]$, we can define the associated ground type context $\|C\|[\bullet^o]$ by adding to Definition 1 the fact $\|\bullet^o\| = \bullet^o$. Hence we can say that $\|t\| = \|C\|[\bar{F} \bar{t}_1 \dots \bar{t}_k \Delta]$.

We see that $\|t\| \rightarrow_{\bar{G}} \|C\|[\bar{e}_{[\forall i x_i \rightarrow \bar{t}_i]} \Delta]$. Claim 10 shows that $\bar{e}_{[\forall i x_i \rightarrow \bar{t}_i]} \Delta = \bar{e}_{[\forall i x_i \rightarrow t_i]} \Delta \rightarrow_{\bar{G}}^* \|e_{[\forall i x_i \rightarrow t_i]}\|$, hence $\|t\| \rightarrow_{\bar{G}} \|C\|[\bar{e}_{[\forall i x_i \rightarrow \bar{t}_i]} \Delta] \rightarrow_{\bar{G}}^* \|C\|[\|e_{[\forall i x_i \rightarrow t_i]}\|] = \|t'\|$. \square

Claim 12. Given a term $t' \in \text{Acc}_{\bar{G}}$ there exists a term $t \in \text{Acc}_G$ such that $t' \rightarrow_{\bar{G}} \|t\|$. *This claim implies property (2).*

Proof of Claim 12. We will divide the relation $\rightarrow_{\bar{G}}$ in two relations: $\rightarrow_{\bar{G}} = \rightarrow_{\Sigma} \uplus \rightarrow_{\mathcal{N}}$ depending of the head symbol of the redex we're rewriting. Let $t \rightarrow_{\bar{G}} t'$ if the rewrite rule applied is $r_{\bar{a}}$ for some $a \in \Sigma$ then $t \rightarrow_{\Sigma} t'$, if the rewrite rule is $r_{\bar{F}}$ with $F \in \mathcal{N}$, then $t \rightarrow_{\mathcal{N}} t'$.

The proof is in four steps:

1. Given a term $t \in \text{Acc}_{\bar{G}}$, there is only a finite number of derivation $t \rightarrow_{\Sigma}^* t'$, furthermore, if $t \rightarrow_{\Sigma}^* t_1$ and $t \rightarrow_{\Sigma}^* t_2$ such that there is no t' such that $t_1 \rightarrow_{\Sigma}^* t'$ or $t_2 \rightarrow_{\Sigma}^* t'$, then $t_1 = t_2$. We name this unique term t^{Σ} , and we notice that if $t \rightarrow_{\Sigma}^* t'$ then $t' \rightarrow_{\Sigma}^* t^{\Sigma}$. Basically this step comes from the fact that the relation \rightarrow_{Σ} strictly decrease the number of occurrences of terms headed by some \bar{a} with $a \in \Sigma$.
2. Let $t = C[\bar{F} t_1 \dots t_{\text{arity}(\bar{F})}]$, then $t^{\Sigma} = C^{\Sigma}[\bar{F} t_1 \dots t_{\text{arity}(\bar{F})}]$, C^{Σ} being defined inductively: if $C[\bullet] = \bullet$ then $C^{\Sigma}[\bullet] = \bullet$, if $C[\bullet] = a t_1 \dots C'[\bullet] \dots t_k$, then $C^{\Sigma}[\bullet] = a t_1^{\Sigma} \dots C'^{\Sigma}[\bullet] \dots t_k^{\Sigma}$ (Claim 9 shows that these are the only possibilities). This step is shown by induction.
3. If $t \rightarrow_{\mathcal{N}} t'$ i.e. $t = C[\bar{F} t_1 \dots t_k]$ and $t' = C[\bar{e}_{[\forall i x_i \mapsto \bar{t}_i]} \Delta]$ with the appropriate \bar{e} , let $t'' = C^{\Sigma}[\bar{e}_{[\forall i x_i \mapsto \bar{t}_i]} \Delta]$, then $t'^{\Sigma} = t''^{\Sigma}$.
4. Finally we prove by induction that for all t' , there exists $t \in \text{Acc}_G$ such that $t'^{\Sigma} = \|t\|$, which proves the claim.

□

□

B The Type System Detailed

We give here a complete proof of Theorem 3 and Proposition 4. We first recall the type system, and the definition of $\vdash (G, t) \triangleright \sigma$.

$$\begin{array}{c}
\frac{\Theta \vdash t \triangleright \theta_1 \quad \dots \quad \Theta \vdash t \triangleright \theta_n}{\Theta \vdash t \triangleright \bigwedge \{ \theta_1, \dots, \theta_n \}} (Set) \\
\frac{}{\Theta, \alpha \triangleright \bigwedge \{ \theta_1, \dots, \theta_n \} \vdash \alpha \triangleright \theta_i} (At) \quad (for\ all\ i) \\
\frac{}{\Theta \vdash a \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_{i \leq \text{arity}(a)} \rightarrow q_{\infty}} (\Sigma) \quad (for\ a\ a \in \Sigma\ and\ \exists j\ \sigma_j = q_{\infty}) \\
\frac{\Theta \vdash t_1 \triangleright \sigma \rightarrow \theta \quad \Theta \vdash t_2 \triangleright \sigma}{\Theta \vdash t_1 t_2 \triangleright \theta} (App) \\
\frac{}{\Theta \vdash t \triangleright q_{\infty} \rightarrow q_{\infty}} (q_{\infty} \rightarrow q_{\infty} I) \quad (if\ t : \tau_1 \rightarrow \tau_2) \\
\frac{\Theta \vdash t_1 \triangleright q_{\infty}}{\Theta \vdash t_1 t_2 \triangleright q_{\infty}} (q_{\infty} I)
\end{array}$$

Remark that:

- Using rule (Set) one can always prove, for any term t , $\Theta \vdash t \triangleright \bigwedge \{ \}$.
- $\Theta \vdash t \triangleright \bigwedge \{ \theta_1, \dots, \theta_k \}$ if and only if, for all i , $\Theta \vdash t \triangleright \theta_i$.
- There is no rules that directly involve q_{\perp} , but that does not mean that no term matches q_{\perp} , since it can appears in Θ . Rules like (At) or (App) may be use to state that a term matches q_{\perp} .

We say that (G, t) matches the conjunctive mapping σ written $\vdash (G, t) \triangleright \sigma$ if there exists an environment Θ , called a witness environment of $\vdash (G, t) \triangleright \sigma$, which verifies the following properties:

1. $\text{Dom}(\Theta) = \mathcal{N}$,
2. $\forall F : \tau \in \mathcal{N} \quad \Theta(F) :: \tau$,

3. if $F x_1 \dots x_k \rightarrow e \in \mathcal{R}$ and $\Theta \vdash F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_{i \leq k} \rightarrow q$ then either there exists j such that $q_\infty \in \sigma_j$, or $i = k$ and $\Theta, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e \triangleright q$,
4. $\Theta \vdash t \triangleright \sigma$.

Lemma 13 (Isolated Non Terminals). *Given a non terminal F that has not ground type. Then if Θ verifies properties 1 to 3, one cannot prove $\Theta \vdash F \triangleright q_\infty$.*

Proof of lemma 13. The proof comes from the fact that Θ verifies property 3. Assume $\Theta \vdash F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_i \rightarrow q_\infty$. Property 3 states that if $i < \text{arity}(F)$ then there exists $j \leq i$ such that $q_\infty \in \sigma_j$ in particular $i \neq 0$. If $i = \text{arity}(F)$ then by hypothesis, $i \neq 0$. Then, one cannot prove $\Theta \vdash F \triangleright q_\infty$. \square

Lemma 14 (Non fully-applied terminals). *Let Θ be an environment that verifies properties 1 to 3, let F be a non terminal that has not ground type and let $t = F t_1 \dots t_i$ with $i < \text{arity}(F)$. If $\Theta \vdash F t_1 \dots t_i \triangleright q_\infty$ then there exists $j \leq i$ such that $\Theta \vdash t_j \triangleright q_\infty$.*

Proof of Lemma 14. We prove by induction on l the following more general result: if $\Theta \vdash F t_1 \dots t_l \triangleright \sigma_{l+1} \rightarrow \dots \rightarrow \sigma_i \rightarrow q_\infty$ with $F \in \mathcal{N}$ and $i < \text{arity}(F)$, then there exists $j \leq i$ such that $\Theta \vdash t_j \triangleright q_\infty$ if $j \leq l$ or $q_\infty \in \sigma_j$ if $j > l$.

If $l = 1$, then the rule we used to prove $\Theta \vdash F t_1 \triangleright \sigma_2 \rightarrow \dots \rightarrow \sigma_i \rightarrow q_\infty$ could not be $(q_\infty I)$ since one cannot prove $\Theta \vdash F \triangleright q_\infty$. If the rule we used were $(q_\infty \rightarrow q_\infty I)$, then $q_\infty \in \sigma_2$. If it is rule (App) then $\Theta \vdash F \triangleright \sigma_1 \rightarrow \sigma_2 \rightarrow \dots \rightarrow \sigma_i \rightarrow q_\infty$ and $\Theta \vdash t_1 \triangleright \sigma_1$, and since $i < \text{arity}(F)$, property 3 states that there exists $j < l$ such that $\Theta \vdash t_j \triangleright q_\infty$ if $j = 1$, $q_\infty \in \sigma_j$ elsewise. These are the only rules we could have applied.

If $l > 1$. If we applied rule $(q_\infty I)$ then $\Theta \vdash F t_1 \dots t_{l-1} \triangleright q_\infty$ by induction hypothesis there exists $j \leq l-1$ such that $\Theta \vdash t_j \triangleright q_\infty$. If we applied rule $(q_\infty \rightarrow q_\infty I)$, then $q_\infty \in \sigma_{l+1}$. If we applied rule (App) then $\Theta \vdash F t_1 \dots t_{l-1} \triangleright \sigma_l \rightarrow \sigma_{l+1} \rightarrow \dots \rightarrow \sigma_i \rightarrow q_\infty$ and $\Theta \vdash t_l \triangleright \sigma_l$ then by induction hypothesis there exists $j \leq i$ such that either $\Theta \vdash t_j \triangleright q_\infty$ if $j < l$ or $q_\infty \in \sigma_j$ if $j \geq l$. If $j \leq l$ then either $j < l$ and then $\Theta \vdash t_j \triangleright q_\infty$, or $j = l$ and $q_\infty \in \sigma_j$ hence $\Theta \vdash t_j \triangleright q_\infty$, if $j > l$ then $q_\infty \in \sigma_j$ if $j \geq l$. \square

Lemma 15 (Redexes and q_∞). (1) *If Θ verifies properties 1 to 3, then given a term t , if $\Theta \vdash t \triangleright q_\infty$ then either t contains a redex r such that $\Theta \vdash r \triangleright q_\infty$. In particular, if t does not contains any redex, then one cannot prove $\Theta \vdash t \triangleright q_\infty$.*

(2) *If Θ verifies properties 1 to 3, then given a term t , if t contains a redex r such that $\Theta \vdash r \triangleright q_\infty$ then one can prove $\Theta \vdash t \triangleright q_\infty$.*

Proof of Lemma 15. We prove (1) by induction on t . Assume that $\Theta \vdash t \triangleright q_\infty$.

If $t : o$ then either $t = F t_1 \dots t_k$ in which case t is the redex r , or $t = a t_1 \dots t_k$ then the only rule we could have applied to prove $\Theta \vdash t \triangleright q_\infty$ is (Σ) , then there exists $t_i : o$ such that $\Theta \vdash t_i \triangleright q_\infty$, and the result comes by induction.

If $t : \tau$ with $\tau \neq o$. We could not have $t = F \in \mathcal{N}$ since one cannot prove $\Theta \vdash F \triangleright q_\infty$ if F has not ground type. If $t = a t_1 \dots t_i$ then again there exists $t_i : o$ such that $\Theta \vdash t_i \triangleright q_\infty$, and the result comes by induction. If $t = F t_1 \dots t_i$, F has not ground type, and $i < \text{arity}(F)$ since t has not ground type. Then Lemma 14 states that there exists t_j such that $\Theta \vdash t_j \triangleright q_\infty$ and the result comes by induction.

To prove (2), assume that there is a redex r such that $\Theta \vdash r \triangleright q_\infty$ and $t = C[r]$. We prove the result by induction on $C[\bullet]$. If $C[\bullet] = \bullet$, then $t = r$ therefore $\Theta \vdash r \triangleright q_\infty$. Assume $t = t_1 t_2$ with $t_1 = C'[r]$ or $t_2 = C'[r]$. If $t_1 = C'[r]$ then by induction hypothesis, one can prove $\Theta \vdash t_1 \triangleright q_\infty$ and then, using rule $q_\infty I$, $\Theta \vdash t_1 t_2 \triangleright q_\infty$. If $t_2 = C'[r]$, by induction hypothesis, one can prove $\Theta \vdash t_2 \triangleright q_\infty$, using rule $(q_\infty \rightarrow q_\infty I)$, we have $\Theta \vdash t_1 \triangleright q_\infty \rightarrow q_\infty$ and then, rule (App) gives us $\Theta \vdash t_1 t_2 \triangleright q_\infty$. \square

Lemma 16 (Ground type terms and q_{\perp}). *if Θ verifies properties 1 to 3, then if $t : \tau$ and $\Theta \vdash t \triangleright \sigma$, $\sigma :: \tau$. In particular, if $\Theta \vdash t \triangleright q_{\perp}$, then $t : o$.*

Proof of Lemma 16. We can assume, without loss of generality that $\sigma = \{\theta\}$ for some atomic mapping θ . We prove this by induction on the structure of t .

If $t = \alpha$ with $\alpha \in \Sigma \uplus \mathcal{N}$, then the only rules we can apply are (At) , (Σ) and $(q_{\infty} \rightarrow q_{\infty} I)$ and they all satisfy the property.

If $t = t_1 t_2$ with $t_1 : \tau_2 \rightarrow \tau$ and $t_2 : \tau_2$, then the rules we can apply are either (q_{∞}) or (App) . If it is (q_{∞}) then we have proven $\Theta \vdash t \triangleright q_{\infty}$ and $q_{\infty} :: \tau$. If it is (App) it means that we have proven $\Theta \vdash t_1 \triangleright \sigma' \rightarrow \theta$ and $\Theta \vdash t_2 \triangleright \sigma'$, and by induction hypothesis, $\sigma' :: \tau_2$ and $\theta :: \tau$. \square

Theorem 17 (Soundness). *Let G be an HORS, and t be term (of any type), if $\vdash (G, t) \triangleright q_{\infty}$ (resp. $\vdash (G, t) \triangleright q_{\perp}$) then $\mathcal{P}_{\infty}(t)$ (resp. $\mathcal{P}_{\perp}(t)$) holds.*

Proof of Theorem 17.

Lemma 18 (Type Preservation). *Let $t : \tau$ be a term. If $\vdash (G, t) \triangleright \sigma$ and $t \rightarrow_{IO} t'$ then $\vdash (G, t') \triangleright \sigma$.*

Proof of Lemma 18. Assume that $\vdash (G, t) \triangleright \sigma$ and $t \rightarrow_{IO} t'$. Let Θ be a witness environment of $\vdash (G, t) \triangleright \sigma$, we will prove that it is also a witness environment of $\vdash (G, t') \triangleright \sigma$ (we only have to check that $\Theta \vdash t' \triangleright \sigma$).

We know that $t = C[F s_1 \dots s_k]$ and $t' = C[e_{[\forall i x_i \rightarrow s_i]}]$ for some context $C[\bullet : o] : \tau$. We proceed by induction on $C[\bullet]$.

If $C[\bullet] = \bullet$, we can assume without loss of generality that $\sigma = q \in Q$. We look at the proof of $\vdash (G, t') \triangleright \sigma$ and remark that either (1) the proof contains $\Theta \vdash F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow \sigma$ and $\Theta \vdash s_i \triangleright \sigma_i$ for all i and the last steps are using the rule (App) , or (2) the proving tree contains $\Theta \vdash F s_1 \dots s_i \triangleright q_{\infty} \rightarrow q_{\infty}$ and $\Theta \vdash s_{i+1} \triangleright q_{\infty}$, the last steps are using the rule (App) once and then only rule $(q_{\infty} I)$. The former case is impossible: since $t \rightarrow_{IO} t'$ is an IO derivation there's no redex in s_i for all i , hence Lemma 15 shows that one cannot have $\Theta \vdash s_{i+1} \triangleright q_{\infty}$.

Hence the proving tree contains $\Theta \vdash F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow \sigma$ and $\Theta \vdash s_i \triangleright \sigma_i$ for all i , then $\Theta, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e \triangleright q$, and if we replace all statements of $x_i \triangleright \sigma_i$ by the proof of $\Theta \vdash s_i \triangleright \sigma_i$, we obtain a proof of $\Theta \vdash e_{[\forall i x_i \rightarrow s_i]} \triangleright \sigma$.

Now we prove the induction step. Assume that $C = C'[\bullet] t_2$ or $C = t_1 C'[\bullet]$, then $t = t_1 t_2$ with either $t_1 = C'[F s_1 \dots s_k]$ or $t_2 = C'[F s_1 \dots s_k]$. Then $t' = t'_1 t'_2$ with respectively, either $t'_1 = C'[e_{[\forall i x_i \rightarrow s_i]}]$ and $t'_2 = t_2$, or $t'_1 = t_1$ and $t'_2 = C'[e_{[\forall i x_i \rightarrow s_i]}]$. Either way, by induction hypothesis, if $\Theta \vdash t_1 \triangleright \sigma_1$ (resp. $\Theta \vdash t_2 \triangleright \sigma_2$), then $\Theta \vdash t'_1 \triangleright \sigma_1$ (resp. $\Theta \vdash t'_2 \triangleright \sigma_2$). Assume we have proven $\Theta \vdash t \triangleright \sigma$. In order to do it, either we have use rule $(q_{\infty} I)$ or rule (App) , either way we could use the same rule to prove $\Theta \vdash t' \triangleright \sigma$. \square

We extend in an intuitive way the properties \mathcal{P}_{\perp} and \mathcal{P}_{∞} to trees: if t is a tree then $\mathcal{P}_{\perp}(t) = "t = \perp"$ and $\mathcal{P}_{\infty}(t) = "t$ is either infinite or contains $\perp"$.

Lemma 19 (Weak Soundness). *Given a term $t : o$, (1) if $\vdash (G, t) \triangleright q_{\perp}$, then $\mathcal{P}_{\perp}(t^{\perp})$ holds, (2) if $\vdash (G, t) \triangleright q_{\infty}$, then $\mathcal{P}_{\infty}(t^{\perp})$ holds.*

Proof of Lemma 19. We can use Lemma 15 to prove (2): if $\vdash (G, t) \triangleright q_{\infty}$ then t contains a redex, hence t^{\perp} contains \perp , therefore $\mathcal{P}_{\infty}(t^{\perp})$ holds.

We prove (1) by induction on the structure of t^{\perp} . If $t^{\perp} = \perp$ then $\mathcal{P}_{\perp}(t^{\perp})$ is true hence (1) holds. If $t^{\perp} = a$ with $a \in \Sigma$, then $t = a$ and there is no rule that we can apply to state $\vdash (G, a) \triangleright q_{\perp}$, hence (1) and (2) holds. If $t^{\perp} = a t'_1 \dots t'_k$ with $k > 0$, then $t = a t_1 \dots t_k$ with $a \in \Sigma$ and $t_i^{\perp} = t'_i$ for all i . For all

environment Θ , we show by induction that for all i , if $\Theta \vdash a t_1 \dots t_i \triangleright \sigma'$ then $\sigma' = \sigma_1 \rightarrow \dots \rightarrow \sigma_l \rightarrow q_\infty$: The term a can only be judge by the rule (Σ) hence it is true if $i = 0$, the term $(a t_1 \dots t_i) t_{i+1}$ can be judge by rules $(q_\infty I)$, $(q_\infty \rightarrow q_\infty I)$ and (App) and by induction hypothesis, in all three cases, we get $\Theta \vdash a t_1 \dots t_i \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_l \rightarrow q_\infty$ for some l . In particular, we don't have $\vdash (G, t) \triangleright q_\perp$, hence (1) holds. \square

Using Lemma 1 and 2, in order to prove Theorem 17 we can assume that $t : o$. We prove it by contradiction. Assume that $\vdash (G, t) \triangleright q_\infty$ but $\mathcal{P}_\infty(t)$ doesn't hold. Then it means that $\|G_t\|$ is finite and contains only terminals. Since it's finite, there exists a finite IO derivation from t that leads to $\|G_t\|$: $t \rightarrow_{IO}^* \|G_t\|$, hence using Lemmas 18 and 19 we can prove $\mathcal{P}_\infty((\|G_t\|)^\perp)$, but since $\|G_t\|$ is a tree, $\|G_t\| = (\|G_t\|)^\perp$, hence $\|G_t\|$ is infinite or contains \perp which raises a contradiction.

We treat the case $\vdash (G, t) \triangleright q_\perp$ the same way: Assume that $\vdash (G, t) \triangleright q_\perp$ but $\mathcal{P}_\perp(t)$ doesn't hold. Then it means that $\|G_t\|$ contains some terminals. Then there exists a finite IO derivation from t that leads to a term t' such that $t'^\perp \neq \perp$: $t \rightarrow_{IO}^* t'$, hence using Lemmas 18 and 19 we can prove $\mathcal{P}_\perp((t')^\perp)$ which is false. \square

Theorem 20 (Completeness). *Let G be an HORS, if $\mathcal{P}_\infty(t)$ (resp. $\mathcal{P}_\perp(t)$) holds then $\vdash (G, t) \triangleright q_\infty$ (resp. $\vdash (G, t) \triangleright q_\perp$).*

Proof of Theorem 20.

Using Lemma 15 we can assume without loss of generality that t has ground type.

We recall the properties that an environment Θ has to satisfy in order to be a witness of $\vdash (G, t) \triangleright \sigma$.

1. $Dom(\Theta) = \mathcal{N}$,
2. $\forall F : \tau \in \mathcal{N} \ \Theta(F) :: \tau$,
3. if " $F x_1 \dots x_k \rightarrow e$ " $\in \mathcal{R}$ and $\Theta \vdash F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_{i \leq k} \rightarrow q$ then either there exists j such that $q_\infty \in \sigma_j$, or $i = k$ and $\Theta, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e \triangleright q$,
4. $\Theta \vdash t \triangleright \sigma$.

Let \mathcal{E} be the set of environment that matches properties 1 and 2. Let $\mathcal{F} : \mathcal{E} \rightarrow \mathcal{E}$ be a mapping such that for all $F : \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o \in \mathcal{N}$, if $F x_1 \dots x_k \rightarrow e \in \mathcal{R}$ then,

$$\begin{aligned} \mathcal{F}(\Theta)(F) = & \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \mid q \in Q \wedge \forall i \ \sigma_i :: \tau_i \wedge \Theta, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e : q \} \\ & \cup \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_{i \leq k} \rightarrow q_\infty \mid \wedge \forall i \ \sigma_i :: \tau_i \wedge \exists j \ q_\infty \in \sigma_j \} \\ & \cup \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q_\perp \mid \forall i \ \sigma_i :: \tau_i \wedge \exists j \ q_\infty \in \sigma_j \}. \end{aligned}$$

Let $\Theta_0 \in \mathcal{E}$ be the environment such that, for all $F : \tau = \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o \in \mathcal{N}$, $\Theta_0(F)$ is defined and contains all atomic mappings $\theta ::_a \tau$. Notice that:

$$\Theta_0(F) = \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \mid q \in Q \wedge \forall i \ \sigma_i :: \tau_i \} \cup \{ \sigma_1 \rightarrow \dots \rightarrow \sigma_{i < k} \rightarrow q_\infty \mid \forall j \ \sigma_j :: \tau_j \}.$$

Lemma 21 (Universal Witness). *There exists $m \in \mathbb{N}$ such that the judgment $\vdash (G, t) \triangleright \sigma$ holds if and only if $\mathcal{F}^m(\Theta_0) \vdash t \triangleright \sigma$ (This is Proposition 4 with $\Theta^* = \mathcal{F}^m(\Theta_0)$).*

Proof of Lemma 21. We define the partial order \sqsubseteq on \mathcal{E} such that $\Theta_1 \sqsubseteq \Theta_2$ if and only if, for all $F \in \mathcal{N}$, $\Theta_1(F) \subseteq \Theta_2(F)$. Note that if $\Theta_1 \sqsubseteq \Theta_2$ and $\Theta_1 \vdash t \triangleright \sigma$ then $\Theta_2 \vdash t \triangleright \sigma$. $\Theta_0(F)$ contains all atomic mappings $\theta ::_a \tau$, hence Θ_0 is a maximum of \mathcal{E} with respect to \sqsubseteq . Note that the mapping \mathcal{F} is monotonic with

respect to \sqsubseteq (i.e. if $\Theta \sqsubseteq \Theta'$ then $\mathcal{F}(\Theta) \sqsubseteq \mathcal{F}(\Theta')$). Given $\Theta \in \mathcal{E}$, we say that Θ is a post-fixpoint of \mathcal{F} if and only if $\Theta \sqsubseteq \mathcal{F}(\Theta)$. Remark that being a post-fixpoint of \mathcal{F} is the same as verifying property 3.

Since Θ_0 is a maximum of \mathcal{E} , and $\mathcal{F}(\Theta_0) \in \mathcal{E}$, then $\Theta_0 \sqsupseteq \mathcal{F}(\Theta_0)$, therefore, since \mathcal{F} is monotonic, $\Theta_0 \sqsupseteq \mathcal{F}(\Theta_0) \sqsupseteq \mathcal{F}^2(\Theta_0) \sqsupseteq \dots$. Because \mathcal{E} is finite, there exists m such that $\mathcal{F}^m(\Theta_0) = \mathcal{F}^{m+1}(\Theta_0)$, in particular $\mathcal{F}^m(\Theta_0)$ is a post-fixpoint of \mathcal{F} , hence it verifies properties 1, 2, and 3.

Take a witness Θ of $\vdash (G, t) \triangleright \sigma$. Θ is a post-fixpoint of \mathcal{F} , and since $\mathcal{F}^m(\Theta_0)$ is the greatest post-fixpoint, $\mathcal{F}^m(\Theta_0) \sqsupseteq \Theta$, hence $\mathcal{F}^m(\Theta_0) \vdash t \triangleright \sigma$, thus $\mathcal{F}^m(\Theta_0)$ is a witness of $\vdash (G, t) \triangleright \sigma$. \square

Let $G^{(m)} = \langle \mathcal{V}, \Sigma, \mathcal{N}^{(m)} \uplus \{\text{Void} : o\}, \mathcal{R}^{(m)}, I \rangle$ be the scheme such that $\mathcal{N}^{(m)} = \bigcup_{0 \leq i \leq m} \{F_i \mid F \in \mathcal{N}\}$. For all $F \ x_1 \dots x_k \rightarrow e \in \mathcal{R}$, $\mathcal{R}^{(m)}$ contains the following rewrite rules:

$$F_i \ x_1 \dots x_k \rightarrow e_{[\forall H \in \mathcal{N} \ H \mapsto H_{i-1}]} \quad \text{for } i > 0$$

$$F_0 \ x_1 \dots x_k \rightarrow e_{[\forall H \in \mathcal{N} \ H \mapsto H_0]}$$

$$F_0 \ x_1 \dots x_k \rightarrow \text{Void}$$

$$\text{Void} \rightarrow \text{Void}$$

Notice that *Void* here is a non-terminal of order 0 that produce itself. Hence applying its rewrite rule to a term would produce the same term. In the following we forbid this rule to be applied. $G^{(m)}$ with this restriction is said to be recursion free, i.e. the graph whose vertices are the non terminals and where there is an edge from F to G if and only if there exist an allowed rewrite rule $F \ x_1 \dots x_k \rightarrow e$ such that e contains an occurrence of G , has no loop. Such non-recursive schemes are known to be strongly normalizing, i.e. for any term t all derivations using only allowed rewrite rules are finite. In particular, there exists a finite IO derivation $t \rightarrow_{IO}^* t'$ such that $(t')^\perp = \|t\|_{IO}$.

We define the environment $\Theta^{(m)}$ on $\mathcal{N}^{(m)} \uplus \{\text{Void}\}$: for all $F \in \mathcal{N}$, for all $i \leq j$, $\Theta^{(m)}(F_i) = \mathcal{F}^i(\Theta_0)(F)$ and $\Theta^{(m)}(\text{Void}) = \bigwedge \{q_\infty, q_\perp\}$.

Lemma 22. *Given a two terms $t, t' \in \mathcal{T}(\Sigma \uplus \mathcal{N}^{(m)})$ such that $t \rightarrow_{IO} t'$ is allowed in $G^{(m)}$. If $\Theta^{(m)} \vdash t' \triangleright \sigma$, then $\Theta^{(m)} \vdash t \triangleright q$.*

Proof of Lemma 22. We proceed by induction on the structure of t . We prove the initial case: $t = F_l \ s_1 \dots s_k$ and $t' = e_{[\forall i \ x_i \mapsto s_i]}$, with $F_l \ x_1 \dots x_k \rightarrow e \in \mathcal{R}$. We assume without loss of generality that σ is atomic, hence $\sigma = q \in Q$. Let σ_i be the union of all mappings assigned to s_i in the proof of $\Theta^{(m)} \vdash e_{[\forall i \ x_i \mapsto s_i]} \triangleright q$. Then we have $\Theta^{(m)}, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e \triangleright q$. Let $\Theta' = \Theta^{(m)}, F_l \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q$. Since $\Theta' \vdash F_l \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q$, and $\Theta' \vdash s_i \triangleright \sigma_i$ (indeed, $\Theta^{(m)} \subseteq \Theta'$), we can prove $\Theta' \vdash t' \triangleright q$. If $l = 0$, by definition, $F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \in \Theta_0(F)$, and since $\Theta^{(m)}(F_0) = \Theta_0(F)$, we have $F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \in \Theta^{(m)}(F_0)$, hence $\Theta' = \Theta^{(m)}$. If $l > 0$, e only contains terminals of the form G_{l-1} , then we can transform the proof of $\Theta^{(m)}, x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e \triangleright q$ to obtain a proof of $\mathcal{F}^{l-1}(\Theta_0), x_1 \triangleright \sigma_1, \dots, x_k \triangleright \sigma_k \vdash e_{[\forall G \in \mathcal{N} \ G_{l-1} \mapsto G]} \triangleright q$. Then by definition of \mathcal{F} , $F \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \in \mathcal{F}^l(\Theta_0)(F)$, and since $\Theta^{(m)}(F_l) = \mathcal{F}^l(\Theta_0)(F)$, we have $\sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \in \Theta^{(m)}(F_l)$, hence $\Theta' = \Theta^{(m)}$. Thus $\Theta^{(m)} \vdash t \triangleright q$.

For the induction step, We assume without loss of generality that $\sigma = \bigwedge \{\theta\}$. Assume that $C = C'[\bullet] \ t_2$ or $C = t_1 \ C'[\bullet]$, then $t = t_1 \ t_2$ with either $t_1 = C'[F \ s_1 \dots s_k]$ or $t_2 = C'[F \ s_1 \dots s_k]$. Then $t' = t'_1 \ t'_2$ with respectively, either $t'_1 = C'[e_{[\forall i \ x_i \mapsto s_i]}]$ and $t'_2 = t_2$, or $t'_1 = t_1$ and $t'_2 = C'[e_{[\forall i \ x_i \mapsto s_i]}]$. Either way, by induction hypothesis, if $\Theta^{(m)} \vdash t'_1 \triangleright \sigma_1$ (resp. $\Theta^{(m)} \vdash t'_2 \triangleright \sigma_2$), then $\Theta^{(m)} \vdash t_1 \triangleright \sigma_1$ (resp. $\Theta^{(m)} \vdash t_2 \triangleright \sigma_2$). Assume we

have proven $\Theta^{(m)} \vdash t' \triangleright \sigma$. In order to do it, either we have used rule $(q_\infty I)$ or rule (App) , either way we could use the same rule to prove $\Theta^{(m)} \vdash t \triangleright \sigma$. \square

Lemma 23 (Terms that contains *Void*). *Given a term t that contains the non terminal *Void*, one can prove $\Theta^{(m)} \vdash t \triangleright q_\infty$.*

Proof of Lemma 23. We prove the result by induction on the structure of t : if $t = \text{Void}$ then we use rule (At) , if $t = G_j t_1 \dots t_i$ with $G_j \in \mathcal{N}^{(m)}$ and t_ℓ contains *Void* for some $\ell \leq i$, then by induction hypothesis, $\Theta^{(m)} \vdash t_\ell \triangleright q_\infty$. Using rule (Set) we have, for all $\ell' \neq \ell$, $\Theta^{(m)} \vdash t_{\ell'} \triangleright \bigwedge\{\}$. We have by construction $\Theta^{(m)} \vdash a \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q_\infty$ for $\sigma_j = q_\infty$ and $\sigma_i = \bigwedge\{\}$ for all $i \neq j$. Then, if we apply k times rule (App) we can prove $\Theta^{(m)} \vdash a t_1 \dots t_k \triangleright q_\infty$. By definition, if $\sigma_{\ell' \neq \ell} = \bigwedge\{\}$ and $\sigma_\ell = q_\infty$, we have $\sigma_1 \rightarrow \dots \rightarrow \sigma_i \rightarrow q_\infty \in \Theta^{(m)}(G_j)$. Hence one can prove $\Theta^{(m)} \vdash t_i \triangleright q_\infty$.

if $t = G_j t_1 \dots t_i$ with $a \in \Sigma$ and t_ℓ contains *Void* for some $\ell \leq i$ the proof is similar except we use rule (Σ) to prove $\Theta^{(m)} \vdash a \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_i \rightarrow q_\infty$ with $\sigma_{\ell' \neq \ell} = \bigwedge\{\}$ and $\sigma_\ell = q_\infty$. \square

Lemma 24 (Weak Completeness). *Given a term $t : o \in \mathcal{T}(\Sigma \uplus \mathcal{N}^{(m)})$, if $\mathcal{P}_\perp(t^\perp)$ (resp. $\mathcal{P}_\infty(t^\perp)$) holds and if there exists no IO allowed rewrite rule we can apply in t , then $\Theta^{(m)} \vdash t \triangleright q_\perp$ (resp. $\emptyset \vdash t^\perp \triangleright q_\infty$).*

Proof of Lemma 24. We prove both results simultaneously by induction on the structure of t . If $t = \text{Void}$ then one can directly prove both result using rule (At) . We know that $t \neq a$ with $a \in \Sigma$ since $\mathcal{P}_\perp(a)$ (resp. $\mathcal{P}_\infty(a)$) does not hold. If $t = a t_1 \dots t_k$, we know that $\mathcal{P}_\perp(t^\perp)$ doesn't hold, assume that $\mathcal{P}_\infty(t^\perp)$ holds. Since t^\perp contains \perp , there exists j such that t_j^\perp contains \perp , i.e. $\mathcal{P}_\infty(t_j^\perp)$. Furthermore there exists no allowed rewrite rule we can apply in t_j , elseway we could apply it in t . Therefore, by induction hypothesis, $\Theta^{(m)} \vdash t_j \triangleright q_\infty$. Using rule (Set) we have, for all $i \neq j$, $\Theta^{(m)} \vdash t_i \triangleright \bigwedge\{\}$. Rule (Σ) gives $\Theta^{(m)} \vdash a \triangleright \sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q_\infty$ for $\sigma_j = q_\infty$ and $\sigma_i = \bigwedge\{\}$ for all $i \neq j$. Then, if we apply k times rule (App) we can prove $\Theta^{(m)} \vdash a t_1 \dots t_k \triangleright q_\infty$.

Assume now that $t = F_l t_1 \dots t_k$ with $F_l \in \mathcal{N}^{(m)}$. Since there exists no IO allowed rewrite rule we can apply in t it means that there exists i such that t_i contains a redex, but this redex can't be applied, in other words, t_i contains *Void*. Using Lemma 23 we have $\Theta^{(m)} \vdash t_i \triangleright q_\infty$. By definition, if $\sigma_{j \neq i} = \bigwedge\{\}$ and $\sigma_i = q_\infty$, we have $\sigma_1 \rightarrow \dots \rightarrow \sigma_k \rightarrow q \in \Theta^{(m)}(F_l)$ for $q \in Q$. Hence one can prove $\Theta^{(m)} \vdash t \triangleright q$. \square

Now, we can prove the Theorem. Given a term $t : o \in \mathcal{T}(\Sigma \uplus \mathcal{N})$, assume that $\mathcal{P}_\perp(t)$ (resp. $\mathcal{P}_\infty(t)$) holds. We define the term $t^{(m)} : o = t_{[\forall F \in \mathcal{N} \ F \mapsto F_i]} \in \mathcal{T}(\Sigma \uplus \mathcal{N}^{(m)})$. Notice that $\|G_t^{(m)}\|_{IO}$ is obtained by turning some subtrees of $\|G_t\|_{IO}$ into \perp . Hence, $\mathcal{P}_\perp(t^{(m)})$ (resp. $\mathcal{P}_\infty(t^{(m)})$) holds. Let $t' : o \in \mathcal{T}(\Sigma \uplus \mathcal{N}^{(m)} \uplus \{\perp\})$ such that $t^{(m)} \rightarrow_{IO}^* t'$ and $(t')^\perp = \|G_t^{(m)}\|_{IO}$ (we have seen previously that such t' exists). Lemma 24 states that $\Theta^{(m)} \vdash t' \triangleright q_\perp$ (resp. $\Theta^{(m)} \vdash t' \triangleright q_\infty$), then, Lemma 22 shows that $\Theta^{(m)} \vdash t^{(m)} \triangleright q_\perp$ (resp. $\Theta^{(m)} \vdash t^{(m)} \triangleright q_\infty$). Since non terminals in $t^{(m)}$ have the form F_m , if we restrict the domain of $\Theta^{(m)}$ only to $\{F_m \mid F \in \mathcal{N}\}$ the proof still holds, furthermore in this proof, if we remove all “ m ” subscripts, we get $\mathcal{F}^m(\Theta_0) \vdash t \triangleright q_\perp$ (resp. $\mathcal{F}^m(\Theta_0) \vdash t \triangleright q_\infty$). Lemma 21 allows us to conclude: $\vdash (G, t) \triangleright q_\perp$ (resp. $\vdash (G, t) \triangleright q_\infty$). \square

C Selfcorrecting Scheme

Proof of Theorem 5.

Lemma 25 (Equality of Trees). *Let $t : o \in \mathcal{T}(\mathcal{V} \uplus \mathcal{N})$ be a term, then $t^\perp = (t^+)^\perp$.*

Proof of Lemma 25. We prove it by induction on the structure of $t : o$. If $t = F t_1 \dots t_k$ with $F \in \mathcal{N}$ then $t^\perp = \perp$ and $t^+ = F \llbracket t_1 \rrbracket, \dots, \llbracket t_k \rrbracket t_1^{\bar{\sigma}_1^{\tau_1}} \dots t_k^{\bar{\sigma}_k^{\tau_k}}$, then $(t^+)^\perp = \perp = t^\perp$.

If $t = a t_1 \dots t_k$ with $a : o^k \rightarrow o \in \Sigma$ and $t_i : o$ for all i , then $t^+ = a \llbracket t_1 \rrbracket, \dots, \llbracket t_k \rrbracket t_1^+ \dots t_k^+$, $t^\perp = a t_1^\perp \dots t_k^\perp$ and $(t^+)^\perp = a (t_1^+)^\perp \dots (t_k^+)^\perp$. By induction hypothesis, for all i $(t_i^+)^\perp = t_i^\perp$, then $(t^+)^\perp = a t_1^\perp \dots t_k^\perp = t^\perp$. \square

Lemma 26 (Label Conservation in Rewrite Rules). *Given a term $t : \tau = \tau_1 \rightarrow \dots \rightarrow \tau_k \rightarrow o \in \mathcal{T}(\Sigma \uplus \mathcal{N})$, such that $t = F s_1 \dots s_k$ with $F \in \mathcal{N}$, and t is an IO-relevant redex. Note that $t^+ = F \llbracket s_1 \rrbracket, \dots, \llbracket s_k \rrbracket s_1^{\bar{\sigma}_1^{\tau_1}} \dots s_k^{\bar{\sigma}_k^{\tau_k}}$. If $F x_1 \dots x_k \rightarrow e \in \mathcal{R}$, let $t' = e_{[\forall i x_i \mapsto s_i]}$ and $s = e_{\Theta^{\mathcal{Y}} \left[\begin{smallmatrix} \bar{\sigma}_i^{\tau_i} & \bar{\sigma}_j^{\tau_j} \\ [x_i & \mapsto s_i^j] \end{smallmatrix} \right]}$ with $\Theta^{\mathcal{Y}}(x_i) = \llbracket s_i \rrbracket$ for all i (in particular, $t \rightarrow t'$ and $t^+ \rightarrow s$).*

We have $s = (t')^+$.

Proof of Lemma 26. Besides the labeling, by construction, s matches $(t')^+$. Take a subterm e' of e , if one can prove $\Theta^* \vdash e'_{[\forall i x_i \mapsto s_i]} \triangleright \sigma$, then one can prove $\Theta^*, \Theta^{\mathcal{Y}} \vdash e' \triangleright \sigma$, hence s is well labeled, therefore $s = (t')^+$. Then $t^+ \rightarrow_{IO}^* (t')^+$. \square

Given two terms t, t' , we write $t \Rightarrow_{IO} t'$ if t' is obtained by applying in parallel all IO rewrite availables in t . Formally, we define it inductively: if t is an IO-relevant redex and t' is the term obtained by rewriting this redex then $t \Rightarrow_{IO} t'$. If t is not an IO redex and $t = t_1 t_2$ then $t \Rightarrow_{IO} t'$ if and only if:

- either there exists t'_1, t'_2 such that $t_1 \Rightarrow_{IO} t'_1$ and $t_2 \Rightarrow_{IO} t'_2$, and $t' = t_1 t_2$,
- or there exists t'_1 such that $t_1 \Rightarrow_{IO} t'_1$ but no t'_2 such that $t_2 \Rightarrow_{IO} t'_2$ and $t' = t'_1 t_2$,
- or there exists t'_2 such that $t_2 \Rightarrow_{IO} t'_2$ but no t'_1 such that $t_1 \Rightarrow_{IO} t'_1$ and $t' = t_1 t'_2$.

Notice that if such t' exists then it is unique, and it exists if and only if t contains a redex. The $\cdot \Rightarrow_{IO} \cdot$ relation is known as parallel rewrite, and from a term $t : o$, the unique associated parallel derivation $t \Rightarrow_{IO} t_1 \Rightarrow_{IO} t_2 \Rightarrow_{IO} \dots$ leads to the tree $\|G_t\|$.

Lemma 27 (Coincidence of Parallel Derivation). *Given a terms $t \in \mathcal{T}(\Sigma \uplus \mathcal{N})$, and some conjunctive mappings $\sigma_1, \dots, \sigma_k$. There exists $t' \in \mathcal{T}(\Sigma \uplus \mathcal{N})$ such that $t \Rightarrow_{IO} t'$, if and only if there exists $s' \in \mathcal{T}(\Sigma' \uplus \mathcal{N}')$ such that $t^{+\sigma_1, \dots, \sigma_k} \Rightarrow_{IO} s'$. Furthermore, if it is true, then $s' = (t')^{+\sigma_1, \dots, \sigma_k}$.*

Proof of Lemma 27. The first part of the result comes from the observation that t contains a redex if and only if $t^{+\sigma_1, \dots, \sigma_k}$ contains a redex. We prove the second part by induction. If t is an IO-relevant redex, $t^{+\sigma_1, \dots, \sigma_k}$ is too, and Lemma 26 proves the result. If $t = t_1 t_2$, t is not an IO-relevant redex a then $t^{+\sigma_1, \dots, \sigma_k} = t_1^{+\sigma_1, \dots, \sigma_k} t_2^{+\bar{\sigma}_1^{\tau_1} \dots \bar{\sigma}_k^{\tau_k}}$ and $t^{+\sigma_1, \dots, \sigma_k}$ is not an IO-relevant redex. Assume that $t \Rightarrow_{IO} t'$ then,

- either there exists t'_1, t'_2 such that $t_1 \Rightarrow_{IO} t'_1$ and $t_2 \Rightarrow_{IO} t'_2$, and $t' = t'_1 t'_2$,
- or there exists t'_1 such that $t_1 \Rightarrow_{IO} t'_1$ but no t'_2 such that $t_2 \Rightarrow_{IO} t'_2$ and $t' = t'_1 t_2$,
- or there exists t'_2 such that $t_2 \Rightarrow_{IO} t'_2$ but no t'_1 such that $t_1 \Rightarrow_{IO} t'_1$ and $t' = t_1 t'_2$.

By induction hypothesis, $t_i \Rightarrow_{IO} t'_i$ if and only if $t_i^+ \Rightarrow_{IO} t'^+_i$ for $i \in \{1, 2\}$, hence

- either there exists t'_1, t'_2 such that $t_1^{+\sigma, \sigma_1, \dots, \sigma_k} \Rightarrow_{IO} t'_1+$ and $t_2^{+\vec{\sigma}_j} \Rightarrow_{IO} t'_2+$ for all j , and $(t')^{+\sigma_1, \dots, \sigma_k} = (t'_1)^{+\sigma, \sigma_1, \dots, \sigma_k} (t'_2)^{+\vec{\sigma}_1} \dots (t'_2)^{+\vec{\sigma}_r}$,
- or there exists t'_1 such that $t_1^{+\sigma, \sigma_1, \dots, \sigma_k} \Rightarrow_{IO} t'_1+$ but no s'_2 such that $t_2^{+\vec{\sigma}_j} \Rightarrow_{IO} s'_2$ for all j , and $(t')^{+\sigma_1, \dots, \sigma_k} = (t'_1)^{+\sigma, \sigma_1, \dots, \sigma_k} (t_2)^{+\vec{\sigma}_1} \dots (t_2)^{+\vec{\sigma}_r}$,
- or there exists t'_2 such that $t_2^{+\vec{\sigma}_j} \Rightarrow_{IO} (t'_2)^{+\vec{\sigma}_j}$ but no s'_1 such that $t_1^{+\sigma, \sigma_1, \dots, \sigma_k} \Rightarrow_{IO} s'_1$, and $(t')^{+\sigma_1, \dots, \sigma_k} = (t_1)^{+\sigma, \sigma_1, \dots, \sigma_k} (t'_2)^{+\vec{\sigma}_1} \dots (t'_2)^{+\vec{\sigma}_r}$.

Therefore, $t^{+\sigma_1, \dots, \sigma_k} \Rightarrow (t')^{+\sigma_1, \dots, \sigma_k}$. \square

Given a term $t : o$ let $t \Rightarrow_{IO} t_1 \Rightarrow_{IO} t_2 \Rightarrow_{IO} \dots$ be the parallel derivation associated to it. Thanks to Lemma 27 we know that the parallel derivation associated to t^+ is $t^+ \Rightarrow_{IO} t_1^+ \Rightarrow_{IO} t_2^+ \Rightarrow_{IO} \dots$, then $\|G'_{t^+}\|_{IO}$ is the limit of $(t_i^+)^{\perp}$ then $(\|G'_{t^+}\|_{IO})^-$ is the limit of $((t_i^+)^{\perp})^- = (t_i)^{\perp}$. Then $\|G'_{t^+}\|_{IO} = \|G_t\|_{IO}$. \square

Proof of $\|G''\|_{IO} = \|G'\|_{IO}$. Take a term $t \in \mathcal{T}(\Sigma \uplus \mathcal{N})$ we define $void(t) \in \mathcal{T}(\Sigma \uplus \mathcal{N} \uplus \{Void\})$ as the set of termes obtained by substituing some redex r in t such that $\|G'_r\|_{IO} = \perp$ by $Void$. From the definition comes that if $t' \in void(t)$ then $(t')^{\perp} = t^{\perp}$.

Given a term $t \in \mathcal{T}(\Sigma \uplus \mathcal{N})$ and an IO derivation associated $t = t_1 \rightarrow_{IO} t_2 \rightarrow_{IO} \dots$ in G' we construct by induction an IO derivation in G'' $t = t'_1 \rightarrow_{IO} t'_2 \rightarrow_{IO} \dots$ such that for all i $t'_i \in void(t_i)$. The initial step is straightforward: $t \in void(t)$, Assume that $t'_i \in void(t_i)$, and assume that $t_i = C[F t_1 \dots t_k]$ and $t_{i+1} = C[e_{[\forall i, x_i \mapsto t_i]}]$ with $F x_1 \dots x_k \rightarrow e \in \mathcal{R}'$. If this redex is a subterm of another one that is transformed by $Void$ in t'_i then we just rewrite this void obtaining $t'_{i+1} = t'_i$ by induction hypothesis, we still have $t'_{i+1} \in void(t_{i+1})$. If this redex is not transformed in t'_i then we rewrite this redex, and either $\|F t_1 \dots t_k\|_{IO} = \perp$, in which case the semantics associated contains q_{\perp} thanks to Theorem 5, and then $e_{[\forall i, x_i \mapsto t_i]}$ is still a redex and is transformed to $Void$ in t'_{i+1} or $\|F t_1 \dots t_k\|_{IO} \neq \perp$ and no more transformation is added to create t'_{i+1} , in both cases $t'_{i+1} \in void(t_{i+1})$.

This result gives that $\|G'\|_{IO} \sqsubseteq \|G''\|_{IO}$. Since G'' is obtained by from G' changing some redex into other that will produce \perp , it is clear that $\|G''\|_{IO} \sqsubseteq \|G'\|_{IO}$, then $\|G'\|_{IO} = \|G''\|_{IO}$. \square

Proof of $\|G''\| = \|G''\|_{IO}$. We already know that $\|G''\|_{IO} \sqsubseteq \|G''\|$, we just have to show that for all t , if there is \perp at node u in $\|G''_t\|_{IO}$ then there is \perp at node u in $\|G''_t\|$. We show this by induction on the size of u .

If $\|G''_t\|_{IO} = \perp$. Then $\|G'_t\|_{IO} = \|G_t\|_{IO} = \perp$, hence $\llbracket t \rrbracket = \perp$, hence the only derivation in G'' from t is $t \rightarrow Void \rightarrow Void \rightarrow \dots$, therefore $\|G''\| = \perp$. If $u = ju'$ then let a be the terminal at the root of $\|G''\|_{IO}$, then there exists an IO derivation $t \rightarrow^* a t_1 \dots t_k$, and $\|G''_j\|_{IO}$ is equal to the subtree of $\|G''_t\|_{IO}$ rooted at node j . Since $t \rightarrow^* a t_1 \dots t_k$, $\|G''_j\|$ is equal to the subtree of $\|G''_t\|$ rooted at node j and by induction hypothesis, since there is \perp at node u' in $\|G''_j\|_{IO}$, there is \perp at node u' in $\|G''_j\|$ hence there is \perp at node u in $\|G''_t\|$. \square