



Photoinductive degradation of two pesticides by natural dissolved organic matter under simulated sunlight

Emilie Caupos, Patrick Mazellier, Jean-Philippe Croue

► To cite this version:

Emilie Caupos, Patrick Mazellier, Jean-Philippe Croue. Photoinductive degradation of two pesticides by natural dissolved organic matter under simulated sunlight. TransCon2010, Sep 2010, Switzerland. <hal-00822551>

HAL Id: hal-00822551

<https://hal-enpc.archives-ouvertes.fr/hal-00822551>

Submitted on 14 May 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Photoinductive degradation of 17 β -estradiol and isoproturon by natural dissolved organic matter under simulated sunlight





Emilie CAUPOS^{a*}, Marie DEBORDE^a, Patrick MAZELLIER^{a,b}, Jean-Philippe CROUE^{a,c}
^a Laboratoire de Chimie et Microbiologie de l'Eau, Poitiers, France
^b Laboratoire de Physico et Toxico Chimie de l'Environnement, LPTC, Bordeaux, France
^c Kaust, Saudi Arabia * contact : emilie.caupos@univ-poitiers.fr

- > Pesticides are common micropollutants of surface waters. Beside them, pharmaceuticals are now detected, coming from natural excretions or WWTPs.
- > Natural Dissolved Organic Matter (DOM) is ubiquitous in surface waters. It is a complex organic mixture formed by natural decomposition of ecosystems.
- > Photodegradation of aquatic micropollutants can be enhanced through DOM sensitization pathway involving the formation of reactive species.

- > The natural photodegradation of a natural hormone (17 β -estradiol, E2) and a pesticide (isoproturon, IPU) has been studied in the presence of DOM.
- > Sunlight irradiation was performed by a Suntest CPS+ solar simulator (290 nm < λ < 700 nm, 250 W/m², 8h). Concentrations used in the experiments were approximately 1 μ M. Photodegradation efficiencies have been measured in the presence of DOM (20 mg/L) isolated from surface waters.
- > The formation of reactive species was investigated and photoproducts were identified through LC-MS analyses.

A – Photodegradation in the presence of DOM


- Pollutants degradation is always faster in the presence of DOM than in purified water
- Enhancement effect strongly depends on the nature of DOM extracts
- An order of DOM efficiency is observed :
 South Platte (69%, E2; 72%, IPU) >>
 Pinail (41%, E2) >
 Suwannee (39%, E2; 37%, IPU)



B – DOM production of reactive species

- Photodegradation only weakly decreases with the addition of inhibitors (azide \rightarrow 1O_2 ; 2-propanol \rightarrow OH^\bullet)
- For E2 :
 Without inhibitor > with NaN₃ > with 2-propanol
- For IPU :
 Without inhibitor \geq with 2-propanol > with NaN₃
- Reactive species involved :
 > 1O_2 : participation in the photodegradation about 4 – 19%
 > OH^\bullet : participation in the photodegradation about 0 – 17%


Pathway of E2



C – Pathways and photoproducts

- 8 and 5 photoproducts have been identified during the degradation of E2 and IPU respectively
- Pathway of E2 :
 1: hydroxylation of aromatic cycle or cycle closed to aromatic one (E21-E24)
 2: E2 quinone methide derivative (E27)
 3: hydroxylation of aromatic cycle and oxidation of phenolic groups (E25 and E26)
- Pathway of IPU :
 1: demethylation of dimethylurea group (I5)
 2: hydroxylation of aromatic cycle (I1 and I4)
 3: simultaneous demethylation and hydroxylation of isopropyl group (I2 and I3)

Pathway of IPU



Conclusions

- > DOM presents an ability to photoinduce the degradation of micropollutants; its efficiency depending on the micropollutant and on its own properties (nature of the extract)
- > During photoinductive degradation, reactive species such as singlet oxygen and hydroxyl radicals are produced by DOM and react with the pollutants. However, the participation of these 2 molecules has only been observed. Excited triplet states may contribute to a large extent in the reaction.
- > Photoproducts have been observed. They come from hydroxylation, oxidation and demethylation/dealkylation mechanisms of the parent compound.