



Reply to comment by Poole et al. on a tropical NAT-like belt observed from space.

Vincent Noel, H el ene Chepfer

► To cite this version:

Vincent Noel, H el ene Chepfer. Reply to comment by Poole et al. on a tropical NAT-like belt observed from space.. Geophysical Research Letters, American Geophysical Union, 2009, 36, pp.L20804. <10.1029/2009GL039689>. <hal-00439699>

HAL Id: hal-00439699

<https://hal.archives-ouvertes.fr/hal-00439699>

Submitted on 8 Dec 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

1 Reply to the Comment by Poole et al. on “A tropical “NAT-like” belt observed from space”

2

3 V. Noel, LMD/IPSL, CNRS, France

4 H. Chepfer, LMD/IPSL, Univ. Paris 06, France

5

6 proposed for publication in Geophysical Research Letters

7

8 Corresponding author:

9 V. Noel

10 Laboratoire de Meteorologie Dynamique

11 Ecole Polytechnique

12 91128 Palaiseau

13 France

14 vincent.noel@lmd.polytechnique.fr

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

1 **Abstract**

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

In their comment, Poole et al. (2009) aim to show it is highly improbable that the observations described in Chepfer and Noel (2009), and described as "NAT-like" therein, are produced by Nitric Acid Trihydrate (NAT) particles. In this reply, we attempt to show why there is, in our opinion, too little evidence to reject this interpretation right away.

1 The main hypothesis in the Comment by Poole et al. (hereafter called P09) is that the
2 optical observations labeled "NAT-like" in Chepfer and Noel 2009 (hereafter called CN09)
3 are most likely a mixture of Liquid Aerosols (LA) and ice crystals, given particle number
4 concentrations from the literature.

5 First of all, we want to emphasize that, contrarily to what might be inferred from reading
6 P09, CN09 did not conclude that "NAT-like" observations should unambiguously be
7 attributed to NAT particles; the term "NAT-like" referred to the optical behavior of the
8 particles under study. Instead, CN09 proposed several possible interpretations, NAT
9 particles being one of them. The present reply aims to show that, while P09 certainly
10 brings cogent arguments that make the NAT interpretation less probable, it should not be
11 dismissed entirely yet.

12 13 **1. On the detection of optically very thin layers with CALIOP**

14
15 P09 shows that NAT particle concentrations of 10^{-4} cm^{-3} would produce a backscatter near
16 $10^{-6} \text{ km}^{-1} \text{ sr}^{-1}$, too low to be observed from CALIOP.

17 CN09 used observations in the lower tropical stratosphere to calibrate the background
18 backscatter (due to molecules), and identified layers as signal fluctuations above this
19 background. By extracting the particulate backscatter this way, we found it was possible to
20 detect layers with backscatter close to $10^{-5} \text{ km}^{-1} \text{ sr}^{-1}$ above the molecular signal in
21 CALIOP data, i.e. to identify layers that backscatter less than the molecular background
22 ($3\text{-}5 \cdot 10^{-4} \text{ km}^{-1} \text{ sr}^{-1}$ as shown in P09). Those layers possess cloud-like qualities: horizontal
23 and vertical extension, consistent geographic distribution over specific areas,
24 homogeneous optical properties (at least as homogeneous as in PSC observations), etc.

25 A recent paper by Immler et al. (2007) shows a very similar layer observed by ground-
26 based lidar, producing similar levels of backscatter (i.e. near $10^{-5} \text{ km}^{-1} \text{ sr}^{-1}$). The authors
27 speculate this layer is a remnant of ice clouds, stabilized by inorganic acids, and conclude
28 it is most likely made of NAT crystals in concentrations near 1 L^{-1} , i.e. 10^{-3} cm^{-3} . The
29 concentration-backscatter relationship is consistent with results from P09, though both
30 quantities are much higher. In our observations, NAT-like layers produce backscatters
31 between 10^{-5} and $5 \cdot 10^{-4} \text{ km}^{-1} \text{ sr}^{-1}$.

32
33 Since the 10^{-4} cm^{-3} concentration used in P09 for NAT is derived from in-situ studies (by
34 definition limited in scope and altitude), the generality of these numbers cannot be
35 assessed. Moreover, given the lower cut-size of the inlets used to bring particle-laden air

1 into the instruments that measure NO_y , a significant amount of small NAT particles might
2 be missed during in-situ measurements. Thus the existence of higher NAT concentrations
3 (e.g. 10^{-3} cm^{-3}) cannot be discounted yet, for example close to lightning events where
4 HNO_3 amounts increase. P09 conducted simulations of optical properties of particle
5 mixtures with such enhanced NAT levels, but results from these computations only appear
6 in their Fig. 2 and are somewhat surprising (see next section). Since their Fig. 1, which
7 shows backscatter levels expected for various particle mixtures, does not include results
8 from enhanced NAT levels, it does not preclude the possibility to detect NAT layers in such
9 concentrations in CALIOP observations.

10

11 **2. On the optical properties of particle mixtures**

12

13 First, P09 argues that mixtures of LA and NAT particles produce depolarization ratio below
14 the limit used in CN09 (0.05) to discriminate between liquid and solid particles: their Fig. 2
15 shows that, even considering enhanced levels of NAT (10^{-3} cm^{-3} as discussed in the
16 previous section), the produced depolarization ratios do not get higher than 0.02.
17 However, in Fig. 2 the enhanced NAT mixture does not produce higher backscatter than
18 the mixture with a 10^{-4} cm^{-3} NAT concentration ($0.3 < 1-1/R_{532} < 0.6$ in both cases),
19 backscatter levels even appear to be lower with the enhanced mixture (near $1-1/R_{532} \sim 0.6$).
20 It is rather surprising that increasing particle concentrations by an order of
21 magnitude would not lead to increased backscatter. This result, and thus the
22 depolarization ratio values, should be investigated further.

23 Second, P09 argues that a local mixture of liquid aerosols and ice crystals would lead to
24 intermediate values of volumic backscatter and depolarization (between those produced
25 by LA and ice crystals). From a theoretical point of view, it seems to us that the shape
26 hypothesis done for ice and NAT particles is a key point. P09 used oblate particles for
27 both, and it is unclear how much these shapes depolarize compared to others, especially
28 at the considered particle sizes. A higher amount of oblate particles might then be needed
29 to produce a given depolarization ratio compared to non-spheroidal particles - the
30 dependence of depolarization ratio on particle shape is a well-known complex problem
31 (e.g. Takano and Liou 1989). As in Fig. 2 of P09 the depolarization is used as a strong
32 discrimination criteria, it may be useful to test others particle shapes to check how they
33 affect the depolarization-to-concentration relationship. From an observational point of view,
34 in CN09 we attempted to extract the particulate backscatter and depolarization from the
35 volumic CALIOP observations by removing the background contribution. This approach

1 was able to retrieve particulate depolarization ratios typical of ice crystals (0.4-0.6) in
2 layers with a much lower volumic depolarization ratio (0-0.2), so we assumed the
3 background depolarization (including LA) was correctly cleaned up. After this processing,
4 the layers labeled NAT-like in CN09 still show a intermediate depolarization (0.1-0.3), while
5 other layers are clearly gathered near 0 (liquid aerosols) or 0.4-0.6 (ice crystals).

6 Finally, regarding the optical properties of particle layers, in addition to the backscatter and
7 depolarization ratios CN09 also used the color ratio, which has not been tested in P09. It
8 maybe be useful to test whether the mixing of ice+liquid aerosols as proposed by P09 is
9 consistent with observed color ratios.

10 Nonetheless, we agree that CN09 NAT-like layers could be due to (1) intense, small-scale,
11 local increases of BSA concentrations mixed with ice crystals (in which case the process
12 mentioned above would fail to totally remove the background contribution to the
13 backscatter); however this hypothesis does not seem necessarily more supported by
14 evidence than NAT particles, (2) more "exotic" processes (e.g. deposition of liquid aerosols
15 on ice crystals leading to particle rounding and weakened depolarization). In those cases,
16 the final particulate depolarization (after background correction) would indeed be
17 intermediate and could be consistent with observations. It would be very interesting to find
18 a way to discriminate between these two hypothesis (NAT particles / ice + liquid aerosols)
19 based on optical observations alone.

21 **3. Conclusion**

22
23 As already stated in CN09, we agree with P09 that there is not enough evidence to
24 unambiguously conclude that NAT-like layers in CN09 are actually composed of NAT
25 crystals. Mixtures of liquid aerosols and ice, as proposed by P09, appear as a convincing
26 possibility. As mentioned in CN09, other potential candidates include very small, non-
27 spherical ice crystals that have been observed in situ in significant concentrations (Gayet
28 et al. 2007); it remains to be seen if such particles are thermodynamically stable in the
29 specific conditions of the TTL. However, this reply tries to show why we also think that it is
30 too soon to definitely rule out a NAT composition for the non-ice / non-liquid layers in
31 CALIOP TTL observations. More studies, including remote sensing and in-situ
32 observations, are required to fully resolve these questions.

34 **References**

35

- 1 • Chepfer H, Noel V 2009: A tropical "NAT-like" belt observed from space. *Geophys. Res. Lett.* **36**, L03813.
- 2
- 3 • Gayet J.-F., Shcherbakov V, Mannstein H, Minikin A, Schumann U, Ström J, Petzold A, Ovarlez J, Immler F 2007: Microphysical and optical properties of midlatitude cirrus clouds observed in the southern hemisphere during INCA. *Q. J. R. Meteorol. Soc.* **132**,
- 4 2719.
- 5
- 6
- 7 • Hopfner M, Luo B, Massoli P, Cairo F, Spang R, Snels M, Don-Francesco G, Stiller G, Clarmann T, Fischer H, Biermann U 2006: Spectroscopic evidence for NAT, STS, and ice in MIPAS infrared limb emission measurements of polar stratospheric clouds. *Atmos. Chem. Phys.* **6**, 1201–1219.
- 8
- 9
- 10
- 11 • Immler F, Krueger K, Tegtmeier S, Fujiwara M, Fortuin P, Verver G, Schrems OJ 2007: Cirrus clouds, humidity, and dehydration in the tropical tropopause layer observed at
- 12 Paramaribo, Suriname (5.8°N, 55.2°W). *Geophys. Res. Lett.* **112**, D03209.
- 13
- 14 • Poole L. R., Pitts M. C. and Thomason, L. W. 2009: Comment on "A Tropical NAT-like belt observed from space by H. Chepfer and V. Noel". *Geophys. Res. Lett.*, this issue.
- 15
- 16 • Popp PJ, Marcy TP, Jensen EJ, Karcher B, Fahey DW, Gao RS, Thompson KH T. L. and Rosenlof, Richard EC, Herman RL, Weinstock EM, Smith JB, May RD, Vomel H, Wilson
- 17 JC, Heymsfield AJ, Mahoney MJ, Thompson AM 2006: The observation of nitric acid-containing particles in the tropical lower stratosphere. *Atmos. Chem. Phys.* **6**, 601-611.
- 18
- 19
- 20 • Takano Y, Liou K. 1989: Solar Radiative Transfer in Cirrus Clouds. Part I: Single-
- 21 Scattering and Optical Properties of Hexagonal Ice Crystals. *J. Atmos. Sci.* **46**, 3–18.
- 22
- 23 • Thampi BV, Sunilkumar SV, Parameswaran KJ 2009: Lidar studies of particulates in the UTLS region at a tropical station over the Indian subcontinent. *Geophys. Res. Lett.* **114**,
- 24 D08204.
- 25
- 26 • Toon OB, Pollack JB, Khare BN 1976: The optical constants of several atmospheric aerosol species: Ammonium sulfate, aluminium oxide, and sodium chloride. *J. Geophys. Res.* **81**(33), 5733-5748.
- 27
- 28 • Voigt C, Schlager H, Roiger A, Stenke A, de Reus M, Borrmann S, Jensen E, Schiller C, Konopka P, Sitnikov N. 2008: Detection of reactive nitrogen containing particles in the tropopause region - evidence for a tropical nitric acid trihydrate (NAT) belt. *Atmos. Chem. Phys.* **8**, 7421-7430.
- 29
- 30
- 31
- 32
- 33
- 34
- 35