



# Oceanic CO<sub>2</sub> outgassing triggered by terrestrial organic carbon fluxes during deglacial flooding

Thomas Extier, Katharina D. Six, Bo Liu, Hanna Paulsen, and Tatiana Ilyina

Max Planck Institute for Meteorology, Hamburg, Germany

**Correspondence to:** Thomas Extier (thomas.extier@mpimet.mpg.de)

## Abstract.

Oceans play a major role on the exchange of carbon with the atmosphere and thereby on past climates with glacial/interglacial variations of the CO<sub>2</sub> concentration. The melting of ice sheets during deglaciations lets the sea level rise which leads to the flooding of coastal land areas resulting in the transfer of terrestrial organic matter to the ocean. However, the consequences of such fluxes on the ocean biogeochemical cycle and uptake/release of CO<sub>2</sub> are poorly constrained. Moreover, this potentially important exchange of carbon at the land-sea interface is not represented in most Earth System Models. We present here the implementation of terrestrial organic matter fluxes into the ocean at the transiently changing land-sea interface in the Max Planck Institute for Meteorology Earth System Model (MPI-ESM) and investigate their effect on the biogeochemistry during the last deglaciation. Our results show that during the deglaciation, most of the terrestrial organic matter inputs to the ocean occurs during Meltwater Pulse 1a (between 15-14 ka) which leads to additional 21.2 GtC of terrestrial origin (mostly originating from wood and humus). Although this additional organic matter input is relatively small in comparison to the global ocean inventory (0.06 %) and thus doesn't have an impact on the global CO<sub>2</sub> flux, the terrestrial organic matter fluxes initiate oceanic outgassing at regional hotspots like in Indonesia for a few hundred years. Finally, sensitivity experiments highlight that terrestrial organic matter fluxes are the drivers of oceanic outgassing in flooded coastal regions during Meltwater Pulse 1a. Furthermore, the magnitude of outgassing is rather insensitive to higher carbon to nutrients ratios of the terrestrial organic matter. Our results provide a first estimate of the importance of terrestrial organic matter fluxes in a transient deglaciation simulation. Moreover, our model development is an important step towards a fully coupled carbon cycle in an Earth System Model applicable for simulations of glacial/interglacial cycles.

## 1 Introduction

Since the middle to late Pliocene (approximately 3 million years ago) the climate has undergone large variations triggered by the glacial and interglacial cycles with associated changes in ice sheets volume, sea level, oceanic circulation and atmospheric CO<sub>2</sub> concentration (e.g. Kohfeld and Ridgwell, 2009; Sosdian and Rosenthal, 2009; Spratt and Lisiecki, 2016). The last deglaciation defined to be from 21 to 10 ka (thousands of years before present) is the most recent manifestation of these changes and has been extensively examined in studies based on proxy records (e.g. Barker et al., 2009; Denton et al., 2010; Clark et al., 2012; Marcott et al., 2014) and Earth System Models (ESMs) with a focus on the timing of the last deglaciation, on



the ocean thermohaline circulation or on the ice sheet dynamics (e.g. Bonelli et al., 2009; Liu et al., 2009; Menviel et al., 2011; Roche et al., 2011; Heinemann et al., 2014; Ivanovic et al., 2016; Klockmann et al., 2016). The melting of ice sheets during the last deglaciation is accompanied by a sea level increase of about 95 m (Spratt and Lisiecki, 2016) resulting in flooding of land coastal areas and changes in the oceanic coastlines. In this case, the carbon and nutrients previously stored on land in vegetation and soil are transferred to the ocean, potentially impacting the global ocean biogeochemistry with implications for the uptake and release of carbon by the ocean. Indeed, the carbon and nutrients bound in the terrestrial material might change the ocean biogeochemistry once they have decomposed. A specificity of this terrestrial organic matter (terrOM) is the higher carbon to nutrients ratio than the marine organic matter. But as of now, the role of this terrestrial organic matter transfer to shelf areas on the glacial/interglacial atmospheric CO<sub>2</sub> variations remains poorly constrained. The representation of these fluxes does not exist in climate models, making a consistent quantification of their role in the context of the global carbon cycle challenging. The CO<sub>2</sub> increase from 188 to 264 ppm observed during the last deglaciation (Bereiter et al., 2015) results from the combination of mechanisms partly associated with ocean outgassing, following changes in the ocean physics and biogeochemistry and could be directly dependent of the land-sea organic matter fluxes during flooding events.

Previous attempts have tried to explain the changes in atmospheric CO<sub>2</sub> from changes in the physical ocean conditions over the course of the deglaciation using Earth System Models of Intermediate Complexity (EMICs) or paleoproxy records. For example, Schmittner et al. (2007) used a coupled climate-carbon cycle model to show that the oceanic carbon content decreases following the AMOC shutdown, explained by the reduced ocean solubility due to higher Southern Ocean temperature. Menviel et al. (2008) found from glacial water hosing simulations that after a small increase in atmospheric CO<sub>2</sub>, the carbon oceanic sink becomes dominant leading to a decrease of atmospheric CO<sub>2</sub> of ~10 ppm, also consistent with the response of a coupled climate-carbon cycle model to freshwater discharge (Obata, 2007). More recent work from Menviel et al. (2014) combined paleodata and climate simulations from two EMICs (LOVECLIM and UVic ESCM) to show that the ocean can act either as a sink or a source of carbon depending on the bottom water transport in the Pacific Ocean. But in the end, all these models only considered the ocean-atmosphere CO<sub>2</sub> gas exchange and carbon fluxes between the land carbon pool and the atmosphere, ignoring direct carbon fluxes across the land-sea continuum and the associated consequences on the biogeochemistry.

There is a lack of global coupled climate models that consider fluxes of terrestrial organic matter induced by flooding, as well as their consequences on the ocean over long time period such as deglaciations. Only few previous studies conducted time-slice sensitivity experiments with EMICs to quantify the impact of changes in marine physical and biogeochemical conditions on ocean outgassing. Running transient simulations with General Circulation Models remain challenging. We present here for the first time a coupled transient simulation over the last deglaciation using MPI-ESM (Max Planck Institute Earth System Model) that takes into account (1) a fully interactive adaptation of the ocean bathymetry with corresponding changes of the land-sea distribution (Meccia and Mikolajewicz, 2018), (2) a transient river routing (Riddick et al., 2018) and the representation of new processes that are (3) the automatic adjustment of marine biogeochemical tracers under changing ocean bathymetry and land-sea distribution and (4) the fluxes of terrestrial organic matter between land and ocean during flooding events. A special focus is placed on the time period of the large freshwater inputs between 15-14 ka and corresponding to



Meltwater Pulse 1a (MWP1a) to address the role of terrestrial organic matter fluxes on the CO<sub>2</sub> fluxes between the ocean and the atmosphere in context of this millennial event.

## 2 Model description

MPI-ESM is a comprehensive Earth system model composed of four components and one coupler: the atmospheric component ECHAM6.3, the ocean dynamics component MPIOM1.6, the ocean biogeochemistry component HAMOCC6, the land, hydrology and dynamic vegetation component JSBACH3.2 and the coupler OASIS3-MCT. It is currently used in the version 1.2 following the latest updates detailed in Mauritsen et al. (2019). Several components of MPI-ESM have been further extended with new developments to take into account a varying land-sea mask during the period of sea level changes (Meccia and Mikolajewicz, 2018) and the river routing (Riddick et al., 2018) as described in the following sections. In addition, we present here new developments that allow it to take into account the terrestrial organic matter fluxes between land and ocean at a transiently changing land-sea interface as well as their effect on the ocean biogeochemistry.

### 2.1 HAMOCC

The global ocean biogeochemical model HAMOCC (HAMBURG Ocean Carbon Cycle) is part of the ocean component MPIOM (Max Planck Institute Ocean Model) which is used to simulate the oceanic physics by resolving primitive equations under the hydrostatic and Boussinesq approximation on a C-grid with a free surface for every time step (1h) (Jungclaus et al., 2013). HAMOCC simulates the biogeochemical tracers and processes in the oceanic water column, the sediment and at the air-sea interface. HAMOCC has been previously described in Ilyina et al. (2013) and then revised in Mauritsen et al. (2019) with additional processes like marine nitrogen fixation by Cyanobacteria (Paulsen et al., 2017) and nitrogen deposition. Recently, additional ocean tracers have been added in HAMOCC to simulate carbon isotopes (Liu et al., 2021). In this paper we use the HAMOCC6 version as described in Mauritsen et al. (2019) with specific new features for the coupling of the land-sea-continuum that are described in the next sections. The grid configuration of HAMOCC is identical to the ocean component MPIOM and consists of a bipolar grid with one pole over Greenland and another pole over Antarctica. The coarse-resolution ocean grid used for paleoclimate purpose is noted GR30 and has a horizontal resolution of about 300 km. The vertical resolution is composed of 40 unevenly spaced layers with level thickness increasing with depth. Details on the coupling between MPIOM and HAMOCC can be found in Maier-Reimer et al. (2005).

The biogeochemistry of the water column is computed based on the extended NPZD (Nutrients, Phytoplankton, Zooplankton and Detritus) model described in Six and Maier-Reimer (1996) and includes the following pools: phytoplankton, zooplankton, cyanobacteria, dissolved organic matter, particulate organic matter, dissolved inorganic phosphate, dissolved inorganic nitrate, dissolved iron, O<sub>2</sub>, dissolved silicate, opal, calcium carbonate and nitrous oxide (N<sub>2</sub>O). The model solves the ocean carbonate system (mocsy 2.0) for total dissolved inorganic carbon (DIC) and total alkalinity according to the protocol of the Ocean Model Intercomparison Project (Orr and Epitalon, 2015). We also use the total pH scale and equilibrium constants recommended by



Dickson et al. (2007) and Dickson (2010). All tracers are mass conserving and follow the variations of temperature and salinity with respect to hydrodynamical processes. In contrast to Mauritsen et al. (2019), we revised the flux of particulate organic matter. Instead of using a linear increasing sinking speed we use a constant sinking speed of  $5 \text{ m day}^{-1}$  and introduce a temperature dependent remineralisation rate with a  $Q_{10}$  of 2.3. The remineralization of organic matter can be aerobic or anaerobic with denitrification or sulfate reduction when  $\text{O}_2$  concentration passes a critical threshold. This version of HAMOCC also uses an updated procedure to account for weathering fluxes entering the ocean. Instead of a globally uniform distribution, weathering fluxes are only distributed along the coast. The water column interacts with atmospheric components by air-sea exchange of gaseous tracers ( $\text{O}_2$ ,  $\text{N}_2$ ,  $\text{N}_2\text{O}$  and  $\text{CO}_2$ ). All atmospheric concentrations are prescribed. Dust deposition from the atmosphere to the ocean is also taken into account based on the time slice estimates of Albani et al. (2016) (see details in Section 2.4).

The simulation of the oceanic sediment follows the approach of Heinze and Maier-Reimer (1999). The biogeochemical tracers of the sediment include DIC, alkalinity as well as dissolved phosphate, nitrate, silicate, oxygen, iron,  $\text{N}_2$  and  $\text{H}_2\text{S}$  for the pore water fraction and detritus, opal, calcium carbonate and clay for the solid fraction. The processes of decomposition of detritus, dissolution of opal and calcium carbonate and the carbon chemistry are similar to those in the water column. There is vertical diffusion of dissolved tracers in pore water within the sediment column, as well as a diffusive exchange with the water column above. The sediment is resolved by 12 layers with increasing thickness and decreasing porosity from top to bottom, the bottom layer being defined as an underlying burial layer. In the burial, no biogeochemical processes are active and there is no pore water diffusion between burial and active sediment layers. This HAMOCC version also includes erosion under higher bottom shear stress. At most half of the first sediment layer volume of the solid components can be eroded within one time step. Due to remineralization and dissolution of solid components as well as erosion, gaps might be produced within the sediment layers. They are removed by shifting the solid sediment constituents downward within the sediment column. To guarantee that each sediment layer is properly filled with solid components according to solid fraction, the sediment column is shifted upward and filled with sand from the burial layer. Sand is an inert component and can not be eroded (Mathis et al., 2019).

## 2.2 Dynamic land-sea mask and hydrological discharge

In context of ice sheets growth or decay and associated changes in freshwater input and sea level, it is necessary to take into account the variations in ocean bathymetry and coastlines when performing transient simulations. Meccia and Mikolajewicz (2018) describe a new automatic method for interactive bathymetry and land-sea mask changes in MPIOM, allowing simulations for transient time periods. To do so, an ice sheet reconstruction is used to compute the time-dependent freshwater fluxes to the ocean and to derive the topography, which is then used to obtain changes in bathymetry and land-sea mask. Changes in ocean bathymetry and the land-sea mask are updated every 10 years. In the case of ocean expansion, new grid cells are flooded by the surrounding water. For HAMOCC, this also includes pore water of new sediment grid cells. The solid fraction of the new sediment column is filled with sand. This automatic bathymetry adjustment conserves mass (i.e. salt content and all other oceanic tracers) at the global and regional scales (changes in ocean volume match the freshwater input and the global inventory of tracers is constant in the absence of sources or sinks).



Changes in ice sheets volume since the Last Glacial Maximum (LGM) induce changes in the river pathways that need to be taken into account to properly estimate the runoff during the last deglaciation. Riddick et al. (2018) presented a new development within the land surface component JSBACH (Jena Scheme for Biosphere-Atmosphere Coupling in Hamburg) (Reick et al., 2013; Kleinen et al., 2020; Reick et al., 2021) and the Hydrological Discharge model (Hagemann and Dümenil, 1997; Hagemann and Gates, 2001) to automatically generate dynamic river directions and flow parameters for past conditions. First the orography is adjusted and corrected. Then river directions and flow parameters are generated and the location of the river mouths are determined. The ice sheet height and isostatic adjustments are taken from an ice sheet reconstruction and the land-sea mask is generated using the technique described above. These changes in freshwater discharge to the ocean have significant impacts on the global oceanic circulation (via the North Atlantic and Arctic Oceans) and as a consequence also affect atmospheric circulation.

### 2.3 Land-sea carbon and nutrients transfer


In this version of MPI-ESM, we account for the fact that due to increasing sea level (following the melting of the ice sheets) coastal areas are flooded. ECHAM6.3 does not include fractional grid boxes, i.e. a single grid box is treated as land (ocean) as long as the land (ocean) fraction is larger than 50 %. Flooding, i.e. conversion from land to ocean, occurs if the land fraction within one grid cell is less than 50 %. Drying of an ocean grid cell in the case of a decrease of the sea level is also considered. In this case, all pore water tracers, i.e. phosphate, nitrate, dissolved inorganic carbon and alkalinity are redistributed to the water column to guarantee mass conservation and the solid parts of the sediment are considered as inactive land pools.

In the land component JSBACH, each grid box represents the diversity of plant functional types (PFTs) depicting various vegetation types (Reick et al., 2021). 11 PFTs are described in the model: 8 for natural vegetation and 3 for land-use types (which are not used in this deglaciation simulation). The land module also accounts for coupled carbon and hydrological cycles. As part of the atmospheric component, the land component JSBACH uses the same coarse-resolution grid noted T31 with an approximate grid spacing of 400 km (different from the GR30 grid of MPIOM). In the case of flooding, the terrestrial organic matter of the flooded grid cell is collected on the T31 grid and remapped to the GR30 grid. The transfer from land to ocean is mass conserving.

As a consequence of flooding events, the land carbon pools (wood, woody litter above and below ground, humus) are transferred to corresponding oceanic pools in the water column and sediment (Figure 1). So in addition to the 25 prognostic tracers already existing in HAMOCC, we added 3 new prognostic tracers to represent wood, woody litter and humus. In case of flooding, woody litter above ground is distributed uniformly over the water column of the corresponding ocean grid cells. Woody litter below ground and humus are transferred to the sediment where they participate in all sedimentary processes and can be eroded (Figure 1). Wood is located at the water-sediment interface. It is not advected and does not participate in erosion. All terrestrial organic particles in the water column sink with the same speed as marine particulate organic matter. All terrestrial organic matter is remineralized using oxygen in case of aerobic remineralization or  $\text{NO}_3$  and  $\text{NO}_2$  if anaerobic remineralization



takes place (see equations below). It is crucial to note that the stoichiometry of terrestrial organic matter differs from that of marine organic matter defined as C:N:P = 122:16:1 (Takahashi et al., 1985). The carbon to nutrients ratio for woody litter (above and below ground) is defined as C:N:P = 7600:51:1 (Goll et al., 2012). The carbon to nutrients ratio for wood is set to C:N:P = 3650:11:1 (Goll et al., 2012). The carbon to nutrients ratio for humus is C:N:P = 465:10:1 (Goll et al., 2012). Remineralization of terrestrial organic matter leads to changes in dissolved inorganic carbon, phosphate, nitrate, dissolved oxygen and alkalinity in the water column or the sediment. In contrast to the long-living terrestrial material that is transferred to the ocean, short-living material from the vegetation (green biomass, non-woody litter above and below ground) is considered to be remineralized during the flooding event and is treated as a CO<sub>2</sub> source for the atmosphere (Figure 1). However, since we used prescribed CO<sub>2</sub> concentrations in this simulation, the carbon emitted to the atmosphere from short-living land sources has no effect on the climate.




**Figure 1.** Scheme of the pre- and post-flooding environments for terrestrial organic matter.

Assuming that terrestrial carbon compounds that are carbohydrates  $(CH_2O)_x$  give the classical organic matter composition defined as  $(CH_2O)_x(NH_3^+)_yH_3PO_4$ , we use the equations of Paulmier et al. (2009) presented below to calculate the oxygen and nitrate demand for remineralization, as well as alkalinity change. Considering that land biomass contains no excess  $H^+$  compared to organic material in the ocean we obtain:


$$\begin{aligned}
 175 \quad c &= a + 4 && \text{for oxygen} \\
 z &= 0 = b - 2c - d + 5 && \text{for } H^+
 \end{aligned}
 \tag{1}$$



We use the equation for aerobic remineralization with an autotrophic shortcut to nitrate as defined in Paulmier et al. (2009):



The change in alkalinity is given by (-d-1) following Eq. (2). For complete denitrification (with conversion of the ammonium produced by anaerobic remineralization into  $N_2$ ) we use the following equation (Paulmier et al., 2009):



For the  $NO_3^-$  change, the value corresponds to (alkalinity change + 1), i.e.  $(\frac{4}{5} a + \frac{1}{5} b - \frac{2}{5} c + 1)$ . This leads to the overall values presented in Table 1 for the different carbon and nutrients compositions of terrestrial organic matter.

Terrestrial OM	C:N:P	Aerobic remineralization		Anaerobic remineralization	
		O <sub>2</sub> demand	ΔAlk	NO <sub>3</sub> <sup>-</sup> demand	ΔAlk
Wood	3650:11:1	3672	-12	2926.6	2925.6
Woody litter	7600:51:1	7702	-52	6110.6	6109.6
Humus	465:10:1	485	-11	378	377

**Table 1.** Compositions of terrestrial organic matter with consumption of oxygen, nitrate and change in alkalinity during remineralization.

## 2.4 Model and experiment setup

Two spin-up runs (between 26-24 ka and 24-21 ka) were first performed to bring the physical, climatic and biogeochemical systems into quasi-equilibrium. We then ran the deglaciation simulation starting at 21 ka. The initial conditions and forcing are described next. The greenhouse gases concentrations, i.e. CO<sub>2</sub>, CH<sub>4</sub> and NO<sub>2</sub> are from Köhler et al. (2017) and the orbital parameters are taken from Berger and Loutre (1991). The model is also forced with the ice sheet reconstruction GLAC1D (Tarasov et al., 2012) and dust deposition from Albani et al. (2016) which is linearly interpolated over the deglaciation since the data is only available for specific time-slices (21 ka, 16 ka, 10 ka, 8 ka, 6 ka, 4 ka, 2 ka). The initial marine nutrients and carbon inventories are set to similar values to those of the present-day ocean with global mean concentrations of 4.2 mmol N m<sup>-3</sup> for nitrate, 0.52 mmol P m<sup>-3</sup> for phosphate, 9.2 mmol Si m<sup>-3</sup> for silicate and 1915 mmol C m<sup>-3</sup> for DIC. The global





195 mean alkalinity concentration is set higher than present-day with  $2300 \text{ mmol m}^{-3}$  to stabilize the marine chemistry to the lower LGM atmospheric  $\text{CO}_2$  concentration. Weathering fluxes are set to a global constant value for the entire simulation to compensate for burial fluxes, based on the previous spin-up runs. The global weathering rates used to calculate the coastal input fluxes correspond to  $1 \text{ kmol P s}^{-1}$  for organic material,  $1720 \text{ kmol C s}^{-1}$  for inorganic material and  $210 \text{ kmol Si s}^{-1}$  for silicate. The weathering fluxes are independent from the river discharge even if they might locally show small variations because the length of the coastline varies. Moreover, we consider that high latitudes coastlines get only 20 % of the global value to avoid excessive inputs of silicate that would occur if the weathering rates were distributed uniformly over the coastlines (Lacroix et al., 2020). Finally, we assume that sea-ice contains only pure water and thus does not carry any biogeochemical tracer or salt. The land-sea mask, bathymetry, river directions and flow parameters are updated every 10 years.

200 Different sensitivity experiments have also been performed during Meltwater Pulse 1a by branching off new simulations at 15 ka from the deglaciation simulation. These sensitivity experiments ran for 1000 years. One sensitivity experiment has been run without terrestrial organic matter fluxes to the ocean to investigate the changes in sea-air carbon flux. Furthermore, a set of two sensitivity experiments has been run with modified carbon to nutrients ratios for terrestrial organic matter entering the ocean (details in Section 3.3).

### 3 Results and Discussion


#### 210 3.1 Ocean and land responses over the deglaciation

The last deglaciation simulation is characterized by several changes in the oceanic physics and biogeochemistry, as well as in the land carbon content. The first part of the deglaciation, i.e. 21-15 ka (Figure 2), is marked by small changes in the global ocean primary production from  $52.1 \text{ GtC y}^{-1}$  to  $52.6 \text{ GtC y}^{-1}$ , as well as in the ocean physics with a relatively constant AMOC (Atlantic Meridional Overturning Circulation). The LGM AMOC strength is  $22.5 \text{ Sv}$  ( $1 \text{ Sv} = 10^6 \text{ m}^3 \text{ s}^{-1}$ ) which is within the range of a multimodel mean LGM maximum AMOC value of  $23 \pm 3 \text{ Sv}$  (Muglia and Schmittner, 2015). Global net  $\text{CO}_2$  flux between the ocean and the atmosphere is mostly negative, except during small intervals like between 18.75-18.39 ka (Figure 2c). This indicates that the ocean is mainly acting as a sink of carbon in this simulation with prescribed  $\text{CO}_2$  forcing. This general uptaking of  $\text{CO}_2$  by the ocean could also be induced by the oceanic circulation with the Southern Ocean being already well ventilated since the beginning of the deglaciation in our simulation. A rapid change is observed in the ocean between 15-14 ka with the global primary production decreasing by  $4.94 \text{ GtC y}^{-1}$  and the ocean uptaking up to  $0.30 \text{ GtC y}^{-1}$  (Figure 2c-d). This abrupt event is correlated to the large freshwater input of  $0.5 \text{ Sv}$  originating from the North Atlantic and Arctic Oceans and is called Meltwater Pulse 1a. As a result the AMOC strength decreases from  $20$  to  $3 \text{ Sv}$  (Figure 2b). The mean Sea Surface Temperature (SST) decreases by more than  $5 \text{ }^\circ\text{C}$  in the North Atlantic and the mean Sea Surface Salinity (SSS) decreases by around  $3 \text{ psu}$ . Globally, the mean SST and SSS decrease by  $0.53 \text{ }^\circ\text{C}$  and  $0.24 \text{ psu}$ , respectively. This decrease in AMOC streamfunction is also observed when looking at a cross-section in the Atlantic Ocean between the Last Glacial Maximum state and the minimum of the streamfunction at  $14.5 \text{ ka}$ . During the LGM, the maximum of the streamfunction in the Atlantic Ocean is at around  $1200 \text{ m}$  depth at  $30^\circ \text{ N}$  (Figure 3). Between 15-14.5 ka the AMOC strength decreases significantly with only weak






circulation remaining for the top 2000 m, the most important export being in the South Atlantic. After this large meltwater pulse and the AMOC streamfunction minimum at 14.5 ka, the Atlantic circulation takes 500 years to return to its original state.  
 230 This decrease in the AMOC strength during MWP1a is also observed in the  $^{231}\text{Pa}/^{230}\text{Th}$  measured in sediment cores and used to infer past oceanic circulation (McManus et al., 2004).



**Figure 2.** Land-ocean-atmosphere comparison during the last deglaciation. The presented outputs start at 21 ka, after the spin-up runs between 26-21 ka. (a) Freshwater input to the global ocean. (b) Atlantic Meridional Overturning Circulation streamfunction. (c) Global net  $\text{CO}_2$  flux between the ocean and the atmosphere. Positive values mean that the ocean is outgassing to the atmosphere and negative values that the ocean is uptaking carbon. (d) Global ocean primary production. (e) Total carbon in all terrestrial carbon pools, i.e. vegetation, soil and litter. The thick darker curves are 500 years running mean for the freshwater input and 50 years running mean for the other panels. A zoom over Meltwater Pulse 1a is presented on the right.




**Figure 3.** Cross-section of the AMOC streamfunction for the Last Glacial Maximum at 21 ka (a) and during MWP1a at 14.5 ka (b).



For the land part, the total carbon in all terrestrial carbon pools (litter, soil and vegetation) increases from 922.9 GtC to 1302.7 GtC between 21-15 ka and reaches 1563.6 GtC at 12 ka (Figure 2e). This increase is explained by the progressively warmer climate during the deglaciation and by the increase of C<sub>3</sub> plants and Gross Primary Productivity (GPP) on land. Even if our simulation currently doesn't continue beyond 12 ka, the total land carbon content evolution since the Last Glacial Maximum is within the range of other modelling studies like Prentice et al. (2011), O'ishi and Abe-Ouchi (2013), and Ganopolski and Brovkin (2017). Furthermore, in a setup with a fully coupled land and ocean carbon cycle it would be the ocean that supplies this additional land carbon, which would demand a constant outgassing over the period during which the land carbon inventory increases. During MWP1a, the C<sub>3</sub> cover fraction and the land GPP decrease, indicating the replacement of trees by grasslands and resulting in a decrease of 62.38 GtC stored in litter and vegetation pools as shown in Fig. 2e.

We can also evaluate the ability of the model to reproduce the biome distribution before the beginning of MWP1a at 15 ka in comparison to pollen data like the BIOME6000 Version 4.2 reconstruction (Harrison, 2017) based on the Palaeovegetation Mapping Project (Prentice and Webb III, 1998; Prentice et al., 2000; Harrison et al., 2001; Bigelow et al., 2003; Pickett et al., 2004). To do so we used the biomisation technique presented in Dallmeyer et al. (2019) to convert the different PFT cover fractions modelled in JSBACH into 9 biomes. Although the LGM conditions were different from those at 15 ka, in absence of other reconstructions, the BIOME6000 pollen record shows a good overall agreement with the modelled biome reconstruction (Figure 4). At high latitudes of the Northern Hemisphere, both tundra and boreal forests are modelled for the areas that are not covered by ice, which is consistent with the few pollen data sets available at these locations. Temperate forest is modelled over part of North America, grassland over Europe and temperate/warm forest over East Asia. This is generally in agreement with the pollen record even if some local discrepancies are observed like in central Asia. At low latitudes the model mostly reproduces the tropical forest (over Eastern South America, West Africa and Indonesia) as observed in the pollen data (Figure 4). The Indonesian region is compared in more detail with other studies since most of the terrestrial organic matter inputs during MWP1a comes from this region (see next sections). Dubois et al. (2014) used isotopic composition of vascular plant fatty acid ( $\delta^{13}C_{FA}$ ) from surface sediments in Indonesian seas to infer past regional vegetation over Indonesia during the LGM. They showed that during this time period the predominant vegetation was characterized by C<sub>3</sub> plants over central Indonesia. Indeed, even if the climate was colder and drier than during the Preindustrial, it was not sufficiently so to alter the vegetation distribution and decrease the rainforest coverage. Our biome reconstruction also agrees with modelling studies like the one of Cannon et al. (2009) or Prentice et al. (2011) which shows using a dynamic vegetation model that tropical forest dominated the Sunda shelf during the Last Glacial Maximum. Recently, Dallmeyer et al. (2019) also evaluated the vegetation reconstruction from 4 different Earth System Models using the same harmonization method for PFT distribution that we used for Fig. 4. All ESMs used in Dallmeyer et al. (2019) model tropical forest over Indonesia during the LGM. All together, these results support the conclusion that tropical forests developed following the LGM continued to do so until at least 15 ka, even if the climate was colder and drier than the Preindustrial.




**Figure 4.** Biome distribution modelled by JSBACH at 15 ka. The superimposed circles are the pollen data from the BIOME6000 Version 4.2 reconstruction at 21 ka (Harrison, 2017).

### 265 3.2 Land-sea carbon fluxes


The transient adaptation of the land-sea mask and bathymetry during the deglaciation allows for flooded coastal land areas to be accounted for when the sea level changes. During the deglaciation, before and after Meltwater Pulse 1a (i.e. 21-15 ka and 14-12 ka), flooding is observed on almost all continental coasts but with a higher number of flooded land coastal regions located above 60° N (Figure 5). During MWP1a, local coastal land surfaces are flooded in East Asia, Indonesia and Australia, but again with a higher number of the flooded grid cells in Northern Europe since the major source of the meltwater is in the Northern Hemisphere (Figure 5). In terms of area, the flooded high latitudes coastal regions represent  $1.03 \times 10^6 \text{ km}^2$ , which is larger than the  $8.04 \times 10^5 \text{ km}^2$  of flooded coastal area in Indonesia.

Organic matter fluxes from land to ocean are computed during the different flooding events and the results obtained from the transient simulation show that terrestrial inputs are frequent over the last deglaciation but most of them happen after 15 ka (Figure 6). There are also differences in the size of the carbon fluxes depending on the origin of the terrestrial organic matter. During MWP1a, 7.7 GtC are emitted to the atmosphere (related to the green vegetation), which corresponds to 26.6 % of the total terrestrial organic matter inputs (Table 2). We however refrain from any discussion since these fluxes don't have an impact on the climate in this simulation with prescribed  $\text{CO}_2$  concentrations. 21.2 GtC goes to the ocean, i.e. 73.4 % of the total terrestrial organic matter inputs (Table 2).



**Figure 5.** Spatial distribution of the coastal flooded areas during the last deglaciation (dark blue) and during the specific interval of MWP1a between 15-14 ka (light blue). Zooms over the Arctic and Indonesian regions are also shown.

Looking at the origin of the terrestrial organic matter inputs to the ocean, wood and humus are the largest contributors with 31.1 and 29.1 % of the total inputs respectively (atmosphere and ocean) and have even higher contribution of 42.5 and 39.6 % for the ocean part only. In comparison to the entire deglaciation between 21-12 ka, the terrestrial organic matter inputs to the atmosphere and to the ocean represent a similar proportion than during MWP1a with respectively 28.4 and 71.6 % for a total of 88.3 GtC (Table 2). Wood and humus are again the major contributors of these land-sea fluxes. We have to emphasize that these numbers are for one millennium and thus the total terrestrial organic matter contribution to the ocean of 21.2 GtC is non negligible compared to the entire deglaciation (63.2 GtC). However, in comparison to the ocean inventory of the Mixed Layer Depth of around 600 GtC or even of the global ocean of around 36000 GtC, we can consider that the terrestrial organic matter fluxes to the ocean are rather small (3.5 % of the Mixed Layer Depth inventory and 0.06 % of the global ocean inventory). Their effects on the ocean biogeochemistry are investigated next in Section 3.3.



**Figure 6.** Time series of terrestrial organic matter inputs during the last deglaciation. Top panel: Terrestrial organic matter (vegetation, green biomass and non-woody litter) inputs to the atmosphere. Middle panel: Terrestrial organic matter (wood and woody litter above ground) inputs to the water column and to the water-sediment interface. Bottom panel: Terrestrial organic matter (woody litter below ground and humus) inputs to the sediment.


	Green biomass and non-woody litter	Wood	Woody litter above ground	Woody litter below ground	Humus	Total
Amount in GtC between 21-12 ka	25.1 (28.4 %)	26.2 (29.7 %)	7.3 (8.3 %)	3.2 (3.6 %)	26.5 (30 %)	88.3 (100 %)
Amount in GtC between 15-14 ka	7.7 (26.6 %)	9.0 (31.1 %)	2.6 (9 %)	1.2 (4.2 %)	8.4 (29.1 %)	28.9 (100 %)
Destination	Atmosphere	Ocean-sediment interface	Water column	Sediment	Sediment	Atmosphere and ocean

**Table 2.** Carbon mass of terrestrial organic matter going to the atmosphere, ocean-sediment interface, water column and sediment after the flooding of the coastal land areas during the last deglaciation and MWP1a.



Among all contributors of terrestrial organic matter during MWP1a, the Indonesian region is the largest one with inputs to the water column that represent 66.4 % of the total inputs (Figure 7a) and explained by wood inputs. The second largest contributor during this meltwater pulse is Australia with 17.3 %, followed by East Asia, the high latitudes of the Northern Hemisphere, South America and Europe with respectively 11.1, 2.2, 1.8 and 1.2 %. All these terrestrial inputs are dominated by wood inputs.

295 As for the part that goes to the sediment, Indonesia and East Asia regions are also major contributors with 43.7 and 18.7 % of the total terrestrial inputs to the sediment during MWP1a (Figure 7b), mostly due to humus. Other regions subject to flooding like Australia, the high latitudes of the Northern Hemisphere and Europe also contribute to terrestrial organic matter inputs to the sediment in smaller amounts. So even if most of the flooded areas are located in the Northern Hemisphere during MWP1a, their contributions of terrestrial inputs to the ocean are less significant than equatorial and low latitudes flooded areas.



**Figure 7.** Contributions of flooded coastal areas to the total terrestrial organic matter inputs (considered as 100 %) that go to the water column (a) and to the sediment (b) during the full deglaciation between 21-12 ka (full bars on the left) and during MWP1a between 15-14 ka (hatched bars on the right).

300 These results are however only representative for a short time period relative to the rapid change induced by the meltwater pulse and could be different from the rest of the deglaciation. To investigate this, we also present the contribution of these same areas during the full deglaciation (21-12 ka). In comparison to the specific case of MWP1a, Europe, South America, East Asia and the high latitudes of the Northern Hemisphere have similar contributions of terrestrial carbon and nutrients to the water column over the entire deglaciation (Figure 7a). In contrast, North America (East and West) and land areas in the

305 high latitudes of the Southern Hemisphere show organic matter inputs of a few percent during the deglaciation, but not during the large meltwater pulse. But the largest contribution of terrestrial organic matter to the water column observed during the deglaciation, i.e. Indonesia, occurs during the large meltwater pulse. In terms of regional contribution to the total terrestrial



inputs to the sediment, Indonesia is once again the largest contributor with 43.7 % (Figure 7b). East Asia contributes with 18.7 %, similar to Australia, but three times more important than Europe and the high latitudes of the Northern Hemisphere with 6.5 %. Among the smallest contributors are North and South America between 1 and 3 %. Once in the sediment, part of the terrestrial organic matter is buried, i.e. 0.042 and 0.168 GtC respectively for the woody litter and humus during the entire deglaciation. Inputs to the water column due to sediment erosion are of the order of 0.0026 and 0.026 GtC for the woody litter and humus respectively. The remaining part is remineralized.

### 3.3 Implications for the ocean biogeochemical cycle

To assess the role of the terrestrial organic matter inputs on ocean biogeochemistry during MWP1a, we performed a sensitivity experiment without terrestrial organic matter fluxes between land and ocean (as described in Section 2.4). The sensitivity experiment is branched off at 15 ka from the reference deglaciation run. Figure 8 shows anomalies of surface alkalinity, dissolved inorganic carbon and phosphate between simulations with and without terrestrial organic matter inputs (i.e. deglaciation reference run - sensitivity experiment) as average over MWP1a.


The surface alkalinity anomaly shows regionally higher values of around 10 mmol m<sup>-3</sup> in central Indonesia, East Asia and up to 22 mmol m<sup>-3</sup> at high latitudes of the Northern Hemisphere where coastal areas are flooded (Figure 8a). The largest difference between the two simulations is observed between Indonesia and Australia with values from 148 to 214 mmol m<sup>-3</sup>. In contrast, lower values are observed in the Atlantic Ocean, in the West Equatorial Pacific and in the Arctic Ocean. A similar pattern is observed for the dissolved inorganic carbon with higher values between 0 and 30 mmol C m<sup>-3</sup> in Indonesia, East Asia and high latitudes of the Northern Hemisphere. Several grid points from North to South Indonesia show higher values between the two simulations with 36 mmol C m<sup>-3</sup> (Northern Indonesia), 90 mmol C m<sup>-3</sup> (central Indonesia) and 228 mmol C m<sup>-3</sup> (Southern Indonesia) (Figure 8b). These positive anomalies (detailed in Fig. 9a for alkalinity and Fig. 9b for DIC) are associated with the input of terrestrial organic matter, mainly wood and humus, during a flooding event with additional carbon and nutrients entering the ocean leading to an increase of the surface alkalinity and DIC once the terrestrial organic matter has been remineralized. At depth, an increase in DIC is observed below 1500 m in the Atlantic Ocean, Nordic Seas, Australia-Indonesia Coastal Province and Sunda-Arafura Shelves Province. For the surface phosphate anomaly, small differences are observed in the Atlantic part of the Southern Ocean and in the equatorial band with values lower by up to 0.02 mmol P m<sup>-3</sup> (Figure 8c). Higher values up to 0.05 mmol P m<sup>-3</sup> are observed in West Indonesia, in the North Atlantic Ocean and at high latitudes of the Northern Hemisphere.

The temporal evolution of the ocean biogeochemistry is also quite similar between the simulations with and without terrestrial organic matter inputs to the ocean. The global surface alkalinity shows similar variations with values ranging from 2247 to 2282 mmol m<sup>-3</sup> between 15-14 ka (Figure 8d). The largest difference observed between the two simulations occurs between 14.69-14.62 ka with the simulation with terrestrial organic matter fluxes being around 4 mmol m<sup>-3</sup> higher than the simulation without these fluxes. This is explained by land-sea fluxes with a total of 3.08 GtC. The global surface DIC and phosphate variations during MWP1a show a similar trend within the two simulations with values ranging respectively from 1916 to 1950






mmol C m<sup>-3</sup> (Figure 8e) and from 0.44 to 0.54 mmol P m<sup>-3</sup> (Figure 8f). The differences in the biogeochemistry within the two simulations highlight the fact that terrestrial organic matter inputs to the ocean only have a relatively small effect of 1-2 % on the global surface alkalinity, dissolved inorganic carbon and phosphate. The land-sea fluxes are indeed relatively small (3.5 % in comparison to the Mixed Layer Depth carbon inventory (upper 200 m of the water column)).




**Figure 8.** Anomaly of the mean surface alkalinity (a), surface dissolved inorganic carbon (b), and surface phosphate (c) between the two simulations with and without terrestrial organic matter fluxes averaged over MWP1a. Time evolution of the two simulations during MWP1a for annual mean global surface alkalinity (d), DIC (e) and phosphate (f).



**Figure 9.** Same as Fig. 8a, b, c but with a focus over Indonesia for the surface alkalinity (a), surface dissolved inorganic carbon (b) and surface phosphate (c) between the two simulations with and without terrestrial organic matter fluxes averaged over MWP1a.

Due to the stoichiometry of the terrestrial organic matter being higher than of the marine organic matter (C:N:P = 3650:11:1 for the wood, 7600:51:1 for the woody litter and 465:10:1 for the humus in comparison to 122:16:1 for marine organic matter) there is an excess of carbon in the ocean once the terrestrial organic matter has been remineralized. Indeed, net primary production uptakes only 122 moles of carbon per mole of phosphate which leaves a large fraction of remineralized terrestrial carbon in the water column. This leads the ocean to outgas  $\text{CO}_2$  to the atmosphere. Similarly to what has been presented above, we investigate the effect of terrestrial organic matter inputs on the  $\text{CO}_2$  fluxes between ocean and atmosphere during MWP1a. Figure 10 shows the anomaly between the two simulations (with and without land-sea fluxes) of the mean surface  $\text{CO}_2$  flux between the ocean and atmosphere. Indonesia shows differences with higher  $\text{CO}_2$  outgassing values of up to  $2.2 \times 10^{-9} \text{ kg C m}^{-2} \text{ s}^{-1}$  for the simulation taking into account the land-sea fluxes. These larger values originate from the remineralization of the wood and humus associated with tropical forest developed at that time, as explained in Section 3.1 and shown in Fig. 4. Other flooded areas during MWP1a also contribute to higher  $\text{CO}_2$  fluxes to the atmosphere for the simulation with terrestrial organic matter fluxes, like in East Asia with  $0.17\text{-}0.22 \times 10^{-9} \text{ kg C m}^{-2} \text{ s}^{-1}$  or in the Northern Australian coast with  $0.36\text{-}0.39 \times 10^{-9} \text{ kg C m}^{-2} \text{ s}^{-1}$ . Overall, the terrestrial organic matter fluxes only have a local influence on the ocean behaviour with higher outgassing in the Indonesian region during the millennial event of MWP1a.



**Figure 10.** Anomaly of the mean surface CO<sub>2</sub> flux between the simulations with and without terrestrial organic matter inputs to the ocean averaged over MWP1a. Negative values indicate flux from the atmosphere to the ocean (uptaking) and positive values indicate flux from the ocean to the atmosphere (outgassing). The black rectangle highlights the area with the largest differences.

360 To investigate the influence of the carbon to nutrients ratio of terrestrial organic matter on the CO<sub>2</sub> fluxes between the ocean and atmosphere during MWP1a we performed two additional sensitivity experiments following the procedure described in Section 2.4. The first sensitivity experiment uses a higher C:N:P ratio of terrestrial organic matter, arbitrarily defined as twice the carbon and nitrogen values of Goll et al. (2012) used in the reference simulation since no higher estimations exist in the literature. The second sensitivity experiment uses a lower C:N:P ratio of terrestrial organic matter based on studies from Lerman et al. (2004) and Cleveland and Liptzin (2007). These changes in the stoichiometry also require an adjustment of the O<sub>2</sub> and NO<sub>3</sub><sup>-</sup> demand and corresponding ΔAlk changes during remineralization of terrestrial organic matter. All values are summarized in Table 3.


		Aerobic remineralization		Anaerobic remineralization		
	Terrestrial OM	C:N:P	O <sub>2</sub> demand	ΔAlk	NO <sub>3</sub> <sup>-</sup> demand	ΔAlk
High stoichiometry sensitivity test	Wood	7300:22:1	7344	-23	5853.2	5852.2
	Woody litter	15200:102:1	15404	-103	12221.2	12220.2
	Humus	930:20:1	970	-21	756	755
Low stoichiometry sensitivity test	Wood	510:4:1	518	-5	410.4	409.4
	Woody litter	186:13:1	212	-14	156.6	155.6
	Humus	140:6.6:1	153.2	-7.6	115.96	114.96

**Table 3.** Terrestrial organic matter stoichiometry with consumption of oxygen, nitrate and change in alkalinity during remineralization for the sensitivity experiments with high and low stoichiometry. For comparison, values for the reference deglaciation simulation are given in Table 1.



We compare the largest outgassing events over Indonesia for the different simulations with and without terrestrial organic matter as well as for the simulations with different carbon to nutrients ratios for this organic matter. As already observed in Fig. 10, the outgassing of the ocean to the atmosphere from North to South Indonesia during MWP1a is driven by the terrestrial organic matter fluxes during flooding events. We observe a CO<sub>2</sub> flux to the atmosphere happening concurrently to the flooding events and then slowly decreasing with time until all the terrestrial organic matter has been remineralized. In Northern Indonesia (12.5° N - 96.5° E), the outgassing starts at 14.64 ka, reaches 3.8 x 10<sup>-9</sup> kg C m<sup>-2</sup> s<sup>-1</sup> and then decreases for 200 years which corresponds to the decay time of the wood material (i.e. stems) in the ocean (Figure 11a). This outgassing event is due to the transfer during the flooding event of terrestrial material to the ocean, dominated by the wood with 1.72 GtC and by the humus with 0.79 GtC. For the simulation without terrestrial organic matter inputs, the ocean behaves differently and rather uptakes carbon with a slightly negative CO<sub>2</sub> flux. This opposite behaviour between the two simulations with and without terrestrial organic matter fluxes is also observed in central and Southern Indonesia and highlights the key role of terrestrial organic matter fluxes in the oceanic outgassing. In central Indonesia (4.5° N - 107.5° E) we observe an outgassing peak of 8.8 x 10<sup>-9</sup> kg C m<sup>-2</sup> s<sup>-1</sup> at 14.54 ka and then a positive CO<sub>2</sub> flux for 300 years (Figure 11b). This outgassing event is primarily the consequence of wood input to the ocean with 1.95 GtC, as well as humus input, the second largest contributor, with 0.69 GtC. In Southern Indonesia (6.5° S - 134.5° E) we also have an outgassing with 5.3 x 10<sup>-9</sup> kg C m<sup>-2</sup> s<sup>-1</sup> to the atmosphere at 14.18 ka due to dominant wood inputs of 2.15 GtC (humus inputs represent 0.67 GtC) (Figure 11c).

For a higher carbon to nutrients ratio the three locations in Indonesia reproduce similar CO<sub>2</sub> fluxes to the atmosphere than those in the reference simulation. For a lower carbon to nutrients ratio, the CO<sub>2</sub> flux is smaller than the reference simulation with only slightly positive values following the flooding event, with maximum of 0.085 x 10<sup>-9</sup> kg C m<sup>-2</sup> s<sup>-1</sup> for Northern Indonesia, 0.67 x 10<sup>-9</sup> kg C m<sup>-2</sup> s<sup>-1</sup> for central Indonesia and 0.14 x 10<sup>-9</sup> kg C m<sup>-2</sup> s<sup>-1</sup> for Southern Indonesia (Figure 11). These sensitivity experiments highlight the fact that even with a very high carbon to nutrients ratio (so larger fraction of remineralized terrestrial carbon in the water column), the outgassing CO<sub>2</sub> flux doesn't increase. However, in the case of the lower carbon to nutrients ratio for terrestrial organic matter, the CO<sub>2</sub> flux to the atmosphere is greatly reduced. But again, these fluxes are only happening at regional hotspots and do not affect the global net CO<sub>2</sub> flux between the ocean and the atmosphere.



**Figure 11.** Evolution of the surface CO<sub>2</sub> flux during MWP1a for flooded grid cells in Northern (a), central (b) and Southern Indonesia (c) for 4 simulations: the reference simulation with the terrestrial organic matter fluxes (dark green), the sensitivity experiment without terrestrial organic matter fluxes (light green), the sensitivity experiment with low stoichiometry for terrestrial organic matter (dark blue) and the sensitivity experiment with high stoichiometry (light blue). 50 years running mean are plotted for each simulation. Positive values indicate an outgassing to the atmosphere and negative values indicate an uptake by the ocean.

## 4 Conclusions

In this study we present for the first time the implementation of terrestrial organic matter fluxes between land and ocean at transiently changing land-sea interface in the global Earth System Model MPI-ESM. This unique setup of MPI-ESM was used to perform a transient deglaciation simulation from 21 to 12 ka accounting for sea level rise induced by meltwater inputs from ice sheets and consequential changes in ocean depth and coast lines. The period between 15-14 ka, which corresponds to Meltwater Pulse 1a, has been highlighted because it is characterized by larger terrestrial organic matter inputs to the ocean than during the first part of the deglaciation. Indeed, a total of 21.2 GtC, mostly arising from Indonesia, goes to the ocean during this millennial event, which represents 34 % of the total amount of terrestrial organic matter entering the ocean over the last deglaciation. This terrestrial carbon is remineralized once in the ocean within a time frame of hundreds of years. The effect of this supplementary carbon brought from land is only observed at regional hotspots with local outgassing to the atmosphere. The carbon input doesn't seem to be large enough to impact the global behaviour of the ocean, considering that it represents



a very small amount in comparison to the global ocean inventory (0.06 %). A sensitivity experiment also emphasizes that the terrestrial organic matter fluxes only have a small effect on the surface alkalinity and dissolved inorganic carbon (around 1-2 % increase). However, the local CO<sub>2</sub> fluxes between the ocean and the atmosphere during MWP1a are driven by the terrestrial organic matter inputs. This regional outgassing to the atmosphere observed in Indonesia is explained by wood inputs and is supported by several lines of evidences suggesting that prior to this meltwater pulse event and even as far back as the Last Glacial Maximum, tropical forest was developed in this region favouring the storage of carbon-rich material on land entering the ocean once the land is flooded. As a complement, an additional set of sensitivity experiments show that the magnitude of outgassing during MWP1a is rather insensitive to higher carbon to nutrients ratio of the terrestrial organic matter. Overall, our simulation is a first step towards a fully coupled ESM including carbon and nutrients fluxes at the land-sea continuum that will be applicable for long transient paleo-simulations over the last glacial/interglacial cycles.

*Author contributions.* TE designed the study and wrote the manuscript. KDS and HP realized the model development. KDS conducted the deglaciation simulation and TE performed the sensitivity experiments. All co-authors contributed to the manuscript with valuable comments and discussions.

415 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We acknowledge support through the project PalMod, funded by the German Federal Ministry of Education and Research (BMBF). Simulations were performed at the German Climate Computing Center (DKRZ). We thank Thomas Riddick for internal review of this manuscript. We also thank Anne Dallmeyer for her help with the biome reconstruction.

420 *Financial support.* This research has been supported by the German Federal Ministry of Education and Research (PalMod initiative, FKZ: grant no. 01LP1925C). The article processing charges for this publication are covered by the Max Planck Society.



## References

- Albani, S., Mahowald, N., Murphy, L., Raiswell, R., Moore, J., Anderson, R., McGee, D., Bradtmiller, L., Delmonte, B., Hesse, P., et al.: Paleodust variability since the Last Glacial Maximum and implications for iron inputs to the ocean, *Geophys. Res. Lett.*, 43, 3944–3954, <https://doi.org/10.1002/2016GL067911>, 2016.
- 425 Barker, S., Diz, P., Vautravers, M. J., Pike, J., Knorr, G., Hall, I. R., and Broecker, W. S.: Interhemispheric Atlantic seesaw response during the last deglaciation, *Nature*, 457, 1097–1102, <https://doi.org/10.1038/nature07770>, 2009.
- Bereiter, B., Eggleston, S., Schmitt, J., Nehrbass-Ahles, C., Stocker, T. F., Fischer, H., Kipfstuhl, S., and Chappellaz, J.: Revision of the EPICA Dome C CO<sub>2</sub> record from 800 to 600 kyr before present, *Geophys. Res. Lett.*, 42, 542–549, <https://doi.org/10.1002/2014GL061957>, 2015.
- 430 Berger, A. and Loutre, M.-F.: Insolation values for the climate of the last 10 million years, *Quaternary Sci. Rev.*, 10, 297–317, [https://doi.org/10.1016/0277-3791\(91\)90033-Q](https://doi.org/10.1016/0277-3791(91)90033-Q), 1991.
- Bigelow, N. H., Brubaker, L. B., Edwards, M. E., Harrison, S. P., Prentice, I. C., Anderson, P. M., Andreev, A. A., Bartlein, P. J., Christensen, T. R., Cramer, W., et al.: Climate change and Arctic ecosystems: 1. Vegetation changes north of 55° N between the last glacial maximum, mid-Holocene, and present, *J. Geophys. Res.-Atmos.*, 108, <https://doi.org/10.1029/2002JD002558>, 2003.
- 435 Bonelli, S., Charbit, S., Kageyama, M., Woillez, M.-N., Ramstein, G., Dumas, C., and Quiquet, A.: Investigating the evolution of major Northern Hemisphere ice sheets during the last glacial-interglacial cycle., *Clim. Past*, 5, <https://doi.org/10.5194/cp-5-329-2009>, 2009.
- Cannon, C. H., Morley, R. J., and Bush, A. B.: The current refugial rainforests of Sundaland are unrepresentative of their biogeographic past and highly vulnerable to disturbance, *P. Natl. Acad. Sci. USA*, 106, 11 188–11 193, <https://doi.org/10.1073/pnas.0809865106>, 2009.
- Clark, P. U., Shakun, J. D., Baker, P. A., Bartlein, P. J., Brewer, S., Brook, E., Carlson, A. E., Cheng, H., Kaufman, D. S., Liu, Z., et al.: Global  
440 climate evolution during the last deglaciation, *P. Natl. Acad. Sci. USA*, 109, E1134–E1142, <https://doi.org/10.1073/pnas.1116619109>, 2012.
- Cleveland, C. C. and Liptzin, D.: C: N: P stoichiometry in soil: is there a “Redfield ratio” for the microbial biomass?, *Biogeochemistry*, 85, 235–252, <https://doi.org/10.1007/s10533-007-9132-0>, 2007.
- Dallmeyer, A., Claussen, M., and Brovkin, V.: Harmonising plant functional type distributions for evaluating Earth System Models, *Clim. Past*, 15, 335–366, <https://doi.org/10.5194/cp-15-335-2019>, 2019.
- 445 Denton, G. H., Anderson, R. F., Toggweiler, J., Edwards, R., Schaefer, J., and Putnam, A.: The last glacial termination, *Science*, 328, 1652–1656, <https://doi.org/10.1126/science.1184119>, 2010.
- Dickson, A. G.: The carbon dioxide system in seawater: equilibrium chemistry and measurements, *Guide to best practices for ocean acidification research and data reporting*, 1, 17–40, 2010.
- 450 Dickson, A. G., Sabine, C. L., and Christian, J. R.: *Guide to best practices for ocean CO<sub>2</sub> measurements.*, PICES Special Publication 3, p. 191, 2007.
- Dubois, N., Oppo, D. W., Galy, V. V., Mohtadi, M., Van Der Kaars, S., Tierney, J. E., Rosenthal, Y., Eglinton, T. I., Lückge, A., and Linsley, B. K.: Indonesian vegetation response to changes in rainfall seasonality over the past 25,000 years, *Nat. Geosci.*, 7, 513–517, <https://doi.org/10.1038/ngeo2182>, 2014.
- 455 Ganopolski, A. and Brovkin, V.: Simulation of climate, ice sheets and CO<sub>2</sub> evolution during the last four glacial cycles with an Earth system model of intermediate complexity, *Clim. Past*, 13, 1695–1716, <https://doi.org/10.5194/cp-13-1695-2017>, 2017.





- Goll, D. S., Brovkin, V., Parida, B., Reick, C. H., Kattge, J., Reich, P. B., Van Bodegom, P., and Niinemets, Ü.: Nutrient limitation reduces land carbon uptake in simulations with a model of combined carbon, nitrogen and phosphorus cycling, *Biogeosciences*, 9, 3547–3569, <https://doi.org/10.5194/bg-9-3547-2012>, 2012.
- 460 Hagemann, S. and Dümenil, L.: A parametrization of the lateral waterflow for the global scale, *Clim. Dynam.*, 14, 17–31, <https://doi.org/10.1007/s003820050205>, 1997.
- Hagemann, S. and Gates, L. D.: Validation of the hydrological cycle of ECMWF and NCEP reanalyses using the MPI hydrological discharge model, *J. Geophys. Res.-Atmos.*, 106, 1503–1510, <https://doi.org/10.1029/2000JD900568>, 2001.
- Harrison, S.: BIOME 6000 DB classified plotfile version 1, available at: <http://researchdata.reading.ac.uk/99/>, 2017.
- 465 Harrison, S., Yu, G., Takahara, H., and Prentice, I.: Diversity of temperate plants in east Asia, *Nature*, 413, 129–130, <https://doi.org/10.1038/35093166>, 2001.
- Heinemann, M., Timmermann, A., Elison Timm, O., Saito, F., and Abe-Ouchi, A.: Deglacial ice sheet meltdown: orbital pacemaking and CO<sub>2</sub> effects., *Clim. Past*, 10, <https://doi.org/10.5194/cp-10-1567-2014>, 2014.
- Heinze, C. and Maier-Reimer, E.: The Hamburg oceanic carbon cycle circulation model version "HAMOCC2s" for long time integrations, 470 DKRZ, [https://doi.org/10.2312/WDCC/DKRZ\\_Report\\_No20](https://doi.org/10.2312/WDCC/DKRZ_Report_No20), 1999.
- Ilyina, T., Six, K. D., Segschneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: Global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth system model in different CMIP5 experimental realizations, *J. Adv. Model. Earth Sys.*, 5, 287–315, <https://doi.org/10.1029/2012MS000178>, 2013.
- Ivanovic, R., Gregoire, L., Kageyama, M., Roche, D., Valdes, P., Burke, A., Drummond, R., Peltier, W. R., and Tarasov, L.: Transient climate 475 simulations of the deglaciation 21-9 thousand years before present (version 1)-PMIP4 Core experiment design and boundary conditions, *Geosci. Model Dev.*, <https://doi.org/10.5194/gmd-9-2563-2016>, 2016.
- Jungclauss, J., Fischer, N., Haak, H., Lohmann, K., Marotzke, J., Matei, D., Mikolajewicz, U., Notz, D., and Von Storch, J.: Characteristics of the ocean simulations in the Max Planck Institute Ocean Model (MPIOM) the ocean component of the MPI-Earth system model, *J. Adv. Model. Earth Sys.*, 5, 422–446, <https://doi.org/10.1002/jame.20023>, 2013.
- 480 Kleinen, T., Mikolajewicz, U., and Brovkin, V.: Terrestrial methane emissions from the Last Glacial Maximum to the preindustrial period, *Clim. Past*, 16, 575–595, <https://doi.org/10.5194/cp-16-575-2020>, 2020.
- Klockmann, M., Mikolajewicz, U., and Marotzke, J.: The effect of greenhouse gas concentrations and ice sheets on the glacial AMOC in a coupled climate model, *Clim. Past*, 12, 1829–1846, <https://doi.org/10.5194/cp-12-1829-2016>, 2016.
- Kohfeld, K. E. and Ridgwell, A.: Glacial-interglacial variability in atmospheric CO<sub>2</sub>, Surface Ocean-Lower Atmosphere Processes, Geo- 485 physical Research Series 37, Washington DC, American Geophysical Union, <https://doi.org/10.1029/2008GM000845>, 2009.
- Köhler, P., Nehrbass-Ahles, C., Schmitt, J., Stocker, T. F., and Fischer, H.: A 156 kyr smoothed history of the atmospheric greenhouse gases CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O and their radiative forcing, *Earth Syst. Sci. Data*, 9, 363–387, <https://doi.org/10.5194/essd-9-363-2017>, 2017.
- Lacroix, F., Ilyina, T., and Hartmann, J.: Oceanic CO<sub>2</sub> outgassing and biological production hotspots induced by pre-industrial river loads of nutrients and carbon in a global modeling approach, *Biogeosciences*, 17, 55–88, <https://doi.org/10.5194/bg-17-55-2020>, 2020.
- 490 Lerman, A., Mackenzie, F. T., and Ver, L. M.: Coupling of the perturbed C–N–P cycles in industrial time, *Aquat. Geochem*, 10, 3–32, <https://doi.org/10.1023/B:AQUA.0000038955.73048.c1>, 2004.
- Liu, B., Six, K. D., and Ilyina, T.: Incorporating the stable carbon isotope <sup>13</sup>C in the ocean biogeochemical component of the Max Planck Institute Earth System Model, *Biogeosciences*, 18, 4389–4429, <https://doi.org/10.5194/bg-18-4389-2021>, 2021.



- 495 Liu, Z., Otto-Bliesner, B., He, F., Brady, E., Tomas, R., Clark, P., Carlson, A., Lynch-Stieglitz, J., Curry, W., Brook, E.,  
et al.: Transient simulation of last deglaciation with a new mechanism for Bølling-Allerød warming, *Science*, 325, 310–314,  
<https://doi.org/10.1126/science.1171041>, 2009.
- Maier-Reimer, E., Kriest, I., Segschneider, J., and Wetzol, P.: The hamburg ocean carbon cycle model hamoc5. 1-technical description  
release 1.1, 2005.
- Marcott, S. A., Bauska, T. K., Buizert, C., Steig, E. J., Rosen, J. L., Cuffey, K. M., Fudge, T., Severinghaus, J. P., Ahn, J.,  
500 Kalk, M. L., et al.: Centennial-scale changes in the global carbon cycle during the last deglaciation, *Nature*, 514, 616–619,  
<https://doi.org/10.1038/nature13799>, 2014.
- Mathis, M., Elizalde, A., and Mikolajewicz, U.: The future regime of Atlantic nutrient supply to the Northwest European Shelf, *J. Marine  
Syst.*, 189, 98–115, <https://doi.org/10.1016/j.jmarsys.2018.10.002>, 2019.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., et al.: Develop-  
505 ments in the MPI-M Earth System Model version 1.2 (MPI-ESM1. 2) and its response to increasing CO<sub>2</sub>, *J. Adv. Model. Earth Sys.*, 11,  
998–1038, <https://doi.org/10.1029/2018MS001400>, 2019.
- McManus, J. F., Francois, R., Gherardi, J.-M., Keigwin, L. D., and Brown-Leger, S.: Collapse and rapid resumption of Atlantic meridional  
circulation linked to deglacial climate changes, *Nature*, 428, 834–837, <https://doi.org/10.1038/nature02494>, 2004.
- Meccia, V. L. and Mikolajewicz, U.: Interactive ocean bathymetry and coastlines for simulating the last deglaciation with the Max Planck  
510 Institute Earth System Model (MPI-ESM-v1. 2), *Geosci. Model Dev.*, 11, 4677–4692, <https://doi.org/10.5194/gmd-11-4677-2018>, 2018.
- Menviel, L., Timmermann, A., Mouchet, A., and Timm, O.: Meridional reorganizations of marine and terrestrial productivity during Heinrich  
events, *Paleoceanography*, 23, <https://doi.org/10.1029/2007PA001445>, 2008.
- Menviel, L., Timmermann, A., Timm, O. E., and Mouchet, A.: Deconstructing the Last Glacial termination: the role of millennial and  
orbital-scale forcings, *Quaternary Sci. Rev.*, 30, 1155–1172, <https://doi.org/10.1016/j.quascirev.2011.02.005>, 2011.
- 515 Menviel, L., England, M. H., Meissner, K., Mouchet, A., and Yu, J.: Atlantic-Pacific seesaw and its role in outgassing CO<sub>2</sub> during Heinrich  
events, *Paleoceanography*, 29, 58–70, <https://doi.org/10.1002/2013PA002542>, 2014.
- Muglia, J. and Schmittner, A.: Glacial Atlantic overturning increased by wind stress in climate models, *Geophys. Res. Lett.*, 42, 9862–9868,  
<https://doi.org/10.1002/2015GL064583>, 2015.
- Obata, A.: Climate–carbon cycle model response to freshwater discharge into the North Atlantic, *J. Climate*, 20, 5962–5976,  
520 <https://doi.org/10.1175/2007JCLI1808.1>, 2007.
- O’ishi, R. and Abe-Ouchi, A.: Influence of dynamic vegetation on climate change and terrestrial carbon storage in the Last Glacial Maximum,  
*Clim. Past*, 9, 1571–1587, <https://doi.org/10.5194/cp-9-1571-2013>, 2013.
- Orr, J. and Epitalon, J.-M.: Improved routines to model the ocean carbonate system: mocsy 2.0, *Geosci. Model Dev.*, 8, 485–499,  
<https://doi.org/10.5194/gmd-8-485-2015>, 2015.
- 525 Paulmier, A., Kriest, I., and Oschlies, A.: Stoichiometries of remineralisation and denitrification in global biogeochemical ocean models,  
*Biogeosciences*, 6, 923–935, <https://doi.org/10.5194/bg-6-923-2009>, 2009.
- Paulsen, H., Ilyina, T., Six, K. D., and Stemmler, I.: Incorporating a prognostic representation of marine nitrogen fixers into the global ocean  
biogeochemical model HAMOCC, *J. Adv. Model. Earth Sys.*, 9, 438–464, <https://doi.org/10.1002/2016MS000737>, 2017.
- Pickett, E. J., Harrison, S. P., Hope, G., Harle, K., Dodson, J. R., Peter Kershaw, A., Colin Prentice, I., Backhouse, J., Colhoun, E. A.,  
530 D’Costa, D., et al.: Pollen-based reconstructions of biome distributions for Australia, Southeast Asia and the Pacific (SEAPAC region) at  
0, 6000 and 18,000 14C yr BP, *J. Biogeog.*, 31, 1381–1444, <https://doi.org/10.1111/j.1365-2699.2004.01001.x>, 2004.



- Prentice, I., Harrison, S., and Bartlein, P.: Global vegetation and terrestrial carbon cycle changes after the last ice age, *New Phytologist*, 189, 988–998, <https://doi.org/10.1111/j.1469-8137.2010.03620.x>, 2011.
- 535 Prentice, I. C. and Webb III, T.: BIOME 6000: reconstructing global mid-Holocene vegetation patterns from palaeoecological records, *J. Biogeog.*, 25, 997–1005, <https://doi.org/10.1046/j.1365-2699.1998.00235.x>, 1998.
- Prentice, I. C., Jolly, D., and Participants, B. .: Mid-Holocene and glacial-maximum vegetation geography of the northern continents and Africa, *J. Biogeog.*, 27, 507–519, <https://doi.org/10.1046/j.1365-2699.2000.00425.x>, 2000.
- Reick, C., Raddatz, T., Brovkin, V., and Gayler, V.: Representation of natural and anthropogenic land cover change in MPI-ESM, *J. Adv. Model. Earth Sys.*, 5, 459–482, <https://doi.org/10.1002/jame.20022>, 2013.
- 540 Reick, C. H., Gayler, V., Goll, D., Hagemann, S., Heidkamp, M., Nabel, J. E., Raddatz, T., Roeckner, E., Schnur, R., and Wilkenskjaeld, S.: JSBACH 3-The land component of the MPI Earth System Model: documentation of version 3.2, 2021.
- Riddick, T., Brovkin, V., Hagemann, S., and Mikolajewicz, U.: Dynamic hydrological discharge modelling for coupled climate model simulations of the last glacial cycle: the MPI-DynamicHD model version 3.0, *Geosci. Model Dev.*, 11, 4291–4316, <https://doi.org/10.5194/gmd-11-4291-2018>, 2018.
- 545 Roche, D. M., Renssen, H., Paillard, D., and Levvasseur, G.: Deciphering the spatio-temporal complexity of climate change of the last deglaciation: a model analysis., *Clim. Past*, 7, <https://doi.org/10.5194/cp-7-591-2011>, 2011.
- Schmittner, A., Brook, E. J., and Ahn, J.: Impact of the ocean’s overturning circulation on atmospheric CO<sub>2</sub>, in: *Ocean Circulation: Mechanisms and Impacts*, edited by: Schmittner, A., Chiang, J. C. H., and Hemming, S. R., *Geophysical Monograph Series*, 173, American Geophysical Union, Washington DC, 315–334, <https://doi.org/10.1029/173GM20>, 2007.
- 550 Six, K. D. and Maier-Reimer, E.: Effects of plankton dynamics on seasonal carbon fluxes in an ocean general circulation model, *Global Biogeochem. Cycles*, 10, 559–583, <https://doi.org/10.1029/96GB02561>, 1996.
- Sosdian, S. and Rosenthal, Y.: Deep-sea temperature and ice volume changes across the Pliocene-Pleistocene climate transitions, *Science*, 325, 306–310, <https://doi.org/10.1126/science.1169938>, 2009.
- Spratt, R. M. and Lisiecki, L. E.: A Late Pleistocene sea level stack, *Clim. Past*, 12, 1079–1092, <https://doi.org/10.5194/cp-12-1079-2016>, 555 2016.
- Takahashi, T., Broecker, W. S., and Langer, S.: Redfield ratio based on chemical data from isopycnal surfaces, *J. Geophys. Res.- Oceans*, 90, 6907–6924, <https://doi.org/10.1029/JC090iC04p06907>, 1985.
- Tarasov, L., Dyke, A. S., Neal, R. M., and Peltier, W. R.: A data-calibrated distribution of deglacial chronologies for the North American ice complex from glaciological modeling, *Earth Plan. Sci. Lett.*, 315, 30–40, <https://doi.org/10.1016/j.epsl.2011.09.010>, 2012.