

Bio-antioxidants Activity: Their Mechanisms and Measurement Methods

Sahari, M. A. *, Berenji Ardestani, S.

*Department of Food Science and Technology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran.

Abstract

Bioactive food components (such as phenolic compounds, carotenoids, anthocyanins, tocopherols, ascorbic acid, flavonoids, minerals, enzymes and pectin) are active ingredients in food or dietary supplements, which proven to have an important role in health, and are safe for human consumption. These compounds exert their antioxidant effects by different mechanisms such as single electron transfer or hydrogen atom transfer, and their efficiencies can be evaluated by several methods such as ferric reducing ability of plasma, oxygen radical absorbance capacity assay, total radical trapping antioxidant parameter, total oxidant scavenging capacity, conjugated dienes, lipoxygenase activity inhibition assay, chemiluminescence, deoxyribose, tocopheroxyl radical attenuating ability, and nitric oxide radicals trapping. In this review, these mechanisms and methods will be discussed in details.

Article Info

Article history

Received 21 Jun 2014

Revised 26 Oct 2014

Accepted 3 Nov 2014

Keywords

Bioactive compounds,
Antioxidant,
Mechanisms of antioxidant action,
Hydrogen atom transfer,
Single electron transfer

Correspondence to

Mohammad Ali Sahari, Tarbiat
Modarres University, Tehran, Iran.
Email address: sahari@modares.ac.ir

1. Introduction

Bioactive food components are active ingredients in food or dietary supplements, which are derived from vegetable, animal and microbial sources, including materials necessary for human major nutritional needs, and materials proven to play a role in the health and safety of human beings. All active components in foods such as micro- and macro-nutrients should be considered as bioactive food compounds. Bioactive compounds are classified into different groups according to their distinctive chemical structures and functions (e.g. phenolic compounds and their subclasses such as flavonoids). The biological activity of a chemical group is affected not only by the differences in chemical composition and structure, but also by such factors as bioavailability, distribution and metabolism. In these experiments, all positive and negative constructive effects of bioactive food components should be investigated in scientific models of risk-benefit; however, toxic components should not be considered as bioactive compounds [1]. Microbial

products including modified proteins and amino acids can be mentioned as natural antioxidants [2, 3]. Functional foods (that are beneficial for health, and help reduce the risk of chronic diseases) include conventional foods containing bioactive natural materials such as fiber, foods enriched with bioactive substances like probiotics and antioxidants, and synthetic commercial nutrients like prebiotics [1, 4]. This study aimed to review the antioxidant properties of bioactive compounds, involved mechanisms and effective factors in bio-antioxidant effects, and at last several practical methods for assessing their antioxidant activity.

2. A Brief about Oxidation Reactions

Oxidizing characteristics of oxygen have a vital role in different biological actions including electron transport in ATP production. While, oxygen is essential for life, it can have destructive effects by means of cellular material oxidation [5]. Auto-oxidation and thermal oxidation of lipids, ce-

llular oxidation pathways, and numerous physiological and biochemical processes in human body under normal circumstances produce two groups of radicals; reactive oxygen species and reactive nitrogen species [6]. Thus, they attempt to obtain or lose electrons in the body, and consequently, cause damages to DNA, proteins, lipids and carbohydrates [4,7,8]. Also, free radicals participate in the mechanism of cytochromes [9]. The lack of anti-oxidative capacity causes imbalance and therefore, high concentrations of reactive oxygen species cause damages to cellular lipids, proteins, and nucleic acids as well as their routine performances [10, 11,12]. Human body always requires antioxidant sources continuously [13]. Vitamins C and E, carotenoids, xanthophylls, tannins, several amino acids, and total phenolic are involved in the antioxidant index of a sample [14, 15, 16].

Antioxidants' functions can be divided into two categories: Chain breaking and path inhibitors. Antioxidants sources in biological systems are categorized as follows: enzymes, large molecules, small molecules, some hormones [17]. Under different conditions and concentrations, antioxidants such as vitamin E and vitamin C can act as antioxidant, peroxidant or pro-oxidant [18, 19].

3. Main Chemical Reactions of Antioxidants

Antioxidants are capable of inactivating radicals with two major mechanisms, hydrogen atom transfer (HAT) and single electron transfer (SET). Methods using HAT mechanism include: oxygen radical absorbance capacity (ORAC), total radical trapping antioxidant parameter (TRAP), total oxidant scavenging capacity (TOSC), chemical luminescence, α -carotene bleaching, low density lipoprotein (LDL) oxidation, and methods using SET mechanism are: ferric reducing antioxidant power (FRAP), TRAP and copper reduction. Total phenolic method (Folin-Ciocalteu method= FC) is among the methods using both HAT and SET mechanisms [17].

4. Factors Affecting Antioxidant Mechanisms and Capacity

Ascorbic acid, scurvy preventive agent, is the most abundant water-soluble antioxidant and a heat sensitive vitamin [20]. The antioxidant effect of ascorbate is attributed to its capacity to remove reactive oxygen species by reaction with superoxide radicals, hydroxyl radicals, hydrogen

peroxide, and single oxygen through the HAT mechanism. Also, this compound removes RNS and prevents nitration reactions. Biological studies of vitamin C have shown its protection role against DNA mutations [21].

Polyphenols are the main plant components with antioxidant activity [13], and the hydrogen donating property of polyphenol compounds is responsible for preventing the production of free radical causing lipid peroxidation [22]. Generally, phenolic compounds in low concentrations show antioxidant behavior. In higher concentrations, they display pro-oxidant behavior, and again by further increasing of the concentration, they show antioxidant behavior [23,24,25]. Flavonoids also chelate metal ions, and therefore, prevent them from participating in the production of free radicals [26,27]. Polyphenolic compounds are implicated in large number of biological effects including anti-bacterial, anti-inflammatory, anti-histamine, anti-hepatitis, anti-viral, anti-cancer, anti-coagulation, and vessel protection [6, 28].

Carotenoids, especially α -carotene, quench single oxygen [12]. They are efficient quenchers against reactive oxygen species, particularly peroxy radical. Gene regulation is one of the most apparent mechanisms in some pathogenic processes, which is influenced by carotenoids. Carotenoids are relatively resistant to thermal decomposition [29].

In human nutrition, selenium, copper, zinc, iron and manganese are essential minerals for optimizing the response of antioxidant enzymes. Prosthetic groups that are present in heme proteins are similar to the antioxidant enzymes of catalase, peroxidase and superoxide dismutase. Research findings have shown that the amount of anthocyanins increases with the increase of zinc level; these compounds prevent membrane lipid peroxidation as a plant protective system against oxidative stress [30].

Vitamin E (α -tocopherol) is the best single oxygen quencher that acts as a chain breaking antioxidant [12].

Antioxidant activities of glutathione and uric acid are exerted by scavenging free radicals and antioxidant abilities of isothiocyanates (including anti-nutritive compounds) inactivate single oxygen [31].

5. Practical Methods for Assessing Antioxidants

Total phenolic content reducing capacity is measured by Folin-Ciocalteu method, which is

well related to the reducing and antioxidant abilities of phenolic compounds. Phenolic compounds react with FCR only in alkaline conditions (sodium carbonate solution at pH=10) [15,20,32].

Flavonoids in food systems act as free radical scavengers and terminate chain radical reactions during the triglyceride oxidation; they also exert various biological effects (such as antioxidant, anti-inflammatory), and protect vascular actions [28]. Flavonoid content was measured using colorimetric method provided by Lenucci *et al.* (2006) with some modifications [21, 23].

Analysis of ascorbic acid in vegetables and fruits was carried out according to the works the Ngee Wen *et al.* (2010) and Sahari *et al.* (2004) [24, 33-34].

Spectrophotometric methods with UV-VIS detectors at 400-600nm wavelengths are usually used for determining peroxide values. These procedures usually use oxidation of Fe (II) to Fe (III) to react with different reagents, and produce colored complexes [35, 36, 37-38].

The FRAP method is related to the reduction of [Fe (III)-(2, 4 and 6-three (2-Pyridyl)-s-Tryazyn=TPTZ)₂] complex to a colored product [Fe (II)-(TPTZ)₂] by an antioxidant and usually in non-physiological conditions at low pH of about 3.6 [24,39,40]. Generally, FRAP mechanism, as compared to the mixed SET and HAT, is based more on electron transfer; so, its combination with other methods for the diagnosis of the dominant mechanisms of various antioxidants is very useful.

The ORAC mechanism is a HAT system, which is monitored by means of spectrophotometer. ORAC is applied to assess the antioxidant capacity of hydrophilic compounds, and with some modifications, it is used to determine the antioxidant capacity of hydrophobic compounds [16, 41, 42]. The ORAC method measures the inhibition of peroxy radical oxidation generated by antioxidant activity, and thus reflects the antioxidants' radical chain breaking activity through H atom transmission. In this method, the peroxy radical reacts with a fluorescent probe to form a non-fluorescent product that can be determined easily by fluorescence assessment.

TRAP method shows the ability of antioxidant compounds to interfere with the reaction between the peroxy radicals derived from AAPH or ABAP (2, 2'-azobis (2-amidinopropane) dihydrochloride) and the target probe. Antioxidants expand the ret-

ardation time to advent oxidized probe, and their ability is measured by percentage of reaction reduction. TRAP values are usually expressed as retardation time or sample reaction time in comparison to the corresponding times for trolox. Copper reduction method is based on the reduction of Cu (II) to Cu (I) by the combined antioxidant action of all samples (reducing agents). CUPRAC (copper reduction antioxidant capacity) values are comparable with total equivalent antioxidant capacity = TEAC values for polyphenols, while FRAP values are usually significantly lower. TOSC method determines the antioxidants absorbance capacity especially against three major oxidants, including hydroxyl radicals, peroxyl radicals and peroxy nitrite. In this method, α -keto- γ -methyl butyric acid is oxidized to form ethylene. Time analysis of ethylene formation is followed by headspace GC analysis, and antioxidant capacity is determined through the antioxidants' ability to prevent ethylene formation towards the control reaction [43]. Fresh LDL is isolated from blood samples, and oxidized by Cu (II) or other substances such as AAPH, forming conjugated dienes that are monitored spectrophotometrically at 234nm wavelength. The effects of antioxidant compounds with HAT mechanism are evaluated by reducing the absorption amounts at 234nm. This method is appropriate for assessing a simple fatty acid, and the primary stage of lipid peroxidation [44,45]. Antioxidants react non-specifically with lipoxygenase through scavenging the radical intermediates and reduce their iron section [13]. The phenolic antioxidant's activity is probably related to their reducing properties, which allow them to act as reducing agents or hydrogen atoms donors. In addition to their ability to chelate metals, they inhibit lipoxygenase and scavenge free radicals [28]. Enzymatic lipid peroxidation is measured by means of spectrophotometer with an increase in the lipid hydroperoxide absorption at 234nm [13]. In a simple and cheap method, deoxyribose sugar decomposes with exposure to the OH[•] generated by radiation or by Fenton systems. If the complex mixture of products is heated under acidic conditions, malone aldehyde is formed, which is identified by its ability to react with thiobarbitonic acid and the formation of pink material. In addition, deoxyribose is used to determine the OH[•] formed in biological systems. It is likely that the decomposition rate of deoxyribose increases in the

presence of reducing agents such as ascorbic acid in the reaction mixture [4, 5, 46].

Tocopheroxyl radical attenuating ability is a quick and easy method for screening the inhibitor's ability to prevent LDL peroxidation. The method is based on the antioxidants' capacity to reduce α -tocopheroxyl radicals' contents. α -tocopherols are produced by UV light, and measured by electron spin resonance spectroscopy. The tocopheroxyl radical attenuating ability method using $\text{CCl}_3\text{O}_2^\cdot$ is expensive because it needs electron spin resonance machine [24].

Nitric oxide radicals trapping method is based on this assumption that sodium nitroprusside, which reacts with environmental oxygen to produce nitrite ions in an aqueous solution at physiological pH, slowly produces nitric oxide. Nitric oxide trapping and competing with oxygen decrease the production of nitrite ions. The absorption at 546nm is read versus the solvent. Quercetin is used as the control for comparison [47]. Proxy nitrite can cause lipid peroxidation, oxidation of methionine and sulfhydryl residues in proteins decrease in antioxidants and damage to DNA.

The antioxidant activity is monitored through the main reaction of HAT by a decrease in the chemical luminescence intensity caused by luminal and superoxide anion radical. Luminol is the most important indicator component for trapping oxidants, and intensifying long lasting and stable emissions. A continuous light output depends on the constant production of free radicals derived from p-iodophenol, luminol and oxygen. Also, the light emission is sensitive to radical scavenger antioxidants interferences. The antioxidant capacity is assessed as the emission light declines [48].

6. Conclusion

Peroxidation of lipids in the cell membrane, caused by free radicals, is critically associated with heart diseases, cancers, aging, etc. This harmful procedure can be inhibited by bio-antioxidants' activities in biosystems. It is well accepted that ascorbic acid, polyphenols, carotenoids, α -tocopherol, minerals as selenium, copper, zinc, iron, manganese and other compounds like glutathione, uric acid and isothiocyanates can exert notable antioxidant properties against free radicals. These antioxidants are capable of inactivating radicals with the two major mechanisms of HAT and SET. HAT-based methods measure the ability

of antioxidants to disable free radicals by hydrogen donating. SET-based methods detect the ability of antioxidants to transfer single electron to reduce any compound, including metals, carbonyls and radicals. Principal mechanisms applied by antioxidant compounds include the reduction of metals such as iron and copper; scavenging free radicals; creating complexes with metal pro-oxidants (chelating); quenching single oxygen; and stimulating anti-oxidative defense enzymatic activities. Some methods are usable to evaluate the activity of bio-antioxidants, which were discussed in this review in details. In brief, methods using HAT mechanism include ORAC, TRAP, TOSC, chemical luminescence, α -carotene bleaching, and LDL oxidation. Methods using SET mechanism include FRAP, TRAP and copper reduction. Total phenolic method (FC) uses both HAT and SET mechanisms.

References

1. Biesalski HK, Dragsted L, Elmadfa I, Grossklaus R, Muller M, Schrenk D, et al. Bioactive compounds: Safety and efficiency. *Nutrition* 2009; 25: 1202-1205.
2. Hall C. Sources of natural antioxidants: Oil seeds, nuts, cereals, legumes, animal products and microbial sources. North Dakota State University. 2001. Available from: <http://www.academia.edu/1072737/>.
3. Takao T, Kitatani F, Watanabe N, Yagi A, Sakata K. A simple screening method for antioxidants and isolation of several antioxidants produced by marine bacteria from fish and shellfish. *Biosci Biotech Biochem* 1994; 58(10): 1780-1783.
4. Meghashri S, Vijay Kumar H, Gopal S. Antioxidant properties of a novel flavonoid from leaves of *Leucas aspera*. *Food Chem* 2010; 122 (1): 105-110.
5. Haghiroalsadat F, Bernard F, Kalantar SM, Sheikhha MH, Hokmollahi F, Azimzadeh M, et al. *Buniumpersicum* (Black Caraway) of Yazd province: Chemical assessment and evaluation of its antioxidant effects. *J Shahid Sadoughi Univ Med Sci* 2010; 18(4): 284-291 [in Persian].
6. Benhammou N, Bekkara FA, Panovska TK. Antioxidant activity of methanolic extracts and some bioactive compounds of *Artiplexhalimus*. *CR Chem* 2009; 12: 1259- 66.
7. Rasooli I, Gachkar L, Yadegarinia D, et al. Relation of anti-oxidative property and free

- radical scavenging capacity to the anti-microbial characteristics of essential oils from *Mentha spicata* L. and *Chenopodium ambrosioides* L. Iranian J Med Arom Plants 2008; 23(4): 492-503 [in Persian].
8. Movahedian I, Dashti Gh, Naderi Gh, *et al.* Anti-atherogenic and antioxidant effects of valine on coronary arteries of hypercholesterolemic animals. Armaghan-e- Danesh J 2008; 13(2): 61-68 [in Persian].
 9. Kamkar A. The study of antioxidant activity of essential oils and extract of Iranian *Anethum graveoloens*. Ofogh-e-Danesh.GMUHS J 2009; 15(2): 11-17 [In Persian].
 10. Kusznierevicz B, Bartoszek A, Wolska L, Drzewiecki J, Gorinstein S, Namiesnik J. Partial characterization of white cabbages (*Brassica oleracea var. Capitata f. alba*) from different regions by glucosinolates, bioactive compounds, total antioxidant activities and proteins. LWT-Food Sci Technol 2008; 41: 1-9.
 11. Jayaprakasha GK, Patil BS. *In vitro* evaluation of the antioxidant activities in fruit extracts from citron and blood orange. Food Chem 2007; 101: 410-418.
 12. Chanwitheesuk A, Teerawutgulrag A, Rakariyatham N. Screening of antioxidant activity and antioxidant compounds of some edible plants of Thailand. Food Chem 2005; 92: 491-497.
 13. Ajila CM, Naidu KA, Bhat SG, Rao UJSP. Bioactive compounds and antioxidant potential of mango peel extract. Food Chem 2007; 105: 982-988.
 14. Gioti E, Fiamegos Y, Salkos D, Stalikas C. Antioxidant activity and bioactive components of the aerial parts of *Hypericum perforatum* L. from Epirus, Greece. Food Chem 2009; 117: 398-404.
 15. Ao C, Li A, Elzaawely AA, Xuan TD, Tawata S. Evaluation of antioxidant and anti-arterial activities of *Ficus microcarpa* L. fil. extract. Food Cont 2008; 19: 940-48.
 16. Perez-Jimenez J, Arranz S, Taberner M, Diaz-Rubio ME, Serrano J, Gono I, *et al.* Updated methodology to determine antioxidant capacity in plant foods, oils and beverages: Extraction, measurement and expression of results. Food Res Int 2008; 41: 274-85.
 17. Prior RL, Wu X, Schaich K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. J Agric Food Chem 2005; 53: 4290-302.
 18. Asgary S, Emami SA, Shams Ardekani MR, Naderi Gh, Aslani S, Kosher C, *et al.* Antioxidant effects of the essential oils from the fruits and twigs of *Juniperus excels* sub sp. *Excels* on several oxidative systems. Iranian J Med Arom Plants 2007; 23(1): 59-72 [in Persian].
 19. Yen G, Hsieh C. Antioxidant activity of extracts from Duzhong (*Eucommia ulmoides*) toward various lipid peroxidation models *in vitro*. J Agric Food Chem 1998; 46: 3952-57.
 20. Rufino M, Alves RE, Brito E, Perez-Jimenez J, Saura-Calixto F, Mancini-Filho J. Bioactive compounds and antioxidant capacities of 18 non-traditional tropical fruits from Brazil. Food Chem 2010; 121(4): 996-1002.
 21. Stratil P, Klejdus B, Kuban V. Determination of phenolic compounds and their antioxidant activity in fruits and cereals. Talanta 2007; 71: 1741-51.
 22. Luo W, Zhao M, Shen G, Rao G. Identification of bioactive compounds in *Phyllanthus emblica* L. fruit and their free radical scavenging activities. Food Chem 2009; 114: 499-504.
 23. Tomaino A, Martorana M, Arcoraci T, Monteleone D, Giovanazzo C, Saija A. Antioxidant activity and phenolic profile of pistachio (*Pistacia vera* L., variety Bronte) seeds and skins. Biochimie 2010; 92(9): 1115-22.
 24. Aruoma OL. Review: Methodological considerations for characterization potential antioxidant actions of bioactive components in plant foods. Mutat Res 2003; 523-24: 9-20.
 25. Amaral S, Mira L, Nogueira JMF, Pereira da Silva A, Felorencio M. Plant extracts with anti-inflammatory properties-A new approach for characterization of their bioactive compounds and establishment of structure antioxidant activity relationships. Bioorg Med Chem 2009; 17: 1876-83.
 26. Cano A, Medina A, Bermejo A. Bioactive compounds in different citrus varieties. Discrimination among cultivars. J Food Compost Anal 2008; 21: 377-381.
 27. Velioglu YS, Mazza G, Gao L, Oomah BD. Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. J Agric Food Chem 1998; 46: 4113-4117.
 28. Barreira GCM, Ferreira ICFR, Oliveira MBPP, Pereira JA. Antioxidant activity and bioactive

- compounds of ten Portuguese regional and commercial almond cultivars. *Food Chem Technol* 2008; 46: 2230-35.
29. Perez-Conesa D, Garcia-Alonso J, Garcia-Valverde V, Iniesta MD, Jacob K, Sanches-Siles, LM, *et al.* Changes in bioactive compounds and antioxidant activity during homogenization and thermal processing of tomato puree. *Innov Food Sci Emerg Technol* 2009; 10: 179-88.
 30. Zare Dehabadi S, Asrar Z. Effect of excess zinc on the concentration of some mineral element and antioxidant responses of spearmint (*Mentha spicata* L.). *Iran J Med Arom Plants* 2009; 24(4): 530-40.
 31. Obon JM, Castellar MR, Cascales JA, Fernandez-Lopez JA. Assessment of the TEAC method for determining the antioxidant capacity of synthetic red food colorants. *Food Res Int* 2005; 38: 843-45.
 32. Ou B, Huang D, Hampsch-Woodili M, Flanagan JA, Deemer EK. Analysis of antioxidant activities of common vegetables employing oxygen radical absorbance capacity (ORAC) and ferric reducing antioxidant power (FRAP) assays: A comparative study. *J Agric Food Chem* 2002; 50: 3122-28.
 33. Ngee Wen T, Prasad KN, Yang B, Ismail A. Bioactive substance contents and antioxidant capacity of raw and blanched vegetables. *Innov Food Sci Emerg Technol* 2010; 11(3): 464-69.
 34. Sahari MA, Boostani M, Hamidi Esfahani Z. Effect of low temperature on the ascorbic acid content and quality characteristics of frozen strawberry. *Food Chem* 2004; 86: 357-363.
 35. Samadloiy HM, Azizi MH, Barzegar M. Anti-oxidative effect of compounds pomegranate seed phenolic components on soybean oil. *J Agric Sci Natur Resour* 2007; 14(4): 193-200 [in Persian].
 36. Hornero-Mendez D, Perez-Galvez A, Minguez-Mosquera MI. A rapid spectrophotometric method for the determination of peroxide value in food lipids with high carotenoid content. *J Am Oil Chem Soc* 2001; 78(11): 1151-55.
 37. Assimopoulou AN, Zlatanov SN, Papageorgiou VP. Antioxidant activity of natural resins and bioactive triterpenes oil substrates. *Food Chem* 2005; 92: 721-27.
 38. Lee OH, Lee BY, Lee J, Lee HB, Son JY, Park CS, *et al.* Assessment of phenolics enriched extract and fractions of olive leaves and their antioxidant activities. *Bioresour Technol* 2009; 100: 6107-13.
 39. Polovka M, Suhaj M. Detection of caraway and bay leaves irradiation based on their extracts antioxidant properties evaluation. *Food Chem* 2010; 119: 391-401.
 40. Tezcan F, Gultekin-Ozguven M, Diken T, Ozcelik B, Bedia Erim F. Antioxidant activity and total phenolic, organic acid and sugar content in commercial pomegranate juices. *Food Chem* 2009; 115: 873-77.
 41. Wu R, Feri B, Kennedy J, Zhao Y. Effects of refrigerated storage and processing technologies on the bioactive compounds and antioxidant capacities of Marion and Evergreen blackberries. *LWT-Food Sci Technol* 2010; 1-12.
 42. Sales J, Resurreccion A. Phenolic profile, antioxidants and sensory acceptance of bioactive-enhanced peanuts using ultrasound and UV. *Food Chem* 2010; 122: 795-803.
 43. Lichtenthaler R, Marx F, Kind OM. Determination of anti-oxidative capacities using an enhanced total oxidant scavenging capacity (TOSC) assay. *Eur Food Res Technol* 2003; 216: 166-73.
 44. Fathi A, Sahari MA, Zangiabadi M, Barzegar M. Application of *Satureja hortensis* L. and *Zataria multiflora* Boiss. essential oils as two natural antioxidants in soybean oil during microwave heating. *J Med Plants* 2011; 10: 12-21.
 45. Vahidyan H, Sahari MA, Barzegar M, Naghdi Badi H. Application of *Zataria multiflora* Boiss. and *Satureja hortensis* L. essential oils as two natural antioxidants in mayonnaise formulated with linseed oil. *J Med Plants* 2012; 11: 69-79.
 46. Halliwell B, Gutteridge JMC, Aruoma OI. The deoxyribose method: A simple test-tube assay for determination of rate constants for reactions of hydroxyl radicals. *Anal Biochem* 1987; 165: 215-19.
 47. Goli Z, Lakzaei M, Pour Amir M. Antioxidant activity of orange peel extract and its effect on lipid oxidation of raw and cooked meat *Hypophthalmichthys molitrix* fish. *Iran J Nutr Sci Food Technol* 2010; 5(2): 19-26.
 48. Alho H, Leinonen, J. Total antioxidant activity measured by chemiluminescence methods. *Method Enzymol* 1999; 299: 3-1.