

Optimizing Submerged Cultivation for the Production of Red Pigments by *Monascus purpureus* on Soybean Meals using Response Surface Methodology

Hadiseh Keivani, Mahshid Jahadi*, Nafiseh Ghasemisepero

Department of Food Science and Technology, Faculty of Agriculture, Isfahan (Khorasgan) Branch, Islamic Azad University, Isfahan, Iran.

Abstract

Background and objective: *Monascus purpureus* is a filamentous fungus with ability to produce pigments with therapeutic properties. Red pigments are especially used as additives, antioxidants, preservatives and substitutions for nitrites in food technology. To decrease fermentation costs, agro-industrial wastes such as soybean meals have been used as substrates. In the current study, red pigment production by *Monascus purpureus* on soybean meals was optimized.

Material and methods: In this study, red *Monascus* pigment production by *Monascus purpureus* ATCC 16362 was carried out under submerged fermentation using soybean meals as nitrogen sources to replace yeast extracts. Central composite design was used to assess the optimum level of soybean meal replacement (0-100%), $ZnSO_4 \cdot 7H_2O$ concentration (0-0.02 g l⁻¹) and thermal stress time of spore suspension at 70°C (50-90 s). Red *Monascus* pigment and biomass productions were assessed as dependent responses.

Results and conclusion: The maximum production of red *Monascus* pigment (4.54 AU ml⁻¹) was achieved under conditions of soybean meal replacement of 79.72%, $ZnSO_4 \cdot 7H_2O$ concentration of 0-0.02 g l⁻¹ and thermal stress time of spore suspension of 81.89 s. The average yield of red *Monascus* pigment, conversion factor of biomass in red pigment $Y_{P/X}$ and cell productivity included 0.324 AU ml⁻¹ day⁻¹, 1.10 AU L g⁻¹ and 0.292 g l⁻¹ day⁻¹, respectively. Results of the current study have demonstrated that combination of soybean meal and yeast extract as nitrogen source is beneficial for the production of red *Monascus* pigment by *Monascus purpureus*.

Conflict of interest: The authors declare no conflict of interest.

How to cite this article

Keivani H, Jahadi M, Ghasemisepero N. Optimizing Submerged Cultivation for the Production of Red Pigments by *Monascus purpureus* on Soybean Meals Using Response Surface Methodology. *Appl Food Biotechnol* 2019; 7(3): 143-151. <http://dx.doi.org/10.22037/afb.v7i3.28931>

Article Information

Article history:

Received 05 Feb 2020
Revised 04 April 2020
Accepted 17 April 2020

Keywords:

- Central composite design
- Monascus purpureus*
- Red pigment
- Response surface method
- Soybean meal

*Corresponding author:

Mahshid Jahadi,
Islamic Azad University,
Isfahan (Khorasgan) Branch,
Arghavanieh, Jey St.,
Khorasgan, Isfahan, Iran.

Tel: +98-34-35342221
Fax: +98-31-35354060
E-mail:
m.jahadi@khuif.ac.ir

1. Introduction

Color is one of the most important characteristics of foods. Use of synthetic dyes can cause carcinogenicity and fetal defects. Therefore, natural pigments have become popular as raw materials in food, pharmaceutical and cosmetics industries. Microbial pigments, as natural pigments (bio pigments) extracted from microorganisms, are produced by a number of bacteria, algae and fungi. Microbial pigment production is preferred over other bio-dye productions because of easy cultivations, production short times and low costs [1,2].

Of the pigment producing microorganisms, *Monascus* (*M.*) *purpureus* is important due to the production of red,

yellow and orange pigments with therapeutic properties such as cancer, inflammation, diabetes and fat prevention and reduction. The *M. purpureus* pigments have been used as antioxidants, additives and preservatives in food products and as substitutions for nitrites in processing of meats. Asian countries monitored citrinin in red yeast rice and established maximum limitations for citrinin in foods for humans [3]. Singgih et al. showed that ethanol extraction of *Monascus* pigments decreased citrinin, which made this pigment safe for human consumption [4].

To decrease costs of fermentation in industrial scales, agricultural and industrial wastes can be used as substrates.

Agro-industrial substrates are considered for the production of secondary metabolites such as production of xanthan from date extracts [5], surfactin from rice straws [6], mycoprotein from date wastes [7], molasses, starches and fruit and vegetable wastes [8], alpha-amylase from wheat bran, rice brans and potato peels [9] and carotenoids from potato skins, mung bean husks, onion peels and pea pods [10]. Recently, studies have been carried out on use of agricultural wastes such as date wastes [11], grape wastes [12], bakery wastes [13], orange wastes [14], sugarcane bagasse [15,16] and bug damaged wheat [17] to produce *Monascus* pigment under submerged fermentation. Submerge fermentation for the production of red *Monascus* pigment can solve problems of low income, capital-intensive, time-consuming and large surface area necessity in solid-state fermentations. Moreover, this method is more appropriate for large-scale industrial production than solid-state production [18].

Several studies have been carried out for red *Monascus* pigment production by *M. purpureus* as a safe microorganism. Production of red *Monascus* pigment under submerged fermentation is affected by various parameters such as pH [11,19], zinc [20], carbon [21,22] and nitrogen sources [19,22-24], salts and environmental stresses [25].

Yeast extract, as a nitrogen source, is a beneficial substrate for *Monascus* pigment production [23]. However, use of other economic sources seems appropriate. Lubricated seeds are rich in proteins and used in dietary enrichments as well as biotechnological processes during fermentation. Soybean meal is a byproduct of oil seeds after lubrication and solvent extraction [26]. Soybean meal contains more than 43% of proteins [27]. Up to date, Soybean meal has been used in various fermentation processes as a substrate to produce secondary metabolites from microorganisms, including fumaric acid by *Rhizopus oryzae* [28], ethanol by *Saccharomyces cerevisiae* and *Zymomonas mobilis* [29] and Monacolin K by *M. purpureus* [30].

In nature, microorganisms are constantly affected by physical and chemical changes in their environments. However, microorganisms include defense mechanisms against a variety of environmental stresses. Numerous secondary metabolites are produced under stressful conditions. Creating environmental stresses such as high temperatures, osmotic pressures and presence of heavy metals such as Zn increases pigment productions. Therefore, pigment production by the microorganisms is increased to adapt to the new conditions [2]. This method can be used to increase production of secondary metabolites from microorganisms [31]. Although Soybean meal residues have been targets of replacing C- and N- sources [26-28], no studies have been carried out to decrease the yeast extract as nitrogen source using Soybean meal in culture media for the production of *Monascus* pigment during fermentation.

Therefore, the aim of this study was to identify significant nutritional and physical factors such as soybean meal replacement (0-100%), $ZnSO_4 \cdot 7H_2O$ concentration (0-0.02 g l⁻¹) and thermal stress time of spore suspension of 70°C (50-90 s) to produce red *Monascus* pigment from *M. purpureus* during submerged fermentation using response surface methodology.

2. Materials and methods

2.1. Microorganism

The *M. purpureus* ATCC 16362 was provided as live culture by the Iranian Scientific and Industrial Research Organization (IROST) (Tehran, Iran). Microorganism was cultured in potato dextrose agar (PDA) media and incubated at 30°C for 7 days. Then, culture was stored at 4°C until use [11].

2.2. Preparation of inoculum and seed culture

The yeast extract powder soluble starch media included 4 g of yeast extract, 1 g of K_2HPO_4 , 0.5 g of $MgSO_4$ and 15 g of soluble starch per liter. Seed culture was prepared using 7-day-old *M. purpureus* spores. The number of spores was adjusted using hemocytometer at 10⁵ (spore ml⁻¹). Spore suspension was heat stressed at 70°C with response surface design (50-90 s). These were then incubated with agitation at 120 rpm for 24-48 h at 30°C [11,25].

2.3. Media preparation and submerge fermentation

The Soybean meal was purchased from a local market and then milled to powder and sieved using 60-mesh net screen. Flavourzyme 1% (w w⁻¹) (Novozyme, Denmark) was added to the Soybean meal and mixed at 100 rpm for 24 h at 40°C. The culture media included 1 g glucose, 0.05 g $MgSO_4 \cdot 7H_2O$, 0.1 K_2HPO_4 , 0.001 g $FeSO_4 \cdot 7H_2O$, 0.05 g KCl, 0.3 g $NaNO_3$, 0.4 g yeast extract (yeast extract contents varied in each treatment and SBM (0-100%) replaced it) and 0-0.002 g of $ZnSO_4 \cdot 7H_2O$ per 100 ml (Table 1). Media was autoclaved at 121°C for 15 min [32,11]. Then, 10-15% (v v⁻¹) of the seed culture added to the media and incubated at 30°C for 14 days using shaker incubator (120 rpm) [11].

Table 1. Levels of the variables in central composite design ($\alpha = 1.68$)

Variables	Range of levels				
	$-\alpha$	-1	0	+1	$+\alpha$
SBMR† (%) (X ₁)	0	20.27	50	79.73	100
$ZnSO_4 \cdot 7H_2O$ (g l ⁻¹) (X ₂)	0	0.004	0.01	0.016	0.02
TST‡ (S) (X ₃)	50	58.11	70	81.89	90

† Soybean meal replacement

‡ Thermal stress time

2.4. Analytical method

2.4.1. Quantitative analysis of pigment

Pigment extraction was carried out according to Kantifedaki et al. Fermented media was mixed with alcohol (70%) and transferred to ultrasonic bath (Parsunic, Iran) for 30 min at 25°C. Then, mixture was filtered using no. 1 Whatman filter papers. The supernatant was used for the red *Monascus* pigment estimation by measuring the absorbance at 500 nm using spectrophotometer (UNICO2100, USA) and expressing the values as AU ml⁻¹ [14].

2.4.2. Biomass assessment

Filtration of the culture media was carried out using no. 1 Whatman filter papers and weighed using digital scale. Filtrates were dried at 75°C until achievement of a constant weight and weighed using digital scale [24].

2.5. Experimental design

In the present study, variables of soybean meal replacement (X_1) (0–100%), ZnSO₄·7H₂O concentration (X_2) (0–0.02 g l⁻¹) and thermal stress time (X_3) (50–90 s) were selected for further optimization by central composite design of response surface methodology using Design Expert Software v.7.0.0. Results of the central composite design experiments for investigating effects of red *Monascus* pigment (Y_1) and biomass (Y_2) productions are presented in Table 2. To investigate effects of the factor and their interactions, a central composite design experiment with 14 axial points and six center points was carried out ($\alpha = 1.68$). After validation of the models, the average yield of red pigment (P_M) (Eq. 1), conversion factor of biomass in red pigment ($Y_{P/X}$) (Eq. 2) and average cell productivity (P_{cells}) (Eq. 3) were calculated after 14 days of incubating [21,33].

$$P_M (\text{AU ml}^{-1} \text{ day}^{-1}) = \frac{P_{max} - P_0}{(t_{P_{max}} - t_{P_0})} \quad \text{Eq. 1}$$

$$Y_{P/X} (\text{AU L g}^{-1}) = \frac{P_{max} - P_0}{X_{max} - X_0} \quad \text{Eq. 2}$$

$$P_{cells} (\text{g l}^{-1} \text{ day}^{-1}) = \frac{X_{max} - X_0}{(t - t_0)} \quad \text{Eq. 3}$$

Where, P_{max} was the maximum rate of red pigment production at time $t_{P_{max}}$ (AU ml⁻¹), P_0 was the quantity of red pigments at t_{P_0} (AU ml⁻¹), $t_{P_{max}}$ was the time of maximum red pigment production (day⁻¹), t_0 was the initial cultivation time, X_{max} was the maximum biomass formation at time t (g l⁻¹) and X_0 was the biomass formation at time t_0 (g l⁻¹).

3. Results and discussion

3.1. Regression model

Results of red *Monascus* pigment and biomass productions were illustrated in Table 2. Red *Monascus* pigment and biomass productions varied 0.5 ±0.28 to 5 ±0.27 AU ml⁻¹ and 1 ±0.11 to 9.7 ±0.43 g l⁻¹, respectively, in various combinations of the 20 treatments. Models, describing behaviors of the responses, were created by finding the best setting of variables to optimize the fermentation process. Second order models of the two responses for coded variables were calculated using Eq. 4 for red *Monascus* pigment and Eq. 5 for biomass productions, respectively:

$$Y_1 = (+2.66) + (0.55 X_1) - (0.82 X_2) - (0.46 X_3) - (0.61 X_2 X_3) + (0.35 X_1^2) \quad \text{Eq. 4}$$

$$Y_2 = (+7.93) + (0.13 X_1) - (0.55 X_2) - (1.43 X_3) - (1.92 X_1 X_2) - (1.13 X_1^2) + (0.60 X_2^2) - (1.27 X_3^2) \quad \text{Eq. 5}$$

Table 2. Central composite design using experimental data and predicted values

Number	Variable			Red <i>Monascus</i> pigment (AU ml ⁻¹)		Biomass production (g l ⁻¹)	
	SBMR (%)	ZnSO ₄ ·7H ₂ O (g l ⁻¹)	TST (s)	Experimental	Predicted	Experimental	Predicted
1	20.27	0.004	58.11	3.2±0.08	2.92	5±0.17	4.86
2	79.73	0.004	58.11	4±0.22	4.02	9.7±0.43	8.98
3	20.27	0.016	58.11	2.3±0.12	2.51	7.78±0.28	7.61
4	79.73	0.016	58.11	3.68±.31	3.60	3.99±0.56	4.03
5	20.27	0.004	81.89	3.6±0.28	3.23	3 ±0.32	1.99
6	79.73	0.004	81.89	5±0.27	4.32	7.02±0.72	6.11
7	20.27	0.015	81.89	0.61±0.17	0.37	4.5±0.12	4.74
8	79.73	0.016	81.89	1.89±0.26	1.46	1.7±0.10	1.16
9	0	0.010	70.00	2.56±0.52	2.74	4.22±0.46	4.52
10	100	0.010	70.00	4.1±0.33	4.58	4.1±0.21	4.97
11	50	0.000	70.00	3.5±0.25	4.03	5.9±0.63	7.16
12	50	0.02	70	1.2±0.39	1.28	5.4±1.05	5.31
13	50	0.01	50	2.98±0.20	2.84	6.5±0.52	6.75
14	50	0.01	90	0.5±0.28	1.30	1±0.11	1.92
15	50	0.01	70	2.3±0.58	2.66	8.67±0.49	7.93
16	50	0.01	70	2.87±0.43	2.66	8±0.68	7.93
17	50	0.01	70	2.65±0.75	2.66	7.6±0.80	7.93
18	50	0.01	70	2.7±0.3	2.66	7±0.18	7.93
19	50	0.01	70	2.4±.29	2.66	8.7±0.16	7.93
20	50	0.01	70	3.08±0.09	2.66	7.83±0.11	7.93

SBMR= soybean meal replacement, TST= thermal stress time,

Table 3. Variance analysis of the response surface design of *Monascus purpureus* red pigment (Y_1) and biomass (Y_2) productions

Source	Red <i>Monascus</i> pigment (Y_1)				Biomass production (Y_2)			
	df	Mean squares	F-value	P-value	df	Mean squares	F-value	P-value
Model	6	3.62	18.15	0.000	7	14.60	21.12	0.000
X_1	1	4.06	20.35	0.001	1	0.25	0.35	0.566
X_2	1	9.17	45.91	0.000	1	4.12	5.96	0.031
X_3	1	2.86	14.33	0.002	1	28.10	40.66	0.000
$X_1 X_2$	-	-	-	-	1	29.64	42.89	0.000
$X_1 X_3$	-	-	-	-	-	-	-	-
$X_2 X_3$	1	2.98	14.91	0.002	-	-	-	-
X_1^2	1	1.83	9.17	0.009	1	18.30	26.48	0.000
X_2^2	-	-	-	-	1	5.21	7.53	0.017
X_3^2	-	-	-	-	1	23.35	37.78	0.000
Residual error	13	0.20	-	-	12	0.69	-	-
Lack of Fit	8	0.27	0.37	0.105	7	0.88	2.08	0.218
Pure Error	5	0.084	-	-	5	0.42	-	-
Corrected total	19	-	-	-	19	-	-	-
$R_1 = 0.89$								
$R_1(\text{adj}) = 0.84$								
$R_2 = 0.92$								
$R_2(\text{adj}) = 0.88$								

Where, X_1 , X_2 and X_3 represented soybean meal replacement, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration and thermal stress time, and Y_1 and Y_2 represented red *Monascus* pigment and biomass productions, respectively. Based on Analysis of variance test (Table 3) coefficient of determination (R^2) and adjusted coefficient of determination (R^2_{adj}) for red *Monascus* pigment and biomass productions were 0.89, 0.84 and 0.92, 0.88, respectively. The R^2 value seemed better between 0 and 1. The closer the R^2 was to 1, the excellent model and the more accurate predicted responses were [34]. This demonstrated good compatibility of the experimental and predicted values for red *Monascus* pigment and biomass productions. Red *Monascus* pigment and biomass production model F -values of respective-ly 18.15 and 21.12 and p -value of less than 0.0001 showed that the model was statistically significant. The lack-of-fit value of red *Monascus* pigment and biomass productions of 0.37 and 2.08 and p -value of 0.1052 and 0.218 showed that lack of fit was not statistically significant. Lack of fit is the variation of the data around the fitted model. If the model does not fit data extremely, the value is significant. Independent variables with no significance were neglected as backward elimination regarding their p -values ($P \leq 0.05$) and the model equation was modified to the new fitted model.

3.2. Interaction between the affecting factors

Of the three variables tested, soybean meal replacement (X_1), $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration (X_2) and thermal stress time (X_3) were highly significant for red *Monascus* pigment production, while interactions between the $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration and the thermal stress time ($X_2 X_3$) and the quadratic effects of soybean meal replacement (X_1) were significant based on the p -values (Table 3). Furthermore, 2D and 3D plots were created for the responses (red *Monascus* pigment production) at any of the two independent variables, while keeping the others at the middle point.

Based on Figure 1A, increases in soybean meal replacement resulted in increases in red *Monascus* pigment production. Amino acids have been reported as the most important N-sources to pigment production by *M. purpureus*. Since Soybean meal is rich in amino acids, it leads to the presence of further amine groups in the amination process of polyketide synthesis [35]. Interaction effects of various $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentrations and thermal stress time on red *Monascus* pigment production are shown in Figure 1B. Increases in $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration led to decreases in red *Monascus* pigment production.


Toxicity of high zinc concentrations could be associated to the creation of magnesium deficiency, which disturbed the microbial growth and hence decreased red pigment production as growth-dependent metabolites [36,37]. The

red *Monascus* pigment production was further sensitive to higher levels of thermal stress time at 70°C and the red pigment content decreased with increasing thermal stress time (Figure 1B). Abrashev et al. showed that temperatures above 50°C for 10-20 min resulted in germination delay in spores of *Aspergillus niger* [38]. Disturbance in fungal metabolism leads to delay germination, resulting in

decreased production of growth-dependent red pigments [37]. Based on Figure 1B, interaction effects of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration and thermal stress time increased red pigment production using levels of both media ($\text{ZnSO}_4 \cdot \text{H}_2\text{O}$ of 0-0.01 g l^{-1} and thermal stress time of 50-70 s).

(A)

Design-Expert@Software
Red Pigment
● Design Points
X1= A: %Soybean meal
Replacement
Actual Factors
B: $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (g l^{-1}) =0.010
C: Thermal Stress (S) =70.00



(B)

Design-Expert@ Software
Red Pigment

Red Pigment
5
0.5

X₁=B: $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (g l^{-1})
X₂=C: Thermal Stress (S)
Actual Factor
A: %Soybean Meal
Replacement=50.00




Figure 1. Effects of various levels of soybean meal replacement (A) and response surface of interaction effects between $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration and thermal stress time (B) on red pigment production by *Monascus purpureus*

In general, 2D and 3D plots were generated for the responses (biomass production) at any of the two independent variables, while keeping the others at their 0 levels. Based on Figure 2A, increases in thermal stress time led to decreases in biomass production. These were further sensitive to higher levels of thermal stress time at 70°C. Therefore, reduction process at higher levels of thermal stress time (greater than 70 s) was significant. This could be linked to delays in germination. Heat stress caused protein denaturation, resulting in polar bond weakening and exposure of hydrophobic groups. Therefore, reactive oxygen species were produced under the heat stress [39]. In


nut shells, reactive oxygen species affects the apical tip growth, hyphal branching and cellular differentiation [40]. Figure 2B reveals that the interaction effects of various $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentrations and soybean meal replacement on biomass production were significant ($P \leq 0.05$) (Table 3). Increases in soybean meal replacement of the media (0-50%) resulted in increases in biomass production up to 8.67 g l^{-1} . However, biomass production decreased by further increases in soybean meal replacement. The soybean meals contains a small quantity of zinc [41]. It could be discussed that the presence of SBM provided a sufficient quantity of zinc to stimulate the microbial growth. Bau et al. reported

that biomass production stopped with the presence of zinc at 2×10^{-3} and 3×10^{-3} M ($0.57\text{-}0.86 \text{ g l}^{-1}$) in liquid media. In addition, the optimum concentration of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ for the growth of *M. purpureus* was 5×10^{-4} M (0.14 g l^{-1}) [20]. Therefore, presence of high quantities of zinc in media

inhibited growth of *M. purpureus*. Gadd reported that a small quantity of zinc was essential for the growth of *M. purpureus*. However, higher quantities of zinc was toxic due to the blockage of functional groups of the enzymes [42].

(A)

Design-Expert® Software
 Biomass
 ● Design Points
 $X_1 = C$: Thermal Stress (S)
 Actual Factors
 A: % Soybean Meal Replacement=50.00
 B: $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (g l^{-1}) =0.010



(B)

Design-Expert® Software
 Biomass
 9.7
 1
 $X_1 = A$: % Soybean Meal Replacement
 $X_2 = B$: $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (g l^{-1})
 Actual Factor
 C : Thermal Stress (S) =70.00




Figure 2. Effects of various levels of thermal stress time (A) and response surface of interaction effects between $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration and soybean meal replacement (B) on biomass contents of *Monascus purpureus*

3.3. Validation of the model under optimized conditions

Validation was carried out under optimized conditions predicted by the model. To validate statistical model and regression equation, optimization program was used with investigation of experiment range using Design Expert Software and the following optimum conditions were achieved: soybean meal replacement of 79.72%, thermal stress time of 81.89 s and $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration of zero. Experimental responses for the red *Monascus* pigment and biomass productions included 4.54 AU ml^{-1} and 4.12 g l^{-1} , respectively. Experimental results were similar to the predicted value of 4.44 AU ml^{-1} . The average yield of red pigment, conversion factor of biomass in red pigment and average cell productivity of the optimized sample included $0.324 \text{ AU ml}^{-1} \text{ day}^{-1}$, 1.10 AU L g^{-1} and $0.292 \text{ g l}^{-1} \text{ day}^{-1}$, respectively. Production of the red *Monascus* pigment at optimized conditions was much higher than those reported by Meinicke et al. [22], Seyedin et al. [23] and Broder and Koehler [43]. Broder and Koehler (2001) produced 1.5 and 3.5 AU ml^{-1} of submerged red pigments using *M. purpureus* and yeast extracts (4 g l^{-1}) [43]. Seyedin et al. achieved red *Monascus* pigment at 2.05 ODU ml^{-1} using yeast extracts. In their study, K_2HPO_4 respectively included 2.75 and 1.5 g l^{-1} using response surface methodology [23].

4. Conclusion

The current study was carried out to assess effects of three important variables (soybean meal replacement, thermal stress time and $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration) on red pigment production by *M. purpureus* using response surface methodology. Results showed that use of soybean meal as a nitrogen source to replace yeast extracts increased red pigment production. Red *Monascus* pigment production was further sensitive to higher levels of thermal stress time at 70°C . Using selected levels of the process variables, a relatively high quantity of the red *Monascus* pigment was achieved with 72% soybean meal replacement and 81.89 s thermal stress time with no $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ concentration in submerge fermentation. Under optimized conditions, quantity of the red Mp, average yield of red *Monascus* pigment (P_M), conversion factor of biomass in red pigments ($Y_{P/X}$) and cell productivity (P_{cells}) were assessed as 4.54 AU ml^{-1} , $0.324 \text{ AU ml}^{-1} \text{ day}^{-1}$, 1.10 AU L g^{-1} and $0.292 \text{ g l}^{-1} \text{ day}^{-1}$, respectively. In general, the lower application of yeast extract may result in decreases in media costs at large scales and can promote commercialization of the red *Monascus* pigment production from SBM. Successful replacement of yeast extract with SBM addresses gaps in other studies for costs of the recently optimized media based on use of yeast extracts for production of red *Monascus* pigment

5. Acknowledgements

The authors thank the laboratory staff within Department of Food Science and Technology, Faculty of Agriculture, Isfahan (Khorasgan) Branch, Isfahan, Islamic Azad University, Isfahan, Iran.

6. Conflict of interest

The authors declare no conflict of interest.

References

1. Panesar R, Kaur S, Panesar PS. Production of microbial pigments utilizing agro-industrial waste: A review. *J Curr Opin Food Sci.* 2015; 1: 70-76. doi: 10.1016/j.cofs.2014.12.002
2. Azlina W, Arul C, Dufoss L. Current perspective of yellowish-orange pigments from microorganisms: A review. *J Clean. Prod.* 2018; 180:168-182. doi: 10.1016/j.jclepro.2018.01.093
3. Agboyibor C, Kong W, Chen D, Zhang A, Niu, S. *Monascus* pigments production, composition, bioactivity and its application: A review. *Biocatal Agric Biotechnol.* 2018 ;16: 433-447. doi: 10.1016/j.bcab.2018.09.012
4. Singgih M, Saraswaty V, Ratnaningrum D, Priatni S, Damayanti S. The influence of temperature and ethanol concentration in monacolin K extraction from *Monascus* fermented rice. *Procedia Chem.* 2014; 9 :242-247. doi: 10.1016/j.proche.2014.05.029
5. Farhadi B, Khosravi-Darani K, Nasernejad B. Enhancement of xanthan production on date extract using response surface methodology. *Asian J Chem.* 2012; 24: 3887-3890.
6. Zhu Z, Zhang F, Wei Z, Ran W, Shen Q. The usage of rice straw as a major substrate for the production of surfactin by *Bacillus amyloliquefaciens* XZ-173 in solid-state fermentation. *J Environ Manage.* 2013; 127: 96-102. doi: 10.1016/j.jenvman.2013.04.017
7. Seyede Reihani FS, Khosravi-Darani K. Mycoprotein production from date waste using *Fusarium venenatum* in a submerged culture. *Appl Food Biotechnol.* 2018; 5: 243-252. doi: 10.22037/afb.v5i4.23139
8. Hashempour-Baltork F, Khosravi-Darani K, Hosseini H, Farshi P, Seyed Reihani SF. Mycoprotein as safe meat substitutes: A review. *J Clean Prod.* 2020; 253: 119958. doi: 10.1016/j.jclepro.2020.119958
9. Mojumdar A, Deka J. Recycling agro-industrial waste to produce amylase and characterizing amylase-gold nanoparticle composite. *Int J Recycle Org Waste Agri.* 2019; 136: 1-7. doi: 10.1007/s40093-019-00298-4
10. Sharma R, Ghoshal G. Optimization of carotenoids production by *Rhodotorula mucilaginosa* (MTCC-1403) using agro-industrial waste in bioreactor: A statistical approach. *J Biotechnol Reports.* 2020; 25: e00407. doi: 10.1016/j.btre.2019.e00407
11. Bakhshi F, Jahadi M. Optimization of the orange pigment production by *Monascus purpureus* from date waste using response surface methodology. *J Food Microbiol.* 2018; 4: 31-39.
12. Silveira ST, Daroit DJ, Brandeli A. Pigment production by *Monascus purpureus* in grape waste using factorial design. *LWT Food Sci Technol.* 2008; 41 :170-174.

- doi: 10.1016/j.lwt.2007.01.013
13. Haque MA, Kachrimanidou V, Koutinas A. Valorization of bakery waste for biocolorant and enzyme production by *Monascus purpureus*. J Biotechnol. 2016; 231 :55-64. doi: 10.1016/j.jbiotec.2016.05.003
 14. Kantifedaki A, Kachrimanidou V, Mallouchos A, Papanikolaou S, Koutinas AA. Orange processing waste valorisation for the production of bio-based pigments using the fungal strains *Monascus purpureus* and *Penicillium purpurogenum*. J Clean Prod. 2018; 185 :882-890. doi: 10.1016/j.jclepro.2018.03.032
 15. Chiu SW, Chan SM. Production of pigments by *Monascus purpureus* using sugar-cane bagasse in roller bottle cultures. World J Microbiol Biotechnol. 1992; 8 :68-70. doi: 10.1007/BF01200689
 16. Silveira ST, Daroit DJ, Sant'Anna V, Brandelli A. Stability modeling of red pigments produced by *Monascus purpureus* in submerged cultivations with sugarcane bagasse. J Food Bioproc Tech. 2013; 6 :1007-1114. doi: 10.1007/s11947-011-0710-8
 17. Duraklı-Velioglu S, Boyacı IH, Şimşek O, Gumuş T. Optimizing a submerged *Monascus* cultivation for production of red pigment with bug damaged wheat using artificial neural networks. J Food Sci Biotechnol. 2013; 22 :1639-1648. doi: 10.1007/s10068-013-0261-z
 18. Yuliana A, Singgih M, Julianti E, Blanc P. Derivates of azaphilone *Monascus* pigments: A review. J Biocatal Agric Biotechnol. 2017; 9: 183-194. doi: 10.1016/j.bcab.2016.12.014
 19. Chen M, Johns MR. Effect of pH and nitrogen source on pigment production by *Monascus purpureus*. J Appl Microbiol Biotechnol. 1993; 40: 132-138. doi: 10.1007/BF00170441
 20. Bau YS, Wong HC. Zinc effects on growth, pigmentation and antibacterial activity of *Monascus purpureus*. J Physiol Plant. 1979; 46: 63-67. doi: 10.1111/j.1399-3054.1979.tb03187.x
 21. da Costa JPV, Vendruscolo F. Production of red pigments by *Monascus ruber* CCT 3802 using lactose as substrate. Biocatal Agric Biotechnol. 2017; 11: 50-55. doi: 10.1016/j.bcab.2017.05.010
 22. Meinicke RM, Vendruscolo F, Moritz DE, de Oliveira D, Schmidell W, Samohyl RW, Ninow JL. Potential use of glycerol as substrate for the production of red pigments by *Monascus ruber* in submerged fermentation. Biocatal Agric Biotechnol. 2012; 1:238-242. doi: 10.1016/j.bcab.2012.03.001
 23. Seyedin A, Yazdian F, Hatamian-Zarmi A, Rasekhi B, Mirderkvand M. Natural red pigment production by *Monascus purpureus*: Bioreactor yield improvement through statistical analysis. J Appl Food Biotechnol. 2015; 2(2): 23-30. doi: 10.22037/afb.v.2i2.7457
 24. Sharmila G, Nidhi B, Muthukumaran C. Sequential statistical optimization of red pigment production by *Monascus purpureus* (MTCC 369) using potato powder. J Ind Crop Prod. 2013; 44: 158-164. doi: 10.1016/j.indcrop.2012.11.007
 25. Babitha S, Soccol CR, Pandey A. Effect of stress on growth, pigment production and morphology of *Monascus* sp. in solid cultures. J Basic Microbiol. 2007; 47: 118-126. doi: 10.1002/jobm.200610261
 26. Thakur SA, Nemade SN, Sharanappa A. Solid state fermentation of overheated soybean meal (waste) for production of protease using *Aspergillus Oryzae*. Int J Innov Res Sci Eng Technol. 2015; 4:18456-18461. doi: 10.15680/IJRSET.2015.0401008
 27. Silva F, Daniel F. Characterization and stability of bioactive compounds from soybean meal. J Food Sci Technol. 2015; 63: 992-1000. doi: 10.1016/j.lwt.2015.04.032
 28. Zhang K, Yu C, Yang S. Effects of soybean meal hydrolysate as the nitrogen source on seed culture morphology and fumaric acid production by *Rhizopus oryzae*. J Process Biochem. 2015; 50(2): 173-179. doi: 10.1016/j.procbio.2014.12.015
 29. Lujan-Rhenals D, Morawicki R, Gbur E, Ricke S. Fermentation of soybean meal hydrolyzates with *Saccharomyces cerevisiae* and *Zymomonas mobilis* for ethanol production. J Food Sci. 2015; 80: 1512-1518. doi: 10.1111/1750-3841.12907
 30. Zhang J, Wang YL, Lu L, Zhang BB, Xu GR. Enhanced production of Monacolin K by addition of precursors and surfactants in submerged fermentation of *Monascus purpureus* 9901. J Biotechnol Appl Biochem. 2014; 61(2): 202-207. doi: 10.1002/bab.1154
 31. Kawahara Y, Kikuchi Y, Kurahashi, O, Kimura E, Nakamatsu T, Goto S. Stress-Resistant Microorganism and Method of Producing Fermentative Product. United States Patent. 2002; 6: 338-956.
 32. Stroher R, Stenzel M, Pereira NC, Zanin GM. Enzymatic extraction of protein from toasted and not toasted soybean meal. J Food Sci. 2011; 1: 463-469. doi: 10.1016/j.profoo.2011.09.071
 33. Kongruang S. Growth kinetic of biopigment production by Thai isolated *Monascus purpureus* in a stirred tank bioreactor. J Ind Micro Biotechnol. 2011; 38: 93-99. doi: 10.1007/s10295-010-0834-2
 34. Jahadi M, Khosravi-Darani K, Ehsani MR, Mozafari MR, Saboury AA, Pourhosseini PS. The encapsulation of flavourzyme in nanoliposome by heating method. J Food Sci Technol. 2015; 52(4): 2063-2072. doi: 10.1007/s13197-013-1243-0
 35. Hajjaj H, Francois J-M, Goma G, Blanc PJ. Effect of amino acids on red pigments and citrinin production in *Monascus ruber*. J Food Sci. 2012; 77(3): 156-159. doi: 10.1111/j.1750-3841.2011.02579.x
 36. Adiga PR, Sastry KS, VenkatasuBramanyan V, Sarma PS. Interrelationships in trace-element metabolism in *Aspergillus niger*. Biochem J. 1961; 8(3): 545-550. doi: 10.1042/bj0810545
 37. Vendruscolo F, Schmidell W, Oliveira D, Ninow J. Kinetic of orange pigment production from *Monascus ruber* on submerged fermentation. J Bioproc Biosyst Eng. 2017; 40(1): 115-121. doi: 10.1007/s00449-016-1679-5
 38. Abrashev R, Dolashka P, Cristova R, Stefanova L, Angelova M. Role of antioxidant enzymes in survival of conidiospores of *Aspergillus niger* 26 under conditions of temperature stress. J Appl Microbiol. 2005; 99(4): 902-909. doi: 10.1111/j.1365-2672.2005.02669.x
 39. Fridovich I. Superoxide radical and superoxide dismutases. J Annu Rev Biochem. 1995; 64: 97-112.

- doi: 10.1146/annurev.bi.64.070195.000525
40. Li Q, Harvey LM, McNeil B. Oxidative stress in industrial fungi. *J Crit Rev Biotechnol.* 2009; 29: 199-213.
doi: 10.1080/07388550903004795
41. Edward G, Perkins. Composition of Soybeans and Soybean Products. In: David R, Erickson. *Practical Handbook of Soybean Processing and Utilization*, 1st Edition. Academic Press and AOCS Press, 1995: pp.9-28.
doi:10.1016/B978-0-935315-63-9.50006-1
42. Gadd GM. 'Interactions of fungip with toxic metals. *J New Phytol.* 1993; 124: 25-60.
doi: 10.1007/978-1-4899-0981-7_28
43. Broder CU, Koehler PE. Pigments produced by *Monascus purpureus* with regard to quality and quantity. *J Food Sci.* 1980; 45: 567-569.
doi: 10.1111/j.1365-2621.1980.tb04102.x

بهینه سازی کشت غوطه‌وری موناسکوس پورپورئوس بر روی کنجاله سویا به منظور تولید رنگدانه قرمز با روش سطح پاسخ

حدیثه کیوانی، مهشید جهادی*، نفیسه قاسمی سپرو

گروه علوم و صنایع غذایی، دانشکده کشاورزی، دانشگاه آزاد اسلامی، واحد اصفهان (خوراسگان)، اصفهان، ایران.

چکیده

سابقه و هدف: موناسکوس پورپورئوس فارچی رشته‌ای با توانایی تولید رنگدانه‌هایی دارای خواص درمانی می‌باشد. رنگدانه‌های قرمز به ویژه به عنوان افزودنی، آنتی‌اکسیدان، نگهدارنده و جایگزین نیترات در صنایع غذایی مورد استفاده قرار می‌گیرند. به منظور کاهش هزینه تخمیر، ضایعات کشاورزی_صنعتی مانند کنجاله سویا به عنوان رشدمایه^۱ مورد استفاده قرار می‌گیرند. در مطالعه حاضر، تولید رنگدانه قرمز توسط موناسکوس پورپورئوس بر روی کنجاله سویا بهینه‌سازی شد.

مواد و روش‌ها: در این مطالعه، تولید رنگدانه قرمز موناسکوس توسط کشت غوطه‌ور موناسکوس پورپورئوس ATCC 16362 با استفاده از کنجاله سویا به عنوان منبع نیتروژن به جای عصاره‌های مخمر انجام شد. طراحی مرکب مرکزی به منظور ارزیابی سطح بهینه جایگزینی کنجاله سویا (۰-۱۰۰٪)، غلظت $ZnSO_4 \cdot 7H_2O$ ($0-0.02 \text{ g l}^{-1}$) و مدت زمان تنش حرارتی سوسپانسیون اسپور در $70^\circ C$ (۵۰-۹۰ s) مورد استفاده قرار گرفت. تولید رنگدانه قرمز موناسکوس و زی‌توده^۲ به عنوان پاسخ‌های وابسته مورد بررسی قرار گرفتند.

یافته‌ها و نتیجه‌گیری: بیشینه تولید رنگ قرمز موناسکوس ($4/54 \text{ AU ml}^{-1}$) تحت شرایط جایگزینی ۷۹/۷۲ درصدی کنجاله سویا، غلظت $0-0.02 \text{ g l}^{-1}$ $7H_2O \cdot ZnSO_4$ و تنش حرارتی سوسپانسیون ۸۱/۸۹ ثانیه به دست آمد. متوسط راندمان رنگدانه قرمز موناسکوس (P_M)، ضریب تبدیل^۳ زی‌توده در رنگدانه قرمز ($Y_{P/X}$)، و بهره‌وری سلولی به ترتیب $0.324 \text{ AU ml}^{-1} \text{ day}^{-1}$ ، $1/10 \text{ AU L g}^{-1}$ و $0.292 \text{ g l}^{-1} \text{ day}^{-1}$ بود. نتایج مطالعه حاضر نشان داده است که ترکیب کنجاله سویا و عصاره مخمر به عنوان منبع نیتروژن برای تولید رنگدانه قرمز موناسکوس توسط موناسکوس پورپورئوس مفید می‌باشد.

واگان کلیدی: طراحی مرکب مرکزی، موناسکوس پورپورئوس، رنگدانه قرمز، روش سطح پاسخ، کنجاله سویا

تعارض منافع: نویسندگان اعلام می‌کنند که هیچ نوع تعارض منافی مرتبط با انتشار این مقاله ندارند.

تاریخچه مقاله

دریافت ۵ فوریه ۲۰۲۰
داوری ۴ آپریل ۲۰۲۰
پذیرش ۱۷ آپریل ۲۰۲۰

واژگان کلیدی

- طراحی مرکب مرکزی
- موناسکوس پورپورئوس
- رنگدانه قرمز
- روش سطح پاسخ
- کنجاله سویا

*نویسنده مسئول

مهشید جهادی

دانشگاه آزاد اسلامی، واحد اصفهان
(خوراسگان)، ارغوانیه، خیابان جی،
خوراسگان، اصفهان، ایران.

تلفن: ۹۸-۳۴-۳۵۳۴۲۲۲۱+

دورنگار: ۹۸-۳۱-۳۵۳۵۴۰۶۰+

پست الکترونیک:

mahshidjahadi@gmail.com

^۱ Substrates هر ماده‌ای که نیاز غذایی و بستر زیست‌ریزاندامگان را فراهم کند.

^۲ Biomass

^۳ Conversion factor