

Research Article

Mechanical Occlusion Chemically Assisted Ablation (MOCA) for Saphenous Vein Insufficiency: A Meta-Analysis of a Randomized Trial

Johanes Nugroho ^{1,2}, Ardyan Wardhana ³, and Cornelia Ghea ³

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

²Dr. Soetomo General Hospital, Surabaya, Indonesia

³PILAR Research and Education, Cambridge, UK

Correspondence should be addressed to Johanes Nugroho; j.nugroho.eko@fk.unair.ac.id

Received 26 October 2019; Revised 24 December 2019; Accepted 13 January 2020; Published 29 January 2020

Academic Editor: Alberto Caggiati

Copyright © 2020 Johanes Nugroho et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Purpose. A previous meta-analysis has conducted nonrandomized trials for mechanochemical ablation (MOCA). Since medium-term follow-up data from randomized clinical trials (RCTs) are becoming available, we chose to perform a meta-analysis of RCTs to assess the efficacy and safety of MOCA for saphenous vein insufficiency. **Methods.** A systematic search of all RCTs comparing the anatomical success of MOCA for saphenous vein insufficiency to thermal ablation was performed using the PubMed and Cochrane Library databases. We employed the Mantel-Haenszel random-effects meta-analysis of outcomes using RevMan 5.3. **Results.** Four studies (615 patients) were included in this meta-analysis. The MOCA group had 93.4% and 84.5%, whereas the thermal ablation group had 95.8% and 94.8% of anatomical success rate at 1 month (short-term) and a period of more than 6 months but less than 1-year follow-up (mid-term), respectively. According to intention-to-treat analysis, there were similar anatomical successes in MOCA and thermal ablation groups at the short-term follow-up (low-quality evidence; relative risk (RR) = 0.98 (95% CI, 0.94–1.03); $P = 0.44$; $I^2 = 53\%$). The estimated effect of MOCA on anatomical success showed a statistically significant reduction at the mid-term follow-up (moderate-quality evidence; RR = 0.89 (95% CI, 0.84–0.95); $P = 0.0002$; $I^2 = 0\%$). MOCA had fewer incidence of nerve injury, deep vein thrombosis, and skin burns compared to the thermal ablation procedure (low-quality evidence; RR = 0.33 (95% CI, 0.09–1.28); $P = 0.11$; $I^2 = 0\%$). **Conclusion.** MOCA offered fewer major complications but lesser anatomical success at the period of more than 6 months but less than 1-year follow-up than thermal ablation. **Trial Registration.** This trial is registered with UMIN Clinical Trial Registry (UMIN ID 000036727).

1. Introduction

Saphenous vein insufficiency is the most common chronic venous incompetence of lower limbs. It affects more than 30% of adults [1]. The first-line treatment of venous incompetence is thermal ablation, e.g., radiofrequency ablation (RFA) and endovenous laser ablation (EVLA) [2]. However, the use of heat energy in thermal ablation was associated with complications such as nerve injury and skin burns [3, 4].

To avoid thermal energy complications, few novel treatments such as sclerotherapy, cyanoacrylate, and mechanochemical ablation (MOCA) have been introduced. One of MOCA device is ClariVein (Merit Medical, South Jordan,

Utah) that combines mechanical injury to the venous endothelium using a rotating wire with simultaneous delivery and dispersion of a liquid sclerosant [5]. The liquid sclerosant causes irreversible damage to the cellular membranes of the endothelium, resulting in the fibrosis of veins [6]. Another technique to produce a nontumescent sclerosant-assisted ablation is by means of Flebogrif (Balton, Poland), a device that is scratching the endothelium lining by a dedicated catheter.

A previous meta-analysis has conducted nonrandomized trials for MOCA [7]. It is important to study the use of MOCA since medium-term follow-up data on randomized clinical trials are now becoming available. Therefore, we

performed a meta-analysis of randomized trials aiming to assess the safety and efficacy of MOCA by comparing thermal ablation for saphenous vein insufficiency.

2. Methods

All RCTs comparing the anatomical success of MOCA for saphenous vein insufficiency (great saphenous vein (GSV), small saphenous vein (SSV), or both) to the success rate of a thermal ablation procedure were eligible for inclusion in this review. All trials comparing MOCA to treatments other than thermal ablation were excluded. A systematic search of the literature was undertaken on July 7, 2019, after receiving an approval from the Institutional Review Board. Two different databases (PubMed and Cochrane Library) were used to perform a systematic search of all the literature without language restriction. Search terms were differently spelled text words or abbreviations, such as “varicose,” “saphenous incompetence,” “saphenous insufficiency,” “saphenous varicose veins,” “saphenous reflux,” “saphenous vein,” “endovenous ablation,” “clarivein,” “MOCA,” “mechanochemical ablation,” “randomized,” and “rct” in the title, abstract, and medical subject heading (MeSH). Reference lists of the included studies were also evaluated to identify additional relevant studies. We followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines for research reporting.

Two investigators independently screened and assessed titles and abstracts before full-text retrieval. The full papers that potentially met the inclusion/exclusion criteria were reviewed by the two authors for final inclusion. Subsequently, two investigators extracted the data, which included authors, year of publication, detailed intervention agent, veins treated (either GSVs or SSVs), postprocedural management, number of patients, primary outcomes, and adverse events. The corresponding author was contacted to obtain any incomplete data. All extracted data were recorded with a dedicated data extraction form on an Excel spreadsheet.

The primary outcome in our meta-analysis was anatomical success that was defined either as complete occlusion of the saphenous vein or at least proximal occlusion (>5 cm proximally occluded with >5 cm distally open) on duplex ultrasound (DUS) imaging. The primary outcome was categorized at the short-term follow-up (1 month) and mid-term follow-up (≥6 months but ≤1 year). Secondary outcomes were phlebitis and major adverse event rates, including nerve injury, DVT, and skin burns. Nerve injury was reported differentially in the studies as sensory disturbance, paresthesia, or numbness. Other minor complications (e.g., hematoma, lumps, and hyperpigmentation) were excluded from the analysis.

Two authors independently assessed the methodological quality of the articles using the Risk of Bias Tool (Cochrane collaboration). They employed a GRADE approach to evaluate the quality of evidence [8, 9]. Factors assessed for the risk of bias were as follows [1]: randomization [2], adherence to intervention [3], measurement [4], missing outcome data, and reporting [5]. We evaluated the GRADE while taking into account the following factors: risk of bias, imprecision,

indirectness of evidence, inconsistency in results among included studies, and reporting bias.

We conducted the meta-analysis using relative risk (RR) for anatomical success rates at short-term and mid-term follow-up periods. We employed the Mantel–Haenszel random-effects method using Review Manager (RevMan v5.3 2014). We conducted an intention-to-treat (ITT) analysis using imputation data according to event rates among completers in the separate groups [10]. Available case analysis was further performed as a sensitivity analysis. We evaluated between and within-study heterogeneity using the I^2 statistic.

3. Results

A total of 100 studies were identified after our initial search was completed (45 citations in PubMed and 55 in Cochrane Library, as presented in Figure 1). Six studies were further retrieved to review the full text. However, two studies were excluded because one lacked a control and one was an in-abstract study [11, 12]. Therefore, four studies were finally included in our analysis [13–16]. Attempts to contact authors to acquire unpublished data were uniformly unsuccessful. As such, only information available from the publication was used. Table 1 summarizes the clinical characteristics of the included studies.

The Cochrane risk of bias analysis is shown in Figure 2. There was a low risk of a randomization process in all included trials. Blinding to participants and the delivering clinicians was not performed in all included trials. There was no placebo included for comparison. However, there was assessor-blinding when evaluating the anatomical success using DUS imaging. One trial was stopped early because reimbursement of MOCA was suspended [13]. Participants who were not followed up ranged from 6% to 34%, and there were no documented reasons for loss to follow-up. All trials reported the results from available case analyses and not ITT analysis.

Table 2 shows the quality of evidence, and the GRADE approach was used for each outcome. We did not downgrade from the risk of bias aspect because most information was from studies that were at low risk of bias. We downgraded one level for the evidence of anatomical success at the short-term follow-up because there was substantial heterogeneity. Indirectness did not appear to be an issue in all outcomes. We downgraded the evidence of major complications for the imprecision aspect because there was a wide CI that included no important effect and RR reduction greater than 25%. The publication bias was assessed for all outcomes because there was asymmetry from a visual inspection of the funnel plot (Figure 3). This finding was not further evaluated by conducting Egger’s test because there were less than 10 included trials.

The MOCA group had 93.4% of anatomical success rate, whereas the thermal ablation group had 95.8% at the 1-month follow-up. According to the meta-analysis using ITT as shown in Table 3, there were similar anatomical successes in the MOCA and thermal ablation groups at the 1-month follow-up (4 trials; RR = 0.98 (95% CI, 0.94–



PRISMA 2009 Flow Diagram




FIGURE 1: PRISMA flow diagram.

1.03); $P = 0.44$); however, there was substantial heterogeneity ($I^2 = 53\%$; $P = 0.09$). At the mid-term follow-up (≥ 6 months but ≤ 1 year), the MOCA group had 84.5% of anatomical success rate, whereas the thermal ablation group had 94.8% (Table 4). The estimated effect of MOCA on anatomical success showed a statistically significant reduction at the mid-term follow-up (3 trials; $RR = 0.89$ (95% CI, 0.84–0.95); $P = 0.0002$), without substantial heterogeneity ($I^2 = 0\%$; $P = 0.52$).

We performed sensitivity analysis using the available case analyses. The direction of effect did not change for anatomical success rates. MOCA did not decrease anatomical success compared to thermal ablation at the 1-month follow-up (4 trials; $RR = 0.98$ (95% CI, 0.93–1.03); $P = 0.46$); however, there was substantial heterogeneity ($I^2 = 56\%$; $P = 0.08$). The RR for anatomical success at the mid-term follow-up (≥ 6 months but ≤ 1 year) was significantly lower after treatment with MOCA than that with thermal ablation (3 trials;

$RR = 0.89$ (95% CI, 0.84–0.96); $P = 0.001$), without substantial heterogeneity ($I^2 = 0\%$; $P = 0.67$).

Table 5 shows the details of adverse events in the four included trials. MOCA had fewer major complications compared to thermal ablation procedures (4 trials, $RR = 0.33$ (95% CI, 0.09–1.28); $P = 0.11$; $I^2 = 0\%$). However, participants in the MOCA group had a higher risk of phlebitis than the thermal ablation group (4 trials, $RR = 1.39$ (95% CI, 0.67–2.86); $P = 0.37$; $I^2 = 0\%$).

4. Discussion

We report that MOCA had a similar anatomical success rate with low-quality evidence at the 1-month follow-up, but there was a reduction of the anatomical success rate with moderate-quality evidence at the period of more than 6 months but less than 1-year follow-up compared with thermal ablation. Previous meta-analyses that included

TABLE 1: The characteristics of the included trials.

Author	Vein insufficiency	Intervention	Control	Postprocedural management	DUS scan finding	Follow-up period (percentage lost-to follow-up in groups MOCA and thermal ablation)	Adverse event identification
Holewijn et al. 2017 [13]	GSV	MOCA using ClariVein and polidocanol 3% 2 ml for the first 10-15 cm and 1.5% for the remainder. Pullback rate 7 s/cm	RFA using Closurefast	Phlebectomy as indicated, stocking compression for 2 weeks	Complete occlusion, partial recanalization >10 cm, complete recanalization	1 month (0% & 0%), 12 months (21.4% & 30.1%), 24 months (26.2% & 21.4%)	Major: DVT, pulmonary embolism, skin burn, saphenous neuralgia Minor: phlebitis, induration, hematoma, hyperpigmentation
Vähäaho et al. 2019 [14]	GSV	MOCA using ClariVein and STS 1.5%. Pullback rate was not defined	RFA using Closurefast or EVLA using ELVes 1470-nm diode radial laser	Phlebectomy for every patient. Other postprocedural management was not defined	Complete occlusion, proximal occlusion, distal occlusion, complete recanalization	1 month (0% & 0%), 12 months (18.2% & 6.1%)	Major: DVT, nerve injury Minor: hyperpigmentation, hematoma, phlebitis, lumps
Bootun et al. 2016 [15]	GSV, SSV	MOCA using ClariVein and STS 2%. Pullback rate was not defined	RFA using Venefit	Phlebectomy as indicated, stocking compression for 2 weeks	Complete occlusion, proximal occlusion, distal occlusion, complete recanalization	1 month (13.3% & 55.9%)	Major: DVT Minor: phlebitis
Lane et al. 2016 [16]	GSV, SSV	MOCA using ClariVein and STS 2%. Pullback rate 7 s/cm	RFA using Venefit	Phlebectomy as indicated, LMWH administration, stocking compression for 2 weeks	Complete occlusion, proximal occlusion, distal occlusion, complete recanalization	1 month (16.9% & 26.8%), 6-months (25.3% & 28%)	Major: DVT, sensory disturbance Minor: phlebitis, hyperpigmentation

	Randomization	Adhering to intervention	Missing outcome data	Measurement	Reporting
Bootun et al. 2016	Low risk	Low risk	Unclear risk	Low risk	Low risk
Holewijn et al. 2017	Low risk	Low risk	Unclear risk	Low risk	Low risk
Lane et al. 2016	Low risk	Low risk	Unclear risk	Low risk	Low risk
Vahaaho et al. 2019	Low risk	Low risk	Unclear risk	Low risk	Low risk



 = Low risk
 = Unclear risk

FIGURE 2: Risk of bias assessment.

TABLE 2: GRADE assessment.

Comparison	Outcomes	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication of bias	GRADE
Anatomical success	Short-term	No serious	Some	No at all	No	Presence	Low
	Mid-term	No serious	No	No at all	No	Presence	Moderate
Complication	Major complication	No serious	No	No at all	Some	Presence	Low
	Phlebitis	No serious	No	No at all	No	Presence	Moderate




FIGURE 3: Funnel plot of comparisons: mechanochemical ablation vs thermal ablation intention-to-treat analysis, outcome: anatomical success short-term.

nonrandomized trials demonstrated that MOCA could be considered an effective treatment for saphenous vein insufficiency; however, these results were based on a pooled analysis of low-quality data [17]. In previous studies, the pooled ana-

tomical success rate at the mid-term follow-up was 91%, 97.1%, and 98.5% for MOCA, RFA, and EVLA, respectively [7, 18]; the rates obtained in our study were lower: 84.5% for MOCA and 94.8% for thermal ablation. Blinding to the

TABLE 3: Summary of findings.

		Studies	Participants	Effect estimate	Heterogeneity
ITT analysis	Anatomical success at short-term	4	615	RR 0.98 [0.94, 1.03]; $P = 0.44$	$P = 0.09$; $I^2 = 53\%$
	Anatomical success at mid-term	3	496	RR 0.89 [0.84, 0.95]; $P = 0.0002$	$P = 0.53$; $I^2 = 0\%$
Available case analysis	Anatomical success at short-term	4	532	RR 0.98 [0.93, 1.03]; $P = 0.46$	$P = 0.08$; $I^2 = 56\%$
	Anatomical success at mid-term	3	386	RR 0.89 [0.84, 0.96]; $P = 0.001$	$P = 0.67$; $I^2 = 0\%$
Adverse events	Major complication	4	615	RR 0.33 [0.09, 1.28]; $P = 0.11$	$P = 0.73$; $I^2 = 0\%$
	Phlebitis	4	615	RR 1.39 [0.67, 2.86]; $P = 0.37$	$P = 0.58$; $I^2 = 0\%$

TABLE 4: Anatomical success and complication rates in each study.

Studies	Anatomical success rates		DVT		Sensory disturbances	
	MOCA	Thermal ablation	MOCA	Thermal ablation	MOCA	Thermal ablation
	Short-term		At the end of study		At the end of study	
Bootun et al. 2016 [15]	91.7%	91.5%	0%	1.7%	0%	0%
Holewijn et al. 2017 [13]	91.3%	99.0%	0%	0%	1.0%	2.9%
Lane et al. 2016 [16]	92.8%	91.5%	1.2%	1.2%	0%	0%
Vähäaho et al. 2019 [14]	100.0%	100.0%	0%	0%	0%	7.6%
Total	93.4%	95.8%	0.7%	1.4%	0.4%	3.2%
	Mid-term					
Lane et al. 2016 [16]	86.7%	92.7%				
Vähäaho et al. 2019 [14]	86.4%	100.0%				
Holewijn et al. 2017 [13]	81.6%	93.2%				
Total	84.5%	94.8%				

TABLE 5: Adverse events in the included trials.

Adverse events	Phlebitis		DVT		Nerve injury	
	MOCA	Thermal ablation	MOCA	Thermal ablation	MOCA	Thermal ablation
Holewijn et al. 2017 [13]	12	8	0	0	1	3
Vähäaho et al. 2019 [14]	1	0	0	0	0	5
Bootun et al. 2016 [15]	0	2	0	1	Not reported	Not reported
Lane et al. 2016 [16]	3	2	1	1	0	0

assessor may have contributed to the lower anatomical success rate observed in our study.

The major benefit of MOCA reported in previous reviews was pain intensity reduction during the procedure [17, 18]. However, we did not perform an analysis of pain intensity because there was a high risk of measurement bias from no participant blinding in the available trials. The potential benefits in reducing the risk of nerve injury might be of considerable clinical importance for MOCA. Our meta-analysis indicated that MOCA had a lower risk of major complications compared with thermal ablation. Nerve injury was seen in 0.4% and 3.2% cases after MOCA and thermal ablation, respectively.

There was disparity in terms of anatomical success measure definitions, postprocedure management, and technical aspects of the MOCA procedure. Holewijn et al.

reported different results from the others as they used a different sclerosant and categorized DUS image findings into 3 categories instead of 4 [13]. The types of sclerosant used by different studies may have influenced anatomical success. There was substantial heterogeneity in anatomical success at the short-term follow-up in our study. Removing those trials improved the heterogeneity without changing the direction of the effect. Therefore, the heterogeneity in our study might be due to various types of sclerosants and definitions of anatomical success.

There were a few limitations in our meta-analysis. First, we evaluated efficacy only from the anatomical success rate aspect, not from pain reduction, clinical success, or quality of life aspects. Second, there was a wide CI around the effect estimate of adverse event outcomes because there were few events in this meta-analysis. Future trials are required. To

date, there is one unpublished trial and one ongoing trial investigating the anatomical success of MOCA compared with thermal ablation [12, 19]. Lastly, the need of data at more than 1 year of outcome measurement at least is suggested.

5. Conclusion

Pooled analysis from limited trials demonstrated that MOCA offered fewer major complications (DVT, nerve injury, and skin burns) but lesser anatomical success at the period of more than 6 months but less than 1-year follow-up than thermal ablation. More RCTs together with proper homogeneous data collection are needed to reach optimal information size.

Data Availability

The data supporting this meta-analysis are from previously reported studies and datasets, which have been cited. The processed data are available in the supplementary files (available here).

Ethical Approval

The study was approved by the Dr. Soetomo General Hospital Surabaya Ethical Committee in Health Research (1286/KEPK/VII/2019).

Conflicts of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be considered as a potential conflict of interests.

Acknowledgments

This work did not receive specific funding but was performed as part of Johannes Nugroho employment at the Department of Cardiology and Vascular Medicine, Universitas Airlangga/Dr. Soetomo General Hospital, East Java, Indonesia.

Supplementary Materials

Forest plot of all outcomes in meta-analysis. (*Supplementary Materials*)

References

- [1] C. J. Evans, P. L. Allan, A. J. Lee, A. W. Bradbury, C. V. Ruckley, and F. G. Fowkes, "Prevalence of venous reflux in the general population on duplex scanning: the Edinburgh vein study," *Journal of Vascular Surgery*, vol. 28, no. 5, pp. 767–776, 1998.
- [2] G. Marsden, M. Perry, K. Kelley, A. H. Davies, and on behalf of the Guideline Development Group, "Diagnosis and management of varicose veins in the legs: summary of NICE guidance," *BMJ*, vol. 347, no. 1, article f4279, 2013.
- [3] J. I. Almeida, J. Kaufman, O. Göckeritz et al., "Radiofrequency endovenous ClosureFAST versus laser ablation for the treatment of great saphenous reflux: a multicenter, single-blinded, randomized study (RECOVERY study)," *Journal of Vascular and Interventional Radiology*, vol. 20, no. 6, pp. 752–759, 2009.
- [4] A. L. Kerver, A. C. van der Ham, H. P. Theeuwes et al., "The surgical anatomy of the small saphenous vein and adjacent nerves in relation to endovenous thermal ablation," *Journal of Vascular Surgery*, vol. 56, no. 1, pp. 181–188, 2012.
- [5] R. L. Mueller and J. K. Raines, "ClariVein mechanochemical ablation: background and procedural details," *Vascular and Endovascular Surgery*, vol. 47, no. 3, pp. 195–206, 2013.
- [6] R. R. van Eekeren, J. L. Hillebrands, K. van der Sloot, J. P. de Vries, C. J. Zeebregts, and M. M. Reijnen, "Histological observations one year after mechanochemical endovenous ablation of the great saphenous vein," *Journal of Endovascular Therapy*, vol. 21, no. 3, pp. 429–433, 2014.
- [7] M. E. Witte, C. J. Zeebregts, G. J. de Borst, M. M. P. J. Reijnen, and D. Boersma, "Mechanochemical endovenous ablation of saphenous veins using the ClariVein: a systematic review," *Phlebology: The Journal of Venous Disease*, vol. 32, no. 10, pp. 649–657, 2017.
- [8] J. P. T. Higgins, J. A. C. Sterne, J. Savović et al., "A revised tool for assessing risk of bias in randomized trials," *Cochrane Database of Systematic Reviews*, vol. 10, no. 1, 2016.
- [9] R. Ryan and S. Hill, *How to GRADE the quality of the evidence Version 3.0*, Cochrane Consumers and Communication Group, 2018, July 2019, https://cc.cochrane.org/sites/cc.cochrane.org/files/public/uploads/how_to_grade.pdf.
- [10] H. JPT and S. Green, *Cochrane handbook for systematic reviews of Interventions Version 5.1.0*, The Cochrane Collaboration, 2011, July 2019, <https://handbook-5-1.cochrane.org/>.
- [11] Y. L. Lam, I. M. Toonder, and C. H. Wittens, "Clarivein mechano-chemical ablation an interim analysis of a randomized controlled trial dose-finding study," *Phlebology: The Journal of Venous Disease*, vol. 31, no. 3, pp. 170–176, 2016.
- [12] C. C. Leung, D. Carradice, T. Wallace, and I. C. Chetter, "Endovenous laser ablation versus mechanochemical ablation with ClariVein® in the management of superficial venous insufficiency (LAMA trial): study protocol for a randomised controlled trial," *Trials*, vol. 17, no. 1, article 421, 2016.
- [13] S. Holewijn, R. R. van Eekeren, A. Vahl et al., "Two-year results of a multicenter randomized controlled trial comparing Mechanochemical endovenous Ablation to RADiOfrequeNcy Ablation in the treatment of primary great saphenous vein incompetence (MARADONA trial)," *Journal of Vascular Surgery: Venous and Lymphatic Disorders*, vol. 7, no. 3, pp. 364–374, 2017.
- [14] S. Vähäaho, O. Mahmoud, K. Halmesmäki et al., "Randomized clinical trial of mechanochemical and endovenous thermal ablation of great saphenous varicose veins," *The British Journal of Surgery*, vol. 106, no. 5, pp. 548–554, 2019.
- [15] R. Bootun, T. R. Lane, B. Dharmarajah et al., "Intra-procedural pain score in a randomised controlled trial comparing mechanochemical ablation to radiofrequency ablation: the Multicentre Venefit versus ClariVein for varicose veins trial," *Phlebology: The Journal of Venous Disease*, vol. 31, no. 1, pp. 61–65, 2016.
- [16] T. Lane, R. Bootun, B. Dharmarajah et al., "A multi-centre randomised controlled trial comparing radiofrequency and mechanical occlusion chemically assisted ablation of varicose veins – final results of the Venefit versus Clarivein for varicose veins trial," *Phlebology: The Journal of Venous Disease*, vol. 32, pp. 89–98, 2016.
- [17] J. J. Sun, M. M. Chowdhury, U. Sadat, P. D. Hayes, and T. Y. Tang, "Mechanochemical ablation for treatment of truncal venous insufficiency: a review of the current literature,"

Journal of Vascular and Interventional Radiology, vol. 28, no. 10, pp. 1422–1431, 2017.

- [18] D. Boersma, V. N. Kornmann, R. R. van Eekeren et al., “Treatment modalities for small saphenous vein insufficiency: systematic review and metaanalysis,” *Journal of Endovascular Therapy*, vol. 23, no. 1, pp. 199–211, 2016.
- [19] D. Boersma, R. R. J. P. van Eekeren, H. J. C. Kelder et al., “Mechanochemical endovenous ablation versus radiofrequency ablation in the treatment of primary small saphenous vein insufficiency (MESSI trial): study protocol for a randomized controlled trial,” *Trials*, vol. 15, no. 1, p. 421, 2014.

International Journal of Vascular Medicine

- Journal overview 
- For authors
- For reviewers
- For editors
- Table of Contents
- Special Issues 

Recently Published

A Review of Risk Factors and Predictors for Hemorrhagic Transformation in Patients with Acute Ischemic Stroke


Sheha E. Thomas | Noorine Plumber | ... | Vasavi Gorantla | + 1 Others

[Read the full article](#)

Journal metrics


Acceptance rate	19%
Submission to final decision	59 days
Acceptance to publication	26 days
CiteScore	2.400
Journal Citation Indicator	0.230
Impact Factor	*

APC **\$900**

[Submit](#) 

Journal profile

International Journal of Vascular Medicine publishes articles focusing on vascular biology and physiology, vascular imaging, endovascular intervention, as well as prevention and treatment of vascular disease.

[About this journal](#) 


Editor spotlight

International Journal of Vascular Medicine maintains an Editorial Board of practicing researchers from around the world, to ensure manuscripts are handled by editors who are experts in the field of study.

[Meet the editorial board](#) 

Special Issues

Do you think there is an emerging area of research that really needs to be highlighted? Or an existing research area that has been overlooked or would benefit from deeper investigation? Raise the profile of a research area by leading a Special Issue.

[Submit your proposal](#) 

Author guidelines

Editorial board

Databases and indexing

Sign up for content alerts


[Sign up](#)



International Journal of Vascular Medicine

COUNTRY

Egypt

 Universities and research institutions in Egypt

SUBJECT AREA AND CATEGORY

Medicine
Cardiology and Cardiovascular
Medicine

PUBLISHER

Hindawi Limited

H-INDEX

27

PUBLICATION TYPE

Journals

ISSN

20902824, 20902832

COVERAGE

2010-2020

INFORMATION

[Homepage](#)

[How to publish in this journal](#)

SCOPE

International Journal of Vascular Medicine is a peer-reviewed, Open Access journal that publishes original research articles, review articles, and clinical studies focusing on vascular biology and physiology, vascular imaging, endovascular intervention, as well as prevention and treatment of vascular disease.

 [Join the conversation about this journal](#)

FIND SIMILAR JOURNALS

options

- 1
Acta Angiologica
POL
58%
similarity
- 2
Vascular Medicine
GBR
56%
similarity
- 3
Sang Thrombose Vaisseaux
FRA
52%
similarity
- 4
Angiology
USA
51%
similarity
- 5
International Angiology
ITA
50%
similarity



International Journal of Vascular Medicine

← Show this widget in your own website

Q3
Cardiology and Cardiovascular Medicine
best quartile

SJR 2020
0.41

powered by scimagojr.com

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" data-bbox="198 683 295 693">
```

SCImago Graphica

Explore, visually communicate and make sense of data with our new free tool.

Get it



Metrics based on Scopus® data as of April 2021

Leave a comment

Name

Email
(will not be published)


Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®





Source details

International Journal of Vascular Medicine

Open Access ⓘ

Scopus coverage years: from 2010 to Present

Publisher: Hindawi

ISSN: 2090-2824 E-ISSN: 2090-2832

Subject area: Medicine: Cardiology and Cardiovascular Medicine

Source type: Journal

CiteScore 2020

2.4 ⓘ

SJR 2020

0.411 ⓘ

SNIP 2020

0.826 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020 ▾

$$2.4 = \frac{121 \text{ Citations 2017 - 2020}}{51 \text{ Documents 2017 - 2020}}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021 ⓘ

$$2.3 = \frac{105 \text{ Citations to date}}{46 \text{ Documents to date}}$$

Last updated on 04 November, 2021 • Updated monthly

CiteScore rank 2020 ⓘ

Category	Rank	Percentile
Medicine		
Cardiology and Cardiovascular Medicine	#175/317	44th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語に切り替える](#)
[切换到简体中文](#)
[切换到繁體中文](#)
[Русский язык](#)

Customer Service

[Help](#)
[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

 RELX

International Journal of Vascular Medicine

- [Journal overview](#)
- [For authors](#)
- [For reviewers](#)
- [For editors](#)
- [Table of Contents](#)
- [Special Issues](#)

International Journal of Vascular Medicine / **About this Journal**

On this page
Aims and scope
Bibliographic information
Open Access
Contact

About this Journal

Aims and scope

International Journal of Vascular Medicine is a peer-reviewed, Open Access journal that publishes original research articles and review articles focusing on vascular biology and physiology, vascular imaging, endovascular intervention, as well as prevention and treatment of vascular disease.

Bibliographic information

ISSN: 2090-2824 (Print)
 ISSN: 2090-2832 (Online)
 DOI: 10.1155/1839

Open Access

International Journal of Vascular Medicine is an open access journal. All articles are immediately available to read and reuse upon publication. More information about our Open Access policy can be found on our [copyright page](#).

Contact

Editorial enquiries should be directed to ijvm@hindawi.com.
 General enquiries should be directed to help@hindawi.com.

Journal metrics	
Acceptance rate	19%
Submission to final decision	59 days
Acceptance to publication	26 days
CiteScore	2.400
Journal Citation Indicator	0.230
Impact Factor	-

APC	\$900
-----	-------

- [Submit](#)
- [Author guidelines](#)
- [Editorial board](#)
- [Databases and indexing](#)
- [Sign up for content alerts](#)

[Sign up](#)

Editorial Board

Academic Editors

- **Antonio Bozzani** , Foundation I.R.C.C.S. Policlinico San Matteo, Italy
- **Alberto Caggiati** , University Sapienza, Italy
- **Aaron S. Dumont**, Tulane University, USA
- **Albert G. Hakaim**, Mayo Clinic, USA
- **John A. Kern** , University of Virginia, USA
- **Mark Morasch**, St. Vincent Healthcare, USA
- **Karlheinz Peter**, Baker IDI Heart and Diabetes Institute, Australia
- **Bhagwan Satiani** , Ohio State University, USA
- **Robert M. Schainfeld** , Harvard University, USA
- **Thomas Schmitz-Rixen**, Goethe-University Medical Center, Germany
- **Michael Schneck**, Loyola University Chicago, USA
- **Mario Zuccarello**, The University of Cincinnati, USA



Table of Contents: 2020

International Journal of Vascular Medicine - Volume 2020 - Article ID 8864056

- Research Article

Long-Term Follow-Up of Contrast-Induced Acute Kidney Injury: A Study from a Developing Country

Ashraf O. Owais | Sameeha A. Alshelleh | ... | Karem H. Alzoubi

21 Dec 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 9796012 - Corrigendum

Corrigendum to "Evaluation of the Correlation between Serum Concentrations of Asymmetric Dimethylarginine and Corrected TIMI Frame Count in Patients with Slow Coronary Flow"

Mahshid Naserifar | Mahshid Ataei | ... | Hesamoddin Hosseinijani

14 Dec 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 6657077

- Research Article

Beneficial Effects of Spirulina Aqueous Extract on Vasodilator Function of Arteries from Hypertensive Rats

Diva M. Villalpando | Carlos M. Verdasco-Martín | ... | Mercedes Ferrer

08 Dec 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 8823039

- Research Article

Comparing and Correlating Outcomes between Open and Percutaneous Access in Endovascular Aneurysm Repair in Aortic Aneurysms Using a Retrospective Cohort Study Design

Peter DeVito | Ali Kimyaghalam | ... | Eugene Vitvitsky

29 Nov 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 2035494

- Research Article

Interesting Scenarios during Radiofrequency Ablation of Varicose Veins at University Hospital of Nepal

S. Vaidya | R. M. Karmacharya | ... | P. Thapa

12 Nov 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 2035494

- Research Article

Interesting Scenarios during Radiofrequency Ablation of Varicose Veins at University Hospital of Nepal

S. Vaidya | R. M. Karmacharya | ... | P. Thapa

12 Nov 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 4592190

- Research Article

Evaluation of the Correlation between Serum Concentrations of Asymmetric Dimethylarginine and Corrected TIMI Frame Count in Patients with Slow Coronary Flow

Mahshid Naserifar | Mahshid Ataei | ... | Hesamoddin Hossainjani

19 Sep 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 5923702

- Research Article

Three-Dimensional Echocardiography in Evaluating LA Volumes and Functions in Diabetic Normotensive Patients without Symptomatic Cardiovascular Disease

Mohamed Hamza | Ahmed Mamdouh | ... | Ahmed Nayel

19 Aug 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 9260812

- Research Article

Serum Endothelin-1 Correlates with Myocardial Injury and Independently Predicts Adverse Cardiac Events in Non-ST-Elevation Acute Myocardial Infarction

Anggoro Budi Hartopo | Indah Sukmasari | ... | Budi Yuli Setianto

04 Aug 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 7624158

- Research Article

The Impact of Coexistence of Smoking and Diabetes on the Coronary Artery Severity and Outcomes following Percutaneous Coronary Intervention: Results from the 1ST Jordanian PCI Registry

Mohamad I. Jarrah | Said Al-Khatib | ... | Nasr Alrabadi

02 Jul 2020

[Download PDF](#)



[Download citation](#)



Atherosclerotic Extension of Carotid Arteries: An Insertion in Clinical Practice

M. L. Furlanetto | E. F. B. Chagas | Payão SLM

23 Jun 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 4219180
- Research Article

Lipoprotein Particle Predictors of Arterial Stiffness after 17 Years of Follow Up: The Malmö Diet and Cancer Study

Jacob Hartz | Ronald M. Krauss | ... | Michele Mietus-Snyder

28 Apr 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 7631958
- Research Article

The Comparison of Nailfold Capillaroscopy between Juvenile Systemic Lupus Erythematosus and Healthy Controls: Correlation with Laboratory and Clinical Parameters

Seyed-Reza Raeeskarami | Navid Namazi | ... | Vahid Ziaee

27 Apr 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 4237204 - Clinical Study

Is Nonmicronized Diosmin 600 mg as Effective as Micronized Diosmin 900 mg plus Hesperidin 100 mg on Chronic Venous Disease Symptoms? Results of a Noninferiority Study

Marcio Steinbruch | Carlos Nunes | ... | Mauro Geller

07 Mar 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 4981092
- Research Article

Effect of Acute Myocardial Infarction on a Disintegrin and Metalloprotease with Thrombospondin Motif 13 and Von Willebrand Factor and Their Relationship with Markers of Inflammation

Abeer A. Al-Masri | Syed Shahid Habib | ... | Hana Al Zamil

11 Feb 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 7942381 - Clinical Study

Automatic Continuous CRT Optimization to Improve Hemodynamic Response: An Italian Single-Center Experience

Gregorio Covino | Mario Volpicelli | Paolo Capogrosso

07 Feb 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 6582742

- Research Article

Safety and Outcomes of Permanent and Retrievable Inferior Vena Cava Filters in the Oncology Population

Saba S. Shaikh | Suneel D. Kamath | ... | Brandon J. McMahon

05 Feb 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 9574734 - Review Article

Systematic Review and Meta: Analysis of Aortic Graft Infections following Abdominal Aortic Aneurysm Repair

O. S. Niaz | A. Rao | ... | P. Somalya

31 Jan 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 8758905

- Research Article

Mechanical Occlusion Chemically Assisted Ablation (MOCA) for Saphenous Vein Insufficiency: A Meta-Analysis of a Randomized Trial

Johanes Nugroho | Ardyan Wardhana | Cornelia Ghea

29 Jan 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 3893261

- Research Article

Hypothermia during Surgical Treatment of Type A Aortic Dissection: A 16 Years' Experience

Francesco Macrina | Maria Cristina Acconcia | ... | Fabio Miraldi

25 Jan 2020

[Download PDF](#)



[Download citation](#)



International Journal of Vascular Medicine - Volume 2020 - Article ID 3673657

- Research Article

Experience in the Management of Patients with Chronic Vascular Ulcers of the Lower Limbs Using Negatively Charged Polystyrene Microspheres

María Teresa Cacia Sánchez

22 Jan 2020

[Download PDF](#)



[Download citation](#)

