

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Papers in Plant Pathology

Plant Pathology Department

2021

Root and crown rot pathogens found on dry beans grown in Mozambique

Suzana Vanessa Ismael Fernandes

G. Godoy-Lutz

Celestina Nhagupana Jochua

Carlos A. Urrea

Kent M. Eskridge

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/plantpathpapers>



Part of the [Other Plant Sciences Commons](#), [Plant Biology Commons](#), and the [Plant Pathology Commons](#)

This Article is brought to you for free and open access by the Plant Pathology Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Plant Pathology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Suzana Vanessa Ismael Fernandes; G. Godoy-Lutz; Celestina Nhagupana Jochua; Carlos A. Urrea; Kent M. Eskridge; James R. Steadman; and Joshua Herr

Root and crown rot pathogens found on dry beans grown in Mozambique

Suzanna Fernandes,^{1,2} Graciela Godoy-Lutz,^{1,3}
Celestina Jochua,^{1,2} Carlos Urrea,⁴ Kent Eskridge,⁵
James R. Steadman,¹ and Joshua R. Herr^{1,6,7}

Suzanna Fernandes and Graciela Godoy-Lutz contributed equally to this work.

1 Department of Plant Pathology, University of Nebraska–Lincoln, Lincoln, NE 68583, USA

2 Present address: Directorate of Agronomy and Natural Resources (DARN), The Agricultural Research Institute of Mozambique (IIAM), Maputo, Mozambique

3 Present address: Instituto Dominicano de Investigaciones Agropecuarias y Forestales (IDIAF), Santo Domingo, Dominican Republic

4 Department of Agronomy & Horticulture, University of Nebraska–Lincoln, Lincoln, NE 68583, USA

5 Department of Statistics, University of Nebraska–Lincoln, Lincoln, NE 68583, USA

6 Center for Plant Science Innovation, University of Nebraska–Lincoln, Lincoln, NE 68588, USA

7 Genetics, Cell, and Molecular Biology Graduate Program, School of Biological Sciences, University of Nebraska, Lincoln, NE 68588, USA

Correspondence – Joshua R. Herr, jherr@unl.edu

Published in *Tropical Plant Pathology* 46 (2021), pp 294–310.

<https://doi.org/10.1007/s40858-021-00422-8>

Copyright © 2021 Sociedade Brasileira de Fitopatologia. Published by Springer Nature. Used by permission.

Submitted 16 March 2019; accepted 16 February 2021; published 25 March 2021.

Abstract

Dry edible beans are a vital food source in Mozambique, East Africa—one that alleviates hunger and malnutrition and adds value to the economy. In recent years, root/crown rot (RCR) pathogens have emerged as limiting constraints in dry bean production. Not much has been characterized concerning the causal agents of RCR in Mozambique. The purpose of this study was to identify the primary pathogen(s) associated with RCR dry bean samples collected at breeder nursery sites and farmer fields in Mozambique using molecular sequencing and culture-based methods. Sequencing revealed, not surprisingly, an increased diversity of fungal/oomycete operational taxonomic units when compared to culture-based methods of diversity. Species of *Fusarium*, mainly *F. oxysporum*, were the dominant taxa detected in RCR dry beans through sequencing the ITS rDNA region and partial EF-1 α gene. Collectively, 333 fungi and/or Oomycetes were isolated in culture during the 2014–2015 growing seasons and tested for pathogenicity on healthy bean seedlings. *Fusarium* species were identified by both morphological and molecular characters. At least 60% of the isolates inoculated on common bean were recognized as potentially pathogenic. From both isolation frequency and pathogenicity testing, *F. oxysporum* and related species play an important role in the bean RCR complex. We found similar results from dry beans grown in the two main bean-growing regions of Mozambique. These findings will allow breeders to screen for resistance to *F. oxysporum* in greenhouse grown bean plants as well as within field grown bean cultivars.

Keywords: Soil-borne fungal pathogens, *Fusarium*, FTA® Cards, High-throughput sequencing, Bean root and crown rot, *Macrophomina*, *Fusarium oxysporum*

Introduction

Dry edible beans (*Phaseolus vulgaris* L.), with a yearly production harvest area of 100,000 ha, are one of the most important food sources in Mozambique (Wortmann et al. 1998). Beans provide an important source of dietary protein, micro-nutrients, and caloric intake for the people of Mozambique, as well as a crucial source of income for the small-scale farmers (Wortmann et al. 1998). Mozambique, which is located on the southeastern coast of Africa, has many challenges for bean production, due in large part to climate variation through cyclical drought and excessive rainfall and plant diseases are exacerbated these changes in climate (Walker et al. 2006). Consequently, diseases such as root/crown rot (RCR) are emerging as limiting factors in bean production enhanced by changing environmental conditions (Farrow et al. 2011; Bodah 2017).

Bean RCR comprises a syndrome attributed to a suite of soil-borne pathogens. These pathogens may be reported either as individual strains

or synergistic infections whose occurrence and severity varies by location, site-specific incidences, soil properties and nutrients, and/or other variable environmental conditions that contribute to disease propagation and spread (Rusuku et al. 1997; Wortmann et al. 1998; Farrow et al. 2011). Most species of soil-borne pathogens have been identified as causal agents using traditional culture methods and pathogenicity tests. The most relevant species reported in Mozambique are *Fusarium solani* (Mart.) Appel, Wollenv. *F. sp. phaseoli* (Burk.) Synd. & Hans, *F. oxysporum* Schlecht. *F. sp. phaseoli* Kendrick & Synder, *Pythium spp.*, *Rhizoctonia solani* Kühn (teleomorph: *Thanatephorus cucumeris* (Tassi) Goid), *Athelia rolfsii* (Curzi) C.C. Tu & Kimbr., and *Macrophomina phaseolina* (Tassi) Goid (Abawi 1989; Rusuku et al. 1997; Paparu et al. 2018). Recently, in eastern and southern Africa, species of *Pythium* have received more attention due to their higher prevalence over other pathogens in disease surveys (Nzungize et al. 2011a, Nzungize et al. 2011b). As a result, breeding programs in some countries have focused on only developing RCR resistance for *Pythium* species.

In Mozambique, there is limited information on the pathogenic species causing, or associated with, bean RCR symptoms. This is despite widespread disease occurrence in the Gurue and Chokwe provinces where beans are most consistently grown. The proper identification of the primary causal pathogen(s) can lead to better disease management strategies and may improve bean production due to the fact that yield loss has not been accurately calculated for each pathogen. Morphological features of pathogens cultured from infected plants provide important information for identification in the laboratory (Narayanasamy 2011), and live pathogenic specimens are vital for disease-resistance inoculations and greenhouse evaluations in breeding programs. In addition to traditional morphological characters, molecular methods, such as high-throughput sequencing, have become commonplace in pathogen detection and species identification (Herr et al. 2015; Hibbett et al. 2016). In the present study, we employed a suite of morphological and molecular identification techniques to identify the diversity of the predominant pathogen (or pathogens) in Mozambique that are associated with bean RCR. We additionally focused on the population structure of *Fusarium oxysporum* isolates recovered from the same bean RCR samples.

Materials and methods

Sample collection and processing

We took advantage of collecting plants from a field test of four Andean Diversity Panels for Bean (Cichy et al. 2015). These panels were situated where root rot screening has already been in effect for at least a decade. The survey locations in the present study included the R10 (North West) region, which has the greatest levels of bean production, and is fed by rainfall; the R3 (South West) region where production is under dry winter irrigation nurseries; and lastly, from two farmer fields: R3 (Chokwe) and R10 (Gurue), respectively, from the Gaza and Zambesia provinces in well-studied agro-ecological zones (Wortmann et al. 1998). The sampling locations in 2014 and 2015 represented contrasting climatic and soil conditions (Fig. 1 and Table S-3). The essential differences of these conditions consisted the sampling locations of Chokwe, where the site is close to sea level, had organic matter-rich Salic Fluvisol soils and recognized as a semi-arid climate, and the sampling locations of Gurue, which is approximately 700 m above sea level, and consist of Ferrossols with less organic matter, and a slightly cooler climate due to the elevation gain of slightly more than 100 m.

Within 30 days after bean plant emergence, we identified and sampled 88 bean plants with RCR symptoms (Fig. 1) along with a healthy control plant in close proximity, typically no more than 1 m away. Plants were harvested in the field, placed in a cooler with ice, and brought to the laboratory for processing. Soil and associated debris were removed from the plants and lesions (approximately 2–4 mm in size) at both the diseased and healthy interface (Fig. 2a) were extracted according to published protocols (Mukuma et al. 2020). Lesions of infected tissue were isolated, flash-frozen, and ground with a micro-pestle (Fig. 2b). Liquid extracts of the lesions were spotted onto the absorbent matrix of Whatman FTA® Cards and matching tissue samples were placed in a coin envelope and labeled with the sample number, bean cultivar name, description of phenotypic symptoms, the location and dates of sampling, and any other field-related data taken during collection, such as unique climate and precipitation notes.




Fig. 1 (a) Bean field affected by root/crown rot (RCR) in Chokwe, Mozambique. (b) Inset photo of row of RCR and wilt of bean plants. (c) Topographical map of Mozambique with specific location on the continent of Africa and the general locations that were sampled in this study.




Fig. 2 (a) Schematic representation of typical bean plant showing infection site and location of where tissues were sampled. (b) Ground RCR tissue for DNA extraction. (c) Tissue extracts blotted onto FTA Card. (d) Entire FTA[®] Card shipped to laboratory. (e) Photo of the pathogenicity test by the “straw method” conducted in the greenhouse.

Fungal/Oomycete cultural isolation

Small segments of 2–4 mm of the RCR diseased plant tissue were surface sterilized by immersing in 10% v/v NaOCl/water for 15–30 s, then transferring to 70% alcohol for 15–30 s, and finally to a distilled water wash for 1–2 min. The infected segments of plant tissue were then blotted on sterile Whatman™ filter paper to air dry, and then transferred using sterile forceps onto 2% water agar where segments reached about 4 mm in length. After this growth period, the resulting isolate hyphae was transferred onto potato dextrose agar (PDA) to allow further growth. Within 2–4 days, 4-mm mycelial plugs were transferred to four PDA




Fig. 3 Photos of Fungi and Oomycetes isolated in this study: (a) *Macrophomina phaseolina*; (b) *Alternaria alternate*; (c) *Pythium ultimum*; (d) *Rhizoctonia solani*; (e) *Fusarium equiseti*; (f) *Fusarium oxysporum*; (g) spores of *Fusarium oxysporum* stained with lactophenol cotton blue; (h) *Fusarium solani*; (i) spores of *Fusarium solani* stained with lactophenol cotton blue.

plates and one water agar (WA) plate for each isolate. These plates were used for morphological observations, future pathogenicity tests, and mycelial DNA extraction for subsequent nucleotide sequencing. Culture characteristics were examined across three different culture media: WA, PDA, and carnation leaf agar (CLA) (Fisher et al. 1982; Leslie and Summerell, 2006). The characteristics observed were growth pattern, colony texture and pigmentation, spore size and shape, and growth rate of the mycelial edge (Fig. 3). Mycelial plugs of a 6-mm diameter were taken from the WA colony edge of each isolate and transferred to PDA plates and incubated under continuous darkness for 9 days at 22–25 °C. Colony diameter was measured on the bottom of the plate. Colony texture and color on PDA were evaluated for each isolate. For fungi and Oomycetes, the genus and species names were assigned using several identification keys (Dhingra and Sinclair 1978; Dugan 2006; Watanabe 2010). Species of *Fusarium* were identified based on the color and growth pattern of colonies grown on PDA, then transferred to CLA for identification using dichotomous keys (Burgess et al. 1994; Leslie and

Summerell 2006) focusing on characters such as the size and morphology of phialides, macro- and micro-conidia, and chlamydo-spores. *Fusarium* isolates grown on CLA media at 20–24 °C with a 12-h light/12-h dark regime for 10 days sporulated on CLA, and spore masses growing on the leaf surface were transferred to a blue drop of lactophenol cotton blue on a glass slide to observe spore size and shape. Diagnostic characters were then photographed using a compound light microscope mounted with a Motic camera (Motic North America, British Columbia, Canada) at × 40 magnification.

DNA extraction from FTA® Cards and plant tissue

In order to extract DNA from field collected samples, the Whatman FTA® Cards and matching tissue samples were sent to the laboratory located at the Department of Plant Pathology at the University of Nebraska-Lincoln, NE, USA, under the USDA-APHIS Permit P526P-17-02138 issued to J.R. Steadman. Samples waiting to be processed were kept in a desiccant chamber at room temperature. Due to a small number of samples from the Whatman FTA® Cards that we were not able to acquire enough DNA for sequencing, as well as the overall proximity and similarities of some of the sampling locations, we evaluated all the data from the closely located geographic locations Chato and Chissano (analyzed further as the Gurue (R10) growing region) along with the larger number of samples from Chokwe (R3) growing region (Fig. 1).

Genomic DNA from the FTA® Cards was recovered from an excised 1-cm² section of the Card by incubation in 200–300 µl of TRIS-EDTA buffer solution (Fluka Analytical, Sigma-Aldrich Co., St. Louis, MO, USA) at 4 °C overnight. After this incubation, 100 µl of the buffer solution was used for DNA extraction with the PowerClean Pro DNA Cleanup Kit (Catalog No. 12997-50 MO BIO Laboratories, Inc., Carlsbad, CA, USA). Genomic DNA was extracted from 100 mg of cryogenic ground RCR diseased plant tissue or isolate mycelia using the PowerPlant Pro DNA isolation kit (Catalog No. 13400-50, MO BIO Laboratories Inc., Carlsbad, CA, USA). Concentration and purity of total DNA from each sample was determined using a DeNovix DS-11 Nanodrop Spectrophotometer (DeNovix Inc., Wilmington, DE). Genomic DNA from the extraction process was stored at –20 °C until sequencing.

PCR amplification and DNA sequencing

Taxonomic identification of the fungal/Oomycete isolates from cultural methods was initially based on morphology, but we subsequently used DNA from each isolate amplified by polymerase chain reaction (PCR) with primer pairs ITS4/ITS5 (White et al. 1990) to verify our identification. To aid in the identification of *Fusarium* species, we sequenced a second marker region, the partial EF-1 α gene using primer pairs EF1/EF2 (O'Donnell et al. 1998). We identified four main pathogen groups associated with diseased RCR plant tissue from sequencing of both the DNA from the FTA® Cards and/or plant tissue extracts. To verify these species complexes, we utilized isolate-specific markers for these pathogens using the following specific primer pairs (Table S-4): FM66/58 COX II for *Pythium* species (Martin 2000); ITSFu F/ITSFu R for *Fusarium* species (Abd-Elsalam et al. 2003); RS4 primers for *Rhizoctonia solani* (Guillemaut et al. 2003) and MpkF1/MpkR1 primers for *Macrophomina phaseolina* (Babu et al. 2007). Isolates of *Macrophomina phaseolina*, *Pythium ultimum*, *Rhizoctonia solani* strain AG- 4, and *F. oxysporum* collected from field grown dry beans from Scottsbluff, NE, USA, were used throughout our tests for PCR quality control. For each amplification, 25 μ l of PCR reaction mixture was prepared by adding 1 μ l of genomic DNA to 24 μ l of a master mix/tube containing 9.5 μ l PCR grade sterile ddH₂O, 12.5 μ l Econotaq® PLUS GREEN 2 \times Master mix (Lucigen, Madison, WI, USA), and using 1 μ l of 0.2 mM/ μ l of each forward and reverse primers. Reactions were performed in a PTC-100 thermal cycler (Bio-Rad Laboratories, Hercules, CA, USA). Amplification fragments were separated by electrophoresis in a 1.5% Ultra-Pure® and Quick Dissolve agarose (Invitrogen, Carlsbad, CA, USA) in TBE buffer and stained with ethidium bromide for visualization under a UV light using a ChemiDoc EQ System with the software package Quantity One (Bio-Rad Laboratories, CA, USA). Amplified fragments for Sanger sequencing were purified with Ultra-Clean PCR Clean-Up Kit (Cat. No. 12500-50 MO BIO Laboratories, Inc., Carlsbad, CA, USA), and sequenced on an ABI 3730XL Sanger Sequencer by ACGT, Inc. (Wheeling, IL, USA).

DNA extracted from the FTA® Cards of field infected RCR plants was isolated with the goal of using both sequence identification methods and phylogenetic analysis of the fungal/ Oomycete composition in the plant lesions (Herr et al. 2015; Hibbett et al. 2016). Sequencing was conducted

at Molecular Research LP (www.mrdnalab.com—Shallowater, TX, USA) on an Illumina MiSeq (Illumina, Inc., San Diego, CA, USA) using PCR primers Euk SSU euk7F/euk570R amplifying the 18S rRNA gene V4 variable region. Sequence data was processed using a taxonomic analysis pipeline (Chiodini et al. 2016) where the raw data set was demultiplexed and barcodes and primers were trimmed from the sequences. The sequence data was subsequently processed by removing sequences <200 bp, as well as reads with homopolymer runs exceeding 10 bp and ambiguous base calls. Sequencing chimeras were detected and removed, and operational taxonomic units (OTUs) were generated by centroid clustering at 3% divergence (using vsearch at 97% similarity; Rognes et al. 2016) followed by removal of true singleton sequences for downstream analysis (Chiodini et al. 2016). Final OTUs were identified using BLASTn against the curated GreenGenes (DeSantis et al. 2006, McDonald et al. 2012), RDP II (Cole et al. 2005), and Silva (Quast et al. 2012) databases, respectively.

Pathogenicity assay

Pathogenicity assays were conducted using 333 fungal and/or Oomycete isolates recovered from the RCR diseased plant tissue. Two-week-old susceptible PINTO 114 bean seedlings, grown in the greenhouse using 6 × 4 cm plastic pots containing steam pasteurized sterilized soil mix, were inoculated using a modified straw test method (Mukuma et al. 2020). Each inoculation was replicated five times and plants were kept in a mist chamber set for humidity ≥ 80% and temperatures approximating 23 °C. Control plants were inoculated with clean PDA plugs containing no visible organismal growth. After 48 h, all plants were moved to greenhouse benches and arranged in a complete randomized block design. Lesion length for each plant was measured and recorded 48 h after removal from the mist chamber (Fig. 2e). Lesion data were analyzed using Statistical Analysis System (SAS) v 9.2 (SAS Institute, Inc. Cary, NY, USA). Mean comparisons were made using Fisher's protected least significance difference at $P = 0.05$. To fulfill Koch's postulates, the fungi were re-isolated from the infected petiole, grown on WA, and subsequently transferred to PDA (as previously described) to re-validate cultural and morphological features.

PCR fingerprinting of *Fusarium oxysporum* with microsatellite markers

The genetic diversity of *F. oxysporum* isolates recovered from RCR diseased plant tissue was determined using PCR fingerprinting analysis with four types of microsatellites. Previously utilized for *Fusarium* taxa (Kang et al. 2002), the universal rice primer (URP) (5'CCCAGCAACT-GATCGCACAC3') was used to amplify *Fusarium* isolates and microsatellites with the repeated motifs (GTG)₅ (Lieckfeldt et al. 1993; Brasileiro et al. 2004), (ACG)₅, and (AGG)₅ (Bahkali et al. 2012). Two primers, the (ACG)_s and (AGG)_s motifs, were chosen from a set of 3 URPs, because they yielded the most consistent and unambiguous fragments and amplified most of the *Fusarium* isolates we obtained. DNA samples from 93 *F. oxysporum* isolates (72 from Gurue and 21 from Chokwe) were amplified with the primers following published protocols (Table S-4). The PCR amplification reactions were prepared for each of the primers and band visualization was done as previously described. Band size and number in the gel were determined in the Band Analysis tools of ImageLab software, version 4.1 (Bio-Rad, Carlsbad, CA, USA). After being scored, the resulting data set was analyzed using the `nei.dist()` function within Poppr (Nei 1973, Kamvar et al. 2014) using R (R Core Team 2018).

Clear and unambiguous bands, amplified by the microsatellite markers, were selected and scored for presence (1) or absence (0) of the corresponding band, which ranged from 300 to 1000 bp. A nonclone-corrected binary data matrix was computed with Genalex v. 6.502 (Peakall and Smouse 2006) and SAS v. 9.2 (SAS Institute, Inc. Cary, NY, USA) utilizing morphological characteristics from the original isolate used in the inoculation assay.

Data visualization and analysis

Sequence chromatograms of ITS rDNA and the partial EF-1 α genes were visualized with Chromat version 2.6.4 (Technelysium Pty., Australia) and after inspection, the resulting FASTA files were exported for further analysis. To infer the identification of the isolates at putative species level, the ITS rDNA sequences were identified using the NCBI BLAST web interface (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) against the NCBI GenBank nt database. Identification of the partial EF-1 α gene to the closest

matching *Fusarium* species was conducted by BLAST analysis against the *Fusarium* EF blast cyber-Infrastructure website (<http://fusariumdb.org/index.php>) available through the Pennsylvania State University, University Park, PA, USA (Park, et al. 2010). For phylogenetic analysis and placement, the FASTA files of the partial EF-1 α gene sequences were subjected to multiple sequence alignment (MSA) using MUSCLE (Edgar 2004) and the resulting alignment was used for the phylogenetic tree construction. Phylogenetic analysis was conducted using Bayesian inference with the BEAST pipeline (Drummond and Rambaut 2007). Sequences of *F. oxysporum* from various *F. oxysporum forma specialis phaseoli* accessions from NCBI GenBank showing 99 to 95% synteny were selected as outgroups and to add resolution beyond those sequences collected for this study. The resultant tree was visualized using iTOL (Interactive Tree of Life—<http://itol.embl.de/>; Letunic and Bork 2016).

The overall abundance of taxa among the samples from the different sampling locations and years were estimated from the identification of OTUs from the Illumina MiSeq sequencing. Heat maps were constructed in R (R Core Team 2018) using the ggplot2 (Wickham 2016) and RColorBrewer packages (Neuwirth, 2014), based on the percent of relative abundance obtained from Illumina MiSeq 18S sequencing for FTA® Cards and RCR diseased plant tissue. Sequences with reads depth less than 10, which we considered putative sequencing artifacts, were removed from further analysis. The overall sample diversity, which was measured as Shannon (Shannon and Wiener, 1963) and Simpson (Simpson 1949) indices, was calculated with the PAST (PAleontological STastics) v 3.12 program (Hammer et al. 2001). Venn diagrams, constructed with Venny 2.1 (<http://bioinfogp.cnb.csic.es/tools/venny/>), were drawn to display similarities (shared reads) and distinctiveness (unique reads) among *F. oxysporum* for the Gurue and Chokwe locations based on Illumina MiSeq 18S and EF-1 α sequence data. Sequencing data from the MiSeq run was used to validate the culture-based and PCR amplified methods for detecting the four main RCR pathogens we identified in this field study. This analysis was conducted by Spearman's rank and rho correlation which was also calculated within the PAST program.

The frequency of the four main fungal/Oomycete genera and species (*Fusarium* spp., *Macrophomina phaseolina*, *Rhizoctonia solani* and *Pythium* spp.) was generated by PCR amplification with specific primers.

A binary matrix was constructed with the color-coded data set for each as presence (1) and absence (0) of the expected fragment. Tables were constructed for each location per year.

Results

Cultural and morphological characterization of isolates obtained from RCE tissue

We obtained 333 fungal and Oomycete isolates in culture, as examined by macro- and microscopic characteristics, from 79% of RCR tissue samples collected in Chokwe and Gurue in 2014 and 2015. Most of the isolates were initially assigned at the genus or species level, based on their characteristic mycelial and spore features, although some isolates did not sporulate or had no diagnostic cultural characteristics. Isolates of *Alternaria* spp., *Fusarium* spp., *Pythium* spp., *M. phaseolina*, and *R. solani* were significantly more abundant than other taxa in the samples. Among the identified isolates, *Pythium* spp. had the fastest growth rate and *M. phaseolina* the slowest. The morphology and size of both phialides and spores, based on growth characters from CLA plates, were used to tentatively identify the *Fusarium* species into 3 morphotypes and/or species complexes: *F. solani*, *F. oxysporum*, and *F. equiseti* (Fig. 3), prior to DNA sequencing analysis.

Identification of cultured isolates

We identified the taxa of cultured isolates of fungi and Oomycetes through DNA sequencing and subsequently compared the recovery of the isolates from Whatman FTA® Cards and RCR tissue. Nucleotide sequence data were identified to closest match using BLAST queries with a cut-off value >98% match and a resultant 42 distinct fungal taxa were identified (Table S-5). *Fusarium oxysporum* was by far the most common genus and species (Figs. 4, 5 and 6). In comparison, the most frequently isolated genus and species in both locations were *F. oxysporum* (34%), *F. equiseti* (13%), *F. solani* (10%), *A. alternata* (9%), *M. phaseolina* (4%), and *R. solani* (2%) (Fig. 4).




Fig. 4 Relative abundance of the 17 most prevalent identified pathogenic fungi and Oomycetes isolated from isolated RCR bean tissue through 18S rDNA Illumina sequencing for years (a) 2014 and (b) 2015, by location of pooled Chokwe and pooled Gurue growing regions.

Pathogenicity assay

The 333 isolates recovered from RCR plant tissues were tested for pathogenicity and 204 were identified as inducing pathogenic-like responses in bean assays (Table S-5) in the greenhouse. We used the criteria of water-soaking followed by necrosis or wilting of the stem of 14-day-old Pinto 114 bean plants observed 3 days after inoculation as exhibiting traits of pathogen damage. Mean lesions ranged from 0.4 to 8 cm in length. The most common isolate recovered from our pathogenicity tests was *Fusarium oxysporum*, followed by *F. equiseti* and *F. solani*

Fig. 5 Heat map of normalized relative abundance of the top 92 fungal operational taxonomic units associated to RCR-symptomatic beans based on 18S rDNA region Illumina sequencing from the pooled Chokwe growing regions in 2014 and 2015.





Fig. 6 Heat map of normalized relative abundance of the top 107 fungal operational taxonomic units associated to RCR-symptomatic beans based on 18S rDNA region Illumina sequencing from the pooled Gurue growing regions in 2014 and 2015.



from both locations of farm fields, with the exception of samples from Chokwe 2015, where *Macrophomina phaseolina* was the most virulent isolate we encountered.

Detection of four primary pathogenic isolates associated with RCR

Microsatellite amplicon sequences from the infected samples and positive controls were consistent with the expected fragment sizes for *Fusarium* spp., *Pythium* spp., *Rhizoctonia solani*, and *Macrophomina phaseolina*. With the primers we utilized, the presence or absence of each genus and/or species per sample was detected (Fig. 7). The frequency of detection varied with both location and year; however, *Fusarium* spp. were detected in highest frequencies and in RCR tissue more than in Whatman FTA® Cards (Fig. 7).

When Illumina sequencing was used for DNA analysis from both FTA® Cards and RCR tissue, not surprisingly, we identified a greater depth of species of fungi/Oomycetes than from the culture-based method (Fig. 7). By location and year, overall, *F. oxysporum* was the most frequently identified isolate in both FTA® Cards and RCR tissue, followed by *F. solani* and other *Fusarium* species, as well as *Rhizoctonia solani* (Figs. 4, 5, and 7). Other fungal/Oomycete species that have been associated with RCR symptoms such as *Pythium ultimum* were less frequently observed. We found it notable that a fungal taxon identified in culture from RCR infected legumes, *M. phaseolina*, was not detected by Illumina amplicon sequencing in either location or year with the Whatman FTA® Cards. Finally, there was a significant correlation between information obtained by FTA® Cards and RCR tissue values based on the Spearman correlation coefficient for Gurue (0.759 and 9.86516, $P < 0.001$) and Chokwe (0.70288 and 0.865, $P < 0.001$) for 2014 and 2015, respectively.

Phylogenetic affinities of Fusarium oxysporum and genetic variation for Fusarium oxysporum assessed with microsatellite primers

To detect the four major pathogen genera and species (*Fusarium* spp., *R. solani*, *P. ultimum*, *M. phaseolina*), we employed Illumina amplicon sequencing, PCR-based microsatellites, and culture-based methods. The Illumina amplicon sequencing identified the most fungi and Oomycetes from FTA® Cards. However, PCR-based methods from RCR tissue were best at detecting *M. phaseolina* (Fig. 7). Every method that we employed




Fig. 7 Detection frequency across the years of 2014 and 2015 for the four main pathogens we identified (all *Fusarium* species, *Macrophomina phaseolina*, all *Pythium* species, and *Rhizoctonia solani*) associated with RCR of beans from sampling sites of (a) Chokwe and (b) Gurue in Mozambique. The methods used for detection in this study were (1) culture-based, (2) Illumina MiSeq FTA Card, (3) Illumina MiSeq tissue sample, (4) primer-specific polymerase chain reaction amplification FTA Card, and (5) primer-specific polymerase chain reaction amplification of tissue sample.

identified *Fusarium* spp. as the most abundant pathogen group in RCR tissues. Other *Pythium* species, as well as *R. solani*, were also detected by amplicon sequencing at higher frequencies than PCR-based or culture-based methods. The frequency of detection of the four most abundant pathogens was higher in the direct assay of RCR diseased plant tissue than in FTA® Cards for samples from both farm field locations in




Fig. 8 Diversity metrics (Simpson and Shannon indices) for the (a) pooled Chokwe growing region RCR bean samples and the (b) pooled Gurue growing region RCR bean samples in 2014 and 2015 using the different diagnostic analysis methods we employed: (1) culture-based methods, (2) Illumina sequencing of FTA Card DNA extractions, and (3) Illumina sequencing of directly sampled RCR tissue DNA extractions directly sampled tissue samples.

Mozambique. The culture-based methods had the lowest detection frequency for species of *Pythium*, *R. solani*, and *M. phaseolina*, but they isolated and identified numerous species of *Fusarium*. The two locations sampled showed differences in community diversity (based on the value of $H' > 2.5$) which revealed the presence of diverse array of isolates associated with RCR disease, whereas Simpson values close to 1 show the predominance of one species or group, which was the case for *Fusarium*, where these taxa were found in the same plant niche for most locations and year tested (Fig. 8). The exception was for the Chokwe location in 2015, where *M. phaseolina* was the predominant fungal species





Fig. 9 Bayesian likelihood phylogenetic tree derived from partial EF-alpha gene region sequences of *Fusarium oxysporum* cultured isolates from Mozambique. Leaf names in green were isolated in from RCR-symptomatic samples from the Gurue region and leaf names in red were isolated from RCR-symptomatic samples from the Chokwe region. Leaf names in black represent 36 *Fusarium oxysporum* partial EF-alpha gene accessions from database of NCBI GenBank.

in culture. Values of Shannon and Simpson diversity indices were similar for both locations and years where *F. oxysporum* had the highest percentage of relative abundance. Of the most predominant species, *F. oxysporum*, was further examined for better taxonomic resolution by sequencing the partial EF-alpha gene and the sequences were used to construct a phylogenetic tree (Fig. 9). The phylogenetic tree revealed at least

seven well-defined clusters within *F. oxysporum* species complex indicating intraspecific variability of the partial EF-1 gene. Additionally, most of the isolate sequences were separated from those of published *F. oxysporum* accessions from the NCBI GenBank database indicating the new isolates from Mozambique are not found in databases. There was no evidence that geographical origins of the isolates influenced the clustering pattern because sequences from both locations were found in species complex clusters. Phylogenetically informative nucleotide positions in the 18S rRNA sequences (generated by the Illumina amplicon sequencing) were used to compare common or shared OTUs in different combinations between the Chokwe and Gurue locations for both years and presented in Venn diagrams (Fig. 10). These common, or shared, amplicon counts are displayed in the overlapping panels and unique OTUs in the non-overlapping areas.

Fig. 10 Euler diagrams depicting shared (intersecting) and unique (not intersecting) operational taxonomic units of *Fusarium oxysporum* identified with Illumina MiSeq analysis of the 18S rDNA region sequences from RCR-symptomatic beans plants in (A) 2014 and (B) 2015 from the pooled samples of the Chokwe and Gurue growing regions in Mozambique.



There were 91 isolates of *F. oxysporum* from Chokwe and Gurue characterized with single microsatellite primers (URP2R, (GTG)₅, (ACG)₅, and (AGG)₅) which generated 16 reproducible fragments, as well as 27 scorable fragments. The fragments (or allele) sizes ranged from 100 to >2000 bp, although for ease of interpretation we only scored those with the range of 200 to 1000 bp. Not all of the isolates amplified using all four primers with the same efficiency. We identified diversity indicators for the Chokwe and Gurue group of isolates utilizing the microsatellite primers URP2R, (GTG)₅, (ACG)₅, and (AGG)₅ (Table S-1), and for example, the percentage of the polymorphisms was highest when we utilized the (AGG)₅ primer set when compared to the rest of the primers. Nei's gene diversity (for heterozygosity estimates), unbiased Nei's gene diversity (a correction for small number of samples), and Shannon's Information Index (extent of genetic variation) (Table S-2) all displayed low values indicating a small genetic distance between the Chokwe and Gurue *F. oxysporum* isolate groups. Analysis of molecular variance revealed that the percentage of total variation within the locations was higher than the variation found between locations. In conclusion, we detected no structural population differences between *F. oxysporum* isolate groups for each location based on the number of isolates and primers tested (Table S-1).

Discussion

The identification of plant pathogens is an important first step to develop preventive management strategies and identify resistant dry edible bean cultivars to root and crown rot (RCR) diseases. In this study we employed three methods to identify the primary causal agent of RCR in dry edible beans in Mozambique: amplicon sequencing of the 18S rRNA and EF1 marker regions, genus and species-specific microsatellite primer amplification paired with morphological identification of cultural material, and DNA sequencing of the ITS rDNA region of the cultured fungi and Oomycetes.

Amplicon sequencing methods revealed the presence of a greater diversity of fungal and Oomycete species than culture-based methods. In all of the methods we employed, species of *Fusarium*, mainly *F. oxysporum*, were the dominant fungal isolates detected either on DNA extracted

from FTA® Cards and RCR diseased plant tissue, or from isolates recovered through conventional culture methods. Other RCR disease-associated fungi, such as *R. solani*, *Pythium* spp., *A. rolsfii*, and *M. phaseolina*, were detected or isolated in cultured samples but were lower in abundance and typically inconsistent in their presence and absence. The exception to this was found in the samples from Chokwe in 2015 where we identified *M. phaseolina* from both the sequencing and culture methods. Other fungal species we detected by amplicon sequencing, widely distributed across the Ascomycota and Basidiomycota, are not known to be associated with RCR and could be considered plant endophytes. The role of these fungi as endophytes in healthy or diseased dry bean plant tissue is yet to be well established (Nair and Padmavathy 2014). Despite the high frequency of isolation of *M. phaseolina* from the 2015 samples collected in Chokwe, the symptoms observed in the collected root/crown samples did not match symptoms typically caused by *M. phaseolina*, which is known to infect the upper stem of more mature plants. We hypothesize that increased rainfall in the Chokwe region during our study resulted in this finding, as *M. phaseolina* has been reported to increase abundance in regions of the tropics and subtropics where crops might be exposed to water stress (Songa and Hillocks 1996). The high isolation frequency of *M. phaseolina* is likely due to climate variability in the 2014 and 2015 growing seasons, which were characterized by low rainfall from October to December of 2014, and the resulting drought in the southern region from January to March of 2015. Later, an El Niño phenomenon was reported in Mozambique in the 2015 growing season which a rainy season in the south of the country was followed by additional pronounced and excessive rain or floods in the northern region (C. Jochua, personal observations). Most of the symptomatic bean samples, which exhibited signs of wilting and necrosis of the stem even before the flowering stage, were associated with infection by species of *Fusarium*. These symptoms were typically associated with recurrent cyclical weather patterns in the country, such as flooding and drought, and have caused environmental conditions which favor the abundance of *Fusarium* species in all locations of Mozambique.

The results of this study are in agreement with previous work (Mukuma et al. 2020) that found *F. oxysporum* was the predominant pathogen associated with RCR of bean in Zambia. However, Rusuku et al. (1997) also studied soil-borne pathogens causing the RCR disease from

culture isolation and found species of *Pythium* to be the most frequently isolated pathogens, over *F. oxysporum*, *M. phaseolina*, and *R. solani* in the RCR complex in Rwanda. To complement the methods used in the previous studies conducted in Zambia and Rwanda, we implemented both amplicon sequencing and microsatellite-based methods as they reveal a finer scale genetic profile of pathogens rather than phenotypic profiles obtained *via* only using cultural methods (Hilton et al. 2016; Mukuma et al. 2020).

In spite of the limitations of not being able to culture all possible fungal isolates, the fungi that we did isolate could be directly tested for pathogenicity to screen dry bean germplasm for disease resistance. Collectively, we identified 333 fungi and/or Oomycetes isolates from Chokwe and Gurue regions in the 2014 and 2015 growing seasons and tested these for pathogenicity against a diversity panel of beans. At least 60% of the isolates were pathogenic to bean, and species of *Fusarium*, primarily *F. oxysporum*, collected from all locations and years produced the longest mean lesion lengths in our pathogenicity assays. Based on the isolation frequency from the field collections and overall pathogenicity observed across the bean diversity panel, we conclude that *F. oxysporum* plays the most predominant role in the bean RCR complex in Mozambique.

More than one pathogen causing RCR of bean was additionally reported by Rusuku et al. (1997) and Mukuma et al. (2020), suggesting a complex of pathogens such as *Pythium* spp., *R. solani*, *M. phaseolina*, *F. oxysporum* f. sp. *phaseoli*, and *A. rolfsii* may be interacting to cause bean RCR. The hypothesis that more than one pathogen causes RCR on beans may have implications in the identification and subsequent breeding in response to broad pathogen-specific RCR disease resistance (Abawi and Pastor-Corrales 1990; Wortmann et al. 1998; Chaudhary et al. 2006; Clare et al. 2010).

With regard to the *Fusarium* species we identified, *F. oxysporum* was the most frequently isolated in our study and identified among the most pathogenic fungi and/or Oomycetes isolated within Mozambique. These results support other studies conducted in Latin America, Spain, and the USA, where *F. oxysporum* has caused RCR outbreaks (Pastor-Corrales and Abawi, 1987; Alves-Santos et al. 2002). In addition, *F. oxysporum* has also been identified as a major pathogen in Central Africa where serious losses on improved climbing bean varieties were reported

(Buruchara and Camacho 2000). This study found that pathogenic isolates of *Fusarium*, with *F. oxysporum* being most notable, were coexisting with other individual *Fusarium* taxa that were not observed to be pathogenic within the same plant. The finding that some of the *Fusarium* species were pathogenic and others were not detrimental to a plant host may be explained by their ubiquitous nature in habitats such as soil, water, and as plant endophytes (Lofgren et al. 2018). Therefore, it is not unusual to have pathogenic and non-pathogenic isolates together in a healthy plant, which contribute to the early development and severity of disease when conditions are unfavorable for bean growth (Leslie and Summerell 2006; Estevez de Jensen et al. 2004; Harveson et al. 2005). It is important to note that *Fusarium* taxa that are typically recognized as opportunistic pathogens and will infect plants under the induction of stressful conditions such as those common in changing climate scenarios (Valverde-Bogantes et al. 2019). We note that, along with *F. oxysporum*, a large portion of the *Fusarium* species complexes identified here were identified as *F. solani* and *F. equiseti* and that we did not investigate in more detail here. We expect these species complexes will be the focus of additional studies inspecting morphotypes and genetic diversity of all RCR bean-associated *Fusarium* taxa.

Based on the amplicon sequencing portion of our study, we identified a diverse community of fungal and Oomycete taxa associated with RCR diseases of bean in two geographically distant regions in Mozambique. A relatively high fungal diversity in different environments has been reported in other studies of the dry bean microbiomes in tropical regions (Pastor-Corrales and Abawi 1987; Alves-Santos et al 2002, Buruchara and Camacho 2000), supporting high-throughput sequencing methods which provide better estimates of the overall fungal diversity of a sample when compared to other identification methods. High-throughput sequencing methods, such as the amplicon sequencing methods we employed in this study, are sensitive methods to identify pathogens associated with RCR when compared to culture-based techniques using morphological approaches that are better suited to only detect individual taxa or those that are easier to culture. Culture-based assessments of diversity can be problematic as some particular species can be isolated with ease in a laboratory setting as they rapidly respond to simple media formulations and are quick to outgrow other organisms. It has been estimated that many fungal taxa are difficult or are unable to be cultured

with our current methodology which hinders the process of identification and experimentation (Hilton et al. 2016). We emphasize that the importance of culturing is vital for establishing cultural material of both fungal and/or Oomycete pathogens which are needed for screening bean lines for RCR resistance.

As previously mentioned, the DNA sequencing methods we used as identification methods were not always in agreement with all the isolated cultures we collected. In general, the combination of molecular- and morphological-based methods, such as the microsatellite and cultural isolation methods we employed here, was better for identifying *Macrophomina phaseolina*, which was not detected by amplicon sequencing from either the FTA® Cards or direct RCR lesions. DNA-based methods, such as high-throughput Illumina amplicon sequencing and PCR-based microsatellite amplification we used in this study, are the most sensitive technique available for the detection of plant pathogens, due in part to the overall depth of sequencing and the specificity of the oligonucleotide primers designed for *Fusarium* taxa (Capote et al. 2012). We hypothesize that the poor amplification of numerous *Macrophomina phaseolina* isolates can be explained by a lack of specificity regarding the oligonucleotide primers and probes, but we have yet to test this hypothesis. Sanger-based sequencing analyses were based on the direct analysis of the genomic DNA of the ITS-2 region which may be variable in the case of *M. phaseolina*. In addition, the high recovery of *M. phaseolina* isolates in samples from the Chokwe region in 2015 may be attributed to environmental conditions in the field, such as high humidity and temperature, where *M. phaseolina* has a high optimal temperature for growth and infection (Songa and Hillocks 1996).

Amplicon sequencing using the Illumina high-throughput platform was used to compare the efficacy of the FTA® Card method for collecting fungi and/or Oomycetes with direct isolation from RCR tissue. The taxonomic diversity assessed by both methods were highly correlated, indicating that FTA® Cards are an acceptable alternative for collection and storage of DNA, particularly in locations where DNA collection *via* direct placement in liquid buffers may be prohibitive, such as field studies. These results are in agreement with the findings obtained by Ndunguru et al. (2005) who used FTA® Cards for sampling and retrieval of DNA and RNA viruses from plant tissues and conducted molecular analysis on viral diversity. Additionally, the proper use of FTA® Cards for DNA

collection and storage largely circumvents the security issues related to the import and export of infected plant tissues. These findings demonstrate the importance of using a combination of diagnostic methods to address identification and phylogenetic placement of primary pathogens associated with host-plants. In this study we used specific primers for the ITS-2 and 18S rDNA regions from RCR fungal pathogen isolates, in addition to species-specific oligonucleotide hybridization, which led to the sequencing and analysis of target DNA suited for both taxonomic and phylogenetic assessment of diversity (Herr et al. 2015; Hibbett et al. 2016). However, we stress the importance of culture-based methods for establishing fungal isolate collections that can be used to test for pathogenicity in field and greenhouse experiments with dry beans and assist in breeding efforts to select for pathogen resistance.

A greater amount of genetic variance within fungal populations from a specific location, rather than between the two regions, was most notable in our microsatellite analysis. Despite using the relatively small suite of microsatellites we tested, our results suggest that either gene flow of the fungi exist between the two regions we surveyed or the fungal isolates are broad in their geographic ranges. Even though the Chokwe and Gurue regions are 1200 km apart, this finding may be explained by a number of factors. For example, many fungi may be seed derived, and an active bean seed exchange between farming locations in Mozambique, as well as with neighboring countries such as Zambia, could be contributing to broad geographic ranges of these fungi. *Fusarium oxysporum* is known to be seed transmitted (Gargouri et al. 2000; Garibaldi et al. 2004; Pires da Silva et al. 2014) and this species complex consists of morphotypes consisting of similar genotypes with diverse lineages (Pires da Silva et al. 2014; Gordon 2017). Additionally, the *F. oxysporum* species complex is not well understood and has no identified sexual reproduction system, and, as a result, the diversity among the isolates may be typical resulting in a limited number of clonal lineages on a large geographic scale (Gordon and Martyn 1997; Valverde-Bogantes et al. 2019).

To our knowledge, this is the first study which utilized both molecular and culture methods to determine the primary pathogens associated with RCR disease of bean in Mozambique. The fact that there are similar pathogen profiles in the two main bean-growing regions of Mozambique allows breeders to broadly screen for resistance to *Fusarium* species, mainly *F. oxysporum*, and cultural methods can provide the pathogen

isolates needed for screening across the country as a whole. Currently, the breeding programs in East Africa are predominantly evaluating *Pythium* spp.; however, the results from this study should provide relevant pathogen information and we hope that it will inform breeding for bean and root rot disease resistance in Mozambique. Lastly, we hope that the findings presented here will provide a basis for future studies of agroecosystem microbiomes (Dundore-Arias et al. 2020) at a finer sampling scale for RCR diseases of dry bean in Mozambique, as well as other tropical and temperate growing regions.



Supplementary Information Tables S-1 through S-5 follow the **References**.

Acknowledgments We recognize the help of many regional farmers, most notably R. Joseph, J. Tenga, and E. Langa, as well as many who remain anonymous, in the field identification and collection of beans with RCR symptoms. We acknowledge the help of Edgar Nieto who provided recommendations on heat map visualization tools using the R statistical framework. We also thank the anonymous reviewers who provided feedback and comments on the initial submission of this article.

Author contribution GG-L, CJ, CU, KE, JRS, and JRH were involved in planning and supervised the work; SF, CJ, and GG-L performed the experiments; GG-L and JRH processed the experimental data and designed the figures; SF, GG-L, JRS, and JRH drafted the manuscript. All authors discussed the results and commented on the manuscript.

Funding This work was completed using the Holland Computing Center of the University of Nebraska, which receives support from the Nebraska Research Initiative. JRS recognizes funding from USDA-NIFA (Grant #11172136) which directly supported this research. JRH acknowledges funding from the US National Air & Space Administration (Grant #80NSSC17K0737), the US National Science Foundation (EPSCoR Grant #1557417), and US National Institute of Justice (Grant #2017-IJ-CX-0025), as well as start-up funding from the University of Nebraska Agricultural Research Division and the University of Nebraska Office of Research and Economic Development, which indirectly supported this research through the support of research facilities. The funding agencies had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Data availability The data sets generated and/or analyzed during the current study are available in the following data repository: https://github.com/HerrLab/Fernandes_et_al_2019

Conflict of interest The authors have declared no competing interests.

References

- Abawi GS (1989) Root rots. In: Bean production problems in the tropics. Centro Internacional de Agricultura Tropical (CIAT), Cali, Colombia, 654 p
- Abawi GS, Pastor-Corrales MA (1990) Root rots of beans in Latin America and Africa: diagnosis, research methodologies, and management strategies. Centro Internacional de Agricultura Tropical (CIAT), Cali, Colombia, 114 p
- Abd-Elsalam KA, Mohamed IN, Abdel-Sattar MA, Khalil MS, Verreet JA (2003) PCR identification of *Fusarium* genus based on nuclear ribosomal-DNA sequence data. *African Journal of Biotechnology* 2:82–85
- Alves-Santos FM, Cordeiro-Rodrigues L, Sayagués JM, Martín- Domínguez R, García-Benavides P, Crespo MC, Díaz-Mínguez JM, Eslava AP (2002) Pathogenicity and race characterization of *Fusarium oxysporum* f. sp. *phaseoli* isolates from Spain and Greece. *Plant Pathology* 51:605–611
- Babu BK, Srivastava AK, Saxena AK, Arora DK (2007) Identification and detection of *Macrophomina phaseolina* by using species-specific oligonucleotide primers and probe. *Mycologia* 99:733–739
- Bahkali AH, Abd-Elsalam KA, Guo JR, Khiyami MA, Verreet JA (2012) Characterization of novel di-, tri-, and tetranucleotide microsatellite primers suitable for genotyping various plant pathogenic fungi with special emphasis on *Fusaria* and *Mycosphaerella graminicola*. *International Journal of Molecular Sciences* 13:2951–2964
- Bodah ET (2017) Root rot diseases in plants: a review of common causal agents and management strategies. *Agricultural Research and Technology* 5:555661
- Brasileiro BTRV, Coimbra MRM, Morais MA Jr, Oliveira NT (2004) Genetic variability within *Fusarium solani* species as revealed by PCR-fingerprinting based on PCR markers. *Brazilian Journal of Microbiology* 35:205–210
- Burgess LW, Summerell BA, Bullock S, Gott KP, Backhouse D (1994) Laboratory manual for *Fusarium* research. The University of Sydney, Sydney, AUS - Royal Botanic Gardens Press, Sydney
- Buruchara R, Camacho L (2000) Common bean reaction to *Fusarium oxysporum* f. sp. *phaseoli*, the cause of severe vascular wilt in Central Africa. *Journal of Phytopathology* 148:39–45
- Capote N, Aguado A, Pastrana AM, Sánchez-Torres P (2012) Molecular tools for detection of plant pathogenic fungi and fungicide resistance. In: Molecular tools for detection of plant pathogenic fungi and fungicide resistance. InTechOpen Press. <https://doi.org/10.5772/38011>
- Chaudhary S, Anderson TR, Park SJ, Yu K (2006) Comparison of screening methods for resistance to *Fusarium* root rot in common beans (*Phaseolus vulgaris* L.). *Journal of Phytopathology* 154:303–308
- Chiodini RJ, Dowd SE, Galandiuk S, Davis B, Glassing A (2016) The predominant site of bacterial translocation across the intestinal mucosal barrier occurs at the advancing disease margin in Crohn's disease. *Microbiology* 162:1608–1619
- Cichy KA, Porch TG, Beaver JS, Cregan P, Fourie D, Glahn RP, Grusak MA, Kamfwa K, Katuramu DN, McClean P, Mndolwa ES, Nchimbi MS, Pastor-Corrales MA, Miklas

- PN (2015) A *Phaseolus vulgaris* diversity panel for Andean bean improvement. *Journal of Crop Science* 55:2149–2160
- Clare MM, Melis R, Dereta J, Laing M, Buruchara RA (2010) Identification of sources of resistance to *Fusarium* root rot among selected common bean lines in Uganda. *Journal of Animal and Plant Sciences* 7:876–891
- Cole JR, Chai B, Farris RJ, Wang Q, Kulam SA, McGarrell DM, Garrity GM, Tiedje JM (2005) The Ribosomal Database Project (RDP-II): sequences and tools for high-throughput rRNA analysis. *Nucleic Acids Research* 33:D294–D296
- Core Team R (2018) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna <https://www.R-project.org/>
- DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen PL (2006) Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and Environmental Microbiology* 72:5069–5072
- Dhingra OD, Sinclair JB (1978) *Biology and pathology of Macrophomina phaseolina*. Imprensa Universitária, Universidade Federal de Viçosa, Viçosa, 166 pp
- Drummond AJ, Rambaut A (2007) BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* 7:214
- Dugan FM (2006) *The identification of fungi: an illustrated introduction with keys, glossary and guide to literature*. APS Press, St Paul, 176 pp
- Dundore-Arias JP, Eloë-Fadrosch EA, Schriml LM, Beattie GA, Brennan FP, Busby PE, Calderon RB, Castle SC, Emerson JB, Everhart SE, Eversole K, Frost KE, Herr JR, Huerta AI, Iyer-Pascuzzi AS, Kalil AK, Leach JE, Leonard J, Maul JE, Prithiviraj B, Potrykus M, Redekar NR, Rojas JA, Silverstein KAT, Tomso DJ, Tringe SG, Vinatzer BA, Kinkel LL (2020) Community-driven metadata standards for agricultural microbiome research. *Phytobiomes Journal* 4:115–121
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32:1792–1797
- Estevez de Jensen CE, Kurle JE, Percich JA (2004) Integrated management of edaphic and abiotic factors limiting yield of irrigation soybean and dry bean in Minnesota. *Field Crops Research* 86:211–224
- Farrow A, Musoni D, Cook S, Buruchara R (2011) Assessing the risk of root rots in common beans in East Africa using simulated, estimated, and observed daily rainfall data. *Experimental Agriculture* 47:357–373
- Fisher NL, Burgess LW, Toussoun TA, Nelson PE (1982) Carnation leaves as a substrate for preserving cultures of *Fusarium* species. *Journal of Phytopathology* 72:151–152
- Gargouri S, Hajlaoui MR, Abdennadher M, Marrakchi M (2000) Isolation and morphological and molecular identification of *Fusarium* spp. transmitted by watermelon seeds. *Bulletin OEPP* 30:217–222
- Garibaldi A, Gilardi G, Gullino ML (2004) Seed transmission of *Fusarium oxysporum* f. sp. *lactucae*. *Phytoparasitica* 32:61–65
- Gordon TR (2017) *Fusarium oxysporum* and the *Fusarium* wilt syndrome. *Annual Review of Phytopathology* 55:23–39
- Gordon TR, Martyn RD (1997) The evolutionary biology of *Fusarium oxysporum*. *Annual Review of Phytopathology* 35:111–128

- Guillemaut C, Edel-Hermann V, Camporota P, Alabouvette C, Richard-Molard M, Steinberg C (2003) Typing of anastomosis groups of by restriction analysis of ribosomal DNA. *Can J Microbiol* 49:556–568
- Hammer O, Harper DAT, Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4:1–9
- Harveson RM, Smith JA, Stroup WW (2005) Improving root health and yield of dry bean on Nebraska panhandle with a new technique for reducing soil compaction. *Plant Disease* 89:279–284
- Herr JR, Öpik M, Hibbett DS (2015) Towards the unification of sequence-based classification and sequence-based identification of host-associated microorganisms. *New Phytologist* 205:27–31
- Hibbett D, Abarenkov K, Kõljalg U, Öpik M, Chai B, Cole J, Wang Q, Crous P, Robert V, Helgason T, Herr JR, Kirk P, Lueschow S, O'Donnell K, Nilsson RH, Oono R, Schoch C, Smyth C, Walker DM, Porras-Alfaro A, Taylor JW, Geiser DM (2016) Sequence-based classification and identification of fungi. *Mycologia* 108:1049–1068
- Hilton SK, Castro-Nallar E, Pérez-Losada M, Toma I, McCaffrey TA, Hoffman EP, Siegel MO, Simon GL, Johnson WE, Crandall KA (2016) Metataxonomic and metagenomic approaches vs. culture-based techniques for clinical pathology. *Frontiers in Microbiology* 7:484
- Kamvar ZN, Tabima JF, Grünwald NJ (2014) Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* 2:e281
- Kang HW, Park DS, Go SJ, Eun MY (2002) Fingerprinting of diverse genomes using PCR with universal rice primers generated from repetitive sequence of Korean weedy rice. *Molecules and Cells* 13:281–287
- Leslie JF, Summerell BA (2006) The *Fusarium* laboratory manual. Wiley-Blackwell Publishing, Hoboken, NJ, pp 388
- Letunic I, Bork P (2016) Interactive tree of life (iTOL) version 3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Research* 44:W242–W245
- Lieckfeldt E, Meyer W, Börner T (1993) Rapid identification and differentiation of yeasts by DNA and PCR fingerprinting. *J Basic Microbiol* 33(6):413–425
- Lofgren LA, LeBlanc NR, Certano AK, Nachtigall J, LaBine KM, Riddle J, Broz K, Dong Y, Bethan B, Kafer CW, Kistler HC (2018) *Fusarium graminearum*: pathogen or endophyte of North American grasses? *New Phytologist* 217:1203–1212
- Martin FN (2000) Phylogenetic relationships among some *Pythium* species inferred from sequence analysis of the mitochondrially encoded cytochrome oxidase II gene. *Mycologia* 92:711–727
- McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL, Knight R, Hugenholtz P (2012) An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *The ISME Journal* 6:610–618
- Mukuma C, Godoy-Lutz G, Eskridge K, Steadman J, Urrea C, Muimui K (2020) Use of culture and molecular methods for identification and characterization of dry bean fungal root rot pathogens in Zambia. *Tropical Plant Pathology* 45:385–396

- Nair DN, Padmavathy S (2014) Impact of endophytic microorganisms on plants, environment, and humans. *The Scientific World Journal* 2014:250693
- Narayanasamy P (2011) Detection of fungal pathogens in plants. In: *Microbial plant pathogens-detection and disease diagnosis*. Springer, Dordrecht
- Ndunguru J, Taylor NJ, Yadav J, Aly H, Legg JP, Aveling T, Fauquet CM (2005) Application of FTA technology for sampling, recovery and molecular characterization of viral pathogens and virus-derived transgenes from plant tissues. *Virology Journal* 2:45
- Nei M (1973) Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences USA* 70:3321–3323
- Neuwirth E (2014) RColorBrewer: ColorBrewer palettes. R package version 1:1–2 <https://CRAN.R-project.org/package=RColorBrewer>
- Nzungize J, Gepts P, Buruchara R, Buah S, Ragama P, Busogoro JP, Baudoin JP (2011a) Pathogenic and molecular characterization of *Pythium* species inducing root rot symptoms of common bean in Rwanda. *African Journal of Microbiological Research* 5:1169–1181
- Nzungize J, Gepts P, Buruchara R, Male A, Ragama P, Busogoro JP, Baudoin JP (2011b) Introgression of *Pythium* root rot resistance gene into Rwandan susceptible common bean cultivars. *African Journal of Plant Sciences* 5:193–200
- O'Donnell K, Kistler HC, Cigelnik E, Ploetz RC (1998) Multiple evolutionary origins of the fungus causing Panama disease of banana: concordant evidence from nuclear and mitochondrial gene genealogies. *Proceedings of the National Academy of Sciences USA* 95: 2044–2049
- Paparu P, Acur A, Kato F, Acam C, Nakibuule J, Musoke S, Nikalubo S, Mukankusi C (2018) Prevalence and incidence of four common bean root rots in Uganda. *Experimental Agriculture* 54:888–900
- Park B, Park J, Cheong KC, Choi J, Jung K, Kim D, Lee YH, Ward TJ, O'Donnell K, Geiser DM, Kang S (2010) Cyber infrastructure for *Fusarium*: three integrated platforms supporting strain identification, phylogenetics, comparative genomics, and knowledge sharing. *Nucleic Acids Research* 39:D640–D646
- Pastor-Corrales MA, Abawi GS (1987) Reaction of selected bean germplasm to infection by *Fusarium oxysporum* f. sp. *phaseoli*. *Journal of Plant Disease* 71:990–993
- Peakall R, Smouse PE (2006) GenA1Ex 6: genetic analysis in Excel - population genetic software for teaching and research. *Molecular Ecology Notes* 6:288–295
- Pires da Silva R, Vechiato MH, Harakava R (2014) EF-1 α gene and IGS rDNA sequencing of *Fusarium oxysporum* f. sp. *vasinfectum* and *F. oxysporum* f. sp. *phaseoli* reveals polyphyletic origin of strains. *Tropical Plant Pathology* 39:64–73
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO (2012) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* 41:D590–D596
- Rognes T, Flouri T, Nichols B, Quince C, Mahé F (2016) VSEARCH: a versatile open-source tool for metagenomics. *PeerJ* 4:e2584
- Rusuku G, Buruchara RA, Gatabazi M, Pastor-Corrales MA (1997) Occurrence and distribution in Rwanda of soil-borne fungi pathogenic to the common bean. *Journal of Plant Disease* 81:445–449

- Shannon CE, Wiener W (1963) *The mathematical theory of communication*. University of Illinois Press, Urbana, IL
- Simpson EH (1949) Measurement of diversity. *Nature* 163:688
- Songa W, Hillocks RJ (1996) Charcoal rot in common bean with special reference to Kenya. *International Journal of Pest Management* 42: 213–219
- Valverde-Bogantes E, Bianchini A, Herr JR, Rose DJ, Wegulo SN, Hallen-Adams HE (2019) Recent population changes of *Fusarium* head blight pathogens: drivers and implications. *Canadian Journal of Plant Pathology* 42:315–329
- Walker TS, Pitoro R, Tomo A, Siteo I, Salencia C, Mahanzule R, Donovan C, Mazuze FM (2006) Priority setting for public-sector agricultural research in Mozambique with the National Agricultural Survey Data (No. 1093-2016-88001)
- Watanabe T (2010) Morphologies of cultured fungi and key to species. In: *Pictorial atlas of soil and seed fungi - Morphologies of cultured fungi and key to species*, 3rd Edition. CRC Press, Florida pp 426. <https://www.routledge.com/Pictorial-Atlas-of-Soil-and-Seed-Fungi-Morphologies-of-Cultured-Fungi-and/Watanabe/p/book/9781439804193>
- White TJ, Bruns T, Lee S, Taylor TW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR protocols: a guide to methods and applications*, Academic Press, Inc., New York. pp. 315–322
- Wickham H (2016) *ggplot2: elegant graphics for data analysis*. Springer-Verlag Press, New York
- Wortmann CS, Kirkby RA, Eledu CA, Allen DJ (1998) *Atlas of common bean (Phaseolus vulgaris L.) production in Africa*. Centro Internacional de Agricultura Tropical (CIAT). CIAT publication number 297, Cali, 131 p

Table S-1 – Table summary of microsatellite AMOVA analysis comparing *Fusarium oxysporum* populations isolated at the Chokwe and Gurue locations.

Table S-2 – Table of within population genetic diversity indicators of *F. oxysporum* isolates from the sampling locations of Chokwe and Gurue based on microsatellite primer amplification.

Table S-3 - Table of plant and soil sampling locations, soil metadata characteristics, and climate information for collection sites in this study.

Table S-4 – Table of oligonucleotide primer sequences used in this study.

Table S-5 – Table of information on each isolate obtained from RCR tissue culture of bean plants collected in Mozambique and the closest database match of the NCBI accession number.

Table S-1

Source of Variation	Degrees of Freedom	SS	Mean Square	Estimated Variance	% Variation
URP2R					
Between populations	1	19.453	19.45	0.596	21
Within populations	62	138.641	2.236	2.236	79
Total	63	158.09		2.82	100
Fst value 0.21					
(AGG) ₅					
Between populations	1	17.942	17.94	0.448	10
Within populations	90	351.797	3.909	3.909	90
Total	91	369.739		4.357	100
Fst value 0.10					
(GTG) ₅					
Between populations	1	12.814	12.814	0.282	7
Within populations	71	251.377	3.541	3.541	93
Total	72	264.192	3.823	3.823	100
Fst value 0.07					
(ACG) ₅					
Between populations	1	2.931	2.937	0.018	1
Within populations	89	208.642	2.344	2.344	99
Total	90	211.560		2.362	100
Fst value 0.01					

P = 0.01

Table S-2

Population	URP2R		(GTG) ₅		(ACG) ₅		(AGG) ₅	
	Chokwe	Gurue	Chokwe	Gurue	Chokwe	Gurue	Chokwe	Gurue
No. bands ^a	16	16	17	21	18	26	19	27
No. bands Freq. >=5% ^b	16	11	17	17	12	15	19	25
No. Private bands ^c	3	3	2	6	1	9	1	9
% of polymorphism	55.17	55.17	58.72	72.41	62.07	89.66	65.2	93.10
Mean h ^d	0.211 ±0.039	0.129 ±0.032	0.225 ±0.040	0.227 ±0.036	0.153 ±0.03	0.160 ±0.027	0.196 ±0.034	0.282 ±0.030
Mean uh ^e	0.241 ±0.044	0.132 ±0.033	0.247 ±0.044	0.232 ±0.037	0.161 ±0.032	0.162 ±0.028	0.207 ±0.035	0.286 ±0.031
I ^f	0.312 ±0.056	0.206 ±0.046	0.33 ±0.056	0.34 ±0.051	0.247 ±0.04	0.268 ±0.03	0.304 ±0.049	0.434 ±0.04

^a = No. of different bands

^b = No. of different bands with a frequency >= 5%

^c = No. of bands unique to a single population

^d = Diversity = $1 - (p^2 + q^2)$

^e = Unbiased diversity = $(N / (N-1)) * h$

^f = Shannon's Information index

= $-1 * (p * \ln(p) + q * \ln(q))$

where for Haploid Binary data, p

= Band Freq. and q = 1 - p

Table S-3

Collection		Coordinates (DMS)			Soils characteristics		
Site	Year	Latitude	Longitude	Altitude (masl)	Soil types	Soil organic matter*	Climate
ADP-NE Root rot nursery in Mutequelesse - Gurue, Zambezia Province	2014, 2015	15° 19' 2.55'' S	36° 42' 8.01'' E	690	Ferrossols	Medium	Cool, sub-humid
ADP-NE Root rot nursery field- Chokwe Research Station, Gaza Province	2014, 2015	24° 30' 15'' S	33° 00' 11'' E	34	Salic Fluvisol (Mananga)	Medium to rich	Semi-arid
Farmer's fields- Chate, Chokwe, Gaza Province	2014, 2015	24° 21' 52'' S	32° 50' 32'' E	43	Eutric Fluvisol (Mananga)	Medium to rich	Semi-arid
Farmer's field- Chissano, Gaza Province	2014	N/D	N/D	N/D	Salic Fluvisol (Mananga)	Medium to rich	Semi-arid
DP-NE Root rot nursery field- Chokwe Research Station, Gaza Province	2015	24° 30' 20'' S	33° 00' 07'' E	33	Salic Fluvisol (Mananga)	Medium to rich	Semi-arid
Farmer's field- Camul-Chokwe, Gaza Province	2015	24° 32' 13'' S	33° 01' 09'' E	31	Salic Fluvisol (Mananga)	Medium to rich	Semi-arid
Farmer's field- Camul-Chokwe, Gaza Province	2015	24° 32' 11'' S	33° 01' 05'' E	31	Salic Fluvisol (Mananga)	Medium to rich	Semi-arid

*Medium = 1.6% - 3.0% organic matter, Rich = 3.1% - 6.0% organic matter

Table S-4

Code	Target Gene	Primer Sequence(5'→3')	Annealing Temperature °C	Source
FM 66 FM 58 COX II	Cytochrome oxidase II	TAGGATTTCAAGATCCTGC CCACAAATTTCACTACATTGA	52° C	Martin, 2000
MpkF1 MpkR1	Internal Transcribed Spacer rRNA	CCGCCAGAGGACTATCAAAC CGTCCGAAGCGAGGTGTATT	56°C	Kishore Babbu <i>et al.</i> , 2007
ITSFuF ITSFuR	Internal Transcribed Spacer rRNA	CAACTCCCAAACCCCTGTGA GCGACGATTACCAGTAACGA	58°C	Ed Elsalam KA <i>et al.</i> , 2003
EF1 EF2	Translation Elongation Factor (1- α)	ATGGGTAAGGA(A/G)GACAA GAC GGA(G/A)GTACCAGT(G/C)AT CATGTT	53°C	O'Donnell <i>et al.</i> , 1998
R1 R4	Internal Transcribed Spacer rRNA	CCTGTGCACCTGTGAGACAG TGTCCAAGTCAATGGACTAT	56°C	Camporota <i>et al.</i> , 2000
ITS5 ITS4	Internal Transcribed Spacer rRNA	GGAAGTAAAAGTCGTAACAA GG TCCTCCGCTTATTGATATGC	55°C	White <i>et al.</i> , 1990
URP2R	Entire genome	CCCAGCAACTGATCGCACAC	45°C	Khan <i>et al.</i> , 2002
(GTG) ₅		GTGGTGGTGGTGGTG	50°C	Brasileiro <i>et al.</i> , 2004
(ACG) ₅		ACGACGACGACGACG	52°C	Bahkali <i>et al.</i> , 2012
(AGG) ₅		AGGAGGAGGAGGAGG	50°C	

Table S-5

Isolate ID	Pathogenicity ^a	Location/ Year of Collection	Source Bean line/Variety	Species	NCBI GenBank accession closest match ^b
M1	NP	Chokwe/2014	Mbulamtwe	<i>Alternaria alternata</i>	KT223359.1
M2	NP	Chokwe/2014	Mbulamtwe	<i>Alternaria alternata</i>	KT223325.1
M3	P	Chokwe/2014	Mbulamtwe	<i>Alternaria alternata</i>	KT274695.1
M4	NP	Chokwe/2014	Mbulamtwe	<i>Alternaria</i> sp.	EF432261.1
M5	NP	Chokwe/2014	Mbulamtwe	No similarity found	
M6	NP	Chokwe/2014	Mbulamtwe	<i>Cladosporium</i> sp.	KC178629.1
M7	NP	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KF881762.1
M8	NP	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KF881759.1
M9	NP	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KC178652.1
M10	NP	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KT192329.1
M11	NP	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KU377991.1
M12	P	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KP278185.1
M13	NP	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KT192329.1
M14	NP	Chokwe/2014	Kisapuri	<i>Alternaria arborescens</i>	KM246282.1
M15	P	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KJ605840.1
M16	NP	Chokwe/2014	Kisapuri	<i>Alternaria arborescens</i>	KJ609138.1
M17	P	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	JQ676197.1
M18	NP	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	KJ739880.1
M19	P	Chokwe/2014	Kisapuri	<i>Alternaria alternata</i>	HQ846574.1
M20	NP	Chokwe/2014	Incomparable	<i>Fusarium oxysporum</i>	KJ439205.1
M21	NP	Chokwe/2014	Incomparable	<i>Alternaria macrospora</i>	DQ156342.1
M22	P	Chokwe/2014	Incomparable	<i>Alternaria alternata</i>	KP271958.1
M23	NP	Chokwe/2014	Maharage Makubwa	<i>Cladosporium</i> sp.	KC178629.1

M24	NP	Chokwe/2014	Maharage Makubwa	<i>Alternaria arborescens</i>	KJ609138.1
M25	NP	Chokwe/2014	Sodan	<i>Chaetomium globosum</i>	KU375642.1
M26	NP	Chokwe/2014	Kablanketi	<i>Mucor circinelloides</i> f. <i>lusitanicus</i>	JF439687.1
M27	NP	Chokwe/2014	P1321094-D	<i>Alternaria alternata</i>	KJ739872.1
M28	P	Chokwe/2014	P1321094-D	<i>Alternaria alternata</i>	KX115415.1
M29	NP	Chokwe/2014	NE34-12-20	<i>Curvalaria lunata</i>	KF498867.1
M30	NP	Chokwe/2014	NE34-12-20	<i>Curvalaria lunata</i>	KF498867.1
M31	NP	Chokwe/2014	NE34-12-20	<i>Curvalaria lunata</i>	KY404178.1
M32	NP	Chokwe/2014	NE34-12-45	<i>Alternaria alternata</i>	KJ605840.1
M33	P	Chokwe/2014	NE34-12-45	<i>Alternaria alternata</i>	KT274695.1
M34	NP	Chate /2014	PAN 127	<i>Cladosporium halotolerans</i>	LN834369.1
M35	P	Chate /2014	PAN 127	<i>Fusarium oxysporum</i>	KJ528881.1
M36	P	Chate /2014	PAN 127	<i>Fusarium oxysporum</i> f. <i>sp. ciceris</i>	HG423346.1
M37	NP	Chate /2014	PAN 127	<i>Fusarium oxysporum</i>	KX196809.1
M38	NP	Chate /2014	PAN 127	<i>Chaetomium globosum</i>	KR063144.1
M39	NP	Chate /2014	PAN 128	<i>Cladosporium cladosporioides</i>	KX664415.1
M40	NP	Chate /2014	PAN 128	<i>Cladosporium tenuissimum</i>	EU272531.1
M41	P	Chate /2014	PAN 128	<i>Fusarium solani</i>	KM235740.1
M42	P	Chate /2014	PAN 128	<i>Fusarium oxysporum</i>	KU872849.1
M43	NP	Chate /2014	PAN 127	<i>Marasmius brunneoaurantiacus</i> voucher	KX149014.1
M44	NP	Chate /2014	PAN 127	<i>Marasmius brunneoaurantiacus</i> voucher	KX149014.1
M45	NP	Chate /2014	PAN 127	<i>Fusarium verticillioides</i>	KF897854.1
M46	P	Chate /2014	PAN 127	<i>Fusarium oxysporum</i>	KU872849.1

M47	P	Chate /2014	PAN 127	<i>Fusarium nyagamai</i>	HF546381.1
M48	NP	Chokwe/2014	PAN 127	<i>Alternaria alternata</i>	KT192402.1
M49	NP	Chokwe/2014	PAN 127	<i>Alternaria alternata</i>	KF881762.1
M50	NP	Chokwe/2014	PAN 127	<i>Alternaria alternata</i>	KT192393.1
M51	P	Chokwe/2014	PAN 127	<i>Alternaria alternata</i>	KJ739880.1
M52	P	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	KU872840.1
M53	P	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	GU724513.1
M54	NP	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	KP050556.1
M55	P	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	KM519660.1
M56	NP	Chissano/2014	PAN 148	<i>Fusarium equiseti</i>	KR094440.1
M57	P	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	KX196807.1
M58	P	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	JF300424.1
M59	NP	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	KX196807.1
M60	P	Chissano/2014	PAN 148	<i>Fusarium oxysporum</i>	GU724513.1
M61	P	Chissano/2014	Icapijão	<i>Fusarium solani</i>	HQ439152.1
M62	P	Chissano/2014	Icapijão	<i>Fusarium solani</i>	EU625405.1
M63	P	Chissano/2014	Icapijão	<i>Fusarium solani</i>	JQ277276.1
M64	P	Chissano/2014	Icapijão	<i>Fusarium solani</i>	KC764913.1
M65	P	Chissano/2014	Icapijão	<i>Fusarium oxysporum</i>	KU056819.1
M66	P	Chissano/2014	Icapijão	<i>Fusarium solani</i>	JN006817.1
M67	NP	Chissano/2014	Icapijão	<i>Rhizoctonia zeae</i>	GQ221863.1
M68	NP	Chissano/2014	Icapijão	<i>Rhizoctonia zeae</i>	GQ221863.1
M69	NP	Chissano/2014	Icapijão	<i>Rhizoctonia zeae</i>	GQ221863.1
M70	NP	Chissano/2014	Icapijão	<i>Fusarium solani</i>	KJ528882.1
M71	NP	Chissano/2014	AP 89	<i>Thielavia terricola</i>	KJ921610.1
M72	NP	Chissano/2014	AP 89	<i>Thielavia terricola</i>	GU966509.1
M73	P	Chissano/2014	AP 89	<i>Fusarium oxysporum</i>	KX196807.1
M74	NP	Chissano/2014	AP 89	<i>Chaetomium funicola</i>	KM979902.1
M75	P	Chissano/2014	AP 89	<i>Fusarium solani</i>	KM235740.1

M76	NP	Chissano/2014	Bonus	<i>Epicocum nigrum</i>	KF881763.1
M77	NP	Chissano/2014	Bonus	<i>Epicocum nigrum</i>	KF881763.1
M78	NP	Chissano/2014	Bonus	<i>Epicocum nigrum</i>	KX664321.1
M79	NP	Chissano/2014	Bonus	<i>Epicocum nigrum</i>	KM519661.1
M80	NP	Chissano/2014	Bonus	<i>Epicocum nigrum</i>	KM519661.1
M81	NP	Chissano/2014	Bonus	<i>Epicocum nigrum</i>	KM519661.1
M82	P	Chissano/2014	LPA 91	<i>Fusarium solani</i>	KF918580.1
M83	NP	Chissano/2014	LPA 91	<i>Fusarium oxysporum</i>	KU872849.1
M84	NP	Chissano/2014	LPA 91	<i>Fusarium oxysporum</i>	KU872849.1
M85	P	Chissano/2014	LPA 91	<i>Fusarium equiseti</i>	KF918580.1
M86	NP	Chissano/2014	LPA 91	<i>Fusarium thapsinum</i>	KX171659.1
M87	NP	Chissano/2014	LPA 91	<i>Fusarium thapsinum</i>	KX171659.1
M88	NP	Chissano/2014	LPA 91	<i>Fusarium thapsinum</i>	KM589051.1
M89	P	Chissano/2014	LPA 91	<i>Fusarium thapsinum</i>	KM589051.1
M90	P	Chissano/2014	LPA 91	<i>Fusarium oxysporum</i>	KU872849.1
M91	P	Chissano/2014	LPA 91	<i>Fusarium equiseti</i>	KR094440.1
M92	NP	Gurue/2014	Kiangwe	<i>Trichoderma harzianum</i>	KR868296.1
M93	NP	Gurue/2014	Kiangwe	<i>Trichoderma atrobrunneum</i>	NR_137298.1
M94	NP	Gurue/2014	Kiangwe	<i>Trichoderma harzianum</i>	KR868296.1
M95	NP	Gurue/2014	Kiangwe	<i>Trichoderma harzianum</i>	HG940486.1
M96	NP	Gurue/2014	Kiangwe	<i>Trichoderma harzianum</i>	KR868309.1
M97	NP	Gurue/2014	Kiangwe	<i>Trichoderma harzianum</i>	KU696482.1
M98	P	Gurue/2014	Goloi	<i>Fusarium proliferatum</i>	KP760063.1
M99	P	Gurue/2014	Goloi	<i>Fusarium circinatum</i>	KC464621.1
M100	P	Gurue/2014	Goloi	<i>Fusarium verticillioides</i>	KX196811.1
M101	P	Gurue/2014	Goloi	<i>Fusarium verticillioides</i>	KX196811.1
M102	P	Gurue/2014	Kablanketi	<i>Fusarium equiseti</i>	EU625404.1

M103	P	Gurue/2014	Kablanketi	<i>Fusarium incarnatum</i>	KM921663.1
M104	P	Gurue/2014	Kablanketi	<i>Fusarium equiseti</i>	KR047055.1
M105	P	Gurue/2014	Kablanketi	<i>Fusarium equiseti</i>	KF863780.1
M106	NP	Gurue/2014	Kablanketi	<i>Fusarium equiseti</i>	KF863780.1
M107	NP	Gurue/2014	Kablanketi	<i>Fusarium equiseti</i>	HM008677.1
M108	NP	Gurue/2014	Kablanketi	<i>Fusarium incarnatum</i>	KU680357.1
M109	NP	Gurue/2014	Kablanketi	<i>Fusarium equiseti</i>	HM008677.1
M110	P	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	KR094440.1
M111	P	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	KR094440.1
M112	P	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	KU856645.1
M113	P	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	KU856645.1
M114	P	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	KU856645.1
M115	NP	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	KU856645.1
M116	P	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	JQ936262.1
M117	P	Gurue/2014	Sodan	<i>Fusarium equiseti</i>	JQ936262.1
M118	P	Gurue/2014	G5087	<i>Nectria haematococca</i>	KJ780750.1
M119	P	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	KP050556.1
M120	P	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	HQ439152.1
M121	P	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	JN006816.1
M122	P	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	HQ439152.1
M123	NP	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	KX196809.1
M124	NP	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	KT223349.1
M125	P	Gurue/2014	G5087	<i>Fusarium solani</i>	EU625405.1
M126	NP	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	KX196809.1
M127	P	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	KC202939.1
M128	P	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	KM817209.1
M129	P	Gurue/2014	G5087	<i>Fusarium oxysporum</i>	KU931543.1

M130	P	Gurue/2014	G5087	<i>Fusarium solani</i>	JN006816.1
M131	P	Gurue/2014	CAL 143	<i>Fusarium solani</i>	KU377510.1
M132	P	Gurue/2014	CAL 143	<i>Fusarium solani</i>	JQ277276.1
M133	P	Gurue/2014	CAL 143	<i>Fusarium oxysporum</i>	KM817213.1
M134	P	Gurue/2014	CAL 143	<i>Fusarium oxysporum</i>	KJ082096.1
M135	P	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	GQ131884.1
M136	P	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	GQ131884.1
M137	P	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	KF498869.1
M138	P	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	KC202939.1
M139	P	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	KU931543.1
M140	P	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	KU931543.1
M141	P	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	KJ562370.1
M142	NP	Gurue/2014	G22246	<i>Fusarium oxysporum</i>	GQ131884.1
M143	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	KU931543.1
M144	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	KU931543.1
M145	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	KU931553.1
M146	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	KM817213.1
M147	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	HG423346.1
M148	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	KJ082096.1
M149	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	KU059956.1
M150	P	Gurue/2014	Badillo	<i>Fusarium oxysporum</i>	KU056819.1
M151	P	Gurue/2014	INIAP480	<i>Fusarium oxysporum</i>	KR047056.1
M152	P	Gurue/2014	INIAP480	<i>Fusarium oxysporum</i>	KU931543.1
M153	P	Gurue/2014	INIAP480	<i>Fusarium oxysporum</i>	KU056819.1
M154	NP	Gurue/2014	INIAP480	<i>Fusarium equiseti</i>	KU926350.1
M155	P	Gurue/2014	INIAP480	<i>Fusarium oxysporum</i> f. <i>sp. vasinfectum</i>	EU849584.1
M156	P	Gurue/2014	INIAP480	<i>Fusarium equiseti</i>	JQ936262.1

M157	P	Gurue/2014	INIAP480	<i>Fusarium oxysporum</i>	GQ131884.1
M158	NP	Gurue/2014	INIAP480	<i>Fusarium equiseti</i>	JF773657.1
M159	P	Gurue/2014	INIAP480	<i>Fusarium oxysporum</i>	GU724514.1
M160	P	Gurue/2014	Canioca/Kibala	<i>Fusarium oxysporum</i>	KU931543.1
M161	P	Gurue/2014	Canioca/Kibala	<i>Fusarium oxysporum</i>	KU931543.1
M162	P	Gurue/2014	Canioca/Kibala	<i>Fusarium oxysporum</i> f. <i>sp. vasinfectum</i>	KU729045.1
M163	P	Gurue/2014	Canioca/Kibala	<i>Fusarium oxysporum</i>	KU931543.1
M164	P	Gurue/2014	Canioca/Kibala	<i>Fusarium oxysporum</i>	KU059956.1
M165	P	Gurue/2014	Canioca/Kibala	No similarity found	
M166	P	Gurue/2014	Canioca/Kibala	<i>Fusarium oxysporum</i>	KF577910.1
M167	NP	Gurue/2014	ND061106	<i>Fusarium equiseti</i>	KF863780.1
M168	P	Gurue/2014	ND061106	<i>Fusarium equiseti</i>	KF863780.1
M169	NP	Gurue/2014	ND061106	<i>Fusarium equiseti</i>	KR047055.1
M170	P	Gurue/2014	ND061106	<i>Fusarium equiseti</i>	KF863780.1
M171	P	Gurue/2014	ND061106	<i>Fusarium oxysporum</i>	KJ019830.1
M172	P	Gurue/2014	ND061106	<i>Fusarium solani</i>	KU382502.1
M173	P	Gurue/2014	ND061106	<i>Fusarium proliferatum</i>	KY590032.1
M174	NP	Gurue/2014	H9659-27-10	<i>Fusarium oxysporum</i>	EU839377.1
M175	P	Gurue/2014	H9659-27-10	<i>Fusarium oxysporum</i>	KT898585.1
M176	NP	Gurue/2014	H9659-27-10	<i>Talaromyces</i> <i>pinophilus</i>	LT558962.1
M177	NP	Gurue/2014	H9659-27-10	<i>Penicillium pinophilum</i>	FJ441618.1
M178	P	Gurue/2014	Krimson	<i>Fusarium</i> <i>brachygibbosum</i>	KF985966.1
M179	P	Gurue/2014	NE34-12-30	<i>Fusarium oxysporum</i> f. <i>sp. momordicae</i>	JN005749.1
M180	P	Gurue/2014	NE34-12-30	<i>Fusarium equiseti</i>	KJ562376.1
M181	NP	Gurue/2014	NE34-12-30	<i>Fusarium oxysporum</i>	KU931553.1

M182	P	Gurue/2014	NE34-12-30	<i>Fusarium oxysporum</i>	KU984712.1
M183	NP	Gurue/2014	NE34-12-30	<i>Fusarium oxysporum</i>	KU931554.1
M184	P	Gurue/2014	NE34-12-30	<i>Fusarium oxysporum</i> f. <i>sp. dianthi</i>	EF421235.1
M185	P	Gurue/2014	NE34-12-30	<i>Fusarium proliferatum</i>	KT207283.1
M186	P	Gurue/2014	NE34-12-30	<i>Fusarium equiseti</i>	KU680356.1
M187	P	Gurue/2014	NE34-12-30	<i>Fusarium oxysporum</i>	HF566400.1
M188	P	Gurue/2014	NE34-12-30	<i>Fusarium oxysporum</i>	KU931543.1
M189	P	Gurue/2014	NE34-12-37	<i>Rhizoctonia solani</i>	KM013470.1
M190	P	Gurue/2014	NE34-12-37	<i>Rhizoctonia solani</i>	KM013470.1
M191	P	Gurue/2014	NE34-12-38	<i>Fusarium oxysporum</i>	KJ528881.1
M192	P	Gurue/2014	NE34-12-38	<i>Fusarium oxysporum</i>	KX196809.1
M193	P	Gurue/2014	NE34-12-38	<i>Fusarium oxysporum</i>	KX196807.1
M194	P	Gurue/2014	NE34-12-38	<i>Fusarium oxysporum</i>	KU056819.1
M195	P	Gurue/2014	NE34-12-48	<i>Fusarium oxysporum</i>	KU056819.1
M196	P	Gurue/2014	NE34-12-48	<i>Fusarium oxysporum</i>	GU181389.2
M197	P	Gurue/2014	NE34-12-48	<i>Fusarium oxysporum</i>	KT207755.1
M198	P	Gurue/2014	NE34-12-48	<i>Fusarium oxysporum</i>	KJ439149.1
M199	NP	Gurue/2014	RR372	<i>Fusarium solani</i>	KJ528882.1
M200	NP	Gurue/2014	Manteiga	<i>Cochliobolus sativus</i>	JQ753975.1
M201	P	Gurue/2014	Manteiga	<i>Fusarium equiseti</i>	KU856645.1
M202	P	Gurue/2014	Manteiga	<i>Fusarium oxysporum</i>	KM817213.1
M203	P	Gurue/2014	Manteiga	<i>Fusarium equiseti</i>	KU856645.1
M204	P	Gurue/2014	Manteiga	<i>Fusarium thapsinum</i>	KU680377.1
M205	NP	Gurue/2014	Manteiga	<i>Fusarium thapsinum</i>	KU680377.1
M206	NP	Gurue/2014	Manteiga	<i>Fusarium equiseti</i>	KU680356.1
M207	NP	Gurue/2014	Manteiga	<i>Fusarium oxysporum</i>	KU056816.1
M208	P	Gurue/2014	Manteiga	<i>Fusarium oxysporum</i>	KU056819.1

M209	P	Gurue/2014	Manteiga	<i>Fusarium oxysporum</i>	KJ544916.1
M210	P	Gurue/2014	Manteiga	<i>Fusarium oxysporum</i>	KF278962.1
M211	P	Gurue/2014	Manteiga	<i>Fusarium oxysporum</i>	KF278962.1
M212	P	Gurue/2014	Manteiga	<i>Fusarium oxysporum</i>	KX196807.1
M213	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium oxysporum</i>	KU056819.1
M214	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium solani</i>	KJ696540.1
M215	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium oxysporum</i>	KT223349.1
M216	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium oxysporum</i>	HF566400.1
M217	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium oxysporum</i>	HM346538.1
M218	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium solani</i>	HQ439152.1
M219	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium solani</i>	KJ696540.1
M220	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium oxysporum</i>	GU724514.1
M221	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium oxysporum</i> f. sp. <i>ciceris</i>	KM817208.1
M222	NP	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium oxysporum</i>	HF566400.1
M223	P	Gurue/2014	VTTTG25 5-1-2	<i>Fusarium solani</i>	KM235740.1
M224	P	Chokwe/2015	INIAP414	<i>Macrophomina phaseolina</i>	KU680393.1
M225	P	Chokwe/2015	INIAP414	<i>Macrophomina phaseolina</i>	KU856652.1
M226	P	Chokwe/2015	INIAP414	<i>Macrophomina phaseolina</i>	KF951750.1
M227	P	Chokwe/2015	INIAP414	<i>Macrophomina phaseolina</i>	KM519193.1
M228	NP	Chokwe/2015	NE34-12-28	<i>Phoma multirostrata</i>	JN542527.1
M229	NP	Chokwe/2015	PI321094-D	<i>Phoma multirostrata</i>	JN542527.1
M230	NP	Chokwe/2015	PI321094-D	<i>Phoma multirostrata</i>	JN542527.1

M231	NP	Chokwe/2015	PI321094-D	<i>Setosphaeria rostrata</i>	KT265240.1
M232	NP	Chokwe/2015	PI321094-D	<i>Peyronellaea glomerata</i>	KR012905.1
M233	NP	Chokwe/2015	PI321094-D	<i>Phoma</i> sp.	KM979987.1
M234	NP	Chokwe/2015	Uyole 98	<i>Phoma multirostrata</i>	KU529840.1
M235	NP	Chokwe/2015	Uyole 98	<i>Phoma multirostrata</i>	KU529840.1
M236	NP	Chokwe/2015	Hutterite	<i>Phoma multirostrata</i>	KU529840.1
M237	NP	Chokwe/2015	Hutterite	<i>Aspergillus calidoustus</i>	HG964947.1
M238	NP	Chokwe/2015	NE34-12-50	<i>Setosphaeria rostrata</i>	KT265240.1
M239	NP	Chokwe/2015	NE34-12-50	<i>Setosphaeria rostrata</i>	KT265240.1
M240	P	Chokwe/2015	NE34-12-50	<i>Fusarium oxysporum</i>	KJ082096.1
M241	P	Chokwe/2015	NE34-12-50	<i>Fusarium solani</i>	KM235740.1
M242	P	Chokwe/2015	PI321094-D	<i>Macrophomina phaseolina</i>	HM990163.1
M243	P	Chokwe/2015	PI321094-D	<i>Macrophomina phaseolina</i>	FJ643531.1
M244	NP	Chokwe/2015	PI321094-D	<i>Phoma multirostrata</i>	JN542527.1
M245	P	Chokwe/2015	PI321094-D	<i>Macrophomina phaseolina</i>	HQ649831.1
M246	P	Chokwe/2015	Kiangwe	<i>Fusarium oxysporum</i>	KM268692.1
M247	P	Chokwe/2015	Kiangwe	<i>Fusarium solani</i>	KM235740.1
M248	NP	Chokwe/2015		<i>Alternaria alternata</i>	KP271958.1
M249	NP	Chokwe/2015	Hutterite	<i>Curvularia hawaiiensis</i>	HG778990.1
M250	NP	Chokwe/2015	Hutterite	<i>Alternaria alternata</i>	KU059951.1
M251	P	Chokwe/2015	RR 375- Local Check	<i>Macrophomina phaseolina</i>	KM519193.1
M252	P	Chokwe/2015	RR 375- Local Check	<i>Fusarium equiseti</i>	EU326202.1
M253	P	Chokwe/2015	RR 375- Local Check	<i>Fusarium oxysporum</i>	KJ207391.1
M254	P	Chokwe/2015	RR 375- Local Check	<i>Fusarium equiseti</i>	KR094440.1

M255	P	Chokwe/2015	RR 375- Local Check	<i>Fusarium solani</i>	KM235740.1
M256	P	Chokwe/2015	G10994	<i>Fusarium equiseti</i>	KR094440.1
M257	P	Chokwe/2015	G10994	<i>Alternaria alternata</i>	KF669893.1
M258	P	Chokwe/2015	G10994	<i>Alternaria alternata</i>	KP003824.1
M259	P	Chokwe/2015	G10994	<i>Fusarium equiseti</i>	KR094440.1
M260	P	Chokwe/2015	G10994	<i>Setosphaeria rostrata</i>	KT265240.1
M261	P	Chokwe/2015	G22246	<i>Phaeosphaeriopsis</i> sp.	JQ936185.1
M262	P	Chokwe/2015	G22246	<i>Phoma multirostrata</i>	JN542527.1
M263	P	Chokwe/2015	G22246	<i>Fusarium oxysporum</i>	KJ082096.1
M264	P	Chokwe/2015	G22246	<i>Phoma</i> sp.	KM516291.1
M265	P	Chokwe/2015	G22246	<i>Phoma multirostrata</i>	KU529840.1
M266	P	Chokwe/2015	G22246	<i>Fusarium oxysporum</i>	KJ082096.1
M267	P	Chokwe/2015	G17913	<i>Macrophomina phaseolina</i>	KU863545.1
M268	P	Chokwe/2015	G10994	<i>Fusarium equiseti</i>	KR094440.1
M269	P	Chokwe/2015	G10994	<i>Macrophomina phaseolina</i>	KU856652.1
M270	P	Chokwe/2015	G10994	<i>Macrophomina phaseolina</i>	HM990163.1
M271	NP	Chokwe/2015	G10994	<i>Chaetomium</i> sp.	HQ608145.1
M272	NP	Chokwe/2015	G10994	<i>Chaetomium</i> sp.	HQ608145.1
M273	P	Chokwe/2015	G22246	<i>Fusarium solani</i>	JN006817.1
M274	P	Chokwe/2015	G22246	<i>Fusarium oxysporum</i>	EU888922.1
M275	P	Gurue/2015	Kiangwe	<i>Fusarium equiseti</i>	KX196808.1
M276	P	Gurue/2015	Kasukanywele	<i>Fusarium oxysporum</i>	KX196809.1
M277	P	Gurue/2015	Kasukanywele	<i>Fusarium solani</i>	KJ620369.1
M278	P	Gurue/2015	SUG-131	<i>Pythium ultimum</i> var. <i>sporangiiferum</i>	AJ628986.1

M279	P	Gurue/2015	SUG-131	<i>Fusarium equiseti</i>	KX196808.1
M280	P	Gurue/2015	SUG-131	<i>Fusarium equiseti</i>	KR094440.1
M281	NP	Gurue/2015	G1375	<i>Fusarium equiseti</i>	KR094440.1
M282	NP	Gurue/2015	G1375	<i>Phoma</i> sp.	JQ388278.1
M283	P	Gurue/2015	G1375	<i>Epicoccum sorghinum</i>	KX289695.1
M284	P	Gurue/2015	G1375	<i>Fusarium oxysporum</i>	EU839378.1
M285	P	Gurue/2015	G10994	<i>Fusarium equiseti</i>	KR364597.1
M286	P	Gurue/2015	G22246	<i>Fusarium solani</i>	EU625405.1
M287	P	Gurue/2015	Larga Commercial	<i>Fusarium equiseti</i>	KX196808.1
M288	NP	Gurue/2015	Larga Commercial	<i>Rhizoctonia solani</i>	KF372651.1
M289	P	Gurue/2015	Larga Commercial	<i>Fusarium oxysporum</i>	KJ082096.1
M290	NP	Gurue/2015	Larga Commercial	<i>Fusarium oxysporum</i>	GU724513.1
M291	P	Gurue/2015	Larga Commercial	<i>Epicoccum sorghinum</i>	KX289695.1
M292	P	Gurue/2015	Larga Commercial	<i>Alternaria alternata</i>	KM519671.1
M293	NP	Gurue/2015	Larga Commercial	<i>Fusarium equiseti</i>	KX196808.1
M294	NP	Gurue/2015	Larga Commercial	<i>Fusarium oxysporum</i>	KF577910.1
M295	P	Gurue/2015	Larga Commercial	No similarity found	
M296	P	Gurue/2015	INIAP-414	<i>Fusarium oxysporum</i>	GU724513.1
M297	P	Gurue/2015	INIAP-414	<i>Rhizoctonia solani</i>	KF372651.1
M298	NP	Gurue/2015	INIAP-414	<i>Rhizoctonia solani</i>	KF372652.1
M299	NP	Gurue/2015	INIAP-414	<i>Phoma herbarum</i>	KU529842.1
M300	NP	Gurue/2015	INIAP-414	<i>Cladorrhinum samala</i>	FM955447.1
M301	NP	Gurue/2015	INIAP-414	<i>Phoma</i> sp.	JQ388278.1
M302	P	Gurue/2015	INIAP-414	<i>Fusarium proliferatum</i>	KJ528883.1
M303	P	Gurue/2015	INIAP-414	<i>Fusarium oxysporum</i>	KX196809.1

M304	P	Gurue/2015	INIAP-414	<i>Fusarium oxysporum</i>	KX196809.1
M305	NP	Gurue/2015	INIAP-414	<i>Phoma herbarum</i>	KU529842.1
M306	P	Gurue/2015	INIAP-414	<i>Fusarium verticillioides</i>	KU680389.1
M307	P	Gurue/2015	INIAP-414	<i>Fusarium oxysporum</i>	KX196809.1
M308	NP	Gurue/2015	INIAP-414	<i>Epicoccum sorghinum</i>	KT310093.1
M309	P	Gurue/2015	INIAP-414	<i>Fusarium verticillioides</i>	KU680389.1
M310	P	Gurue/2015	INIAP-414	<i>Fusarium oxysporum</i>	GU724514.1
M311	P	Gurue/2015	INIAP-414	<i>Fusarium oxysporum</i>	KX196809.1
M312	P	Gurue/2015	PI321094-D	<i>Fusarium solani</i>	KU377510.1
M313	P	Gurue/2015	PI321094-D	<i>Fusarium oxysporum</i> f. sp. <i>ciceris</i>	KR364590.1
M314	NP	Gurue/2015	NE34-12-28	<i>Fusarium oxysporum</i>	GU724513.1
M315	P	Gurue/2015	NE34-12-28	<i>Fusarium equiseti</i>	KF863780.1
M316	NP	Gurue/2015	NE34-12-28	<i>Epicoccum sorghinum</i>	KX289695.1
M317	NP	Gurue/2015	NE34-12-28	<i>Alternaria alternata</i>	KT223359.1
M318	P	Gurue/2015	NE34-12-28	<i>Fusarium oxysporum</i>	GU724514.1
M319	P	Gurue/2015	A222	<i>Fusarium oxysporum</i> f. sp. <i>vanillae</i>	KM005080.1
M320	P	Gurue/2015	A222	<i>Fusarium oxysporum</i>	JF440593.1
M321	NP	Gurue/2015	A222	<i>Fusarium oxysporum</i>	EU326216.1
M322	P	Gurue/2015	A222	<i>Fusarium oxysporum</i> f. sp. <i>ciceris</i>	KR364590.1
M323	P	Gurue/2015	Magnum	<i>Fusarium oxysporum</i>	KX834820.1
M324	P	Gurue/2015	Magnum	<i>Fusarium solani</i>	JN006817.1
M325	P	Gurue/2015	Magnum	<i>Macrophomina</i> <i>phaseolina</i>	HQ649832.1
M326	NP	Gurue/2015	Magnum	<i>Rhizoctonia solani</i>	JF701745.1
M327	P	Gurue/2015	Magnum	<i>Macrophomina</i> <i>phaseolina</i>	KM979991.1

M328	NP	Gurue/2015	Magnum	<i>Pleosporaceae</i> sp.	HQ832799.1
M329	P	Gurue/2015	Kiangwe	<i>Fusarium proliferatum</i>	KU680369.1
M330	P	Gurue/2015	Kiangwe	No similarity found	
M331	P	Gurue/2015	NE34-12-50	<i>Fusarium falciforme</i>	KC254047.1
M332	P	Gurue/2015	NE34-12-50	<i>Fusarium oxysporum</i>	EF495235.1
M333	P	Gurue/2015	NE34-12-50	<i>Fusarium solani</i>	FJ478114.1

^aP-Pathogenic; NP-Non-pathogenic

^bQuery cover >99%. Identity 97-1