


SPECIAL ISSUE ARTICLE

Overexpression of RNA-binding bacterial chaperones in rice leads to stay-green phenotype, improved yield and tolerance to salt and drought stresses

Rajasheker Guddimalli¹ | Anil Kumar Somanaboina² | Surender Reddy Palle¹ |
 Sujatha Edupuganti³ | Divya Kummari⁴ | Sudhakar Reddy Palakolanu⁴ |
 Jalaja Naravula² | Jawahar Gandra⁵ | Insaf A. Qureshi⁶ | Nagaraju Marka⁷ |
 Rathnagiri Polavarapu⁸  | Polavarapu Bilhan Kavi Kishor²

¹Department of Genetics, Osmania University, Hyderabad, India

²Department of Biotechnology, Vignan's Foundation for Science, Technology and Research, Guntur, India

³Department of Botany, Osmania University, Hyderabad, India

⁴Cell, Molecular & Genetic Engineering Lab, International Crops Research Institute for the Semi-Arid Tropics, Hyderabad, India

⁵Department of Life Sciences, School of Sciences B-II, Jain University, Bengaluru, India

⁶Department of Biotechnology and Bioinformatics, School of Life Sciences, University of Hyderabad, Hyderabad, India

⁷Biochemistry Division, ICMR-National Institute of Nutrition, Hyderabad, India

⁸Research & Development Division, Genomix CARL, Pulivendula, India

Correspondence

Polavarapu Bilhan Kavi Kishor, Department of Genetics, Osmania University, Hyderabad 500 007, India.
 Email: pbkavi@yahoo.com

Funding information

SERB (DST), Grant/Award Number: SERB/SB/SO/PS/104/2013

Edited by: V. Kumar

Abstract

Genes encoding bacterial cold shock proteins A (*CspA*, 213 bp) and B (*CspB*, 216 bp) were isolated from *Escherichia coli* strain K12, which showed 100% homology with gene sequences isolated from other bacterial species. In silico domain analysis showed eukaryotic conserved cold shock domain (CSD) and ribonuclease-binding domain (RBD) indicating that they bind to RNA and are involved in temperature stress tolerance. Overexpression of these two genes in *E. coli* resulted in higher growth in presence of 200 mM NaCl and 300 mM mannitol. Western blot confirmed the translational products of the two genes. Seedlings of indica rice were transformed with *Agrobacterium tumefaciens* containing pCAMBIA1301 *CspA* and *CspB* genes. Transgene integration was confirmed by β -glucuronidase (GUS) histochemical assay, polymerase chain reaction (PCR) amplification, and gene copy number by Southern blotting. Chlorophyll, proline, Na⁺, and K⁺ contents were higher in transgenics exposed to 150 mM NaCl and drought (imposed by withholding water) stresses during floral initiation stage. Catalase (CAT), superoxide dismutase (SOD), and guaiacol peroxidase (GPX) activities increased, while malondialdehyde (MDA) content was low in transgenics. Transgenics displayed increased root, shoot, and panicle lengths, root dry mass, and a distinct stay-green (SGR) phenotype. Higher transcript levels of *CspA*, *CspB*, *SGR*, *chlorophyllase*, isopentenyl adenine transferase 1 (*IPT1*), 9-*cis*-epoxycarotenoid dioxygenase (*NCED*), *SOD*, and sirtuin 1 (*SIRT1*) genes were observed in transgenics compared to wild type plants (WT) under multiple stresses. Present work indicates that bacterial chaperone proteins are capable of imparting SGR phenotype, salt and drought stress tolerance alongside grain improvement.

1 | INTRODUCTION

Salt and drought stresses are the most devastating environmental factors that result in severe crop losses throughout the world. While salt

imposes ionic stress, drought rapidly depletes underground water tables. Often salt and drought stresses are coupled with high temperature stress. Combined effects of major abiotic stresses cause over 50% losses in agricultural crop production (Bray et al., 2020).

Consequently, this entails the generation of crop plants that use water with higher efficiencies (Morison et al., 2008). Breeding methods helped to improve several crop plants for salt and drought stress tolerance, but potential exists for further improvement of crops with better yields and grain quality under different degrees of stress conditions or their combined effects (Fukai & Cooper, 1995; Sreenivasulu et al., 2015). Rice is sensitive to salt and drought stresses (Ismail & Thomson, 2011; Sreenivasulu et al., 2015). Salt stress leads to ion (Na^+ and Cl^-) toxicity, alters cellular metabolism, lowers osmotic potential and photosynthetic efficiency resulting in reduced plant growth rate and yield (Acosta-Motos et al., 2017). Response to drought stress depends upon timing of the stress during early seedling growth, vegetative stage, at the time of flowering or during grain maturity. Terminal drought stress severely impairs the number of filled grains and grain quality (Sreenivasulu et al., 2015). Genetic engineering approaches, such as incorporation of microbial and plant genes associated with biosynthetic pathways of compatible solutes like mannitol, proline, and glycine betaine into crop plants, resulted in the development of abiotic stress tolerant plants (Huang et al., 2000; Kishor et al., 1995; Kishor et al., 2005; Tarczynski et al., 1993; Xu et al., 1996). Further, overexpression of yeast trehalose-6-phosphate synthase gene in tobacco ensued in less dehydration and necrosis (Romero et al., 1997). Similarly, modulation of the polyamine biosynthetic pathway genes in transgenic rice conferred drought tolerance (Capell et al., 2004). Sato and Yokoya (2008) noticed increased drought stress tolerance in rice by overexpressing a small heat-shock protein sHSP17.7. Gu et al. (2013) observed enhanced drought tolerance in transgenic rice overexpressing genes that encode C_4 photosynthetic pathway enzymes. Maheshwari et al. (2017) reported combined drought and high temperature stress tolerance in rice by overexpressing APETALA-type of transcription factor *SbAP37*. Amara et al. (2013) reported enhanced water stress tolerance of transgenic maize overexpressing *LEA Rab28* gene. However, such transgenics did not exhibit greater tolerance to multiple stresses and final productivity under terminal drought stress conditions. Therefore, it is necessary to explore and exploit the extensive genetic diversity that exist in bacteria, fungi, and plants for improving plant abiotic stress tolerance especially during flowering and grain development stages of plant growth. Since, rice is highly susceptible to salt and water stresses, imparting stress tolerance is vital for stabilizing its yields.

In *Escherichia coli*, nine members of the cold shock protein (CSP) gene family *cold shock protein A - cold shock protein I (CspA-CspI)* have been identified, which function as RNA chaperons and help the cells to facilitate proper transcription as well as translation under low temperature stress (Graumann & Marahiel, 1998; Jiang et al., 1997). It has been observed that *CspD* inhibits DNA replication (Uppal et al. 2014), though Csp's in general have been found to play important roles during osmotic, starvation, ethanol, and pH stresses (Keto-Timonen et al., 2016). *CspB* and *CspC* have been found to play critical roles in NaCl, pH, and ethanol stresses, but not *CspA* (Derman et al., 2015). Schmid et al. (2009) found that *CspA*, *CspB*, and *CspD* genes are dispensable for cold and NaCl stress adaptations in *Listeria monocytogenes*. Thus, conflicting roles for CspA have been noticed in literature. CspD protein of *E. coli* has been found to modulate growth-

phase-dependent nutrient-stress adaptation (Yamanaka & Inouye, 1997). On the other hand, in *B. subtilis*, *CspA*, *CspB*, and *CspD* have been found to be associated with normal as well as growth-phase-dependent stress adaptation responses (Weber & Marahiel, 2002). Csp's appear to be essential for growth of bacteria under cold stress conditions and also bind to single stranded RNA and DNA molecules. Of the several Csp's, *CspA* has been shown to unwind the secondary structures of partially double stranded RNA molecules generated during low temperature stress conditions (Jiang et al., 1997). Premature transcription termination occurs due to RNA secondary structure formation in several prokaryotic systems, but *CspA*, *CspC*, and *CspE* have the transcription anti-termination activity (Bae et al., 2000). Jiang et al. (1997) demonstrated increased translation by *CspA* under cold stress conditions through the removal of stabilized RNA secondary structures. Castiglioni et al. (2008) expressed *CspA* and *CspB* genes in maize, which conferred abiotic stress tolerance with improved grain yield. Likewise, improved drought stress tolerance in wheat was noticed with the overexpression of synthetic bacterial cold shock protein gene *SeCspA* (Yu et al., 2017). Transgenic Arabidopsis overexpressing *SeCspA* and *SeCspB* genes displayed better seed germination and enhanced root length under water-deficit conditions when compared to wild-type (WT) plants (Yu et al., 2017). They also noticed important desirable characters like low levels of MDA, Na^+ , higher chlorophyll, and proline levels under salt and water stress conditions. But they did not notice any change in the phenotype under abiotic stress conditions. Field experiments with transgenic wheat containing *SeCspA* gene (but not *SeCspB* transgenics) revealed increased 1000-grain weight as well as grain yield under drought stress (Yu et al., 2017). Accordingly, in the present study, two Csp genes (*CspA* and *CspB*) were cloned and overexpressed in rice for testing against salt and drought stresses during the flowering stage. We report the generation of transgenic rice tolerant to drought and salt stresses (exposed during the floral initiation stage) with significant increase in panicle length and improved yield. Further, for the first time, we report the occurrence of a stay-green (SGR) phenotype as a result of the overexpression of both *CspA* and *CspB* genes. Since a SGR phenotype was noticed, we looked at the expression levels of some of the associated genes like *STAY GREEN* (SGR, linked to drought stress), *CHLOROPHYLLASE* (involved in chlorophyll catabolism), *ISOPENENTYLTRANSFERASE 1 (IPT1)*, associated with cytokinin biosynthesis and control of senescence), *9-CIS-EPOXYCAROTENOID DIOXYGENASE (NCED)*, involved in ABA biosynthesis), copper-zinc chloroplastic *SUPEROXIDE DISMUTASE (SOD)*, associated with scavenging superoxide radicals), *sirtuin 1 (SIRT1)* gene (implicated in leaf senescence and regulation of photosynthetic activity), besides *CspA* and *CspB* (Cucurachi et al., 2012) and linked their expressions with abiotic stresses.

2 | MATERIALS AND METHODS

2.1 | Isolation and vector construction of *CspA* and *CspB* genes

Full length primers were designed using IDT OligoAnalyzer online tool (<https://eu.idtdna.com/calc/analyzer>) for gene sequences available at the NCBI GenBank. The following forward 5'ATGTCCGGTAAAATG

ACTGGTATCG3', 5'ATGTCAAATAAAATGACTGGTTAGTA3' and reverse 5'TTACAGGCTGGTTACGTTACCAG3', 5'TTAATCAGTAATGATGACATTTGCT3' primers were used for *CspA* and *CspB* gene amplifications, respectively, from the *E. coli* K12 strain. PCR conditions used were initial denaturation at 94°C for 5 min, followed by 35 cycles at 94°C for 30 s, annealing at 57°C for *CspA* gene and 54°C for *CspB* for 30 s and extension at 72°C for 30 s; final extension step at 72°C for 5 min using the Biorad C1000 Touch™ Thermal Cycler PCR System. The amplicons were cloned into TA cloning vector pTZ57R/T (Thermo Fischer Fermentas #K0691) separately using T₄ DNA ligase (Fermentas), then transferred into the *E. coli* Top10 strain using 0.1 M CaCl₂ and heat shock at 42°C. The transformed colonies were grown on Luria-Bertani (LB) medium (Bertani, 1951) containing 100 mg L⁻¹ ampicillin, and blue/white colonies were screened with isopropyl β-D-1-thiogalactopyranoside (IPTG)/5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal). Transformed white colonies were picked; the plasmid was isolated by alkaline lysis method, confirmed by polymerase chain reaction (PCR), and the genes were sequenced (Eurofins). The DNA sequences were blasted against NCBI GenBank reference genes and submitted to the National Center for Biotechnology Information (NCBI). Both *CspA* and *CspB* genes were ligated into the intermediate PRT101 vector (3.3 Kb) containing the cauliflower mosaic virus 35S (CaMV35S) promoter and CaMV polyA terminator and transferred into *E. coli* using the *KpnI* and *BamHI* restriction sites. The construct was confirmed with *HindIII* restriction enzyme digestion. The released construct was transferred into the plant binary vector pCAMBIA1301 (13.1 Kb) containing β-glucuronidase (*GUS*) as reporter gene and hygromycin phosphotransferase II (*hptII*) as selection marker. The pCAMBIA1301 plasmid harboring *CspA* and *CspB* genes were separately mobilized into *Agrobacterium tumefaciens* LBA4404 strain. The recombinant colonies were selected on yeast extract peptone (YEP) medium containing 50 mg L⁻¹ kanamycin, and 20 mg L⁻¹ rifampicin.

2.2 | Rice genetic transformation by in planta method

Seeds of indica rice variety BPT5204 were dehusked and sterilized with 0.1% mercuric chloride for 5 min. Seeds were washed thoroughly with sterile distilled water for 3 times. For germination, sterile seeds were soaked in autoclaved water for 2 days in dark at 26–28°C. For each construct (*CspA* and *CspB*), 2000 seedlings were infected with *Agrobacterium*. Once shoots and roots emerged, they were pricked with 0.5 × 25 mm sterile needles. Immediately, wounded seedlings were incubated in half-strength Murashige and Skoog's (MS) basal liquid medium (Murashige & Skoog, 1962) containing 100 μM acetosyringone and *Agrobacterium* containing *CspA* and *CspB* genes separately. Seedlings were agitated for 20 min at 100g and inoculated onto MS medium containing 100 μM acetosyringone, and incubated for 2–3 days at 28°C. For elimination of bacterial infection, seedlings were treated with 250 mg L⁻¹ cefotaxime. Seedlings growing on 50 mg L⁻¹ hygromycin containing medium were finally transferred into pots filled with garden soil. After one-month, putative transformants

were transplanted to new pots. Transgenic lines developed from *CspA* and *CspB* genes were labeled as A and B lines, respectively.

2.3 | Screening of transgenics for GUS assay and molecular analysis

Leaf material from 45-day-old putative transgenic plants was taken, rinsed twice with sterile distilled water, and incubated at 37°C in an incubator in presence of *GUS* staining buffer [2 mM 5-bromo-4-chloro-3-indolyl-β-D-glucuronic acid (X-Gluc)], 2 mM potassium ferricyanide and 2 mM potassium ferrocyanide in 0.1 M sodium phosphate buffer, pH 7.0. The samples were incubated for 16–20 h and destained by incubating in 90% ethanol. The procedure was repeated several times to visualize the blue color. Genomic DNA was isolated from *CspA*, *CspB* transgenics and WT plants by cetyltrimethyl ammonium bromide (CTAB) method (Dellaporta et al., 1983). All the in planta transformed plants along with WT were screened by PCR using gene-specific primers for the presence of transgenes (*CspA* and *CspB*) and *hptII* marker. Gene copy number analysis was performed by following the procedure laid down by Sambrook and Russell (2001). *CspA* and *CspB* cassettes were released using *HindIII* restriction enzyme and probed with *hptII* gene. Nitrocellulose membrane was developed by chemiluminescence method (Roche DIG DNA labeling kit).

2.4 | Bacterial growth study by pET expression

CspA and *CspB* genes were cloned into bacterial expression vector pET32a(+) using *EcoRI* and *HindIII* restriction enzymes and transformed into *E. coli* expression compatible cells BL21 (DE3). For functional analysis, recombinant *CspA* and *CspB* along with plain vector cells were grown in 200 ml LB medium with and without 200 mM NaCl and 300 mM mannitol. Corresponding controls were also maintained. The bacterial growth was measured by UV-Visible spectrophotometer (Thermo Scientific). Cells were induced with 1 mM IPTG at OD₆₀₀ nm at 30°C. Whole protein was isolated from 4-h-induced culture, using 6 M urea, 1% sodium dodecyl sulfate (SDS), and 0.1 M dithiothreitol (DTT), the cell lysate was heat denaturated (90°C) for 5 min with 1× SDS sample buffer. Samples were resolved in 12% SDS-PAGE gel with prestained protein ladder (Genei), blotted on nitrocellulose membrane, AntiHis tag primary antibody (Invitrogen 100 mg per 200 μl, 1:1000 dilution) was added. Membrane was washed twice with washing buffer, secondary antibody (Invitrogen, 1 mg) conjugated with horse radish peroxidase (HRP) (1 mg ml⁻¹, 1:10 000 dilution) was added and the blot developed by adding 3,3', 5,5'-tetramethylbenzidine (TMB) solution (Thermo Scientific).

2.5 | Segregation analysis of *hptII* in T₁ and T₂ generations

The T₁ seeds obtained from single copy selfed T₀ seeds were sterilized and transferred onto Petri dishes containing MS medium with

50 mg L⁻¹ hygromycin. After 10 days of growth, transgenics were screened for segregation analysis by studying hygromycin resistant and sensitive seedlings. Hemizygous seedlings were later transferred to the pots containing garden soil. Similarly, T₂ transgenic lines obtained from selfing of hemizygous lines were grown on MS medium with 50 mg L⁻¹ hygromycin along with WT plants and screened for their homozygous nature. Homozygous T₂ lines were used for subsequent analysis.

2.6 | Salt and drought stress treatments

Before floral initiation stage, homozygous A₁₅₋₁₋₁, A₅₉₋₁₋₁, A₆₂₋₁₋₃, B₂₋₁₋₄, B₁₆₋₁₋₂, and B₄₂₋₁₋₃ transgenic lines alongside WT plants growing in the pot conditions were used to assess salt and drought stress tolerance. Salt treatment was given by adding 250 ml of 150 mM NaCl solution to each pot on the first day. The drought stress was imposed by withholding water for 8 days. The corresponding controls were maintained for all the treatments. After 8 days of stress treatment, all plants were watered with normal tap water. The same treatment was used for all experiments unless otherwise mentioned. After stress treatment, transgenic lines and WT plants were evaluated for salt and drought stress respectively.

2.7 | Estimation of chlorophyll, proline, malondialdehyde (MDA), relative water content (RWC), and antioxidant enzyme activities

Salt and drought-treated transgenic lines and WT plants were used for the estimation of chlorophyll, proline, MDA content, and antioxidant enzyme activities. Chlorophyll *a* and *b* were estimated following the method of Arnon et al. (1974). Quantities of chlorophyll *a* and *b* were calculated and represented as mg g⁻¹ fresh weight. Proline was estimated following the method of Bates et al. (1973) and expressed as μmol proline g⁻¹ fresh weight of leaf tissue. Lipid peroxidation was determined by measuring the amount of MDA produced by thiobarbituric acid (TBA) reaction as described by Hodges et al. (1999). Lipid peroxidation is expressed in MDA μmol g⁻¹ fresh weight of leaf tissue. The RWC was measured by taking fresh weight, turgid weight, and dry weight of leaves (Van Heerden & de Villiers, 1996). Activity of catalase (CAT) enzyme was assessed by following the procedure of Aebi (1984). One unit of enzyme activity is defined as 1 μmol of H₂O₂ decomposed to water min⁻¹ mg⁻¹ of protein. The reaction of superoxide dismutase (SOD) was carried out by the method of Beauchamp and Fridovich (1971). SOD activity is defined as that amount of enzyme required to inhibit the reduction of nitrobluetetrazolium by 50% under the specified conditions. Activity is expressed as units mg⁻¹ protein min⁻¹. Guaiacol peroxidase (GPX) was estimated according to Eglely et al. (1983). Enzyme activity is expressed as μmol H₂O₂ reduced mg⁻¹ of protein min⁻¹. Protein was quantified by Bradford assay (Bradford, 1976) using bovine serum albumin as standard.

2.8 | Estimation of ions in transgenics and WT plants

Transgenic lines A₆₂₋₁₋₃ and B₄₂₋₁₋₃ (selected based on the performance under stress) and WT plants after salt and drought treatments were used for measuring ions. Na⁺ and K⁺ ions were quantified using atomic absorption spectrometer. Samples were dried in an oven at 104°C, powdered, digested with 8 ml of nitric acid and peroxide, filtered and diluted with 50 μl deionized water.

2.9 | Anatomy of WT and transgenic rice roots

One-month-old seedlings of rice WT, CspA (transgenic line A₆₂₋₁₋₃), and CspB (transgenic line B₄₂₋₁₋₃) were subjected to 150 mM NaCl and drought stress (withholding water) stresses for 48 h. Anatomical sections of the roots were taken with a sharp blade, stained with toluidine blue (5 mg ml⁻¹) for 15 min, destained with water and observed under brightfield microscope (Olympus model CH20iBIMF).

2.10 | Root morphology, shoot, and panicle lengths, seed number per panicle in transgenics and WT plants

Before seed setting, salt treatment was given by adding 250 ml of 150 mM NaCl solution to each pot on the first day. The drought stress was imposed by withholding water for 8 days. After 8 days of stress treatment, all plants were watered with normal tap water. The corresponding controls were maintained for all the treatments. Transgenic lines and WT plants were analyzed for root length, root dry mass, shoot, panicle lengths, and number of seeds per panicle.

2.11 | Gene expressions using quantitative real-time PCR (qRT-PCR) under multiple stress conditions

The relative quantification of gene expressions was studied in 1-month-old seedlings of T₂ transgenic lines A₆₂₋₁₋₃ and B₄₂₋₁₋₃. Seedlings were treated with salt (150 mM), drought (200 mM mannitol), cold (4°C), and high temperature (42°C) stresses. After 4 h of treatment, total RNA was isolated from root and leaf samples using guanidine thiocyanate method. Concentration of RNA was measured by Nano Drop and diluted to the final RNA concentration of 1 μg. It was converted to cDNA using Thermo Revert Aid cDNA Kit (Thermo Scientific). The cDNA was used as template for the gene expression studies (Applied Biosystem Fast PCR 7500) using SYBR green method. Actin gene served as an internal control. In 96 well plates, reaction volume was adjusted to final volume of 20 μl with Tm 60°C. The 2^{-ΔΔCt} values calculated are represented in the form of a graph. The relative expression levels of the genes *CspA*, *CspB*, *SGR*, *chlorophyllase*,

IPT1, *NCED*, *SOD*, and *SIRT1* were studied under diverse abiotic stress conditions in the transgenic lines and WT plants.

2.12 | Statistical analysis of data

All experiments were repeated with biological triplicates and also technical triplicates. For gene expression studies, data from biological triplicates along with technical replicates were collected. Average values along with standard deviations are represented in graphs/tables. All the data were subjected to one-way analysis of variance (ANOVA), since *CspA* and *CspB* lines were developed separately. *P*-value <0.5 was determined and the level of significance indicated in the graphs/tables.

3 | RESULTS

3.1 | *CspA* and *CspB* gene cloning and *in planta* transformation

In the present study, both *CspA* and *CspB* genes were cloned from the *E. coli* strain K12. The length of the *CspA* (MH029279) and *CspB*

(MH029280) gene sequences were 213 and 216 bp, respectively (Figure 1A). *In silico*, studies of domain analysis showed eukaryotic conserved cold shock domain (CSD) and ribonuclease-binding domain (RBD) in *CspA* (Figure S1) and *CspB* (Figure S2) indicating that they are involved in temperature stress. Presence of the gene cassettes were confirmed by restriction digestion (Figure 1B). *A. tumefaciens* LBA4404 pCAMBIA1301 *CspA* (Figure 1C) and pCAMBIA1301 *CspB* (Figure 1D) gene constructs driven by CaMV35S promoter and CaMV polyA terminator were used for *in planta* transformation. The *in planta* transformed seedlings were grown to maturity in pot conditions.

3.2 | Characterization of transgenics for transgene integration

All the T₀ generation, plants were tested for GUS histochemical activity alongside WT plants. GUS histochemical activity was noticed in transgenics, but not in WT plants (Figure 2). A total of six *CspA* and six *CspB* transgenic lines were found PCR positive. Thus, the transformation frequency was only 0.33%. Putative transgenic lines were named as A₁₂, A₁₅, A₄₅, A₅₀, A₅₉, A₆₂ and B₂, B₁₆, B₃₅, B₄₂, B₈₂, B₉₈ for *CspA* and *CspB* genes, respectively. PCR analysis showed an amplification




FIGURE 1 Gene amplification, vector construction, and recombinant vector map. (A) Amplification of bacterial gene 1 = *CspA* (213 bp), 2 = *CspB* (216 bp), M = 100 bp DNA marker, (B) recombinant construct harboring binary vector pCAMBIA1301 with *HindIII* restriction enzyme digestion, (C) recombinant vector map of pCAMBIA1301 containing hygromycin phosphotransferase (*hptII*), CaMV35S, cauliflower mosaic virus 35S (*CaMV35S*) promoter; *CspA* gene, CaMV poly A terminator, (D) recombinant vector map of pCAMBIA1301 containing *hptII*, CaMV35S, *CspB* gene, CaMV poly A terminator. M = 1Kb DNA marker, 1 = pCAMBIA 1301 + *CspA* construct (11 kb), 2 = pCAMBIA 1301 + *CspB* construct, CaMV35S promoter, *CspA* gene, *CspB* gene, CaMV poly A terminator

size of 213 bp for *CspA* (Figure 3A), and 216 bp for *CspB* (Figure 3B). Similarly, amplification of 750 bp for *hptII* gene was observed in both the transgenic lines (Figure 3C,D). Gene amplification by PCR was not detected in the WT plants (Figures 3A–D). Gene copy number by Southern blotting confirmed the integration of *CspA* and *CspB* genes in the transgenic lines (Figure 3E,F). Out of 12 transgenics, four lines from *CspA* (*A*₁₅, *A*₄₅, *A*₅₉, *A*₆₂) and four from *CspB* (*B*₂, *B*₁₆, *B*₄₂, *B*₉₈) were confirmed to have a single gene copy insertions (Figure 3E, F).




FIGURE 2 Histochemical GUS assay in *T*₀ transgenic plants. *CspA*, *CspA* putative transgenic plant; *CspB*, *CspB* putative transgenic plant; WT, wild type plant

On the other hand, in WT plants, no band was noticed, indicating that there was no transgene integration (Figure 3E,F). Subsequently, transgenics with single gene copy insertions were used for salt and drought stress tolerance experiments.

3.3 | Bacterial expression of *CspA* and *CspB* under NaCl and mannitol stresses

CspA and *CspB* genes were transferred into pET-32a(+) expression vector (Figure 4A). The presence of approximately 25.4 KD recombinant proteins along with the tags was noticed (Figure 4B), and protein expressions confirmed by blot with Anti-6× His Tag antibodies (Figure 4C). Once the expression of the recombinant proteins was confirmed, the growth of the recombinants was measured with and without salt and mannitol stresses. Growth of the untransformed cells decreased slightly in 200 mM NaCl and drastically under 300 mM mannitol (Figure 5A,B). Contrarily, *CspB* cells showed more resistance to mannitol than *CspA* (Figure 5B) but growth of *CspA* was better under salt stress treatment (Figure 5A).

3.4 | Mendelian inheritance pattern of *hptII* in *T*₁ and *T*₂ generations

*T*₁ transgenic line progenies segregated in a ratio of 3 tolerant:1 susceptible (Table S1). All *T*₂ progenies obtained from selfed hemizygous lines (*A*₁₅₋₁₋₁, *A*₅₉₋₁₋₁, *A*₆₂₋₁₋₃, *B*₂₋₁₋₄, *B*₁₆₋₁₋₂, and *B*₄₂₋₁₋₃) segregated in




FIGURE 3 Molecular characterization of transgenics. (A) Amplification of *CspA* gene in *T*₀ transgenics. (B) Amplification of *CspB* gene in *T*₀ transgenics. (C) *hptII* selection marker gene (750 bp) amplification from *T*₀*CspA* transgenics. (D) *hptII* selection marker gene (750 bp) amplification from *T*₀ *CspB* transgenics. (E) *CspA* gene copy number as analyzed by Southern blot (genomic DNA was digested with *HindIII* and probed by *hptII* marker). (F) *CspB* gene copy number as analyzed by Southern (genomic DNA was digested with *HindIII* and probed by *hptII* marker). *A*₁₂, *A*₁₅, *A*₄₅, *A*₅₀, *A*₅₉, *A*₆₂, *T*₀*CspA* transgenics; *B*₂, *B*₁₆, *B*₃₅, *B*₄₂, *B*₈₂, *B*₉₈, *CspB* transgenics; +C, positive control; M, 100 bp DNA marker; WT, wild type




FIGURE 4 Expression of *CspA* and *CspB* genes in pET. (A) *CspA* and *CspB*, pET-32a (+) vector was double digested with *EcoRI* and *HindIII* enzymes. (B) SDS-PAGE, 1 mM IPTG was used for induction of recombinant protein in *E. coli* BL21. (C) Western blot analysis of recombinant *CspA* and *CspB* protein expression with pET-32a(±) vector. *CspA*, *CspA* + pET-32a(±); *CspB*, *CspB*+ pET-32a(±); M, protein marker; UC, untransformed control

a 1:2:1 ratio (Table S2). Homozygous dominant lines displayed 100% germination on the selection medium. On the other hand, WT seeds failed to germinate at all on the hygromycin medium.

3.5 | Salt and drought stress tolerance in T₂ generation

Transgenic lines exhibited enhanced tolerance to salt and drought stresses in comparison with WT plants. Transgenic leaves remained relatively green in color with less yellowing, less leaf rolling, delayed senescence and higher tillering. Conversely, WT leaves were found sensitive to salt and drought stresses, exhibited wilting, increased leaf rolling, early senescence, and lower tillering (Figure 6). Transgenic lines overexpressing *CspA* and *CspB* exhibited SGR phenotype and delayed senescence compared to WT plants (Figure 6).

3.6 | Chlorophyll, proline, MDA contents, and antioxidative enzyme activities

No significant decrease in chlorophyll *a* content was observed in transgenic lines. However, chlorophyll *b* decreased slightly under the treatments. Content of chlorophyll *a* declined in WT plants under salt and drought stress conditions from 2.1 to 1.41 mg g⁻¹ and 2.1 to 1.45 mg g⁻¹, respectively. Similarly, chlorophyll *b* in WT declined from 0.75 to 0.46 mg g⁻¹ and 0.75 to 0.53 mg g⁻¹ under salt and drought stresses, respectively (Table 1). Proline content significantly increased in all transgenic lines compared to WT when treated with drought and salt stresses. Proline content increased to 2.1- and 2.7-folds in *CspA* and *CspB* transgenics treated with salt and drought stresses respectively (Figure 7A). Contrarily, levels of MDA were higher in WT than the transgenics under stress conditions. Similarly, under salt stress, MDA content was enhanced from 0.481 to 0.707 μmol g⁻¹ fresh weight in *CspA* and 0.482 to 0.690 μmol g⁻¹ fresh weight in *CspB*

transgenics (Figure 7B). Under drought stress, MDA content increased from 0.481 to 0.570 μmol g⁻¹ fresh weight in *CspA* and 0.482 to 0.536 μmol g⁻¹ fresh weight in *CspB* transgenic lines (Figure 7B). *CspA* and *CspB* transgenics exhibited significant increase in RWC compared to WT under salt and drought stresses (Figure 7C). After treatment with salt and drought, CAT specific activity was increased in both the transgenic lines and WT plants (Figure 7D). Under salt stress, specific CAT activity increased from 58.6 to 69 units in *CspA* and 65.6 to 66.3 units mg⁻¹ protein min⁻¹ in *CspB* transgenic lines (Figure 7D). Similarly, under drought stress, CAT activity leaped from 58.6 to 82 units mg⁻¹ protein min⁻¹ in *CspA* and 65.6 to 94.3 units mg⁻¹ protein min⁻¹ in *CspB* (Figure 7D). Under salt stress, the SOD activity was elevated from 10.85 to 26.6 units mg⁻¹ protein min⁻¹ in *CspA* and from 12.3 to 26.5 units mg⁻¹ protein min⁻¹ in *CspB* transgenics (Figure 7E). Under drought stress, SOD activity increased from 10.85 to 26.7 units mg⁻¹ protein min⁻¹ in *CspA* and 12.3 to 26.5 units mg⁻¹ protein min⁻¹ in *CspB* (Figure 7E). Specific activity of GPX increased in both the transgenic lines and WT plants, respectively (Figure 7F). Under salt stress, GPX activity increased from 297.3 to 312 units mg⁻¹ protein min⁻¹ in *CspA* and 298 to 343 units mg⁻¹ protein min⁻¹ in *CspB* transgenics (Figure 7F). Under drought stress, activity of GPX increased from 297.3 to 358 units in *CspA* and 298 to 374 units mg⁻¹ protein min⁻¹ in *CspB* (Figure 7F).

3.7 | Estimation of Na⁺ and K⁺ under stress conditions

No significant accumulation of Na⁺ and K⁺ was noticed in WT root and leaf tissues (Figure 8). On the contrary, significant accumulation of Na⁺ was recorded in transgenic lines under salt and drought stresses. Under salt stress, Na⁺ levels in root tissues increased 3.6- and 2.2-folds in *CspA* and *CspB* lines, respectively. Likewise, under drought stress, 2.4- and 1.7-folds higher Na⁺ levels were noticed in leaf tissues of *CspA* and *CspB* lines, respectively (Figure 8A). Similarly,




FIGURE 5 Recombinant BL21 strain with pET-32a(±) vector containing *CspA* and *CspB* genes and the growth of the strain with and without stress. (A) Effect of salt stress on bacterial growth, (B) effect of mannitol stress on bacterial growth. MNCspA, recombinant *CspA* strain grown in presence of 300 mM mannitol; MNCspB, recombinant strain *CspB* grown in presence of 300 mM mannitol; MNUC, UC grown in presence of 300 mM mannitol; STCspA, recombinant *CspA* strain grown in 200 mM NaCl; STCspB, recombinant *CspB* strain grown in 200 mM NaCl; STUC, UC grown in presence of 200 mM NaCl; WSCspA, *CspA* recombinant strain grown without stress; WSCspB, *CspB* recombinant strain grown without stress; WSUC, untransformed cells (UC) grown without stress (WS). OD was measured at 600_{nm} after growing for a duration of 16 h. Statistical significance is indicated as star (*) (P -value <0.05)

significant increase in Na^+ was noticed in transgenic root tissues after the imposition of salt and drought stresses (Figure 8B). Under stress conditions, enhanced accumulation of K^+ content was recorded in transgenics and WT leaf and root tissues compared to their corresponding controls. Accumulation of K^+ was higher in transgenics compared to WT (Figure 8C,D).

3.8 | Root, shoot, panicle lengths, root biomass, and number of seeds per panicle

After stress treatments, the root traits varied in transgenics and WT plants (Figure 9A). Devoid of stress, transgenic lines showed better

root length (32.5 cm) than WT (26.02 cm) plants (Figure 9A). Root lengths decreased under salt stress conditions (Figure 9B). The decrease in root lengths was more pronounced in salt-treated transgenics compared to roots grown under drought (Figure 9B). With the imposition of drought stress, root length decreased significantly in both the transgenic lines compared to WT (Figure 9B). Root dry mass decreased under salt and drought stresses in WT in comparison with transgenics (Figure 9C). After treatment with salt and drought stresses, significant decrease in shoot length was observed in WT and transgenic lines. The decrease in shoot length was higher under salt stress in comparison with drought stress-treated plants (Figure 10A). Devoid of stresses, the average length of the panicle in WT and transgenic lines was 14.21 ± 0.61 cm, and 18.54 ± 0.44 cm, respectively (Figure 10B). Panicle length decreased in WT plants (13.50 ± 0.83 cm and 10.03 ± 2.13 cm) as well as in transgenics under salt and drought stresses (15.39 ± 0.42 cm and 15.83 ± 0.67 cm), respectively (Figure 10B). Thus, the decrease in panicle length was more drastic under drought compared to salt stress. After stress imposition, a higher decrease in the number of seeds per panicle was noticed in WT compared to transgenics. A number of seeds per panicle in WT plants were reduced from 115.3 (devoid of stress) to 100.6 and 92.3 under salt and drought stresses, respectively. However, in *CspA* transgenic line, the reduction in seed number was less. In the line A_{15-1-1} , seed number decreased from 140 to 120.5 and 132.6, and in A_{59-1-1} , from 129.3 to 125.5 and 124.6 under salt and drought stresses, respectively. In the lines B_{2-1-4} and B_{42-1-3} , number of seeds decreased from 140.16 and 147.3 to 124.3, 107.1, 115.5 and 129.3 under salt and drought stress conditions, respectively (Table 2).

3.9 | Anatomy of WT and transgenic roots

One-month-old seedlings of WT, *CspA*, and *CspB* were exposed to 150 mM NaCl and also to drought stress (by withholding water) for 48 h along with appropriate controls. Anatomical sections of the WT roots revealed thin-walled sclerenchyma cells and loosely organized aerenchyma (Figure S3). In contrast, transgenic plants (*CspA* and *CspB* lines) were characterized by thick-walled sclerenchyma cells. Further, compactly arranged cortical cells with narrow intercellular spaces were the characteristic feature of the transgenic roots (Figure S3).

3.10 | *CspA* and *CspB* transcript levels in T_2 transgenics and expression of SGR associated genes under salt, drought, cold, and heat stresses

Transcript levels were higher in *CspA* (RTA) leaves and roots than in *CspB* (RTB) leaves and roots. Induction of the *chlorophyllase* gene in both *CspA* and *CspB* was markedly higher under the four abiotic stress conditions, than without stress. All the four stresses influenced the *IPT1* gene, more so under cold and drought compared to salt and high temperature stresses (Figure 11). *NCED* was




FIGURE 6 Evaluation of T_2 transgenic lines subjected to salt and drought stresses. A_{15-1-1} , A_{59-1-1} , A_{62-1-3} , T_0 *CspA* transgenic lines; B_{2-1-4} , B_{16-1-2} , B_{42-1-3} , T_0 *CspB* transgenic lines; drought, drought stress imposed by water withholding; salt, salt stress imposed by 150 mM NaCl; WS, without stress; WT, wild type

TABLE 1 Chlorophyll *a* and *b* contents in WT and transgenic lines grown under salt and drought stress conditions

Stress	WT	A_{15-1-1}	A_{59-1-1}	A_{62-1-3}	B_{2-1-4}	B_{16-1-2}	B_{42-1-3}
Chlorophyll <i>a</i>							
WS	2.10 (± 0.001)	2.11 (± 0.001)	2.15 (± 0.001)	2.14 (± 0.005)	2.03 (± 0.002)	2.16 (± 0.001)	2.15 (± 0.001)
Salt	1.41 (± 0.47)	1.88 (± 0.093) ^a	1.84 (± 0.10) ^a	1.88 (± 0.05) ^a	1.95 (± 0.22)	1.93 (± 0.13)	2.01 (± 0.04) ^a
Drought	1.45 (± 0.54)	1.94 (± 0.12)	1.93 (± 0.11)	1.89 (± 0.12)	1.94 (± 0.13)	1.94 (± 0.12)	2.02 (± 0.05)
Chlorophyll <i>b</i>							
WS	0.75 (± 0.001)	0.88 (± 0.001)	0.80 (± 0.002)	0.80 (± 0.008)	0.75 (± 0.006)	0.79 (± 0.004)	0.81 (± 0.01)
Salt	0.46 (± 0.47)	0.58 (± 0.09) ^a	0.501 (± 0.10)	0.505 (± 0.05)	0.75 (± 0.17)	0.775 (± 0.09) ^a	0.687 (± 0.032)
Drought	0.53 (± 0.54)	0.71 (± 0.12)	0.60 (± 0.11) ^a	0.64 (± 0.12)	0.88 (± 0.10)	0.89 (± 0.06)	0.75 (± 0.07)

Note: Data represent means \pm SD ($n = 9$; three biological replicates).

Abbreviations: A_{15-1-1} , A_{59-1-1} , A_{62-1-3} , B_{2-1-4} , B_{16-1-2} , B_{42-1-3} , transgenic lines; drought, drought by withholding water; salt, 150 mM NaCl; WS, without stress, WT, wild-type plants.

^aIndicates significant at $P < 0.05$ as analyzed by one-way ANOVA.

also induced by the abiotic stresses and the expression was higher under salt, cold, and high temperatures than under drought stress. Both *CspA* and *CspB* were strongly upregulated by cold (but not in *CspB* roots), followed by salt and drought. Expressions were low under high temperature stress, especially in *CspB* roots. *SGR* gene expression was enhanced steeply by drought and cold in comparison with salt and high temperature stresses (Figure 11). *SOD* expression was prompted by salt and high temperature stresses in comparison with cold and drought. Comparatively, cold and drought stresses promoted *SIRT1* gene expressions more than salt and high temperatures (Figure 11).

4 | DISCUSSION

4.1 | Gene cloning, validation of *CspA* and *CspB* genes under salt and drought (mannitol) stresses

Csps have been reported to contain a CSD, which is generally composed of 65 to 70 amino acid residues in bacteria, as well as in higher organisms including plants (Horn et al., 2007). *Csp* proteins are 7 to 10 kDa in size and contain the nucleic acid-binding activities. Jiang et al. (1997) pointed out that in *E. coli*, CSD acts as an RNA chaperone and converts double-stranded RNA into single-stranded RNA. Several




FIGURE 7 Estimation of proline, malondialdehyde (MDA), relative water content (RWC), activities of catalase (CAT), superoxide dismutase (SOD), and guaiacol peroxidase (GPX) in transgenic rice exposed to salt stress and drought conditions (A) Proline, (B) MDA, (C) RWC, (D) CAT (units mg⁻¹ protein min⁻¹), (E) SOD (units mg⁻¹ protein min⁻¹), (F) GPX (units mg⁻¹ protein min⁻¹). Salt and drought stresses were imposed by adding 150 mM NaCl water and withholding respectively. A₁₅₋₁₋₁, A₅₉₋₁₋₁, A₆₂₋₁₋₃, B₂₋₁₋₄, B₁₆₋₁₋₂, B₄₂₋₁₋₃, transgenic lines; FW, leaf fresh weight; WS, without stress; WT, wild type. Statistical significance is indicated as star (*) (P -value < 0.05)

of the Csp proteins may stimulate growth during stress acclimation. Hunger et al. (2006) found out that Csps work in concert with a DEAD box helicase to rescue misfolded mRNA and help in transcription (El-Sharoud & Graumann, 2007). Bae et al. (2000) showed that CspA, CspC, and CspE genes act as antiterminators and regulate the expression of cold-inducible genes. In the present study, bacterial CspA and CspB genes were cloned which shared 100% homology with the *E. coli* sequences and displayed both CSD as well as RBD. Plants also have CSD proteins, which differ from that of Csps known to occur in prokaryotes (Sasaki & Imai, 2012). Several of the bacterial Csps and plant CSDs were found to be induced under cold stress (Jung et al., 2010; Sasaki et al., 2007). Interestingly, though *E. coli* Csps are responsive to cold stress and function as RNA chaperones (Graumann & Marahiel, 1998), they share a domain with AtCSP3, which plays a pivotal role in low temperature tolerance (Kim et al., 2009). In the present study, both CspA and CspB genes were validated for the growth profile of *E. coli* under mannitol (drought) and salt

stress conditions. Growth of untransformed *E. coli* cells decreased drastically in presence of mannitol unlike that of CspA and CspB containing bacterial cells indicating that these genes are associated with mannitol/drought stress besides cold stress tolerance. *E. coli* containing both CspA and CspB displayed almost similar responses in terms of growth. Genetically altered *E. coli* also displayed a similar growth pattern under salt stress indicating that both CspA and CspB protect bacteria against salt stress. Western blot results confirmed the presence of translational product in transformed *E. coli* inferring functional expression of the genes. Nakaminami et al. (2006) determined the importance of C-terminal region of a plant cold shock domain protein (CSDP) and showed that deletion of all C-terminal zinc fingers in wheat cold shock protein 1 (WCSP1) abolished the growth stimulatory activity in *E. coli* during cold stress indicating that the CCHC-type zinc fingers in CSDPs are highly vital for growth. These experiments were not performed in the present study, but such a possibility cannot be ruled out.




FIGURE 8 Ion analysis of leaf and root tissues under salt and drought stresses. (A) Na⁺ content in leaf mg g⁻¹ DW, (B) Na⁺ content in root mg g⁻¹ DW, (C) K⁺ content in leaf mg g⁻¹ DW, (D) K⁺ content in root mg g⁻¹ DW. A₄₂₋₁₋₃ and B₆₂₋₁₋₃, transgenic lines; DW, dry weight; WT, wild type. Statistical significance is indicated as star (*) (P -value < 0.05)

4.2 | Generation and characterization of transgenics

Characterization of both *CspA* and *CspB* genes by PCR amplifications and Southern blot analysis indicated that these genes were incorporated stably into the host plant. qRT-PCR analysis showed that both *CspA* and *CspB* genes are expressed in leaf, root, and internodal tissues at the transcriptional level under drought stress. Fang et al. (1997) found that *E. coli CspA* does not increase the transcription but increases the stability of mRNA during cold stress. Protein folding becomes highly inefficient and the function of ribosomes is impeded under stress conditions (Keto-Timonen et al., 2016). However, the Csp family of proteins counteracts and prevents the formation of secondary structures in mRNA and help in the initiation of translation.

4.3 | Chlorophyll, proline, MDA, ion analysis, and antioxidative enzyme activities under stresses

Chlorophyll (both *a* and *b*) levels did not decline much under stress conditions in *CspA* and *CspB* lines in comparison with WT plants. A higher chlorophyll content in wheat transgenics (in comparison with WT plants) containing *SeCspA* was reported by Yu et al. (2017). These results indicate that *CspA* and *CspB* genes prevent faster degradation of chlorophyll or slow down the process of senescence under stress by an unknown mechanism. Both the transgenic lines exhibited a SGR phenotype in rice, but have not been reported earlier in maize and wheat (Castiglioni et al., 2008; Yu et al., 2017). The higher chlorophyll content in the transgenics (*CspA* and *CspB*) supports the SGR

phenotype that generally maintains higher chlorophyll content than the non-SGR genotype. SGR is an important agronomic trait that permits plants to maintain active photosynthesis and subsequently improve the grain-filling even under adverse conditions (Borrell et al., 2014; Jaeggli et al., 2017). Proline levels were higher in both the transgenics in comparison with WT plants under stress. Accumulation of compatible osmolytes such as proline under stress conditions has been registered in many transgenics (Anjaneyulu et al., 2014; Reddy et al., 2015). Proline helps in the conversion of O₂⁻ to H₂O₂ and O₂ in the chloroplasts, mitochondria, and peroxisomes (Fridovich, 1989; Yiu & Tseng, 2005). Further, it has been noticed that proline protects plants against the oxidative damage caused by abiotic stress conditions (Molinari et al., 2007). Studies on transcriptomic analysis of gene expressions between SGR and senescing lines of sorghum revealed enrichment of genes associated with the “response to osmotic stress” (Jhonson et al., 2015). They noticed high expression of the delta-pyrroline-5-carboxylate synthase 2 (*P5CS2*) gene (involved in proline biosynthesis) in SGR compared to senescent line. Further, the expression of *P5CS2* was correlated with high levels of proline accumulation. Surprisingly, *P5CS2* has been found to lie within the *Stg1* (stay-green) quantitative trait loci. Also, polymorphisms have been identified in known *cis*-elements in the promoter regions of *P5CS2* (Jhonson et al., 2015). Thus, these elements could be responsible perhaps for the differences in the expression of *P5CS2* gene between SGR and senescent lines. Such a finding has a bearing in our understanding of drought tolerance mechanism in crop plants. Lipid peroxidation levels (MDA) were reduced in *CspA* and *CspB* transgenics compared to WT plants, an indication of the consequence of *CspA* and *CspB* genes in transgenics (Semchuk et al., 2012). Significantly higher ionic levels




FIGURE 9 Root phenotype, root length, and root biomass in transgenics and WT. (A) Root morphology, (B) root length (cm), (C) root dry mass (g) in transgenic lines under salt and drought stresses. A₁₅₋₁₋₁, A₅₉₋₁₋₁, A₆₂₋₁₋₃, *CspA* transgenic lines; B₂₋₁₋₄, B₁₆₋₁₋₂, B₄₂₋₁₋₃, *CspB* transgenic lines; drought, withholding water; salt, 150 mM NaCl; WS, without stress; WT, wild type. Statistical significance is indicated as star (*) (P -value < 0.05)

(Na⁺ and K⁺) were observed in the transgenics in comparison with WT plants upon exposure to water-deficit and salt stress conditions. Na⁺ inhibits the influx of K⁺ into cytosol under high salt stress conditions leading to the accumulation high levels of Na⁺ inside the cells (Hanin et al., 2016). Higher Na⁺ levels inside the cytoplasm is generally accompanied by loss of K⁺. So, enhanced uptake of K⁺ under NaCl stress is difficult because of competition from Na⁺ for K⁺-binding sites on transport systems (Chen et al., 2007). But, being a macronutrient, K⁺ plays a crucial role in protein synthesis and phloem sugar loading in

plants (De Schepper et al., 2013). In line with this, Chen et al. (2007) demonstrated that cytosolic K⁺ to Na⁺ ratio is an important determinant of plant salinity tolerance. So, preventing the loss of K⁺ and acquisition of K⁺ under salt stress is crucial for maintaining a proper cytosolic K⁺ concentration (Assaha et al., 2017; Chen et al., 2007). In transgenic *CspA* and *CspB*, accumulation of K⁺ was higher in comparison with Na⁺, which is vital for stress tolerance. Though uptake of K⁺ is through activation of HAK transporters, how *Csps* help to maintain K⁺ acquisition is obscure at the moment. Overall, the specific activities of CAT, SOD, and GPX were enhanced in transgenics compared to WT plants, perhaps an outcome of the transgenes. SOD catalyzes superoxide radicals into H₂O₂, which can be converted subsequently by CAT and GPX into water and O₂. A large body of information exists about the involvement and effective purging of superoxide radical and in stress protection in transgenics (Reddy et al., 2015; Tseng et al., 2007; Yiu & Tseng, 2005). Both CAT and GPX help in the conversion of H₂O₂ and therefore prevent oxidative damage (Esfandiari & Shekari, 2007; Willekens et al., 1995). Increased CAT activity assists in cell membrane stability by decreasing H₂O₂ levels under stress. Increased levels of CAT were also recorded under abiotic stress conditions in maize and *Sesamum* (Koca et al., 2007; Neto et al., 2006). Enhanced GPX was noticed in transgenic finger millet and sorghum under salt stress (Anjaneyulu et al., 2014, Reddy et al., 2015). Antioxidative enzyme activities are vital since they can modulate redox homeostasis of cells (Bhavanath et al., 2011; Huang et al., 2013). Thus, a clear correlation between increased proline (implicated in SGR phenotype), decreased lipid peroxidation and higher activities of antioxidative enzyme activities in *CspA* and *CspB* rice transgenics support that bacterial chaperones help to protect the activities and prevail over salt and drought stress-caused damages.

4.4 | Anatomy of WT differs from transgenic rice plants

Root morphology in transgenics differed from that of the WT plants, which led us to investigate the root anatomy. Roots in WT plants when exposed to abiotic stresses, displayed thin-walled sclerenchymatous cells and loosely organized aerenchyma, an indication of poor defense against stress. This may allow transport of ions such as Na⁺ and Cl⁻ under salt stress or dehydration to occur under drought. Transport of ions into cortex and phloem tissues might damage the cells in all root zones before initiating the process of any structural defense. Contrary to this, transgenic roots (*CspA* and *CspB* lines) under stress conditions exhibited thick-walled sclerenchyma cells indicating a strong defense against salinity and water-limited conditions. Compactly arranged cortical cells with narrow intercellular spaces may reduce the transport of ions into inner phloem cells. Overall, it appears that cell wall integrity features (unlike that of WT) in transgenic roots help the cell viability, which then trigger the biosynthesis and accumulation of polyphenolic substances or deposition of lignin in the epidermal, hypodermal and intercellular spaces of outer cortical cells (Zagorchev et al., 2014, Gall et al., 2015, Kishor et al., 2015). Mourasobczak et al. (2011) and Chun et al. (2019) have provided




FIGURE 10 Analysis of shoot length and panicle length. (A) Shoot length (cm), (B) panicle length (cm) in transgenic lines under salt and drought stresses. Salt and drought stresses were imposed by 150 mM NaCl and water withholding water before booting stage. *A*₁₅₋₁₋₁, *A*₅₉₋₁₋₁, *A*₆₂₋₁₋₃, *CspA* transgenic lines; *B*₂₋₁₋₄, *B*₁₆₋₁₋₂, *B*₄₂₋₁₋₃, *CspB* transgenic lines; WS, without stress; WT, wild type. Statistical significance is indicated as star (*) (*P*-value < 0.05)

TABLE 2 Number of seeds/panicle in WT and transgenics under salt and drought stress conditions


Stress	WT	<i>A</i> ₁₅₋₁₋₁	<i>A</i> ₅₉₋₁₋₁	<i>A</i> ₆₂₋₁₋₃	<i>B</i> ₂₋₁₋₄	<i>B</i> ₁₆₋₁₋₂	<i>B</i> ₄₂₋₁₋₃
WS	115.3 (±4.92)	140 (±1.63)	129.3 (±0.68)	141 (±0.81)	140.16 (±0.95)	137.8 (±2.21)	147.3 (±2.72)
Salt	100.6 (±6.59)	120.5 (±2.85) ^a	125.5 (±0.4) ^a	125.6 (±4.64) ^a	124.3 (±0.27) ^a	116.3 (±1.24) ^a	115.5 (±4.49) ^a
Drought	92.3 (±9.39)	132.6 (±0.54) ^a	124.6 (±1.8) ^a	121.3 (±0.54) ^a	107.1 (±8.47) ^a	121.6 (±2.68) ^a	129.3 (±0.54) ^a

Note: Data represent means ± SD (*n* = 9 for each line).

Abbreviations: *A*₁₅₋₁₋₁, *A*₅₉₋₁₋₁, *A*₆₂₋₁₋₃, *B*₂₋₁₋₄, *B*₁₆₋₁₋₂, *B*₄₂₋₁₋₃, transgenic lines; drought, drought by withholding water; salt, 150 mM NaCl; WS, without stress; WT, wild-type plants.

^aIndicates significant at *P* < 0.05.

FIGURE 11 Transcript analysis of transgenes and stay-green (SGR) associated gene expression of transgenics against different abiotic stresses using qRT-PCR. Relative gene expressions were measured under abiotic stress (48 h) conditions in transgenic rice seedlings (*CspA*₆₂₋₁₋₃) (*CspB*₄₂₋₁₋₃) under salt (150 mM NaCl), drought (200 mM mannitol), cold (4°C), and high temperature (42°C) stresses. Expression (qRT-PCR) of *chlorophyllase*, *isopentenyltransferase 1 (IPT1)*, *9-cis-epoxycarotenoid dioxygenase (NCED)*, bacterial *CspA* transgene (*RTA*), bacterial *CspB* transgene (*RTB*), *SGR*, mitochondrial *SOD* and *SIRT* genes. Actin gene was used as an internal control



molecular and genetic evidence indicating the importance of enhanced lignin accumulation in the plant cell wall during the responses to salt stress. Also, expression of genes of the lignin biosynthetic pathway was

positively correlated with drought stress tolerance (Hu et al., 2009). Associated with this phenomenon, increased lignin biosynthesis has been noticed under abiotic stress conditions (Mourasobczak et al., 2011),

which can reduce penetration and transpiration of cell wall water leading to osmotic balance during abiotic stress conditions (Monties & Fukushima, 2001). Lignins are the cross-linked phenolic polymers, and the enzyme peroxidase enhances the content of phenolics (Kim et al., 2008). In the present study, an increase in peroxidase activity (glutathione peroxidase) has been found in transgenics, which may lead to lignin deposition in the cell walls. Anatomical sections of roots in deed show increased lignin deposition. In line with this, double mutants (*atprx2/atprx25*, *atprx2/atprx71* and *atprx25/atprx71*) with reduced peroxidase activity have been shown to contain lower levels of lignin than single mutants (Shigeto et al., 2015). Taken together, it appears that lignin deposition in cell walls is critical under salt stress conditions.

4.5 | *CspA* and *CspB* overexpression increased root length, shoot length, panicle length, and seed number per panicle

Roots in transgenic plants were longer compared to WT plants indicating that bacterial chaperones by some mechanism improve root length. Such a morphological variation in root length was noticed earlier in tobacco and sorghum transgenics (Kishor et al., 1995; Reddy et al., 2015). Average lengths of transgenic (*CspA* and *CspB*) rice shoots were found to be slightly higher than in WT plants. Since *Csps* are involved in many cellular processes to promote normal growth, protect RNA structure and stress adaptation responses (Keto-Timonen et al., 2016), it is possible that they might play a role on the increased lengths of both roots and shoots in the transgenic plants. WT plants matured 20 days earlier than the transgenics which displayed SGR phenotype indicating that *CspA* and *CspB* proteins delay chlorophyll degradation or slow down the process of senescence, though the mechanism is unclear. Higher chlorophyll content in transgenics is an indication for drawing such an inference. It has been suggested by Sasaki and Imai (2012) that CSDPs regulate embryo development, flowering time and fruit development indicating their diverse roles. On the other hand, *SeCspA* and *SeCspB* did not improve cold stress in transgenic wheat, but synthetic *CspA* gene improved drought stress under the field conditions. In the present study, transgenic rice exhibited better drought and salt (150 mM NaCl) stress tolerance under pot conditions compared to the WT plants. In contrast, Sasaki et al. (2015) showed that *Arabidopsis AtCSDP2* negatively regulates freezing tolerance. Further, they demonstrated that overexpression of *AtCSP2* resulted in reduced salt stress tolerance in *Arabidopsis*, indicating that it is a negative regulator of salt stress. Park et al. (2009) demonstrated that CSDPs affect seed germination and growth of *Arabidopsis* plants under abiotic stress, thus inferring the implication of CSDPs during seed germination. *AtCSP3* overexpression in *Arabidopsis* resulted in improved salt and drought stress tolerance by upregulating the expression of stress-related proteins (Kim et al., 2013). Yu et al. (2017) showed that overexpression of *CspA* and *CspB* genes caused the upregulation of *TaCDPK3* transcription factor in wheat. It is known that CDPKs play crucial roles in stress signal transduction and regulate the downstream genes that can be activated in

turn by ABA (Sanders et al., 2002). Bacterial chaperones (*CspA* and *CspB*) upon expression conferred abiotic stress tolerance in maize in the field conditions (Castiglioni et al., 2008). These experiments indicate that bacterial chaperones have the potential to combat abiotic stresses in higher plants and hence need to be exploited further, especially a combination of *Csp* genes (*CspA* through *CspI*) for obtaining superior transgenics with higher levels of abiotic stress tolerance. Melencion et al. (2017) showed that an RNA chaperone functioned as a universal stress protein in *Arabidopsis* and displayed increased cold stress tolerance. *E. coli CspA* and *CspB* synthetic genes also enhanced the cold tolerance when overexpressed in *Arabidopsis thaliana* (Yu et al., 2017), suggesting that synthetic genes display identical functions in *Arabidopsis* and impart cold stress tolerance (Yu et al., 2017). The above experiments prove that bacterial *Csps* play crucial roles under abiotic stresses in plants also. But, so far, the SGR phenotype has not been reported upon the expression of bacterial *Csp* genes in crop plants. Transgenic rice plants expressing *CspA* and *CspB* genes revealed enhanced drought tolerance as seen by better panicle lengths. Both salt and drought stresses reduced the number of seeds per panicle in transgenics in comparison with the plants not exposed to stress conditions. The yield loss under salt and drought stresses ranged from 13% to 18% when compared with non-stressed conditions in WT plants. In contrast, only 3% loss was recorded in the line *A59-1.1* irrespective of salt and drought stress imposition. In the transgenic line *B2-1.4*, 10% and 25% loss in yield were noticed in salt and drought stress conditions, respectively, in comparison with unstressed environments. Thus, yield penalty was noticed in transgenics in comparison with stressed conditions, but not devoid of stress. Yield was higher in transgenics when compared to WT plants under both salt and water-limited conditions. Similarly, transgenic maize displayed yield benefits of up to 15% under drought stress in the field, with no yield penalty even under non-stressed conditions (Castiglioni et al., 2008). These data suggest that effective RNA chaperone activity is indispensable to plants growing under water deficit conditions since they improve panicle length and seed weight as pointed out by Castiglioni et al. (2008). They noticed that a functional RNA-binding site is critical to confer improved yields under drought stress in maize.

4.6 | Enhanced *CspA*, *CspB*, and SGR-associated transcript levels in T_2 transgenics

Upregulation of both *CspA* and *CspB* genes was noticed under normal as well as abiotic stress conditions. Breakdown of chlorophyll is common during leaf senescence, seed maturation and fruit ripening. Consistent with this observation, expression of the gene *chlorophyllase* was high under all the four abiotic stresses. Chlorophyll is broken down to products via the pheophorbide A oxygenase pathway (Hortensteiner, 2013). However, in the present study, breakdown of chlorophyll during seed maturation stage is delayed and *CspA* and *CspB* transgenics displayed SGR phenotype consistently. This is quite surprising, unexpected and has not been reported earlier in *Csp* expressed crop plants like maize and wheat (Castiglioni et al., 2008;

Yu et al., 2017). It has been suggested that the SGR gene discovery is an association between stress-induced leaf senescence and stability of chlorophyll metabolism (Sakuraba et al., 2014; Thomas & Ougham, 2014). Further, the SGR phenotype is a result of alterations in chlorophyll metabolism. It could be either due to delayed degradation or over-production of chlorophyll under stress conditions (Hortensteiner & Kräutler, 2011). SGRs can play a role in stress tolerance as pointed out by Jagadish et al. (2015). It has been reported that some Csp genes are inducible by abiotic stresses other than cold (Keto-Timonen et al., 2016). Cytokinin is an important phytohormone and is associated with alterations of source/sink relationships (Leopold & Kawase, 1964), counteraction of high temperature stress (Caers et al., 1985), and also stimulation of chlorophyll synthesis (Arnold & Fletcher, 1986). In line with this, expression of the *IPT1* gene (associated with cytokinin biosynthesis) was found higher in transgenics especially under cold and drought stresses. *IPT1* may stimulate chlorophyll biosynthesis and thus help to maintain the SGR phenotype. Expression of *NCED* was also higher in transgenics under different abiotic stresses when compared to non-stress conditions. This is a crucial gene associated with ABA biosynthesis. Its expression was induced by NaCl, cold, high temperature as well as drought, suggesting the critical role that it plays in response to the multiple abiotic stress tolerance in rice. Earlier, Huang et al. (2018) also noticed induction of *NCED* by NaCl, PEG, and H₂O₂ in rice. *PYL9*, a member of the ABA receptors, increases leaf senescence, but promotes drought tolerance in *Arabidopsis* (Zhao et al., 2016). This is in contrast to the other studies, which show that delayed leaf senescence or the SGR type increases tolerance to drought stress by regulating the *IPT* gene (Rivero et al., 2007). *PYL9* is an ABA-dependent pathway and hence plants get adapted to drought. On the other hand, *IPT* involves ABA-independent signal transduction. *IPT* overexpressed plants increase cytokinin synthesis, become greener, and thus adapt to drought stress (Clark et al., 2004; Merewitz et al., 2010; Rivero et al., 2007). Phytohormones like jasmonic acid, salicylic acid, and ABA promote senescence (Liang et al., 2014; Qi et al., 2015). Contrarily, cytokinin has an antipodal function and is known to delay leaf senescence. *IPT* overexpression in *Agrostis stolonifera* alleviated drought stress-induced inhibition of root growth by activating ROS-scavenging systems (Xu et al., 2016). In the present study, cold and drought stresses induced higher expression of the plant *SIRT1* gene (a human homolog) than salt and high temperature, indicating its association with these stresses. In plants, *SIRT* genes have been implicated against genome instability and cell oxidative damage (Huang et al., 2007). *SIRT*s are also associated with leaf senescence, regulation of photosynthetic activity, transcription, metabolism, and DNA damage repair (Cucurachi et al. 2012, Lagunas-Rangel, 2019, Zheng, 2020). Thus, several of these genes might be associated with the SGR phenotype to protect the plants against abiotic stress conditions. Also, Csp genes have been shown to contribute to osmotic and oxidative stress tolerance as pointed out by Keto-Timonen et al. (2016). Our results reveal that transgenic rice plants containing both *CspA* and *CspB* genes display stronger abiotic stress tolerance and improved yields compared to WT plants.

5 | CONCLUSIONS

We demonstrate here for the first time that overexpression of bacterial chaperone genes (*CspA* and *CspB*) in rice results in a SGR phenotype, late maturity, and improved yields under salt and drought stress conditions in comparison with that of WT plants. Higher expression of *IPT1*, *NCED*, *SGR*, and *SOD* genes under diverse abiotic stress conditions in transgenics illustrates that these genes play a pivotal role in imparting SGR character in rice. Thus, there is an urgent need to explore diverse bacterial Csp genes for generating abiotic stress-tolerant crop plants.

ACKNOWLEDGMENTS

GR is grateful to the DST-SERB for providing fellowship. PBK is thankful to Vignan's Foundation for Science, Technology and Research, Vadlamudi, Guntur, for providing emeritus fellowship. This research is supported by a grant from the Department of Science and Technology (DST)-SERB, New Delhi, through the grant number SERB/SB/SO/PS/104/2013.

AUTHOR CONTRIBUTIONS

P. B. K. Kishor conceived and designed the experiments and analyzed the data. Guddimalli Rajasheker cloned the genes, carried out the experiments and analyzed the data. Guddimalli Rajasheker, Somanaboina A. Kumar, Palle S. Reddy, Divya Kummari, Sudhakar R. Palakolanu, Gandra Jawahar, Rathnagiri Polavarapu, and P. B. K. Kishor analyzed and interpreted the data and wrote the manuscript. Dr. Insaf A. Qureshi helped in carrying out the western blot. Guddimalli Rajasheker, Palle S. Reddy, Somanaboina A. Kumar, Divya Kummari, Sujatha Edupuganti, Nagaraju Marka, Sudhakar R. Palakolanu, Gandra Jawahar, Rathnagiri Polavarapu, Jalaja Naravula, Insaf A. Qureshi, and P. B. K. Kishor critically analyzed the manuscript. All authors read and approved the manuscript.

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the supplementary material of this article

ORCID

Rathnagiri Polavarapu  <https://orcid.org/0000-0002-9587-7547>

REFERENCES

- Acosta-Motos, J.R., Ortuno, M.F., Bernal-Vicente, A., Diaz-Vivancos, P., Sanchez-Blanco, M.J. & Hernandez, J.A. (2017) Plant responses to salt stress: adaptive mechanisms. *Agronomy*, 7, 18. <https://doi.org/10.3390/agronomy7010018>.
- Aebi, H. (1984) Catalase in vitro. In: Packer, L. (Ed.) *Methods in enzymology*. Orlando: Academic Press, pp. 121–126.

- Amara, I., Capellades, M., Ludevid, M.D., Pagès, M. & Goday, A. (2013) Enhanced water stress tolerance of transgenic maize plants overexpressing LEA Rab28 gene. *Journal of Plant Physiology*, 170, 864–873.
- Anjaneyulu, E., Reddy, P.S., Sunita, M.S., Kavi Kishor, P.B. & Meriga, B. (2014) Salt tolerance and activity of antioxidative enzymes of transgenic finger millet overexpressing a vacuolar H⁺-pyrophosphatase gene (*SbVPPase*) from *Sorghum bicolor*. *Journal of Plant Physiology*, 171, 789–798.
- Arnold, V. & Fletcher, R.A. (1986) Stimulation of chlorophyll synthesis by benzyladenine and potassium in excised and intact cucumber cotyledons. *Physiologia Plantarum*, 68, 169–174.
- Arnon, D.I., McSwain, B.D., Tsujimoto, H.Y. & Wada, K. (1974) Photochemical activity and components of membrane preparation from blue-green algae. I. Coexistence of two photosystems in relation to chlorophyll a and removal of phycocyanin. *Biochimica et Biophys Acta-Bioenergetics*, 357, 231–245.
- Assaha, D.V.M., Ueda, A., Saneoka, H., Al-Yahyai, R. & Yaish, M.W. (2017) The role of Na⁺ and K⁺ transporters in salt stress adaptation in glycophytes. *Frontiers in Physiology*, 8, 509.
- Bae, W., Xia, B., Inouye, M. & Severinov, K. (2000) *Escherichia coli* CspA-family RNA chaperones are transcription antiterminators. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 7784–7789.
- Bates, L.S., Waldren, R.P. & Teare, I.D. (1973) Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39, 205–208.
- Beauchamp, C. & Fridovich, I. (1971) Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Analytical Biochemistry*, 44, 276–287.
- Bertani, G. (1951) Studies on lysogenesis. I. the mode of phage liberation by lysogenic *Escherichia coli*. *Journal of Bacteriology*, 62, 293–300.
- Bhavanath, J., Anubha, S. & Avinash, M. (2011) Expression of *SbGSTU* (tau class glutathione S-transferase) gene isolated from *Salicornia brachiata* in tobacco for salt tolerance. *Molecular Biology Reports*, 38, 4823–4832.
- Borrell, A.K., Mullet, J.E., Jaeggli, B.G., van Oosterom, E.J., Hammer, G.L., Klein, P.E., et al. (2014) Drought adaptation of stay-green cereals associated with canopy development, leaf anatomy, root growth and water uptake. *Journal of Experimental Botany*, 65, 6251–6263.
- Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72, 248–254.
- Bray, E.A., Bailey-Serres, J. & Weretilnyk, E. (2020) Responses to abiotic stresses. In: Buchanan, B., Gruissem, W. & Jones, R. (Eds.) *Biochemistry and molecular biology of plants*, 2nd edition. Rockville, MD: American Society of Plant Physiologists, pp. 1051–1100.
- Caers, M., Rudelsheim, P., Van Onckelen, H. & Horemans, S. (1985) Effect of heat stress on photosynthetic activity and chloroplast ultrastructure in correlation with endogenous cytokinin concentration in maize seedlings. *Plant and Cell Physiology*, 26, 47–52.
- Capell, T., Bassie, L. & Christou, P. (2004) Modulation of the polyamine biosynthetic pathway in transgenic rice confers tolerance to drought stress. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 9909–9914.
- Castiglioni, P., Warner, D., Bensen, R.J., Anstrom, D.C., Harrison, J., Stoecker, M., et al. (2008) Bacterial RNA chaperones confer abiotic stress tolerance in plants and improved grain yield in maize under water-limited conditions. *Plant Physiology*, 147, 446–455.
- Chen, Z., Pottosin, I.I., Cuin, T.A., Fuglsang, A.T., Tester, M., Jha, D., et al. (2007) Root plasma membrane transporters controlling K⁺/Na⁺ homeostasis in salt-stressed barley. *Plant Physiology*, 145, 1714–1725.
- Chun, H.J., Baek, D., Cho, H.M., Lee, S.H., Jin, B.J., Yun, D.-J., Hong, Y.-S., et al. (2019) Lignin biosynthesis genes play critical roles in the adaptation of Arabidopsis plants to high-salt stress. *Plant Signaling & Behavior*, 14(8), 1625697. <http://doi.org/10.1080/15592324.2019.1625697>.
- Clark, D.G., Dervinis, C., Barrett, J.E., Klee, H. & Jones, M. (2004) Drought-induced leaf senescence and horticultural performance of transgenic PSAG12-IPT petunias. *Journal of the American Society for Horticultural Science*, 129, 193–199.
- Cucurachi, M., Busconi, M., Morreale, G., Zanetti, A., Bavaresco, L. & Fogher, C. (2012) Characterization and differential expression analysis of complete coding sequences of *Vitis vinifera* L. sirtuin genes. *Plant Physiology and Biochemistry*, 54, 123–132.
- De Schepper, V., De Swaef, T., Bauweraerts, I. & Steppe, K. (2013) Phloem transport: a review of mechanisms and controls. *Journal of Experimental Botany*, 64, 4839–4850.
- Dellaporta, S.L., Wood, J. & Hicks, J.B. (1983) A plant DNA mini preparation: version II. *Plant Molecular Biology Reporter*, 1, 19–21.
- Derman, Y., Söderholm, H., Lindström, M. & Korkeala, H. (2015) Role of csp genes in NaCl, pH, and ethanol stress response and motility in *Clostridium botulinum* ATCC3502. *Food Microbiology*, 46, 463–470.
- Egley, G.H., Paul, R.N., Vaughn, K.C. & Duke, S.O. (1983) Role of peroxidase in the development of water impermeable seed coats in *Sida spinosa* L. *Planta*, 157, 224–232.
- El-Sharoud, W.M. & Graumann, P.L. (2007) Cold shock proteins aid coupling of transcription and translation in bacteria. *Science Prog.*, 90, 15–27.
- Esfandiari, E. & Shekari, F. (2007) The effect of salt stress on antioxidant enzymes activity and lipid peroxidation on the wheat seedlings. *Notulae botanicae Horti Agrobotanici Cluj-Napoca*, 35, 48–56.
- Fang, L., Jiang, W., Bae, W. & Inouye, M. (1997) Promoter-independent cold-shock induction of CspA and its derepression at 37 degrees by mRNA stabilization. *Molecular Microbiology*, 23, 355–364.
- Fridovich, I. (1989) Superoxide dismutases. An adaptation to a paramagnetic gas. *The Journal of Biological Chemistry*, 264, 7761–7764.
- Fukai, S. & Cooper, M. (1995) Development of drought-resistant cultivars using physiomorphological traits in rice. *Field Crops Research*, 40, 67–86.
- Gall, H.L., Philippe, F., Domon, J.M., Gillet, F., Pelloux, J. & Rayon, C. (2015) Cell wall metabolism in response to abiotic stress. *Plants*, 2015, 112–166.
- Graumann, P. & Marahiel, M.A. (1998) A superfamily of proteins that contain the cold-shock domain. *Trends in Biochemical Sciences*, 23, 286–290.
- Gu, J.F., Qiu, M. & Yang, J.C. (2013) Enhanced tolerance to drought in transgenic rice plants overexpressing C₄ photosynthesis enzymes. *Journal of Crop*, 1, 105–114.
- Hanin, M., Ebel, C., Ngom, M., Laplaze, L. & Masmoudi, K. (2016) New insights on plant salt tolerance mechanisms and their potential use for breeding. *Frontiers in Plant Science*, 7, 1787.
- Hodges, D.M., DeLong, J.M., Forney, C.F. & Prange, R.K. (1999) Improving the thiobarbituric acid-reactive-substances assay for estimating lipid peroxidation in plant tissues containing anthocyanin and other interfering compounds. *Planta*, 207, 604–611.
- Horn, G., Hofweber, R., Kremer, W. & Kalbitzer, H.R. (2007) Structure and function of bacterial cold shock proteins. *Cellular and Molecular Life Sciences*, 64, 1457–1470.
- Hortensteiner, S. (2013) Update on the biochemistry of chlorophyll breakdown. *Plant Molecular Biology*, 82, 505–517.
- Hortensteiner, S. & Kräutler, B. (2011) Chlorophyll breakdown in higher plants. *Biochimica et Biophys Acta-Bioenergetics*, 1807, 977–988.
- Hu, Y., Li, W.C., Xu, Y.Q., Li, G.J., Liao, Y. & Fu, F.L. (2009) Differential expression of candidate genes for lignin biosynthesis under drought stress in maize leaves. *Journal of Applied Genetics*, 50, 213–223.
- Huang, J., Hirji, R., Adam, L., Rozwadowski, K.L., Hammerlind, J.K., Keller, W.A., et al. (2000) Genetic engineering of glycine betaine production toward enhancing stress tolerance in plants: metabolic limitations. *Plant Physiology*, 122, 747–756.
- Huang, L., Sun, Q., Qin, F., Li, C., Zhao, Y. & Zhou, D.X. (2007) Down-regulation of a silent information regulator2-related histone deacetylase gene, OsSRT1, induces DNA fragmentation and cell death in rice. *Plant Physiology*, 144, 1508–1519.

- Huang, X.S., Wang, W., Zhang, Q. & Liu, J.H. (2013) A basic helix-loop-helix transcription factor, PtrbHLH, of *Poncirus trifoliata* confers cold tolerance and modulates peroxidase-mediated scavenging of hydrogen peroxide. *Plant Physiology*, 162, 1178–1194.
- Huang, Y., Guo, Y., Liu, Y., Zhang, F., Wang, Z., Wang, H., et al. (2018) 9-cis-epoxycarotenoid dioxygenase 3 regulates plant growth and enhances multi-abiotic stress tolerance in rice. *Frontiers in Plant Science*, 9, 162.
- Hunger, K., Beckering, C.L., Wiegeshoff, F., Graumann, P.L. & Marahiel, M. A. (2006) Cold-induced putative DEAD box RNA helicases CshA and CshB are essential for cold adaptation and interact with cold shock protein B in *Bacillus subtilis*. *Journal of Bacteriology*, 188, 240–248.
- Ismail, A. & Thomson, M. (2011) Molecular breeding of Rice for problem soils. In: Varshney, R.K. & Costa de Oliveira, A. (Eds.) *Root genomics*. Berlin: Springer, pp. 289–311.
- Jaeggli, B.G., Mortlock, M.Y. & Borrell, A. (2017) Bigger is not always better: reducing leaf area helps stay-green sorghum use soil water more slowly. *Environmental and Experimental Botany*, 138, 119–129.
- Jagdish, K.S., Kavi Kishor, P.B., Bahuguna, R.N., von Wirén, N. & Sreenivasulu, N. (2015) Staying alive or going to die during terminal senescence—an enigma surrounding yield stability. *Frontiers in Plant Science*, 6, 1070.
- Jhonson, S.M., Cummins, I., Lim, F.L., Slabas, A.R. & Knight, M.R. (2015) Transcriptomic analysis comparing stay-green and senescent *Sorghum bicolor* lines identifies a role for proline biosynthesis in the stay-green trait. *Journal of Experimental Botany*, 66, 7061–7073.
- Jiang, W., Hou, Y. & Inouye, M. (1997) CspA, the major cold-shock protein of *Escherichia coli*, is an RNA chaperone. *The Journal of Biological Chemistry*, 272, 196–202.
- Jung, Y.H., Yi, J.Y., Jung, H.J., Lee, Y.K., Lee, H.K., Naicker, M.C., et al. (2010) Overexpression of cold shock protein A of *Psychromonas arctica* KOPRI 22215 confers cold-resistance. *The Protein Journal*, 29, 136–142.
- Keto-Timonen, R., Hietala, N., Palonen, E., Hakakorpi, A., Lindstrom, M. & Korkeala, H. (2016) Cold shock proteins: a minireview with special emphasis on Csp-family of enteropathogenic *Yersinia*. *Frontiers in Microbiology*, 7, 1151.
- Kim, M.H., Sasaki, K. & Imai, R. (2009) Cold shock domain protein 3 regulates freezing tolerance in *Arabidopsis thaliana*. *The Journal of Biological Chemistry*, 284, 23454–23460.
- Kim, M.H., Sato, S., Sasaki, K., Saburi, W., Matsui, H. & Imai, R. (2013) Cold shock domain protein 3 is involved in salt and drought stress tolerance in *Arabidopsis*. *FEBS Open Bio*, 3, 438–442.
- Kim, Y., Kim, C.Y., Song, W., Park, D., Kwon, S., Lee, H., et al. (2008) Overexpression of sweet potato swpa4 peroxidase results in increased hydrogen peroxide production and enhances stress tolerance in tobacco. *Planta*, 227, 867–881.
- Kishor, P.B.K., Hima Kumari, P., Sunita, M.S.L. & Sreenivasulu, N. (2015) Role of proline in cell wall synthesis and plant development and its implications in plant ontogeny. *Frontiers in Plant Science*, 6, 544.
- Kishor, P.B.K., Hong, Z., Miao, G.H., Hu, C. & Verma, D.P.S. (1995) Overexpression of D-pyrroline-5-carboxylate synthetase increases proline production and confers osmotolerance in transgenic plants. *Plant Physiology*, 108, 1387–1394.
- Kishor, P.B.K., Sangam, S., Amrutha, R.N., Sri Laxmi, P., Naidu, K.R., Rao, K. R.S.S., et al. (2005) Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: its implications in plant growth and abiotic stress tolerance. *Current Science*, 88, 424–438.
- Koca, H., Bor, M., Ozdemir, F. & Turkan, I. (2007) The effect of salt stress on lipid peroxidation, antioxidant enzymes and proline content of sesame cultivars. *Environmental and Experimental Botany*, 60, 344–351.
- Lagunas-Rangel, F.A. (2019) Current role of mammalian sirtuins in DNA repair. *DNA Repair (Amst)*, 80, 85–92.
- Leopold, A.C. & Kawase, M. (1964) Benzyladenine effects on bean leaf growth and senescence. *American Journal of Botany*, 51, 294–298.
- Liang, C., Wang, Y., Zhu, Y., Tang, J., Hu, B., Liu, L., et al. (2014) OsNAP connects abscisic acid and leaf senescence by fine-tuning abscisic acid biosynthesis and directly targeting senescence-associated genes in rice. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 10013–10018.
- Maheshwari, P., Kiran, B., Punita, D.L. & Kishor, P.B.K. (2017) Overexpression of SbAP37 in rice alleviates concurrent imposition of combination stresses and modulates different sets of leaf protein profiles. *Plant Cell Reports*, 36, 773–786.
- Melencion, S.M.B., Chi, Y.H., Pham, T.T., Paeng, S.K., Wi, S.D., Lee, C., et al. (2017) RNA chaperone function of a universal stress protein in *Arabidopsis* confers enhanced cold stress tolerance in plants. *International Journal of Molecular Sciences*, 18, 2546.
- Merewitz, E., Gianfagna, T. & Huang, B. (2010) Effects of SAG12-ipt and HSP18.2-ipt expression on cytokinin production, root growth and leaf senescence in creeping bentgrass exposed to drought stress. *Journal of the American Society for Horticultural Science*, 135, 230–239.
- Molinari, H.B.C., Marur, C.J., Daros, E., MKF, C., JFRP, C., JCB, F., et al. (2007) Evaluation of the stress-inducible production of proline in transgenic sugarcane: osmotic adjustment, chlorophyll fluorescence and oxidative stress. *Plant Physiology*, 130, 218–229.
- Monties, B. & Fukushima, K. (2001, 2001) Occurrence, function and biosynthesis of lignins. In: Hofrichter, M. & Steinbuechel, A. (Eds.) *Biopolymers. Lignin, humic substances and coal*, Vol. 1. Weinheim, Germany: Wiley, pp. 1–64.
- Morison, J.I.L., Baker, N.R., Mullineaux, P.M. & Davies, W.J. (2008) Improving water use in crop production. *Philosophical transactions of the Royal Society of London*, 363, 639–658.
- Mourasobczak, J., Souza, U. & Mazzafera, P. (2011) Drought stress and changes in the lignin content and composition in *Eucalyptus*. *BMC Proceedings*, 2011, 5.
- Murashige, T. & Skoog, F. (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15, 473–497.
- Nakaminami, K., Karlson, D.T. & Imai, R. (2006) Functional conservation of cold shock domains in bacteria and higher plants. *Proceedings of the National Academy of Sciences of the United States of America*, 103, 10122–10127.
- Neto, A.D.A., Prisco, J.T., Eneas-Filho, J., Abreu, C.E.B. & Gomez-Filho, E. (2006) Effect of salt stress on antioxidative enzymes and lipid peroxidation in leaves and roots of salt-tolerant and salt-sensitive maize genotypes. *Environmental and Experimental Botany*, 56, 87–94.
- Park, S.J., Kwak, K.J., Oh, T.R., Kim, Y.O. & Kang, H. (2009) Cold shock domain proteins affect seed germination and growth of *Arabidopsis thaliana* under abiotic stress conditions. *Plant and Cell Physiology*, 50, 869–878.
- Qi, T., Wang, J., Huang, H., Liu, B., Gao, H., Liu, Y., et al. (2015) Regulation of jasmonate-induced leaf senescence by antagonism between bHLH subgroup IIIe and III d factors in *Arabidopsis*. *Plant Cell*, 27, 1634–1649.
- Reddy, P.S., Jageswar, G., Rasineni, G.K., Maheswari, M., Reddy, A.R., Varshney, R.K., et al. (2015) Proline over-accumulation alleviates salt stress and protects photosynthetic and antioxidant enzyme activities in transgenic sorghum [*Sorghum bicolor* (L) Moench]. *Plant Physiology and Biochemistry*, 94, 104–113.
- Rivero, R.M., Kojima, M., Gepstein, A., Sakakibara, H., Mittler, R., Gepstein, S., et al. (2007) Delayed leaf senescence induces extreme drought tolerance in a flowering plant. *Proceedings of the National Academy of Sciences of the United States of America*, 104, 19631–19636.
- Romero, C., Belles, J.M., Vaya, J.L., Serrano, R. & Cullianez-Maria, F.A. (1997) Expression of the yeast trehalose-6-phosphate synthetase gene in transgenic tobacco plants: pleiotropic phenotypes include drought tolerance. *Planta*, 201, 293–297.
- Sakuraba, Y., Park, S.Y., Kim, Y.S., Wang, S.H., Yoo, S.C., Hörtensteiner, S., et al. (2014) *Arabidopsis* STAY-GREEN2 is a negative regulator of chlorophyll degradation during leaf senescence. *Molecular Plant*, 7, 1288–1302.

- Sambrook, J. & Russell, S.W. (2001) *Molecular cloning: a laboratory manual*, 3rd edition. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Sanders, D., Pelloux, J. & Brownlee, C. (2002) Calcium at the crossroads of signalling. *Plant Cell*, 14, 401–417.
- Sasaki, K. & Imai, R. (2012) Pleiotropic roles of cold shock domain proteins in plants. *Frontiers in Plant Science*, 2, 116–119.
- Sasaki, K., Kim, M.H. & Imai, R. (2007) Arabidopsis cold shock domain protein2 is a RNA chaperone that is regulated by cold and developmental signals. *Biochem Biophys Res Com*, 364, 633–638.
- Sasaki, K., Liu, Y., Kim, M. & Imai, R. (2015) An RNA chaperone, AtCSP2, negatively regulates salt stress tolerance. *Plant Signaling and Behavior*, 10, e1042637.
- Sato, Y. & Yokoya, S. (2008) Enhanced tolerance to drought stress in transgenic rice plants overexpressing a small heat-shock protein smHSPs177. *Plant Cell Reports*, 27, 329–334.
- Schmid, B., Klumpp, J., Raimann, E., Loessner, M.J., Stephan, R. & Tasara, T. (2009) Role of cold shock proteins in growth of *Listeria monocytogenes* under cold and osmotic stress conditions. *Applied and Environmental Microbiology*, 75, 1621–1627.
- Semchuk, N.M., Vasylyk, Y.V., Lushchak, O.V. & Lushchak, V.I. (2012) Effect of short-term salt stress on oxidative stress markers and antioxidant enzymes activity in tocopherol deficient *Arabidopsis thaliana* plants. *Ukrainskii Biokhimičeskii Zhurnal*, 84, 41–48.
- Shigeto, J., Itoh, Y., Hirao, S., Ohira, K., Fujita, K. & Tsutsumi, Y. (2015) Simultaneously disrupting *AtPrx2*, *AtPrx25* and *AtPrx71* alters lignin content and structure in *Arabidopsis* stem. *Journal of Integrative Plant Biology*, 57, 349–356.
- Sreenivasulu, N., Butardo, V.M., Misra, G., Cuevas, R.P., Anacleto, R. & Kavi Kishor, P.B. (2015) Designing climate-resilient rice with ideal grain quality suited for high-temperature stress. *Journal of Experimental Botany*, 66, 1737–1748.
- Tarczynski, M.C., Jensen, R.G. & Bohnert, H.J. (1993) Stress protection of transgenic tobacco by production of the osmolyte mannitol. *Science*, 259, 508–510.
- Thomas, H. & Ougham, H. (2014) The stay-green trait. *Journal of Experimental Botany*, 65, 3889–3900.
- Tseng, M.J., Liu, C.W. & Yiu, J.C. (2007) Enhanced tolerance to sulfur dioxide and salt stress of transgenic Chinese cabbage plants expressing both superoxide dismutase and catalase in chloroplasts. *Plant Physiology and Biochemistry*, 45, 822–833.
- Uppal, S., Shetty, D.M., & Jawali, N. (2014) Cyclic amp receptor protein regulates *cspD*, a bacterial toxin gene, in *Escherichia coli*. *Journal of Bacteriology*, 196(8), 1569–1577. <http://doi.org/10.1128/jb.01476-13>.
- Van Heerden, P.D.R. & de Villiers, O.T. (1996) Evaluation of the relative water content and the reduction of 2, 3, 5-triphenyltetrazoliumchloride as indicators of drought tolerance in spring wheat cultivars. *South Af J Plant Soil*, 13, 131–135.
- Weber, M.H. & Marahiel, M.A. (2002) Coping with the cold: the cold shock response in the Gram-positive soil bacterium *Bacillus subtilis*. *Philosophical transactions of the Royal Society of London*, B357, 895–907.
- Willekens, H., Inze, D., Van Montagu, M. & Van Camp, W. (1995) Catalases in plants. *Molecular Breeding*, 1, 207–228.
- Xu, D., Duan, X., Wang, B., Hong, B., Ho, T. & Wu, R. (1996) Expression of a late embryogenesis abundant protein gene, HVA1, from barley confers tolerance to water-deficit and salt stress in transgenic rice. *Plant Physiology*, 110, 249–257.
- Xu, Y., Burgess, P., Zhang, X. & Huang, B. (2016) Enhancing cytokinin synthesis by overexpressing *ipt* alleviated drought inhibition of root growth through activating ROS-scavenging systems in *Agrostis stolonifera*. *Journal of Experimental Botany*, 67, 1979–1992.
- Yamanaka, K. & Inouye, M. (1997) Growth-phase-dependent expression of *cspD*, encoding a member of the CspA family in *Escherichia coli*. *Journal of Bacteriology*, 179, 5126–5130.
- Yiu, J.C. & Tseng, M.J. (2005) Manipulation of superoxide dismutase and catalase to enhance sulfur dioxide tolerance in transgenic Chinese cabbage. *Acta Horticulturae*, 692, 91–100.
- Yu, T., Xu, Z., Guo, J., Wang, Y., Abernathy, B., Fu, J., et al. (2017) Improved drought tolerance in wheat plants overexpressing a synthetic bacterial cold shock protein gene *SeCspA*. *Scientific Reports*, 7, 44050.
- Zagorchev, L., Kamenova, P. & Odjakova, M. (2014) The role of plant cell wall proteins in response to salt stress. *The Scientific World Journal*, 2014, 764089, 9 pages.
- Zhao, Y., Chan, Z., Gao, J., Xing, L., Cao, M., Yu, C., et al. (2016) ABA receptor *PYL9* promotes drought resistance and leaf senescence. *Proceedings of the National Academy of Sciences of the United States of America*, 113, 1949–1954.
- Zheng, W. (2020) The plant sirtuins. *Plant Science*, 293, 110434.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Guddimalli R, Somanaboina AK, Palle SR, et al. Overexpression of RNA-binding bacterial chaperones in rice leads to stay-green phenotype, improved yield and tolerance to salt and drought stresses. *Physiologia Plantarum*. 2021;1–18. <https://doi.org/10.1111/ppl.13369>