

1 **Production and use of two marine zooplanktons,**

2 ***Tigriopus japonicus* and *Diaphanosoma celebensis*, as live food for**

3 **red sea bream *Pagrus major* larvae**

4
5 **ATSUSHI HAGIWARA¹, HEE-JIN KIM^{1*}, HOKUTO MATSUMOTO¹, YUSUKE OHTA¹,**

6 **TETSURO MORITA², AKIMASA HATANAKA², RISA ISHIZUKA², YOSHITAKA SAKAKURA¹**

7
8 ¹*Graduate School of Fisheries and Environmental Sciences, Nagasaki University, Bunkyo 1-14,*
9 *Nagasaki 852-8521, Japan.*

10 ²*Central Research Laboratory, Nippon Suisan Kaisha, Ltd., 1-32-3 Nanakuni, Hachioji-shi, Tokyo*
11 *192-0991, Japan.*

12
13
14 * Corresponding author: Tel/Fax: (+81) 95-819-2830. Email: heejin@nagasaki-u.ac.jp

26 **ABSTRACT:** We evaluated the effectiveness of two representative marine zooplanktons, harpacticoid copepod
27 *Tigriopus japonicus* and euryhaline cladoceran *Diaphanosoma celebensis* as live food for red sea bream *Pagrus major*
28 larvae. Chicken-dropping extract (CDE) was applied to both zooplankton cultures for improving population growth.
29 Population growth of both animals was significantly enhanced by CDE supplementation (at 1 or 2 ml/l). The highest
30 amount of DHA and higher DHA/EPA ratio was detected in *T. japonicus*, whereas *D. celebensis* showed similar values
31 to that of *Artemia*. Effectiveness of both animals as live food was tested by rearing red sea bream larvae for 28 days
32 and compared with that of *Artemia*. There were no significant differences in total length (8.6 ± 1.1 - 8.7 ± 0.7 mm) and
33 wet weight (8.2 ± 0.3 - 9.4 ± 0.1 mg) among fish larvae received three different zooplanktons. Survival rate was
34 significantly higher with *T. japonicus* ($39.4\pm 3.1\%$) than *D. celebensis* ($20.8\pm 3.8\%$) and *Artemia* ($16.7\pm 9.8\%$).
35 Viability was significantly higher in fish fed with *T. japonicus* ($60.0\pm 27.8\%$) and *D. celebensis* ($60.0\pm 32.2\%$) than those
36 with *Artemia* ($44.4\pm 12.3\%$). Fish fed with *T. japonicus* contained higher n-3 highly unsaturated fatty acids than those
37 with *D. celebensis* and *Artemia*. It is concluded that *T. japonicus* and *D. celebensis* have high potential as live food for
38 marine fish larviculture.

39
40 **KEY WORDS:** copepoda, cladocera, red sea bream, larviculture, growth, survival.

41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57 **INTRODUCTION**

58

59 Food web of hydrosphere utilizes various zooplanktons as energy transporter from photosynthetic sources to series of
60 consumers, e.g. larval animals. Among the zooplanktons in the marine ecosystem, monogonont rotifer *Brachionus*
61 *plicatilis* sp. complex are widely applied to the commercial hatchery facilities as an initial live food mainly because of
62 their small size which is suitable for the larvae, rapid population growth, and ease to be cultured and nutritionally
63 fortified. Once the larvae are in the advanced stage, brine shrimp (*Artemia* spp.) is also generally supplied to larval
64 animals associated with development [1, 2] because of its convenience in use (i.e., cyst usage to reduce labor-intensive
65 live food availability) and good nutritional value [3]. In spite of the long history of using *Artemia*, many challenges
66 still remain. At present, the most marketed cysts of *Artemia* are from the Great Salt Lake (GSL), and thus its provision
67 is unpredictable in terms of demand, harvest, cost and nutritional values [4]. Based on these issues, there is a growing
68 interest on the use of other zooplanktons and the need to establish a method of mass culturing them in the hatchery [5].

69 Copepods are major part of the diet for larval animals in the pelagic food chain and are generally known to match the
70 nutritional requirements of the predators, and have higher nutritional value compared to rotifers (*Brachionus* spp.) and
71 *Artemia* [6-8]. Interest in copepod as a live food for aquaculture has grown since 1980's. Harpacticoid copepod
72 *Tigriopus japonicus* can be cultured at higher density compared to other copepod species and thrive in harsh
73 environmental conditions [9-11]. In addition to these biological characteristics, relatively small size (1 mm of adult
74 body length) zooplanktons attracts attentions for usability as a live food [12, 13], while its epibenthic habitat remained a
75 problem to extend for aquaculture facilities targeted marine fish larvae [14].

76 Cladocerans comprised the natural diet for many brackish and freshwater larval animals [15] and due to their
77 parthenogenetic reproduction, rapid propagation is possible. The brackishwater cladoceran, *Diaphanosoma celebensis*
78 has the similar size distribution to *Artemia* and strong tolerance to salinity variations [16, 17]. Based on these
79 perspectives, the studies for its application to the larviculture have been tried extensively [18-20].

80 Seedling production in aquaculture is usually confronted by the cost of producing enough and highly nutritious live
81 foods e.g. zooplankton. To answer this issue, organic fertilizers such as animal manures were suggested as a booster
82 of zooplankton population growth [21, 22] and this method is proven to be useful in many developing countries [23].
83 Among animal manures, chicken manure is preferred because it is easily soluble and contains high level of nitrogen,
84 phosphorus and potassium [24, 25]. It is indeed known to enhance population growth of zooplankton populations in
85 fishpond setting [26]. In this study, we tested the use of chicken-dropping extracts (CDE) to enhance the population
86 growth of two zooplanktons: *T. japonicus* and *D. celebensis* which have high potential as a live food for intensive
87 larviculture. The mass cultured zooplanktons were fed to red sea bream *Pagrus major* larvae, and growth and survival

88 of the larvae was compared to those fed on *Artemia franciscana* to elucidate the qualification of those zooplanktons as a
89 live food.

90

91

92 **MATERIALS AND METHODS**

93

94 **CDE effects on the population growth of zooplanktons**

95

96 The copepod *T. japonicus* was cultured in 800 ml of glass beaker containing 800 ml of natural sea water (34 ppt) with
97 initial density at 0.4 ind./ml (total 320 individuals consisted of 160 nauplii and 160 copepodites). Copepods were fed
98 on vitamin B₁₂ enriched *Chlorella vulgaris* (Super fresh chlorella-V12, Chlorella Industry Co. Ltd., Fukuoka, Japan) at
99 2.5×10^6 cells/ml every 3 days. The culture of cladoceran *D. celebensis* was initiated in 200 ml of glass beaker
100 containing 200 ml of diluted seawater (22 ppt) with 100 individuals (at 0.5 ind./ml), which was randomly selected from
101 a preliminary culture maintained under the same environmental conditions as experimental cultures. Because of the
102 difficulty to conduct small scale batch culture using *Chlorella*, the animals were fed on *Nannochloropsis oculata* at
103 7×10^6 cells/ml every 2 days to maintain their conditions without external stresses like aeration. The microalgae *N.*
104 *oculata* was cultured in the modified Erd-Schreiber medium [27] under continuous light with gentle aeration. Prior to
105 feeding, the culture medium of *N. oculata* was centrifuged at $3968 \times g$ for 10 min and collected cells were re-suspended
106 in the zooplankton culture medium. Photoperiod and temperature were in the two set up were adjusted at optimal
107 conditions for each species by preliminary tests (personal information): at 18L:6D, 25 °C for *T. japonicus* and under
108 total darkness at 28 °C for *D. celebensis*. Culture media for zooplanktons were prepared by GF/C (CAT No. 1822-047,
109 Whatman) filtration of natural seawater followed by autoclave sterilization at 121 °C for 20 min.

110 Chicken-dropping extract (CDE) was prepared by the following procedure: 1 kg of fermented chicken droppings
111 (Shitama Inc., Fukuoka, Japan) were mixed with 10 g of fossil coral powder (Coral international Co. Ltd., Okinawa,
112 Japan). The mixture was boiled in 5 l of tap water for 40-50 minutes and then kept overnight at room temperature. The
113 resulting supernatant was filtered in plankton net (150-200 µm of mesh) and mixed with extracted liquid from sludge by
114 the same filtration. The solution (CDE) was preserved at 5 °C until use. To determine the effect of CDE on the
115 population growth of *T. japonicus* and *D. celebensis*, CDE concentration was adjusted in the culture media at 0 (without
116 CDE: control), 1, or 2 ml/l with three replicates and the population density was estimated every 3 days for copepod and
117 every 2 days for cladoceran during culture periods. The culture period lasts for 30 days for copepod, and 18 days for
118 cladoceran.

119 **Potential as live food for marine fish larvae**

120 Different zooplankton species were cultured in order to determine their potential as live food for *Pagrus major*
121 larviculture. L-type rotifer *Brachionus plicatilis* sensu stricto (Makishima strain) was cultured in 50 l of artificial
122 seawater (Marine Art Hi, Tomita Pharmaceutical Co. Ltd., Naruto, Japan) adjusted at 22 ppt and 25 °C under 12L:12D
123 of photoperiod with gentle aeration. The rotifers were daily fed with *C. vulgaris* (Super Fresh Chlorella-V12) at
124 2.5×10^6 cells/ml twice a day. For *Artemia* feeding, the cyst of *A. franciscana* was incubated in 5 l of 34 ppt artificial
125 sea water at 22 °C of water temperature under 12L:12D of photoperiod with strong aeration. From day 2 after
126 hatching, the nauplii were fed daily with Super Fresh Chlorella-V12 at 2.5×10^6 cells/ml for 24 hours. The copepod *T.*
127 *japonicus* was semi-continuously cultured in 100 l of 34 ppt artificial sea water at 25 °C with 80 ml/min of aeration
128 under 12L:12D of photoperiod. Food supplement was daily performed at 5.0×10^6 cells of *Chlorella*/ml. The
129 cladoceran *D. celebensis* was semi-continuously cultured in 30 l of 22 ppt artificial seawater at 25 °C with gentle
130 aeration and twice supplementation of *Chlorella* at 2.5×10^6 cells/ml a day. The CDE was supplied to each culture of *T.*
131 *japonicus* and *D. celebensis* at 1 and 2 ml/l, respectively, due to the limited amount of the prepared CDE.

132 To compare size distribution of the three zooplanktons tested, 100 individuals of each species were fixed with 5% of
133 neutral formalin for *Artemia* and Lugol solution for copepods and cladocera. The body length of fixed individuals
134 were measured using microscopic measurement system including stereomicroscope (SteREO Discovery V8, ZEISS,
135 Germany) equipped with a digital camera (Axio Cam HSm, ZEISS) and an image-analysis software (Axio Vision
136 Release 4.8.2., ZEISS). The measurement was performed under $\times 20$ of magnification. The fatty acid composition of
137 the cultured zooplanktons was analyzed by the following procedure. Mass cultured zooplanktons were collected by
138 plankton net (45 μ m of mesh) and rinsed with distilled water at several days intervals. After removal of remaining
139 water with a paper tissue, the samples were preserved at -40 °C until analysis. Fatty acid analysis was performed at
140 Oita Marine Biological Technology Center, Nippon Suisan Kaisha Ltd., Oita, Japan, and the detailed procedure is same
141 as that used for the fish larvae.

142 For larviculture trials, fertilized eggs of red sea bream *P. major* were obtained from a local fish farmer hatchery
143 (Ogata Suisan, Kumamoto, Japan). The eggs were incubated in a 100 l of polycarbonate tank containing 34 ppt
144 artificial seawater at 18 °C with 100 ml/min of aeration. Newly hatched larvae (0 days post hatch, dph) were transferred
145 into 9 aquaria each containing 100 l of 34 ppt artificial seawater in a temperature controlled room with 5 l of ceramic
146 sand (grain size: 0.3-0.6 mm, Micros ceramic, NORRA Co. Ltd., Kyoto, Japan) covering the bottom of the tank.
147 Larvae were stocked in each tank following the procedure of Kim et al. [28] in which larval density was adjusted to 10
148 ind./l. Water temperature in the tank was gradually raised from 18 to 22 °C by daily acclimation of 1°C. Laval
149 rearing was performed at a light-dark cycle of 12L:12D. The microalgae (Super Fresh Chlorella-V12) at 5×10^5

150 cells/ml was added into the prepared aquaria on 4 dph and this density was maintained until 28 dph [29, 30]. The
151 feeding scheme is shown in Fig. 1. Fish from 4 (mouth opening) to 23dph were fed on the rotifers twice a day and the
152 density was maintained at 10 ind./ml in the larval rearing tanks. On 16 dph, 400 fish were newly transferred into each
153 experimental tank to sort the fish number prior to switch the food items from rotifer to the targeted species. The tanks
154 were assigned to each diet treatment with triplicates. When larvae reached notochord flexion phase (from 20 dph),
155 *Artemia*, copepod, and cladoceran were fed in triplicates at 0.01 ind./ml 3 to 5 times a day according to their growth.
156 To estimate the effectiveness of the zooplanktons as live food for fish larvae, the 6 following parameters were
157 conducted [28, 31].

158

159 ***Hatching and Survival activity index***

160 The fertilized fish eggs (30 eggs) were incubated in a 500 ml of glass beaker containing 500 ml of artificial seawater at
161 18°C under total darkness to calculate hatching rate and survival activity index (SAI) with triplicate observations. The
162 hatching rate was determined by the number of hatchlings after 24 h. The SAI was estimated by the following
163 equation [31]:

$$SAI = \frac{1}{N} \sum_{i=1}^K (N - hi) \times i$$

164 where N is the total number of examined larvae, hi is the cumulated mortality by i -th day, K is the number of days
165 elapsed until all fish larvae died due to starvation.

166

167 ***Survival***

168 The survival rate was calculated from the mean number of surviving fish larvae in three aquaria for each zooplankton
169 species on the last day of larviculture (28 dph). The adjusted density at 400 individuals on 16 dph was applied as an
170 initial number of fish larvae.

171

172 ***Viability***

173 On the last day of larviculture, air exposure test was conducted to compare the viability of the larvae fed on each
174 zooplankton. We caught fish larvae on a net (130×345 mm, Super net M, SANY co., Ltd., Kanagawa, Japan) from
175 each tank (n=3) and exposed them to air for 1 minute. After this, the fish were immediately returned to seawater and
176 their survival was observed every 2 hours for 24 hours.

177

178 ***Growth***

180 Larval growth was determined by measuring the total length and wet weight. On 20, 23, 26, and 28 dph, 20 larvae
181 were randomly collected from each aquarium, and anaesthetized with MS222 followed by 5% neutral formalin fixation.
182 The total length was measured with all the fixed larvae under digital microscope (VH-6300, Keyence, Japan). Wet
183 weight of fish on 28 dph was measured using an analytical balance (AB204-S, Mettler-Toledo International Inc., United
184 Kingdom). Using these data, the total biomass of fish larvae (i.e., production) was calculated with the number of
185 surviving larvae on the last day of the experiment. To estimate optimal prey size, upper jaw length (JL) was measured
186 using the larvae of 20 dph, and the mouth diameter was determined by the following equation: $\sqrt{2}(JL)$, where the
187 assumption is that the mouth opens to an angle of 90° during prey capture [32].

188

189 ***Fatty acid analysis***

190 For the fatty acid composition of fish larvae fed on three different diets, the reared larvae were sampled at the end of the
191 experiment and preserved at -40 °C until analysis. The analysis was performed at Central Research Laboratory of
192 Nippon Suisan Kaisha, Ltd. with the following detailed method. Pooled cultured zooplankton or fish larvae (on 20 and
193 28 dph) homogenates were precisely weighted in glass tubes. An internal standard consisting of 20 µg tricosanoic-
194 acid (C23:0) and 50 µg butylated hydroxytoluene dissolved in 2 ml of methanol-hexane 4:1 (v/v) was added to
195 biological samples and methylated in the presence of 200 µl acetyl chloride at 80 °C for 1 h, based on the method of
196 Lepage and Roy [33]. After cooling on ice, 5 ml of 6% (w/v) aqueous potassium carbonate was added to each tube to
197 stop the methylation reactions, and centrifuged at 2000 rpm for 5 minutes. The upper organic phase containing the
198 fatty acid methyl ester was collected, and analyzed on a DB-Wax column (30 m length, 0.32 mm id, 0.25 µm film)
199 (Agilent Technologies) coupled to a GC System 6850N (Agilent Technologies). The gas chromatography oven
200 temperature was 180 °C and increased at a rate of 3 °C/min to a final temperature of 230 °C.

201

202 **Statistical analysis**

203

204 The CDE effects on the population growth of the both zooplanktons related to its concentrations and culture day were
205 analyzed by two-way repeated-measures ANOVA using Statview version 5.0 (SAS Institute Inc., USA). When
206 significant differences were detected ($P<0.05$), Tukey's HSD test was performed by R version 3.1.2 [34]. For the
207 fish larviculture, the mean body length of the three zooplankton species, and survival, wet weight and biomass of fish
208 larvae on 28 dph associated with the targeted live food species were compared with one-way ANOVA followed by
209 Tukey-Kramer *post hoc* test as the first test showed significant differences ($P<0.05$). To compare the viability of fish
210 larvae, Log-rank test were performed. The variation of larval total length was analyzed by two-way repeated-

211 measures ANOVA followed by Tukey-Kramer *post hoc* test associated with the food types and culture days (20, 23, 26,
212 and 28 dph). These analyses for fish larviculture were performed by Statview (SAS institute).

213

214

215 **RESULTS**

216

217 **Effects of CDE on zooplankton populations**

218

219 The population growth of copepods was observed with developmental stages: nauplius, copepodite, and
220 nauplius+copepodite (Fig. 2). The population growth of each treatment increased with the culture days ($P<0.0001$)
221 but the pattern was different among CDE concentrations ($P<0.0001$). At 2 ml/l of CDE, active population growth was
222 obtained regardless of developmental stages ($P<0.0001$) and population growth decreased at lower CDE concentration.
223 On the last day of culture, three developmental groups showed the highest count at 2 ml/l ($P<0.0001$): 4408.9±321.1 ind.
224 of nauplii, 7768.9±635.5 ind. of copepodites, and 12177.8±694.6 ind. of total population.

225 The population growth of cladocera varied with the two parameters: culture days and CDE concentrations (Fig. 3). The
226 cladocera population maintained steady growth until day 12 but sharply decreased thereafter. The highest density
227 (14.0±2.6 ind./ml) on day 12 was observed with 2 ml/l of CDE supplementation and it was decreased with the reduction
228 of CDE concentration: 10.3±1.5 ind./ml at 1 ml/l and 8.7±3.2 ind./ml in the control group ($P<0.05$).

229 Total fatty acid level of each animal was described as follows: 0.76% of *Artemia* wet weight, 1.16% of copepod, and
230 1.04% of cladocera. The highest proportion of n-3 HUFA and DHA (C22:6n-3) / EPA(C20:5n-3) ratio were in
231 copepods cultured with CDE (Table 2).

232

233 **Fish larviculture**

234

235 The employed fish eggs showed 94.4±5.1% of hatching rate and 13.6±6.6 of SAI, respectively. Fish larvae from
236 these eggs showed significantly higher survival rate with the copepods compared to those reared with cladocerans or
237 *Artemia* (Table 1) ($P<0.05$). The larval diets of copepod and cladoceran induced higher viability compared to *Artemia*
238 (Table 1) ($P<0.05$).

239 Total length of fish larvae on 28 dph was shown as 8.7±0.7 mm with *Artemia*, 8.5±1.1 mm with copepod, and
240 8.8±0.7 mm with cladoceran, respectively without significant differences among diet treatments. The calculated
241 mouth diameter of 20 dph fish larvae was 0.97±0.08 mm, therefore the size of optimal prey was ranged from 0.5 to 0.7

242 mm which is similar to the mean size of each zooplankton species: 0.8 ± 0.1 mm for *Artemia*, 0.9 ± 0.1 mm for copepods,
243 and 0.7 ± 0.0 mm for cladoceran (Fig. 5). There were no significant differences in wet weight of fish (Table 1) and the
244 total biomass (i.e., production of 28 dph, Fig. 6) of fish larvae among those with three larval diets. The larval
245 production on the last day of larviculture was shown as follows: 552.0 ± 325.4 mg with *Artemia* feeding, 1395.6 ± 564.2
246 mg with copepod, and 780.3 ± 134.4 mg with cladoceran.

247 The total fatty acids were estimated to compose 1.3% of larval wet weight with *Artemia* and copepod diet, and 1.5%
248 with cladoceran which value was slightly higher than initial rotifer-fed larvae (1.4%) on the last day of larviculture.
249 Among these fish larvae (on 28 dph), only the copepod-fed one showed higher proportion of total n-3 HUFA and
250 DHA/EPA ratio compared to the initial larvae on 20 dph (Table 3).

251

252

253 **DISCUSSION**

254

255 Intensive larviculture system of red sea bream has been stably established with rotifers and *Artemia* as live foods related
256 to the developmental stages [35, 36]. Previous studies were conducted in small scale larviculture and obtained high
257 survival, viability and growth with copepod diet, but it has not been applied to intensive culture in larger scale.
258 Feasibility of the tested zooplanktons: *T. japonicus* and *D. celebensis* depends on the competitive cost to culture them at
259 higher densities for the intensive larviculture system. This study made an attempt to mass culture of these
260 zooplanktons with the addition of CDE, because we further aimed to promote cost-effective method of mass production
261 of these zooplanktons to fish culturists. Our results clearly showed the efficiency of CDE to enhance production
262 achieve high density culture of the employed species with proper feeding dosages (Fig. 2 and 3). Many studies on the
263 use of animal manure showed that indeed, CDE could enhance zooplankton population growth [37-39]. The chicken
264 manure is known to contain water-soluble natural 17β -estradiol (E2) [40, 41]. In addition, supplementation of
265 synthetic E2 (10-1000 μ g/l) increased reproduction of *D. celebensis* [42], but not that of *T. japonicus* [43]. The increase
266 in the population growth in *D. celebensis* with the addition CDE should be viewed as direct effect, but other
267 mechanisms may be involved in *T. japonicus*. *T. japonicus* is known as omnivorous and shift feeding resources to
268 detritus when living particles become limited [44]. Detritus contains bacteria which are important decomposer of
269 organic matters, and its population increased with chicken manure dosage [45]. The accelerated bacterial growth with
270 the CDE is expected to induce the better growth of copepod population in this study. Not only to the copepods,
271 several studies also reported that bacteria contribute to the diet of cladoceran [46, 47]. Therefore, the high population
272 growth of cladoceran is probably due to the build-up of bacterial populations by CDE.

273 Hatching rate of fertilized eggs and SAI are used to estimate initial larval quality [48]. These values in this study
274 are comparable to those reported by Kim et al. [28] and are higher than those of other fishes [49]. SAI of fish larvae is
275 influenced by ambient environmental conditions and *Epinephelus akaara* larvae exhibited about 12 of SAI under
276 optimal conditions for stable growth and development [50]. It should indicate that the set-up conditions for the present
277 larval rearing are suitable for the targeted larval fish. Under this condition, the effective larval rearing of red sea
278 bream was constructed with copepod and cladoceran cultured with CDE, and these fish larvae showed higher survival
279 and viability (Table 1) compared to those with *Artemia* in the present study.

280 The survival rate associated with the dietary sources was the highest with the copepod *T. japonicus* (Table 1) even
281 though it has epibenthic features. Influences of turbulence by aeration and predator presence may also change
282 swimming behavior of copepods (e.g., frequency and speed) [14, 51] and should be examined. The size distribution of
283 the employed cladoceran was estimated most favorable for the targeted fish larvae (Fig. 5) to induce the high capture
284 efficiency. The larval capture efficiency with copepod is lower than with cladocerans, although, the earlier stage
285 larvae prefer the copepod adults and nauplii because copepods yield substantial energy to larval fish because of the
286 minimal handling time [52]. This phenomenon is expected to induce the highest survival rate with copepod diet
287 (Table 1). All the mass cultures of zooplanktons were maintained with *C. vulgaris* which has similar cell component
288 to *N. oculata* [53]. The cultivated copepods with CDE contained the highest amount of fatty acids compared to the
289 other zooplanktons (Table 2), although, the level was comparatively lower than the natural one [54]. However, this
290 amount is enough to maintain the stable growth and development of red sea bream larvae [55], especially n-3 HUFAs
291 was highly contained in copepods. It was expected that the copepod feeding induced stable survival of the targeted
292 fish larvae [56]. In addition, the parameter of DHA/EPA ratio is regarded as an effective factor to determine food and
293 larva quality. The optimal ratio of marine fish larval food is estimated more than 1 by comparing with natural diet
294 [57] and the ratio of reared fish larvae is more than 5 which was determined by grunt *Plectorhynchus cinctus* [58].
295 The effects of DHA/EPA ratio on the larval survival was clearly observed in this study and the copepod-fed larvae
296 showed the highest survival rate (Table 1) associated with the higher ratio of it (5.6 in Table 3).

297 *P. major* larvae fed with *T. japonicus* and *D. celebensis* showed the better resistance than those fed with *Artemia*
298 (Table 1). Successful larval rearing generally depends on first feeding regimes with live food species and its nutritional
299 quality. The dietary lipids especially the essential fatty acid (EFA) is recognized as one of the most important
300 nutritional factor that influence larval growth and survival [59, 60] and its deficiency will result to various symptoms
301 including decrease of larval health, poor growth, low feed efficiency, anaemia and high mortality [61-63]. It was also
302 reported that red sea bream during larval development especially utilized the neutral lipids 16:0, 18:1(n-9), and 22:6(n-3)
303 to maintain their growth and survival [64]. While DHA has an important role in stress resistance of mahimahi

304 *Coryphaena hippurus* [65], DHA content with larval stress resistance is not demonstrated in the present study (Table 1).
305 Free amino acids (FAA) is mainly used as metabolic fuel and for body protein synthesis, thus it is regarded as important
306 nutritional components influencing the viability of early stage marine fish [66]. In the wild, various copepods contain
307 more than twice of FAA per gram of wet mass than *Artemia* [6, 67, 68]. Thus, it is expected that FAA composition of
308 the cultured copepods and cladocerans affects the higher viability of fish larvae.

309 The growth of the *P. major* was not significantly influenced by the zooplanktons tested (Fig. 4). To date, studies
310 have shown that larvae fed with copepods or cladocerans achieved better growth than those fed with *Artemia*. This is
311 in the cases of sea bass *Lates calcarifer* [69], yellowtail clownfish *Amphiprion clarkia* [70], barber goby *Elacatinus*
312 *Figaro* [71], mangrove killifish *Kryptolebias marmoratus* [72] and kuruma prawn *Marsupenaeus japonicus* [18].
313 Pandey et al. [13] and Grageda et al. [72] detected that the larval growth was significantly related to feeding behavior of
314 the mangrove killifish larvae in terms of food size preference. Until 26 dph, copepod-fed *P. major* larvae showed low
315 morphometric growth compared to cladoceran and *Artemia*-fed, but on 28 dph the larvae grew comparatively fast
316 caused by the expected reason; the shifting food size preference to more than 0.8 mm (Fig. 5). Similar trends were
317 found with former studies which determined food size selectivity by larval gut analyses [73, 74]. The supply of fatty
318 acids is expected the other reason why the reared fish larvae showed no differences in larval wet weight (individual) and
319 growth (Table 1; Fig. 4, 6). The supplied n-3 HUFA which is consisted of EPA (20:5n-3), DHA (22:6n-3), has activities
320 in the fatty acid metabolism [75]. The requirement of these fatty acids was estimated as 0.5% of diet dry weight for
321 juvenile [76] and 0.4% of diet wet weight for larvae of red sea bream [77]. The total n-3 HUFA contained in *Artemia*,
322 *T. japonicus*, *D. celebensis* were calculated to be 0.03, 0.46, and 0.05% of total fatty acids and thus, the level of *T.*
323 *japonicus* only satisfies the minimum requirement of the targeted fish larvae (Table 2).

324 The present study showed the enhanced survival and viability of red sea bream *P. major* larvae fed on copepod *T.*
325 *japonicus* and cladoceran *D. celebensis* which were mass-cultured with CDE. We opined that these were due to the
326 optimum nutritional contents of these live foods and their appropriate size that stimulate appetite of the larvae. Thus,
327 we recommend the use of *T. japonicus* and *D. celebensis* as substitute of *Artemia* for intensive marine larviculture.

328

329

330 **ACKNOWLEDGMENTS**

331

332 This research was supported by a Ministry of Education, Culture, Sports, Science and Technology Grant-in-Aid for
333 Scientific Research (B) (2012-2014, No. 24380108) to A. H. The authors deeply appreciate Dr. Helen S. Marcial and
334 anonymous reviewers for improving the manuscript.

335 **REFERENCES**

336

- 337 1. Toledo JD (2001) Research on marine fishes. In L. M. B. Garcia (Ed.), Responsible aquaculture development in
338 Southeast Asia. Proceedings of the seminar-workshop on aquaculture development in Southeast Asia organized by
339 the SEAFDEC aquaculture department, 12-14 Oct. 1999, Iloilo city, Philippines, pp 173-184
- 340 2. Sui L-Y, Wille M, Su X-G, Cheng Y-X, Sorgeloos P (2008) Effect of feeding scheme and prey density on survival
341 and development of Chinese mitten crab *Eriocheir sinensis* zoea larvae. *Aquac Res* 39: 568-576
- 342 3. Léger P, Sorgeloos P(1992) Optimized feeding regimes in shrimp hatcheries. In: Fast, A.W., Lester, L. J. (Eds.),
343 Marine shrimp culture: Principles and practices. Elsevier, New York, pp 225-244
- 344 4. Lavens P, Sorgeloos P (2000) The history, present status and prospects of the availability of *Artemia* cysts for
345 aquaculture. *Aquaculture* 181: 397-403
- 346 5. Pagano M, Saint-Jean L, Arfi R, Bouvy M, Shep H (2000) Population growth capacities and regulatory factors in
347 monosepecific cultures of the cladocerans *Moina micrura* and *Diaphanosoma excisum* and the copepod
348 *Thermocyclops decipiens* from Côte d'Ivoire (West Africa). *Aquat Living Resour* 13: 163-172
- 349 6. Evjemo JO, Reitan KI, Olsen Y (2003) Copepods as live food organisms in the larval rearing of halibut larvae
350 (*Hippoglossus hippoglossus* L.) with special emphasis on the nutritional value. *Aquaculture* 227: 191-210
- 351 7. Marte CL (2003) Larviculture of marine species in Southeast Asia: current research and industry prospects.
352 *Aquaculture* 227: 293-304
- 353 8. Ajiboye OO, Yakubu AF, Adams TE, Olaji ED, Nwogu NA (2011) A review of the use of copepods in marine fish
354 larviculture. *Rev Fish Biol Fish* 21: 225-246
- 355 9. Kitajima C (1973) Experimental trials on mass culture of copepods. *Bull Plankton Soc Japan* 20: 54-60
- 356 10. Fukusho K (1991) Review of the research status of zooplankton production in Japan. Rotifer and microalgae culture
357 systems. Proceedings of a U. S. – Asia workshop. Honolulu, pp 55-60
- 358 11. Hagiwara A, Lee C-S, Shiraishi DJ (1995) Some reproductive characteristics of the broods of the harpacticoid
359 copepod *Tigriopus japonicus* cultured in different salinities. *Fish Sci* 61: 618-622
- 360 12. Park HG, Hur SB, Kim CW (1998) Culturing method and dietary value of benthic copepod, *Tigriopus japonicus*. *J*
361 *Aquac* 11: 261-269
- 362 13. Pandey BD, Hagiwara A, Sakakura Y (2008) Feeding behavior, feed selectivity and growth studies of mangrove
363 killifish, *Kryptolebias marmoratus*, larvae using various live and formulated feeds. *Environ Biol Fish* 82: 365-375
- 364 14. Støttrup JG (2006) A review on the status and progress in rearing copepods for marine larviculture. Advantages and
365 disadvantages. Among calanoid, harpacticoid and cyclopoid copepods. *Avances en Nutrición Acuicola VIII*. VIII

- 366 Symposium Internacional de Nutrición Acuícola, pp 62-83
- 367 15. Alam MJ, Ang KJ, Cheah SH (1993) Use of *Moina microura* (Kurz) as an *Artemia* substitute in the production of
- 368 *Macrobrachium rosenbergii* (de Man) post-larvae. *Aquaculture* 109: 337-349
- 369 16. Korovchinsky NM (1989) Redescription of *Diaphanosoma celebensis* Stingelin, 1900 (Crustacea, Cladocera).
- 370 *Hydrobiologia* 184: 7-22
- 371 17. Achuthankutty CT, Shrivastava Y, Mahambre GG, Goswami SC, Madhupratap M (2000) Parthenogenetic
- 372 reproduction of *Diaphanosoma celebensis* (Crustacea: Cladocera): influence of salinity on feeding, survival,
- 373 growth and neonate production. *Mar Biol* 137: 19-22
- 374 18. Nakamoto T, Maruyama I, Kimura H, Inada Y, Hagiwara A (2008) Two cladoceran species *Moina macrocopa* and
- 375 *Diaphanosoma celebensis*, as live feed for larval prawn, *Penaeus japonicus*. *Aquacult Sci* 56: 31-36
- 376 19. Park JC, Park HG (2010) Optimum salinity and temperature condition for mass culture of the brackish water flea,
- 377 *Diaphanosoma celebensis*. *Kor J Fish Aquat Sci* 43: 139-145 (in Korean with English abstract)
- 378 20. Khatton H, Banerjee S, Yusoff FM, Shariff M (2013) Use of microalgal-enriched *Diaphanosoma celebensis*
- 379 stingelin, 1900 for rearing *Litopenaeus vannamei* (Boone, 1931) postlarvae. *Aquacult Nutr* 19: 163-171
- 380 21. Knud-Hansen CF (1998) Pond fertilization: ecological approach and practical application. Pond
- 381 dynamics/aquaculture collaborative research support program. Oregon State University, Corvallis, OR.
- 382 22. Bwala RL, Omoregie E (2009) Organic enrichment of fish ponds: application of pig dung vs. tilapia yield. *Pak J*
- 383 *Nutr* 8: 1373-1379
- 384 23. Swift DR (1993) *Aquaculture training manual*, 2nd edn. Fishing News Books, Garden-Walk Farnham, Surrey.
- 385 24. Knud-Hansen CF, Batterson TR, McNabb CD, Harahat IS, Sumantadinata K, Eidman HM (1991) Nitrogen input,
- 386 primary productivity and fish yield in fertilized freshwater ponds in Indonesia. *Aquaculture* 94: 49-63
- 387 25. Adewumi AA, Adewumi IK, Olalege VF (2011) Livestock waste-menace: fish wealth-solution. *Afr J Environ Sci*
- 388 *Technol* 5: 149-154
- 389 26. Kang'ombe J, Brown JA, Halfyard LC (2006) Effect of using different types of organic animal manure on plankton
- 390 abundance and on growth and survival of *Tilapia rendalli* (Boulenger) in ponds. *Aquac Res* 37: 1360-1371.
- 391 27. Hagiwara A, Hamada K, Hori S, Hirayama K. (1994) Increase sexual reproduction in *Brachionus plicatilis* (Rotifera)
- 392 with the addition of bacteria and rotifer extracts. *J Exp Mar Biol Ecol* 181: 1-8
- 393 28. Kim H-J, Sakakura Y, Maruyama I, Nakamura T, Takiyama K, Fujiki H, Hagiwara A (2014) Feeding effect of
- 394 selenium enriched rotifers on larval growth and development in red sea bream *Pagrus major*. *Aquaculture* 432:
- 395 273-277
- 396 29. Naas K, Huse I (1996) Illumination in first feeding tanks for marine fish larvae. *Aquacult Eng* 15: 291-300

- 397 30. Naas KE, Níss T, Harboe T (1992) Enhanced first feeding of halibut larvae (*Hippoglossus hippoglossus* L.) in green
398 water. *Aquaculture* 105: 143-156
- 399 31. Shimma H, Tsujigado A (1981) Some biochemical quality of bred scorpaenoid fish, *Sebastes marmoratus*, and
400 activities of their larvae. *Bull Natl Res Inst Aquacult* 2: 11–20 (in Japanese with English abstract)
- 401 32. Shirota A (1970) Studies on the mouth size of fish larvae. *Nippon Suisan Gakkaishi* 36: 353-368
- 402 33. Lepage G, Roy CC (1986) Direct transesterification of all classes of lipids in a one-step reaction. *J Lipid Res* 27: 114-
403 120.
- 404 34. R Core Team (2012). R: A language and environment for statistical computing. R Foundation for
405 Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>
- 406 35. Watanabe T, Kitajima C, Fujita S (1983) Nutritional value of live organisms used in Japan for mass propagation of
407 fish: a review. *Aquaculture* 34: 115-143
- 408 36. Takeuchi T (2001) A review of feed development for early life stages of marine finfish in Japan. *Aquaculture* 200:
409 203-222
- 410 37. Damle DK, Chari MS (2011) Performance evaluation of different animal wastes on culture of *Daphnia* sp. *J Fish*
411 *Aquat Sci* 6: 57-61
- 412 38. Rapatsa MM, Moyo NAG (2013) Performance evaluation of chicken, cow and pig manure in the production of
413 natural fish food in aquadams stocked with *Oreochromis mossambicus*. *Phys Chem Earth* 66: 68-74
- 414 39. Hyder A, War M, Saquib N, Kareem A (2014) Utilization of poultry waste (chicken manure) for cost effective and
415 high density culture of two freshwater cyclopoid copepods *Thermocyclops decipiens* and *Mesocyclops aspericornis*.
416 *Res J Biotech* 9: 86-89
- 417 40. Hakk H, Millner P, Larsen G (2005) Decrease in water-soluble 17 β -estradiol and testosterone in composted poultry
418 manure with time. *J Environ Qual* 34: 943-950
- 419 41. Lorenzen A, Hendel JG, Conn KL, Bittman S, Kwabiah AB, Lazarovitz G, Massé D, McAllister TA, Topp E (2004)
420 Survey of hormone activities in municipal biosolids and animal manures. *Environ Toxicol* 19: 216-225
- 421 42. Marcial HS, Hagiwara A (2007) Effect of diazoin on life stages and resting egg hatchability of rotifer *Brachionus*
422 *plicatilis*. *Hydrobiologia* 593: 219-225
- 423 43. Marcial HS, Hagiwara A, Snell TW (2003) Estrogenic compounds affect development of harpacticoid copepod
424 *Tigriopus japonicus*. *Environ Toxicol Chem* 22: 3025-3030
- 425 44. Poulet SA (1983) Factors controlling utilization of non-algal diets by particle-grazing copepods. A review.
426 *Oceanological Acta* 6: 221-234
- 427 45. Moriarty DJW (1986) Bacterial productivity in ponds used for culture of penaeid prawns. *Microb Ecol* 12: 259-269

- 428 46. Geller W, Müller H (1981) The filtration apparatus of Cladocera: Filter mesh-sizes and their implications on food
429 selectivity. *Oecologia* 49: 316-321
- 430 47. Hart RC, Jarvis AC (1993) *In situ* determinations of bacterial selectivity and filtration rates by five cladoceran
431 zooplankters in a hypertrophic subtropical reservoir. *J Plakton Res* 15: 295-315.
- 432 48. Mushiake K, Fujimoto H, Shimma H (1993) A trial of evaluation of activity in yellow tail, *Seriola quinqueradiata*
433 larvae. *Suisanzoshoku* 41: 339-344 (in Japanese with English abstract)
- 434 49. Vassallo-Agius R, Imaizumi H, Watanabe T, Yamazaki T, Satoh S, Kiron V (2001) The influence of astaxanthin
435 supplemented dry pellets on spawning of striped jack. *Fish Sci* 67: 260-270
- 436 50. Wang H, Fang Q, Zheng L (2002) Effects of salinity on hatching rates and survival activity index of the larvae of
437 *Epinephelus akaara*. *Shuichan Xuebao* 26: 344-350
- 438 51. Fleeger JW (2005) The potential to mass-culture harpacticoid copepods for use as food for larval fish. In: Lee CS et
439 al. (eds) *Copepods in aquaculture*. Blackwell publishing, Australia, pp 11-24
- 440 52. Graeb BDS, Dettmers JM, WahldH, Cáceres CE (2004) Fish size and prey availability affect growth, survival, prey
441 selection, and foraging behavior of larval yellow perch. *T Am Fish Soc* 133: 504-514
- 442 53. Maruyama I., Nakao T., Shigeno I., Ando Y., Hirayama K (1997) Application of unicellular algae *Chlorella*
443 *vulgaris* for the mass-culture of marine rotifer *Brachionus*. *Hydrobiologia* 358: 133-138
- 444 54. McEvoy LA, Naess T, Bell JG, Lie Ø (1998) Lipid and fatty acid composition of normal and malpigmented Atlantic
445 halibut (*Hippoglossus hippoglossus*) fed enriched *Artemia*: a comparison with fry fed wild copepods. *Aquaculture*
446 163: 237-250
- 447 55. Takeuchi T, Shiina Y, Watanabe T (1992) Suitable levels of n-3 highly unsaturated fatty acids in diet for fingerlings
448 of red sea bream. *Nippon Suisan Gakkaishi* 58: 509-514
- 449 56. Shields RJ, Bell JG, Luiz FS, Gara B, Bromage NR, Sargent JR (1999) Natural copepods are superior to enriched
450 *Artemia* nauplii as feed for halibut larvae (*Hippoglossus hippoglossus*) in terms of survival, pigmentation and
451 retinal morphology: relation to dietary essential fatty acids. *J Nutr* 129: 1186-1194
- 452 57. Sargent JR, McEvoy LA, Bell JG (1997) Requirements, presentation and sources of polyunsaturated fatty acids in
453 marine fish larval feeds. *Aquaculture* 155: 117-127
- 454 58. Li Y-Y, Chen W-Z, Sun Z-W, Chen J-H, Wu K-G (2005) Effects of n-3 HUFA content in broodstock diet on
455 spawning performance and fatty acid composition of eggs and larvae in *Plectorhynchus cinctus*. *Aquaculture* 245:
456 263-272
- 457 59. Watanabe T, Ohta M, Kitajima C, Fujita S (1982) Improvement of dietary value of brine shrimp *Artemia salina* for
458 fish larvae by feeding them on ω 3 highly unsaturated fatty acids. *Nippon Suisan Gakkaishi* 48: 1775-1782

- 459 60. Sargent J, McEvoy L, Estevez A, Bell G, Bell M, Henderson J, Tocher D (1999) Lipid nutrition of marine fish
460 during early development: current status and future directions. *Aquaculture* 179: 217-229
- 461 61. Olivotto I, Cardinali M, Barbaresi L, Maradonna F, Carnevali O (2003) Coral reef fish breeding: the secrets of each
462 species. *Aquaculture* 224: 69-78
- 463 62. Olivotto I, Rollo A, Sulpizio R, Avella M, Tosti L, Carnevali O (2006) Breeding and rearing the sunrise dottyback
464 *Pseudochromis flavivertex*: the importance of live prey enrichment during larval development. *Aquaculture* 255:
465 480-487
- 466 63. Faulk CK, Holt GJ (2005) Advances in rearing cobia *Rachycentron canadum* larvae in recirculating aquaculture
467 systems: live prey enrichment and greenwater culture. *Aquaculture* 249: 231-243
- 468 64. Kimata M (1983) Changes of chemical composition during early development in the red sea bream *Chrysophrys*
469 *major* egg and larvae. *J Fac Mar Sci Technol Tokai Univ* 16: 213-223
- 470 65. Kraul S, Brittain K, Cantrell R, Nagao T, Ogasawara A, Ako H, Kitagawa H (2007) Nutritional factors affecting
471 stress resistance in the larval mahimahi *Coryphaena hippurus*. *J World Aquac Soc* 24: 186-193
- 472 66. Rønnestad I, Thorsen A, Finn RN (1999) Fish larval nutrition: a review of recent advances in the roles of amino
473 acids. *Aquaculture* 177: 201-216
- 474 67. Næss T, Germain-Henry M, Naas KE (1995) First feeding of Atlantic halibut (*Hippoglossus hippoglossus*) using
475 different combinations of *Artemia* and wild plankton. *Aquaculture* 130: 235-250
- 476 68. Helland S, Terjesen BF, Berg L (2003). Free amino acid and protein content in the planktonic copepod *Temora*
477 *longicornis* compared to *Artemia franciscana*. *Aquaculture* 215: 213-228
- 478 69. Ganzon-Naret ES, Fermin AC (1994) Effect of delayed feeding of *Artemia salina* and partial replacement by *Moina*
479 *macrocopa* on growth and survival of sea bass, *Lates calcarifer* (Bloch), larvae. *Isr J Aquac Bamidgeh* 46: 48-52
- 480 70. Olivotto I, Buttino I, Borroni M, Piccinetti CC, Malzone MG, Carnevali O (2008) The use of the Mediterranean
481 calanoid copepod *Centropages typicus* in Yellowtail clown fish (*Amphiprion clarkii*) larviculture. *Aquaculture* 284:
482 211-216
- 483 71. Côrtes G de F, Tsuzuki MY (2012) Effect of different live food on survival and growth of first feeding barber goby,
484 *Elacatinus figaro* (Sazima, Moura & Rosa 1997) larvae. *Aquac Res* 43: 831-834
- 485 72. Grageda MVC, Kotani T, Sakakura Y, Hagiwara A (2008) Effects of feeding copepod and *Artemia* on early growth
486 and behavior of the self-fertilizing fish, *Rivulus marmoratus*, under laboratory conditions. *Aquaculture* 281: 100-
487 105
- 488 73. Akazawa A, Sakakura Y, Hagiwara A (2008) Feeding selectivity of marine fish larvae, *Verasper variegatus*, *Seriola*
489 *quinqueradiata* and *Platycephalus* sp. on different sizes and shape of three rotifer strains. *Nippon Suisan Gakkaishi*

- 490 74: 380-388
- 491 74. Stenly W, Sakakura Y, Hagiwara A (2011) Application of the minute monogonont rotifer *Proales similis* de
492 Beauchamp in larval rearing of seven-band grouper *Epinephelus septemfasciatus*. *Aquaculture* 315: 355-360
- 493 75. Yone Y, Fujii M (1975) Studies on nutrition of red sea bream-XI Effect of ω 3 fatty acid supplement in a corn oil
494 diet on growth rate and feed efficiency. *Nippon Suisan Gakkaishi* 41: 73-77
- 495 76. Fujii M, Nakayama H, Yone Y (1976) Effect of ω 3 fatty acids on growth, feed efficiency and fatty acid composition
496 of red sea bream (*Chrysophrys major*). *Rep Fish Res Lab Kyushu Univ* 3: 65-86
- 497 77. Izquierdo MS, Watanabe T, Takeuchi T, Arakawa T, Kitajima C (1989) Requirement of larval red sea bream
498 *Pagrus major* for essential fatty acids. *Nippon Suisan Gakkaishi* 55: 859-867
- 499

500 **Table 1** Larval characteristics of red sea bream *Pagrus major* (28 dph) fed with three different diets

Diet	Survival (%)	Viability (%)	Wet weight (mg/ind.)
<i>A. franciscana</i>	16.7±9.8 ^b	44.4±12.3 ^B	8.2±0.3
<i>T. japonicus</i>	39.4±3.1 ^a	60.0±27.8 ^A	8.8±3.4
<i>D. celebensis</i>	20.8±3.8 ^b	60.0±32.2 ^A	9.4±0.1

501 Values are mean±SD of triplicate observations ($n=3$). Different alphabetical letters in a same column represent
 502 significant differences among three different diets ($a>b$, Tukey-Kramer *post hoc* test, $P<0.05$, $n=3$; $A>B$, Long-rank test,
 503 $P<0.05$, $n=3$).

504

505 **Table 2** Total fatty acids (mg/g WW) and fatty acid composition (%) of the three employed zooplanktons: *Artemia*
 506 *franciscana*, *Tigriopus japonicus*, and *Diaphanosoma celebensis* under the experimental conditions

	<i>A. franciscana</i>	<i>T. japonicus</i>	<i>D. celebensis</i>
Total	7.6	16.1	10.4
C14:0	0.6	0.6	2.0
C16:0	10.4	12.1	13.9
C16:1n-7	3.4	2.5	7.6
C18:0	5.6	3.3	4.4
C18:1n-9	17.6	8.3	5.7
C18:1n-7	8.6	1.8	3.5
C18:2n-6	5.5	16.2	22.6
C18:2n-4	0.0	0.1	0.1
C18:3n-6	0.4	0.2	0.4
C18:3n-4	0.1	0.1	0.1
C18:3n-3	25.0	3.9	6.6
C18:4n-3	3.3	0.2	0.4
C20:0	0.1	0.1	0.1
C20:1n-9	0.5	0.3	0.1
C20:2n-6	0.2	0.1	0.0
C20:3n-6	0.2	1.0	0.4
C20:4n-6	1.9	0.9	1.4
C20:3n-3	0.7	1.2	0.2
C20:4n-3	0.5	0.4	0.0
C20:5n-3 (EPA)	3.6	4.9	4.8
C22:1n-9	0.2	0.3	0.1
C21:5n-3	0.0	0.9	0.1
C22:5n-3	0.0	1.5	0.1
C22:6n-3 (DHA)	0.3	22.3	0.3
unknown	11.3	17.0	25.2
DHA/EPA	0.1	4.6	0.1
Σ n-3HUFA	3.9	28.7	5.2

507
508

509 **Table 3** Total fatty acids (mg/g WW) and fatty acid composition (%) of the fish larvae on 20 (initial) and 28 days post
 510 hatch (dph) with three different diets: *Artemia franciscana*, *Tigriopus japonicus*, and *Diaphanosoma celebensis*

	20 dph (Initial)	28 dph		
		<i>A. franciscana</i>	<i>T. japonicus</i>	<i>D. celebensis</i>
Total	13.8	13.3	13.0	14.7
C14:0	0.6	0.5	0.4	0.8
C16:0	14.5	15.2	15.1	15.4
C16:1n-7	1.4	1.7	1.6	2.6
C18:0	8.9	10.3	8.9	9.3
C18:1n-9	3.9	7.8	6.0	4.8
C18:1n-7	1.2	2.7	1.6	2.9
C18:2n-6	13.7	8.7	11.3	13.9
C18:2n-4	0.0	0.0	0.2	0.1
C18:3n-6	0.1	0.2	0.1	0.1
C18:3n-4	0.1	0.1	0.0	0.0
C18:3n-3	2.4	4.5	1.5	2.2
C18:4n-3	0.0	0.4	0.1	0.4
C20:0	0.2	0.2	0.2	0.2
C20:1n-9	1.5	0.8	0.5	0.5
C20:2n-6	0.2	0.2	0.2	0.2
C20:3n-6	2.4	1.3	1.3	1.2
C20:4n-6	0.9	1.8	1.2	1.0
C20:3n-3	0.9	0.6	0.7	0.4
C20:4n-3	0.5	0.4	0.2	0.2
C20:5n-3 (EPA)	5.4	4.5	3.9	7.4
C22:1n-9	0.2	0.2	0.2	0.2
C21:5n-3	0.5	0.2	0.4	0.2
C22:5n-3	6.5	5.4	3.4	3.9
C22:6n-3 (DHA)	13.0	13.2	21.8	10.8
unknown	20.9	19.0	19.25	21.5
DHA/EPA	2.4	2.9	5.6	1.5
Σ n-3HUFA	24.9	23.2	29.1	22.1

511

512 Figures

513

514 **Fig. 1** Experimental scheme for the larviculture of red sea bream *Pagrus major* with the three targeted zooplanktons:
515 *Artemia franciscana*(A), *Tigriopus japonicus*(T), *Diaphanosoma celebensis*(D) associated with the developmental stage
516 of fish larvae and rearing days (dph, days post hatch).

517

518 **Fig. 2** Population growths of three different developmental groups (a) nauplius,(b) copepodite,and (c)
519 nauplius+copepodite at different concentrations of chicken droppings extract (0, 1, and 2 ml/l) in *Tigriopus japonicus*.
520 Each plot and error bar indicates the mean and standard deviation of triplicate, respectively. Different alphabetical
521 letters represent significant differences (a>b>c>d>e>f>g>h>i>j>k>l, Tukey's HSD test, $P<0.05$, $n=3$).

522

523 **Fig. 3** Population growths of *Diaphanosoma celebensis*at different concentrations of chicken droppings extract (0, 1,
524 and 2 ml/l). Each plot and error bar indicates the mean and standard deviation of triplicate, respectively. Different
525 alphabetical letters represent significant differences (a>b>c>d>e>f, Tukey's HSD test, $P<0.05$, $n=3$).

526

527 **Fig. 4** Larval growth of red sea bream *Pagrus major* with the three live zooplanktonic diets: *Artemia franciscana*,
528 *Tigriopus japonicus*, and *Diaphanosoma celebensis* associated with rearing days. Each plot and error bar indicates the
529 mean and standard deviation of triplicates, respectively.

530


531 **Fig. 5** Size distribution of the three employed zooplanktons: *Artemia franciscana* (a), *Tigriopus japonicus* (b),
532 *Diaphanosoma celebensis* (c). The arrow indicates the estimated range of optimal food size [32]which can be eaten by
533 the targeted fish larvae of red sea bream *Pagrus major* on 20 days post hatch. A superscript letter on the mean size of
534 each zooplankton indicates significant difference (a>b>c, Tukey-Kramer *post-hoc* test, $P<0.0001$, $n=100$).

535

536


537 **Fig.6** Total biomass (i.e., production) of red sea bream *Pagrus major* larvae with the three live zooplanktonic diets:
538 *Artemia franciscana*, *Tigriopus japonicus*, and *Diaphanosoma celebensis*. Each column and error bar indicates the
539 mean and standard deviation of triplicates, respectively.

540



541
 542
 543
 544

Fig. 1.



545
 546
 547
 548

Fig. 2.






Fig. 3.

549
 550
 551
 552
 553



554
555
556
557

Fig. 4.



558

559


560

561

562

Fig. 5.

563



564

565

566

567

568

Fig. 6.