

Two New Maltol Glycosides and Cyanogenic Glycosides from *Elsholtzia rugulosa* Hemsl.

HAIZHOU LI · TATSUYA NAKASHIMA · TAKASHI TANAKA · YING-JUN ZHANG ·

CHONG-REN YANG · ISAO KOUNO

HAIZHOU LI · TATSUYA NAKASHIMA · TAKASHI TANAKA* · ISAO KOUNO

Graduate School of Biomedical Sciences,

Nagasaki University, Bunkyo-Machi 1-14, Nagasaki 852-8521, Japan

E-mail: t-tanaka@nagasaki-u.ac.jp

Tel.:+81-95-819-2433

Fax.:+81-95-819-2477

YING-JUN ZHANG · CHONG-REN YANG

Laboratory of Phytochemistry, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, P. R. China

Abstract Two new maltol glycosides, maltol 6'-*O*- β -D-apiofuranosyl- β -D-glucopyranoside and maltol 6'-*O*-(5-*O*-*p*-coumaroyl)- β -D-apiofuranosyl- β -D-glucopyranoside, were isolated from *Elsholtzia rugulosa* Hemsl. along with eleven known compounds including prunasin and amygdalin. The structures were determined on the basis of spectroscopic and chemical evidence. This is the second example of isolation of cyanogenic glycosides from Lamiaceous plants.

Keywords *Elsholtzia rugulosa* · Lamiaceae · maltol glycoside · cyanogenic glycoside

Introduction

Elsholtzia rugulosa Hemsl., which belongs to the Labiatae family, is distributed in the Yunnan, Sichuan and Guizhou provinces of China. It is one of the honey plants in this region, and has also been used as herbal tea or folk medicine for the treatment of colds, coughs, headaches, pharyngitis and fever [1]; however, the chemical constituents of this plant have not been clarified. This paper describes the isolation and structural determination of two new maltol glycosides together with eleven known compounds including two cyanogenic glycosides.

Results and discussion

An aqueous acetone extract of the air-dried plant was subjected to a combination of Diaion HP-20, Sephadex LH-20, silica gel and Chromatorex ODS column chromatography with various solvent systems to furnish 13 compounds. Known compounds maltol 3-*O*- β -D-glucoside [2], luteolin [3], luteolin 7-*O*- β -D-glucoside [4], luteolin 3'-*O*- β -D-glucuronide [5], quercetin 3-*O*- β -D-glucoside [6], tuberonic acid β -D-glucoside [7], benzyl alcohol β -D-glucoside [8], prunasin [9], amygdalin [10], caffeic acid [11], and rosmarinic acid [12] were identified based on comparison of their spectral data with those of authentic samples cited in the literature. This is only the second example of isolation of cyanogenic glycosides, prunasin and amygdalin, from Lamiaceous plants: the first case was from an Australian plant *Clerodendrum grayi* [13].

Compound **1** was obtained as a white amorphous powder and its molecular formula $C_{17}H_{24}O_{12}$ was determined by ^{13}C NMR spectral analysis, MALDI-TOF-MS (m/z 443, $[M+Na]^+$), and elemental analysis. The UV and IR spectral data of **1** showed the presence of an α , β -unsaturated carbonyl group (255 nm and 1650 cm^{-1}). The 1H and ^{13}C NMR spectra of **1** resembled those of maltol 3-*O*- β -D-glucoside and showed signals due to a maltol group [δ_H 8.00 (1H, d, $J=5.6$ Hz, H-6), 6.45 (1H, d, $J=5.6$ Hz, H-5), and 2.45 (3H, s, H-7); δ_C 164.6 (C-2), 143.3 (C-3), 177.0 (C-4), 117.2 (C-5), 157.1 (C-6), and 15.9 (C-7)]. However, two sugar anomeric proton signals appeared at δ_H 4.78 (d, $J=7.6$ Hz, H-1') and 4.95 (d, $J=2.2$ Hz, H-1''). The corresponding anomeric carbon resonances were observed at δ_C 105.0 and 110.7, respectively, in the ^{13}C NMR spectrum. Chemical shift comparison of the remaining carbon signals (Table 1) with those of known 6-*O*-apiosyl glucoside suggested the presence of glucose and apiose in **1** [14]. Appearance of a set of doublets with geminal coupling attributable to an isolated methylene (apiose H-4'') supported the presence of apiose. The relatively large J values (7.6 Hz) of the anomeric proton at δ_H 4.78 indicated a β -configuration for the glucopyranosyl moiety. The ^{13}C NMR shift of the anomeric carbon of the apiofuranosyl unit at δ_C 110.7 also indicated a β -configuration [14]. The location of the β -apiose moiety at the C-6' of β -glucose was apparent from the large low field shift of the C-6' (δ 68.8, $\Delta\delta$ 6.3) compared with that of maltol 3-*O*- β -D-glucoside (δ 62.5) [2, 14]. The absolute configurations of the glucose and apiose were determined to be D-glucose and D-apiose by reversed-phase HPLC after conversion of the sugars to thiocarbamoyl-thiazolidine derivatives [15].

Compound **2** was obtained as a white amorphous powder and the 1H and ^{13}C NMR spectra were closely related to those of **1**, except for appearance of additional 9 sp^2

carbon signals arising from a *p*-coumaroyl ester group [δ_{H} 7.59 (d, $J=16.0$ Hz, H-7'''), 6.30 (d, $J=16.0$ Hz, H-8'''), 7.41 (2H, d, $J=8.5$ Hz, H-2''', H-6'''), 6.76 (2H, d, $J=8.5$ Hz, H-3''', H-5'''); δ_{C} 127.0 (C-1'''), 131.1 (C-2''', C-6'''), 116.8 (C-3''', C-5'''), 161.2 (C-4'''), 146.8 (C-7'''), 114.7 (C-8'''), 168.7 (C-9''')] [11]. The formula $\text{C}_{26}\text{H}_{30}\text{O}_{14}$, which was determined by MALDI-TOF-MS (m/z 589, $[\text{M}+\text{Na}]^+$) and elemental analysis, is consistent with **2** being a *p*-coumaroyl ester of **1**. The ^{13}C NMR spectrum showed sugar carbon signals similar to those of **1** except for the low field shift of apiose C-5''' (δ_{C} 67.4, $\Delta\delta$ 1.9). In addition, a large low field shift of the apiose H-5''' signal at δ 4.17 (d, $J = 11.4$ Hz) and 4.21 (d, $J = 11.4$ Hz) compared to those of **1** [δ 3.54 (2H, s)] unequivocally indicated the acylation of this position. Acid hydrolysis of **2** yielded *p*-coumaric acid and maltol along with D-apiose and D-glucose. Thus the structure of **2** was concluded to be maltol 6'-*O*-(5-*O*-*p*-coumaroyl)- β -D-apiofuranosyl- β -D-glucopyranoside.

This study revealed the presence of two cyanogenic glycosides, prunasin and amygdalin, and the total isolation yield was 0.08 % from dry weight. Since Armeniacae semen (apricot kernel) which contains these compounds has been used for cough remedies in Europe and China, this result chemically supports the use of *E. rugulosa* for the treatment of colds and coughs in China. In addition, luteolin 3'-*O*- β -D-glucuronide [16-19] and rosmarinic acid [20-22] were found to be major phenolic constituents and these compounds are known to have anti-inflammatory activity.

Experimental

General

UV spectra were obtained with a Jasco V-560 UV/Vis spectrophotometer and optical rotations were measured with a Jasco DIP-370 digital polarimeter. ^1H and ^{13}C NMR spectra were measured in CD_3OD at 27 °C with a Varian Unity plus 500 spectrometer operating at 500 MHz for ^1H and 125 MHz for ^{13}C . Coupling constants are expressed in Hz and chemical shifts are given on a δ (ppm) scale. MS were recorded on a Voyager DE-PRO (Applied Biosystems) and JEOL JMS-700N spectrometer, 2,5-dihydroxybenzoic acid and glycerol were used as the matrix for MALDI-TOF-MS and FAB-MS measurements, respectively.

Plant material

The aerial parts of *E. rugulosa* Hemsl. were collected in Yunnan Province, China and identified by Professor Yang Chongren. A voucher specimen was deposited in the Kunming Institute of Botany, CAS, Kunming, China.

Extraction and isolation

The air-dried aerial parts of *Elsholtzia rugulosa* (3.0 kg) were extracted with 70% aqueous acetone (10 l) at room temperature three times. The acetone was removed by evaporation under reduced pressure (*ca.* 40 °C), and the resulting dark green precipitates consisting mainly of chlorophylls and waxes were removed by filtration. The filtrate

was concentrated to give an extract (170 g), which was directly subjected to Diaion HP-20 column chromatography (8 cm i.d. × 45 cm) with water containing increasing proportions of MeOH (20% step-wise elution from 0 – 100 %) and finally with 50 % aqueous acetone, and fractionated into 4 fractions. The first fraction (67.6 g) contained highly water soluble compounds and was mainly composed of sugars and the last fraction (19.6 g) contained non-polar substances including chlorophylls. Fraction 3 (21.3 g) was separated into two fractions by Diaion HP-20 (0 – 100 % MeOH in H₂O, 10% step-wise elution) and the first fraction was combined with fraction 2 because these fractions contained similar compounds. Fraction 3-2 was subjected to Sephadex LH-20 column chromatography with 60 % MeOH to yield luteolin (76.3 mg). Fraction 2, after mixing with fraction 3-1, (35.0 g) was chromatographed over Diaion HP-20 (0 – 100 % MeOH in H₂O, 10% step-wise elution) to yield 4 fractions: 2-1 (2.0 g), 2-2 (17 g), 2-3 (3.7 g) and 2-4 (10.0 g). Fraction 2-1 was crystallized from water to give luteolin 3'-*O*-β-D-glucuronide (1.72 g). Fraction 2-2 was separated by a combination of column chromatography using Sephadex LH-20 (40 – 80 % MeOH), Chromatorex ODS (0 – 80 % MeOH) and silica gel 60 (CHCl₃-MeOH-H₂O, 80 : 20 : 2 – 70 : 30 : 5, v/v/v) to furnish **1** (56.4 mg), **2** (44.2 mg), maltol 3-*O*-β-D-glucoside (52.2 mg), tuberonic acid β-D-glucoside (35.2 mg), benzyl alcohol β-D-glucoside (512.1 mg), prunasin (2.18 g), amygdalin (178.0 mg) and caffeic acid (60.4 mg). Fraction 2-3 was successively separated by Sephadex LH-20 (40 – 60 % MeOH), Chromatorex ODS (0 – 80 % MeOH) and then silica gel 60 (CHCl₃-MeOH-H₂O, 70 : 30 : 5) to yield luteolin 7-*O*-β-D-glucoside (50.0 mg) and quercetin 3-*O*-β-D-glucoside (84.0 mg). Chromatography of fraction 2-4 over Sephadex LH-20 (60 – 80 % MeOH),

Chromatorex ODS (20 – 60 % MeOH) and Sephadex LH-20 (EtOH) afforded rosmarinic acid (874.5 mg) and caffeic acid (67.5 mg).

Maltol 6'-O- β -D-apiofuranosyl- β -D-glucopyranoside (1)

White amorphous powder. $[\alpha]_D^{25}$ -6.2° (*c* 0.7, MeOH). UV $\lambda_{\max}^{\text{EtOH}}$ ($\log \epsilon$): 255 (3.97).

IR ν_{\max} cm^{-1} : 3376, 1650. MALDI-TOF-MS *m/z*: 443 $[\text{M}+\text{Na}]^+$. *Anal. Calcd* for

$\text{C}_{17}\text{H}_{24}\text{O}_{12} \cdot 3\text{H}_2\text{O}$: C, 43.04; H, 6.37. Found: C, 43.01; H, 6.41. ^1H and ^{13}C NMR see

Table 1.

Maltol 6'-O-(5-O-*p*-coumaroyl)- β -D-apiofuranosyl- β -D-glucopyranoside (2)

White amorphous powder. $[\alpha]_D^{25}$ -81.5° (*c* 0.6, MeOH). UV $\lambda_{\max}^{\text{EtOH}}$ ($\log \epsilon$): 258 (4.09),

313 (4.35). IR ν_{\max} cm^{-1} : 3402, 1703, 1647, 1613. MALDI-TOF-MS *m/z*: 589 $[\text{M}+\text{Na}]^+$.

Anal. Calcd for $\text{C}_{26}\text{H}_{30}\text{O}_{14} \cdot 4\text{H}_2\text{O}$: C, 48.90; H, 6.00. Found: C, 48.58; H, 6.12. ^1H and

^{13}C NMR see Table 1.

Acid hydrolysis of 2

A solution of **2** (3 mg) in 1M HCl (0.5 ml) was heated at 90 – 100 °C in a screw-capped vial for 5 h. The mixture was partitioned with EtOAc (0.5 ml) and the EtOAc layer was analyzed by HPLC [Cosmosil 5C₁₈ AR II (250 × 4.6 mm i.d.); gradient elutions of CH₃CN in 50 mM H₃PO₄ from 4-30% in 39 min and 30-75% in 15 min; flow rate, 0.8 mL/min; detection with a Jasco MD-910 photodiode array detector].

The t_R and UV absorptions of the two peaks at 14.61 min and 29.55 min was coincided with those of maltol (UV λ_{\max} : 223 nm) and *p*-coumaric acid (UV λ_{\max} : 310 nm), respectively.

Determination of aldose configuration

A solution of **1** (0.8 mg) in 1M HCl (0.2 ml) was heated at 90 – 100 °C in a screw-capped vial for 5 h. The mixture was neutralized by addition of Amberlite IRA400 (OH⁻ form) and filtered. The filtrate was dried *in vacuo*, dissolved in 0.2 ml of pyridine containing L-cysteine methyl ester (10 mg/ml) and reacted at 60 °C for 1 h. To this mixture a solution (0.2 ml) of *o*-torylisothiocyanate in pyridine (10 mg/ml) was added and it was heated at 60 °C for 1 h. The final mixture was directly analyzed by HPLC [Cosmosil 5C₁₈ AR II (250 × 4.6 mm i.d., Nacalai Tesque Inc.); 25% CH₃CN in 50 mM H₃PO₄; flow rate, 0.8 mL/min; detection, 250 nm]. The t_R of the two peaks at 17.48 and 28.57 min coincided with those of D-glucose and D-apiose, respectively. The standard samples of D- and L-apiose were synthesized from L- and D-ribose, respectively [23]. The aqueous layer remaining after partition with EtOAc on the acid hydrolysis of **2** was treated in the manner similar to that described for **1** and the HPLC analysis showed the presence of D-glucose and D-apiose.

Acknowledgements

The authors are grateful to Mr. K. Inada and Mr. N. Yamaguchi (Nagasaki University) for NMR and MS measurements. This work was supported by a Grant-in-aid for Scientific Research No. 18510189 from the Japan Society for the Promotion of Science.

References

1. Jiangshu New College of Medicine (1985) In: The Dictionary of Chinese Medicine, Shanghai, Shanghai Press of Science and Technology, p 2132
2. Tanaka N, Sakai H, Murakami T, Saiki Y, Chen CM, Iitaka Y (1986) Chemical and chemotaxonomic studies on pterophyten. LXII. Chemical studies on the substance from *Arachniodes maximowiczii* Ohwi. Chem Pharm Bull 34:1015-1022
3. Ulubelen A, Doguc T (1974) Flavonoid compounds from the flowers of *Genista lydia*. Planta Med 25:39-42
4. Kikuchi M, Yamauchi Y (1985) Studies on the constituents of *Osmanthus* species. IV. On the components of the leaves of *Osmanthus fortunei* Carr. Yakugaku Zasshi 105:542-546
5. Ulubelen A, Miski M, Neuman P, Mabry TJ (1979) Flavonoids of *Salvia tomentosa* (Labiatae). J Nat Prod 42:261-263
6. Markham KR, Ternai B, Stanley R, Geiger H, Mabry TJ (1978) Carbon-13 NMR studies of flavonoids—III : Naturally occurring flavonoid glycosides and their acylated derivatives. Tetrahedron 34:1389-1397
7. Ueda M, Okazaki M, Ueda K, Yamamura S (2000) Leaf-Closing Substance of *Albizia julibrissin* Durazz. Tetrahedron 56:8101-8105
8. Miyase T, Ueno A, Takizawa N (1987) Studies on the glycosides of *Epimedium grandiflorum* Morr. var. *thunbergianum* (Miq.) Nakai. I. Chem Pharm Bull 35:1109-1117

9. Ling SK, Tanaka T, Kouno I (2002) New cyanogenic and alkyl glycoside constituents from *Phyllagathis rotundifolia*. *J Nat Prod* 65:131-135
10. Tanaka T, Zhang H, Jiang ZH, Kouno I (1997) Relationship between hydrophobicity and structure of hydrolysable tannins, and association of tannins with crude drug constituents in aqueous solution. *Chem Pharm Bull* 45:1891-1897
11. Jiang ZH, Hirose Y, Iwata H, Sakamoto S, Tanaka T, Kouno I (2001) Caffeoyl, coumaroyl, galloyl, and hexahydroxydiphenoyl glucoses from *Balanophora japonica*. *Chem Pharm Bull* 49:887-892
12. Tanaka T, Morimoto S, Nonaka G, Nishioka I, Yokozawa T, Chung HY, Oura H (1989) Magnesium and ammonium-potassium lithospermates B, the active principles having a uremia-preventive effect from *Salvia miltiorrhiza*. *Chem Pharm Bull* 37:340-344
13. Miller RE, McConville MJ, Woodrow IE (2006) Cyanogenic glycosides from the rare Australian endemic rainforest tree *Clerodendrum grayi* (xLamiaceae). *Phytochemistry* 67:43-51
14. Jiang ZH, Tanaka T, Kouno I (1999) New phenylpropanoid glycosides from the fruits of *Illicium anisatum*. *Chem Pharm Bull* 47:421-422
15. Tanaka T, Nakashima T, Ueda T, Tomii T, Kouno I (2007) *Chem Pharm Bull*, 55:899-901.
16. Gutierrez-Venegas G, Kawasaki-Cardenas P, Rita Arroyo-Cruz S, Maldonado-Frias S (2006) Luteolin inhibits lipopolysaccharide actions on human gingival fibroblasts. *Eur J Pharmacol* 541:95-105
17. Kim JS, Lee HJ, Lee MH, Kim J, Jin C, Ryu JH (2006) Luteolin inhibits LPS-stimulated inducible nitric oxide synthase expression in BV-2 microglial cells. *Planta Med* 72:65—68
18. Odontuya G, Hoult JRS, Houghton PJ (2005) Structure-activity relationship for antiinflammatory effect of luteolin and its derived glycosides. *Phytother Res* 19: 782-786
19. Hougee S, Sanders A, Faber J, Graus YMF, van den Berg WB, Garssen J, Smit HF, Hoijer MA (2005) Decreased pro-inflammatory cytokine production by LPS-stimulated PBMC upon in vitro incubation with the

- flavonoids apigenin, luteolin or chrysin, due to selective elimination of monocytes/macrophages. *Biochem Pharmacol* 69:241-248
20. Englberger W, Hadding U, Etschenberg E, Graf E, Leyck S, Winkelmann J, Parnham MJ (1988) Rosmarinic acid: a new inhibitor of complement C3-convertase with anti-inflammatory activity. *Int J Immunopharmac* 10:729-737
 21. Osakabe N, Yasuda A, Natsume M, Yoshikawa T (2004) Rosmarinic acid inhibits epidermal inflammatory responses: anticarcinogenic effect of *Perilla frutescens* extract in the murine two-stage skin model. *Carcinogenesis* 25:549—557
 22. Osakabe N, Takano H, Sanbongi C, Yasuda A, Yanagisawa R, Inoue K, Yoshikawa T (2004) Anti-inflammatory and anti-allergic effect of rosmarinic acid (RA); inhibition of seasonal allergic rhinoconjunctivitis (SAR) and its mechanism. *BioFactors* 21:127—131
 23. Yun M, Moon HR, Kim HO, Choi WJ, Kim Y, Park CS, Jeong, LS (2005) A highly efficient synthesis of unnatural L-sugars from D-ribose. *Tetrahedron Lett* 46:5903-5905

Figure Legend

Figure 1 Structures of compounds **1** and **2**




Figure 1

Table 1. ^1H - and ^{13}C -NMR data for compounds **1** and **2** (^1H ; 500 MHz, ^{13}C ; 125 MHz, CD_3OD)

	1		2	
	^{13}C	^1H	^{13}C	^1H
maltol				
2	164.6		164.4	
3	143.3		143.3	
4	177.0		176.9	
5	117.2	6.45 (d, 5.6)	117.2	6.38 (d, 5.6)
6	157.1	8.00 (d, 5.6)	157.0	7.89 (d, 5.6)
7	15.9	2.45 (s)	15.9	2.39 (s)
D-glucose				
1'	105.0	4.78 (d, 7.6)	105.2	4.72 (d, 7.6)
2'	74.9	3.27-3.45 (m)	74.9	3.25-3.40 (m)
3'	77.9	3.27-3.45 (m)	78.4	3.25-3.40 (m)
4'	71.3	3.27-3.45 (m)	71.3	3.25-3.40 (m)
5'	77.4	3.27-3.45 (m)	77.4	3.25-3.40 (m)
6'	68.5	3.94 (dd, 2.0, 11.2) 3.60 (dd, 6.1, 11.2)	68.5	3.93 (d, 11.2) 3.57 (dd, 6.0, 11.2)
D-ribose				
1''	110.7	4.95 (d, 2.2)	110.4	4.94 (d, 2.2)
2''	77.8	3.84 (d, 2.2)	77.9	3.92 (d, 2.2)
3''	80.4		78.9	
4''	75.3	3.89 (d, 9.5) 3.73 (d, 9.5)	75.3	3.90 (d, 9.8) 3.78 (d, 9.8)
5''	65.5	3.54 (2H, s)	67.4	4.21 (d, 11.4) 4.17 (d, 11.4)
<i>p</i> -coumaroyl group				
1'''			127.0	
2''', 6'''			131.1	7.41 (d, 8.5)
3''', 5'''			116.8	6.76 (d, 8.5)
4'''			161.2	
7'''			146.8	7.59 (d, 16.0)
8'''			114.7	6.30 (d, 16.0)
9'''			168.7	

 Coupling constants (J in Hz) are given in parentheses