

Protection by Polaprezinc Against Radiation-Induced Apoptosis in Rat Jejunal Crypt Cells

Mutsumi MATSUU-MATSUYAMA^{1*}, Kazuko SHICHIJO¹, Kumio OKAICHI²,
Toshiyuki NAKAYAMA¹, Masahiro NAKASHIMA³, Takashi UEMURA⁴,
Daisuke NIINO⁵ and Ichiro SEKINE¹

Radiation-induced apoptosis/Polaprezinc/p53/Rats/Intestine.

Polaprezinc, an anti-ulcer drug, is a chelate compound consisting of zinc and L-carnosine. Polaprezinc has been shown to prevent gastric mucosal injury. The anti ulcer effects of polaprezinc have been ascribed to its antioxidative property. The effect of polaprezinc on ionizing radiation-induced apoptosis was studied in the jejunal epithelial crypt cells of rats. Seven-to eight week-old Wistar rats, which were treated with 100 mg/kg of polaprezinc orally 1h before irradiation or 2% carboxymethyl cellulose sodium in controls, were exposed to whole body X-ray irradiation at 2 Gy. The number of apoptotic cells per jejunum crypt was counted in haematoxylin and eosin stained sections at 0–6 h after irradiation. TUNEL positive cells and immunopositive cells for active caspase-3 per crypt were also counted. Accumulation of p53, p21^{WAF1/CIP1} and Bax expression in the jejunum after irradiation were examined by Western blot analyses. Polaprezinc treatment given prior to radiation resulted in a significant reduction in numbers of apoptotic cells, TUNEL positive cells and active caspase-3 immunopositive cells in jejunal crypt cells. Polaprezinc treatment resulted in decreases of p53 accumulation, p21^{WAF1/CIP1} and Bax expression after irradiation. Polaprezinc has a protective effect against ionizing radiation induced apoptosis in rat jejunal crypt cells.

INTRODUCTION

Polaprezinc [*N*-(3-aminopropionyl)-L-histidinato zinc], a chelate compound developed as an anti-cancer drug consists of zinc ion and L-carnosine, has been commonly used in the treatment of gastric ulcer.¹⁾ Polaprezinc has been effective in preventing gastric lesions in a wide of variety of experimental models without affecting gastric acid secretion and endogenous prostaglandins.^{2,3)} The protection of the gastric mucosa by polaprezinc may be attributed to its antioxidant activity,⁴⁾ membrane-stabilizing action⁵⁾ and stimulation of mucus production.⁶⁾

Exposure of the intestine to ionizing radiation results in

the rapid, apoptotic death of the stem cells.⁷⁾ Crypt disturbance results from the destruction of the stem cells, which are responsible for repopulating the lining of the intestine. At suitable high doses, damage to the crypt is accompanied by functional changes, such as malabsorption, which is expressed clinically as acute bowel reactions.^{8,9)} Apoptosis is a programmed process of active cell death that involves gross morphological alterations including condensation of nuclear chromatin, compaction of cytoplasmic organelles, membrane blebbing, and cellular fragmentation into apoptotic bodies.¹⁰⁾ Identification and counting of apoptotic bodies in haematoxylin and eosin (H&E)-stained sections has been used extensively to study apoptosis induced by ionizing radiation and other genotoxic stimuli in the intestine.^{11,12)}

Previously, we reported that sucralfate and kefir have a protective effect on radiation-induced apoptosis, presumably in protecting the stem cell region of the rat colonic crypt cells from radiation.^{13,14)} Sucralfate, an aluminum hydroxide complex of sulfated sucrose, is an anti-ulcer agent with a protective effect of radiation-induced apoptosis in the rat colon following decreases of p53 accumulation, p21^{WAF1/CIP1} expression and Bax/Bcl-2 ratio.¹⁴⁾ The p53 tumour suppressor gene is one of the primary cellular factors which determine the nature of growth arrest and/or cell death in response to ionizing radiation.^{15–18)} An increase in the level of the p53

*Corresponding author: Phone: +81-95-819-7107,
Fax: +81-95-819-7108,
E-mail: mutsumi@net.nagasaki-u.ac.jp

¹⁾Division of Tumor and Diagnostic Pathology; ²⁾Department of Radiation Biophysics; ³⁾Tissue and Histopathology Section, Atomic Bomb Disease Institute, Nagasaki University Graduate School of Biomedical Sciences, 1-12-4 Sakamoto, Nagasaki, 852-8523, Japan; ⁴⁾Clinical Research Center, National Hospital Organization (NHO) Nagasaki Medical Center, Nagasaki, Kubara 2-1001-1, Omura, Nagasaki 856-8562, Japan; ⁵⁾Department of Pathology, School of Medicine, Kurume University, Asahi-machi 67, Kurume, 830-0011, Japan.
doi:10.1269/jrr.07114

protein in irradiated cells and the consequent up-regulation of p21^{WAF1/CIP1}, the cyclin dependent kinase inhibitory protein, appear to be critical components of G1 arrest.^{19,20} Radiation-induced intestinal apoptosis in mice has also been shown to be regulated by several members of the bcl-2 family of proteins, including bax.²¹

The aim of this study was to evaluate whether polaprezinc was capable of preventing acute X-ray radiation-induced apoptosis and to examine the mechanism by which polaprezinc exerts its effect on radiation-induced apoptosis in the jejunal crypt cells of rats. Routine histological study with haematoxylin and eosin staining (H&E) confirmed the presence of apoptosis. Identification of apoptosis was confirmed using a TUNEL technique. Active caspase-3, which is a key effector of the apoptotic pathway, was then studied in the jejunal crypt cells by immunohistochemistry. The effects of polaprezinc on the expression of p53, p21^{WAF1/CIP1} and Bax after irradiation were also examined.

MATERIALS AND METHODS

Animals and treatment of polaprezinc

Seven-to eight week-old male Wistar rats (230–310 g) were purchased from Charles River Japan (Atsugi, Japan). A total 60 of rats were used for examination. The rats were housed in groups of 2 to 3 per cage in an air-conditioned room at 24°C (lights on from 7 a.m. to 9 p.m.), and allowed free access to food (laboratory chow F2, Japan CLEA, Tokyo) and tap water at the Laboratory Animal Center of Nagasaki University. All animals were kept in a specific pathogen-free facility at the Animal Center in accordance with the rules and regulations of the Institutional Animal Care and Use Committee.

Rats were given of 100 mg/kg polaprezinc (Zeria Pharmaceutical Co. Ltd., Tokyo) dissolved in 2.5 ml carboxymethyl cellulose sodium (CMC) orally 1 h before irradiation. Control rats were given only 2.5 ml 2% CMC solution.

Irradiation

Irradiation was performed between 9:00 a.m. and 12:00 p.m. Rats received whole-body X-ray irradiation using a Toshiba EXS-300 X-ray, 200 KV, 15mA apparatus with 0.5 mm Al filter at a dose-rate of 0.864 Gy/min.

Histology and assessing apoptosis

Rats in each group were sacrificed by deep anesthesia at 1 h, 2 h, 4 h and 6 h after 2 Gy irradiation. A 4 cm segment of the jejunum was resected at 2 cm from the ligament of Trietz. The colon segment was also resected. Tissue samples were fixed overnight in 4% neutral-buffered formalin, and embedded in paraffin blocks from which 3- μ m sections were cut and stained with H&E. Identification of apoptosis was confirmed using a TUNEL technique (Apoptag Peroxidase *In Situ* Apoptosis Detection Kit, Chemicon, Temecula, CA).

This method detects apoptosis-associated DNA fragmentation by labeling of 3'-OH termini with digoxigenin nucleotides using terminal deoxynucleotidyl transferase.²² Active caspase-3 was detected by anti-active caspase-3 polyclonal antibody (R&D Systems, Abingdon, UK) staining. This antibody specifically recognizes amino acids 163–175 of caspase-3, but does not detect the precursor form. Immunohistochemical staining for active caspase-3 was performed with a DAKO LSAB-2 Kit, and peroxidase for use on rat specimens (DAKO Cytomation, Carpinteria, CA) according to the manufacturer's instructions.

Fifty crypts per group from complete jejunal crypts that had been cut in the longitudinal plane were selected for analysis. The incidence of cell death (apoptotic index) in the jejunum and the colon was quantified by counting the number of dead cells in each crypt in H&E-stained sections at $\times 400$ magnification by light microscopy as described previously.¹⁴ The distinctive morphological features of apoptosis, as described by Kerr *et al.*²³ and Walker *et al.*,²⁴ were used to recognize apoptotic cells, and any doubtful cells were disregarded. Small dead cell fragments occurring in tightly knit groups were assessed as originating from one cell, and given a single count.²⁵ TUNEL positive cells and immunopositive cells for active caspase-3 in at least thirty jejunal crypts per animal, and the average value was calculated.

Western blotting

The jejunum tissue of control and polaprezinc-treated rats were removed at 1 h, 2 h, 4 h and 6 h after 2 Gy irradiation and frozen immediately. The tissues were then suspended in RIPA buffer (50 mM Tris, 150 mM NaCl, 1% sodium deoxycholate and 0.05% SDS, pH7.4), broken into pieces on ice and subjected to three freeze-thaw cycles. The insoluble cell debris was removed by centrifugation. Supernatants were collected and the protein concentration was determined using a protein assay reagent (Bio-Rad Laboratories, Hercules, CA). Protein samples (30 μ g) were subjected to 10% or 15% SDS-polyacrylamide gel electrophoresis, then transferred electrophoretically to Hybond ECL Nitrocellulose Membranes (Amersham, Arlington heights, IL). Membranes were incubated with mouse monoclonal anti-p53 (Pab 421) antibody (Oncogene Science Inc., Uniondale, NY), rabbit polyclonal anti-p21^{WAF1/CIP1} antibody (Santa Cruz Biotechnology, Santa Cruz, CA), Bax (BD pharmingen, San Diego, CA) or rabbit polyclonal anti-actin antibody (Sigma, St. Louis, MO). This was followed with a horseradish peroxidase-conjugated anti-mouse IgG antibody (Zymed Labs. Inc., San Francisco, CA) or a horseradish peroxidase-conjugated anti-rabbit IgG (Amersham). Chemiluminescence (ECL Plus Amersham) was used for analyzing levels of protein according to the manufacturer's protocol. Blots were exposed to Hyperfilm ECL (Amersham). NIH image 1.61 software was used in measuring the densities of each of the protein bands. The level of protein after irradiation

was determined using the level of actin as a standard.

Statistical evaluation of data

All values were expressed as the mean \pm SEM of results obtained from three to five animals per data point. Differences between groups were examined for statistical significance using the Student's *t*-test. A $P < 0.05$ was considered to be statistically significance.

RESULTS

Apoptotic index of control rats and polaprezinc-treated rats after irradiation

Figure 1 shows histologic sections stained with H&E of jejunal crypts from control and polaprezinc-treated rats (Fig. 1A, B), TUNEL (Fig. 1C, D) and active caspase-3 antibody (Fig. 1E, F). At 2 h after 2 Gy irradiation, a large number of apoptotic cells were observed in the base of the jejunal crypts of control rats in H&E-stained sections and with TUNEL staining (Fig. 1A, C). In contrast, there was a sharp decrease in the number of apoptotic cells in the jejunal crypts of polaprezinc-treated rats (Fig. 1B, D). Active caspase-3 immunoreactivity was largely confined to apoptotic bodies towards the base of the intestinal crypts in the area where apoptotic bodies are most often seen in H&E-stained sections (Fig. 1E). Active caspase-3 immunoreactivity was also greatly decreased in polaprezinc-treated rats (Fig. 1F) when compared with control rats.

Figure 2 compares the time course of the apoptotic index detected using H&E staining, TUNEL staining and active caspase-3 expression in the jejunum and the colon in control and polaprezinc-treated rats up to 6 h after irradiation with 2 Gy. The apoptotic index of the jejunum in control rats increased by 2 h after irradiation, then decrease gradually at 4 h and 6 h. Polaprezinc treatment significantly reduced the number of apoptotic jejunal crypt cells at 2 h, 4 h and 6 h after 2 Gy irradiation to 13% ($p < 0.001$), 44% ($p < 0.05$) and 41% ($p < 0.01$) of the control rat values, respectively. There was no significant difference in the background (0 h) levels of apoptosis in both groups (Fig. 2A). Figure 2B shows the pattern of TUNEL positive cells following irradiation of the jejunum in control and polaprezinc treated rats. The number of TUNEL positive cells in control rats increased at 2 h and 4 h, and then decreased slightly at 6 h after 2 Gy irradiation. The number of TUNEL positive cells in control rats was not higher than the apoptotic index at each sampling time after irradiation. Polaprezinc treatment significantly reduced the number of TUNEL positive cells at 2 h, 4 h and 6 h after irradiation to 17% ($p < 0.001$), 49% ($p < 0.05$) and 47% ($p < 0.05$) of the control rat values, respectively. Figure 2C illustrates the pattern of caspase-3 activation following irradiation of the jejunum in control and polaprezinc treated rats. The number of active caspase-3 immunopositive cells in control rats peaked at 2 h, and then




Fig. 1. Histologic sections of the jejunum in control rats (A, C and E) and polaprezinc-treated rats (B, D and F) at 2 h after 2 Gy irradiation, stained by the H&E (A, B), the TUNEL method (C, D) to show the cells containing DNA fragments and active caspase-3 (E, F) ($\times 400$).

decreased sharply at 4 h after irradiation. Treating with polaprezinc resulted in a significant decrease of active caspase-3 positive cells at 2 h and 6 h after irradiation to 17% ($p < 0.001$) and 33% ($p < 0.001$) of the control rat values, respectively. Figure 2D shows the apoptotic index following irradiation of the colon in control and polaprezinc treated rats.




Fig. 2. Apoptotic Index (A), TUNEL (B) and active caspase-3 (C) positive cells per jejunal crypt cells, and apoptotic index of colonic crypt cells (D) in control rats (□) and polaprezinc-treated rats (●) at 0 to 6 h after 2 Gy irradiation. Data are the mean ± SEM values of 3–5 rats per data point. **p* < 0.05, ***p* < 0.01 and ****p* < 0.001 vs. control rats.




Fig. 3. Western blotting analysis and kinetics of p53 (A), p21^{WAF1/Cip1} (B) and Bax (C) expression in the jejunum of control and polaprezinc-treated rats. The amounts of p53, p21^{WAF1/Cip1} and Bax were quantified by densitometric analysis. Each of the protein levels is expressed as ratios to non-irradiated rats (0 h) in control (□) or polaprezinc-treated rats (●). Data represent mean ± SEM values of three to four separate experiments from two rats. **p* < 0.05 and ***p* < 0.01 vs. control rats.

The apoptotic index of the colon in both groups increased by 4 h and then decreased. There was no significant difference between control rats and polaprezinc treated rats.

Western blotting for p53, p21^{WAF1/CIP1} and Bax

In an effort to determine how polaprezinc treatment was interfering with the pathways leading to radiation-induced apoptosis, the expressions of p53, p21^{WAF1/CIP1} and Bax after 2 Gy irradiation were examined by Western blot analyses and kinetic diagrams are shown in Fig. 3. p53 accumulation in control rats after irradiation increased 1.2-fold, 2.8-fold, 2.9-fold and 2.6-fold at 1 h, 2 h, 4 h and 6 h respectively over non-irradiated control rats. p53 accumulation of polaprezinc treated rats after irradiation increased 1.2, 1.5 and 1.4-fold at 1 h, 2 h and 4 h respectively, and then decreased to 0.9-fold at 6 h over non-irradiated polaprezinc treated rats. There was a significant difference at 6 h ($p < 0.05$) between control and polaprezinc-treated rats (Fig. 3A). The expression of p21^{WAF-1/CIP1} in control rats after irradiation increased 1.3, 2.4, 3.0 and 3.4-fold at 1 h, 2 h, 4 h and 6 h respectively over non-irradiated control rats, while that in polaprezinc treated rats increased 1.4, 1.8 and 2.3-fold at 1 h, 2 h and 4 h respectively, and then decreased to 1.3-fold at 6 h over non-irradiated polaprezinc treated rats. There was a significant difference at 6 h ($p < 0.05$) between control and polaprezinc-treated rats (Fig. 3B). The expression of Bax in control rats after irradiation increased 1.5, 2.5, 1.9 and 3.7-fold at 1 h, 2 h, 4 h and 6 h respectively over non-irradiated control rats, while that in polaprezinc treated rats increased 1.5 and 1.3-fold at 1 h and 4 h respectively, and decreased to 0.9 and 1.0-fold at 2 h and 6h over non-irradiated polaprezinc treated rats. There was a significant difference at 2 h ($p < 0.05$), 4 h ($p < 0.01$) and 6 h ($p < 0.05$) between control and polaprezinc-treated rats (Fig. 3C).

DISCUSSION

Intestinal stem cells in the crypts are rapidly proliferating cells *in vivo* which are most sensitive to radiation-induced damage. Apoptosis is the main cause of irradiation-induced intestinal damage responsible for many of the side effects of aggressive cancer therapy.^{26,27} Nonalimentary pelvic tumors are usually treated with doses in the range 50–70 Gy, over 6–7 weeks, administered 5 days per week in about 2 Gy fractions on a once-daily basis. Such a treatment schedule frequently results in acute radiation-induced colitis (ARC).^{28,29} The most common acute symptoms of ARC are diarrhea, tenesmus, and rectal bleeding.³⁰ In spite of the high incidence of these problems in a select group of patients, there is no well established therapy. Polaprezinc is an anti-ulcer agent that is comprised of zinc and carnosine. Zinc is an antioxidant that probably acts by preventing oxidation of a specific sulfhydryl group and by substituting for pro-oxidant metals, such as Cu and Fe, at membrane binding

sites.³¹ Clinically, it has been suggested that zinc supplementation used in conjunction with radiotherapy could postpone the development of severe mucositis and dermatitis for patients with cancers of the head and neck, and can also alleviate the degree of mucositis and dermatitis.³² The aim of this study was to evaluate whether polaprezinc was capable of preventing X-ray radiation-induced apoptosis in the jejunum.

Our data demonstrated that polaprezinc treatment before whole body X-ray irradiation resulted in a significant suppression of apoptotic cells in the rat jejunal crypt at 2 Gy. The increased incidence of apoptosis detected by H&E staining compared with background (0 h) levels peaked at 2 h and then decreased gradually by 6 h in the jejunum. The significant suppression of apoptotic cells by polaprezinc treatment was observed at 2 h, 4 h and 6 h using H&E and TUNEL staining methods in jejunal crypt cells (Fig. 2A, B). TUNEL positive cells in controls reached a plateau at 4 h after irradiation, but the sensitivity of TUNEL assays in the controls was not greater than that seen in H&E staining. This may be because the TUNEL assay is greatly dependent on technical details, mostly related to DNA strand-breaks associated with excessive levels of proteinase digestion, with fixation and processing procedures, or with the action of section cutting or other various pretreatments.³³ Caspase-3 activation is a major effector in the apoptotic process following the irradiation of intestinal cells.³⁴ In this study, the apoptosis index in H&E-stained sections of the control groups increased up to 2 h and decreased gradually by 6 h. In contrast, caspase-3 activation peaked at 2 h and then was sharply reduced at 4 h, with no further increase at 4 h and 6 h. A significant suppression of active caspase-3 positive cells was observed in the polaprezinc treated rats at 2 h and 6 h after irradiation (Fig. 2C). These results suggest that polaprezinc can protect radiation-induced apoptosis and inhibit the caspase-3 activation. The number of active caspase-3 positive cells was lower as compared to the apoptotic index in control groups at 4 h and 6 h. The difference in the numbers of apoptotic cells and active caspase-3 positive cells may reflect the time lag of caspase activation, apoptotic body processing, and degradation because caspase-3 was activated earlier in the apoptotic process than DNA cleavage and cell fragmentation, or it may be due to a caspase-independent apoptosis pathway.³⁵ Irradiation induces DNA damage leading to the activation of p53, p21^{WAF-1/CIP1} and Bax in the murine intestinal epithelium.³⁶ To examine the radiation-induced apoptosis pathways suppressed by polaprezinc treatment, p53, p21^{WAF-1/CIP1} and Bax expressions after X-ray irradiation were examined. p53 accumulation, p21^{WAF-1/CIP1} and Bax expression in polaprezinc-treated rats showed significant decreases compared to control rats after X-ray irradiation at 6 h (Fig. 3).

The antioxidant ability of polaprezinc has also been demonstrated by *in vivo* and *in vitro* studies. Yoshikawa *et al.* suggested that the increase in area of the erosions on the

stomach after ischemia-reperfusion and the increase in lipid peroxides in the gastric mucosa were significantly inhibited by the oral administration of zinc N-(3-aminopropionyl)-L-histidine (Z-103).³⁷⁾ In addition Z-103 inhibits superoxide generation by polymorphonuclear leukocytes, and hydroxyl radical production by the Fenton reaction.³⁸⁾ Z-103 is also found to scavenge superoxide as well.³⁹⁾ On the other hand, it was reported in an *in vitro* study that polaprezinc inhibits apoptosis via inhibition of caspase-3 activation in indomethacin treated RGM1 cells, a rat gastric mucosal cell line but did not prevent mitochondrial cytochrome *c* release.⁴⁰⁾ That paper indicated that the molecular target of polaprezinc lies downstream of cytochrome *c* efflux, yet upstream of caspase-3 activation in indomethacin induced apoptosis. In our study the suppression of p53 accumulation, p21^{WAF-1/CIP1} and Bax expression in polaprezinc-treated rats occurred after the peak of the apoptotic index. Our data demonstrated that the mechanism of the protective effect of polaprezinc treated rats from radiation-induced apoptosis at 2 h and 4 h may be dependent on a p53 independent pathway and the possibility that polaprezinc reacts as an anti-oxidant *in vivo* and its anti-ulcer action may be considered in part by the protective effect on the radiation-induced apoptosis mechanism. But in our study, we didn't investigate the effect of polaprezinc on antioxidant activity in rat intestine after irradiation. Further studies are required to elucidate the mechanism of antiapoptotic effect induced by polaprezinc treatment.

A recent study suggested that polaprezinc protects the small intestine (in either animals or humans) from a nonsteroidal anti-inflammatory drug (NSAID) induced injury, and stimulated cell migration and proliferation *in vitro*.⁴¹⁾ Furthermore, it is suggested that zinc L-carnosine administration protects the colonic mucosa against acetic acid by induction of HSP72 and suppression of NF-kappa B activation.⁴²⁾ In this study, polaprezinc administered orally 1 h before irradiation did not suppress apoptotic cells in the colon (Fig. 2D). This may be because polaprezinc was not present in the colon during radiation in this study design. It may be important the polaprezinc mechanism of membrane-stabilizing action and stimulation of mucus production in the protection of radiation-induced apoptosis in the intestine. Pretreatment with polaprezinc and its presence in the lower digestive tract during radiation may be important for the therapy. It may be possible to prevent acute radiation proctitis with polaprezinc pretreatment during radiotherapy for pelvic malignancies, but further studies are needed. In conclusion, polaprezinc has shown a protective effect of radiation-induced apoptosis in the rat jejunum crypt cells, and decreases of p53 accumulation, p21^{WAF-1/CIP1} and Bax expression were demonstrated.

ACKNOWLEDGEMENTS

This work was partly supported by the 21st Century Cen-

ter of Excellence (COE) Program from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES

1. Ueki, S., Seiki, M., Yoneta, T., Omata, T., Hori, Y., Ishikawa, M. and Tagashira, E. (1989) Effect of Z-103 on compound 40/80-induced gastric lesions in rats. *Scand. J. Gastroenterol.* **24**: 202–205.
2. Arakawa, T., Satoh, H. and Nakamura, A. (1990) Effects of zinc L-carnosine on gastric mucosal and cell damage caused by ethanol in rats. *Dig. Dis. Sci.* **35**: 559–566.
3. Seiki, M., Ueki, S., Hori, Y., Aita, H., Yoneta, T. and Tagashira, E. (1991) Effects of a new compound, Z-103 on various acute experimental models of gastric and duodenal lesions in rats. *Therapeutic Research* **12**: 3243–3251.
4. Yoshikawa, T., Yamaguchi, T., Yoshida, N., Yamamoto, H., Kitazumi, S., Takahashi, S., Naito, Y. and Kondo, M. (1997) Effect of Z-103 on TNB-induced colitis in rats. *Digestion* **58**: 464–468.
5. Cho, C. H., Luk, C. T. and Ogle, C. W. (1991) The membrane-stabilizing action of zinc carnosine (Z-103) in stress-induced gastric ulceration in rats. *Life Sci* **49**: 189–194.
6. Arakawa, T., Satoh, H., Nakamura, A., Fukuda, T., Sakuma, H., Nakamura, H., Ishikawa, M., Seiki, M. and Kobayashi, K. (1990) Effect of zinc-L-carnosine on gastric mucosal and cell damage caused by ethanol in rats. Correlation with endogenous prostaglandin E2. *Dig. Dis. Sci.* **35**: 559–566.
7. Potten, C. S. (1991) Extreme sensitivity of some intestinal crypt cells to X and gamma irradiation. *Nature* **269**: 518–521.
8. Potten, C. S., Merritt, A., Hickman, J., Hall, P. and Faranda, A. (1994) Characterization of radiation-induced apoptosis in the small intestine and its biological implications. *Int. J. Radiat. Biol.* **65**: 71–78.
9. Keefe, D. M., Brealey, J., Golland, G. J. and Cummins, A. G. (2000) Chemotherapy for cancer causes apoptosis that precedes hypoplasia in crypts of the small intestine in human. *Gut* **47**: 632–637.
10. Kerr, J. F. R., Wyllie, A. H. and Currie, A. R. (1972) Apoptosis: a basic biological phenomenon with wide ranging implications in tissue kinetics. *Br. J. Cancer* **26**: 239–257.
11. Ijiri, K. and Potten, C. S. (1983) Response of intestinal cells of differing topographical and hierarchical status to ten cytotoxic drugs and five sources of radiation. *Br. J. Cancer* **47**: 175–185.
12. Potten, C. S. (1992) The significance of spontaneous and induced apoptosis in the gastrointestinal tract of mice. *Cancer Metastasis Rev.* **11**: 179–195.
13. Matsuu, M., Shichijo, K., Okaichi, K., Wen, C. Y., Fukuda, E., Nakashima, M., Nakayama, T., Shirahata, S., Tokumaru, S. and Sekine, I. (2003) The protective effect of fermented milk kefir on radiation-induced apoptosis in colonic crypt cells of rats. *J. Radiat. Res.* **44**: 111–115.
14. Matsuu-Matsuyama, M., Shichijo, K., Okaichi, K., Ishii, K., Wen, C. Y., Fukuda, E., Nakayama, T., Nakashima, M., Okumura, Y. and Sekine, I. (2006) Sucralfate protects intestinal epithelial cells from radiation-induced apoptosis in rats. *J. Radiat. Res.* **47**: 1–8.
15. Kastan, M. B., Onyekwere, O., Sidransky, D., Vogelstein, B.

- and Craig, R. W. (1991) Participation of p53 protein in the cellular response to DNA damage. *Cancer Res.* **51**: 6304–6311.
16. Kuerbitz, S. J., Plunkett, B. S., Walsh, W. V. and Kastan, M. B. (1992) Wild-type p53 is a cell cycle check point determinant following irradiation. *Proc. Natl. Acad. Sci. USA.* **89**: 7491–7495.
 17. Tishler, R. B., Calderwood, S. K., Coleman, N. C. and Price, B. D. (1993) Increase in sequence specific DNA binding by p53 following treatment with chemotherapeutic and DNA damaging agents. *Cancer Res.* **53**: 2212–2216.
 18. Zhan, Q., Carrier, F. and Fornace, A. J. (1993) Induction of cellular p53 activity by DNA-damaging agents and growth arrest. *Mol. Cell. Biol.* **13**: 4242–4250.
 19. El-Deiry, W. S., Harper, J. W., O'Connor, P. M., Velculescu, V. E., Canman, C. E., Jackman, J. *et al.* (1994) Waf1/Cip1 is induced in p53 mediated G1 arrest and apoptosis. *Cancer Res.* **54**: 1169–1174.
 20. Bae, I., Fan, S., Bhatia, K., Kohn, K. W., Fornace, A. J. and O'Connor, P. M. (1995) Relationships between G1 arrest and stability of the p53 and p21^{waf1/cip1} proteins following gamma-irradiation of human lymphoma cells. *Cancer Res.* **55**: 2387–2393.
 21. Pritchard, D. M., Potten, C. S., Korsmeyer, S. J., Roberts, S. and Hickman, J. A. (1999) Damage-induced apoptosis in intestinal epithelia from bcl-2-null and bax-null mice: investigations of the mechanistic determinants of epithelial apoptosis in vivo. *Oncogene* **18**: 7287–7293.
 22. Wijsman, J. H., Jonker, R. R., Keijzer, R., Van De Velde, C. J. and Van Dierendonck, J. H. (1993) A new method to detect apoptosis in paraffin sections: in situ end labeling of fragmented DNA. *J. Histochem. Cytochem.* **41**: 7–12.
 23. Kerr, J. F., Winterford, C. M. and Harmon, B. V. (1994) Apoptosis. Its significance in cancer and cancer therapy. *Cancer* **73**: 2013–2026.
 24. Walker, N. I., Harmon, B. V., Gobe, G. C. and Kerr, J. F. (1988) Patterns of cell death. *Methods. Achiev. Exp. Pathol.* **13**: 18–54.
 25. Arai, T., Kida, Y., Harmon, B. V. and Gobe, G. C. (1996) Expression and localization of clusterin mRNA in the small and large intestine of the irradiated rat: its relationship with apoptosis. *Int. J. Radiat. Biol.* **69**: 547–553.
 26. Potten, C. S. (1996) What is an apoptotic index measuring? A commentary. *Br. J. Cancer* **74**: 1743–1748.
 27. Potten, C. S., Wilson, J. W. and Booth, C. (1997) Regulation and significance of apoptosis in the stem cells of the gastrointestinal epithelium. *Stem Cells* **15**: 82–93.
 28. Nussbaum, M. L., Campama, T. J. and Weese, J. L. (1993) Radiation-induced intestinal injury. *Clin. Plast. Surg.* **20**: 573–580.
 29. Hovdenak, N., Fajardo, L. F., and Hauer-Jensen, M. (2000) Acute radiation proctitis: A sequential clinicopathologic study during peric radiotherapy. *Int. J. Radiat. Oncol. Biol. Phys.* **48**: 1111–1117.
 30. Gilinsky, N. H., Burns, D. G., Barbezat, G. O., Levin, W., Myers, H. S. and Marks, I. N. (1983) The natural history of radiation-induced proctosigmoiditis. *Q. J. Med.* **205**: 40–53.
 31. Bray, T. M. and Bettger, W. J. (1990) The physiological role of zinc as an antioxidant. *Free Radic. Biol. Med.* **8**: 281–291.
 32. Lin, L.-C., Que, J., Lin, L.-K. and Lin, F.-C. (2006) Zinc supplementation to improve mucositis and dermatitis in patients after radiotherapy for head-and-neck cancers: A double-blind, randomized study. *Int. J. Radiat. Oncol. Biol. Phys.* **65**: 745–750.
 33. Duan, W. R., Gamer, D. S., Williams, S. D., Funckes-Shippy, C. L., Spath, I. S. and Blomme, E. A. G. (2003) Comparison of immunohistochemistry for activated caspase-3 and cleaved cytokeratin 18 with the TUNEL method for quantification of apoptosis in histological sections of PC-3 subcutaneous xenografts. *J. Pathology* **199**: 221–228.
 34. Marshman, E., Ottewell, P. D., Potten, C. S. and Watson, A. J. (2001) Caspase activation during spontaneous and radiation-induced apoptosis in the murine intestine. *J. Pathol.* **195**: 285–292.
 35. Inagaki-Ohara, K., Takamura, N., Yada, S., Alnadjim, Z., Liu, E., Yu, X., Yoshida, H. and Lin, T. (2002) Radiation-induced crypt intestinal epithelial cell apoptosis *in vivo* involves both caspase-3 dependent and -independent pathways. *Dig. Dis. Sci.* **47**: 2823–2830.
 36. Wilson, J. W., Pritchard, D. M., Hickman, J. A. and Potten, C. S. (1998) Radiation-induced p53 and p21^{waf1/cip1} expression in the murine intestinal epithelium. Apoptosis and cell cycle arrest. *Am. J. Pathol.* **153**: 899–909.
 37. Yoshikawa, T., Naito, Y., Tanigawa, T., Yoneta, T., Yasuda, M., Ueda, S., Oyamada, H. and Kondo, M. (1991) Effect of zinc-carnosine chelate compound (Z-103), a novel antioxidant, on acute gastric mucosal injury induced by ischemia-reperfusion in rats. *Free Radic. Res. Commun.* **14**: 196–289.
 38. Naito, Y., Yoshikawa, T., Tanigawa, T., Oyamada, H., Takemura, T., Yoshida, N., Sugino, S., Kondo, M. and Yoneta, T. (1989) Protection by zinc N-(3-aminopropionyl)-L-histidine against stress-induced gastric mucosal lesions in rats and its antioxidative effects. In *Medical, Biochemical and Chemical Aspects of Free Radicals* (ed. O. Hayaishi, E. Niki, M. Kondo, T. Yoshikawa), Elsevier, Amsterdam, pp. 473–476.
 39. Yoshikawa, T., Naito, Y., Tanigawa, T., Yoneta, T., Oyamada, H., Ueda, S., Takemura, T., Sugino, S. and Kondo, M. (1988) Effect of the zinc-carnosine chelate Z-103 on the generation of superoxide anion radicals from hypoxanthine-xanthine oxidase system. *J. Clin. Exp. Med.* **147**: 927–928.
 40. Fujii, Y., Matura, T., Kai, M., Kawasaki, H. and Yamada, K. (2000) Protection by polaprezinc, an anti-ulcer drug, against indomethacin induced apoptosis in rat gastric mucosal cells. *Jpn. J. Pharmacol.* **84**: 63–70.
 41. Mahmood, A., FitzGerald, A. J., Marchbank, T., Ntatsaki, E., Murray, D., Ghosh, S. and Playford, R. J. (2007) Zinc carnosine, a health food supplement that stabilises small bowel integrity and stimulates gut repair processes. *Gut* **56**: 168–175.
 42. Odashima, M., Otaka, M., Jin, M., Wada, I., Horikawa, Y., Matsushashi, T., Ohba, R., Hatakeyama, N., Oyake, J. and Watanabe, S. (2006) Zinc L-carnosine protects colonic mucosal injury through induction of heat shock protein 72 and suppression of NF-kappa B activation. *Life sci.* **79**: 2245–2250.

Received on November 13, 2007

Revision received on January 14, 2008

Accepted on February 6, 2008

J-STAGE Advance Publication Date: April 15, 2008