

Kent Academic Repository

Full text document (pdf)

Citation for published version

Mifsud, Duncan V., Hailey, Perry A., Traspas Muiña, Alejandra, Auriacombe, Olivier, Mason, Nigel J. and Ioppolo, Sergio (2021) The role of terahertz and far-IR spectroscopy in understanding the formation and evolution of interstellar prebiotic molecules. *Frontiers in Astronomy and Space Sciences*, 8. ISSN 2296-987X.

DOI

<https://doi.org/10.3389/fspas.2021.757619>

Link to record in KAR

<https://kar.kent.ac.uk/92262/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

The Role of Terahertz and Far-IR Spectroscopy in Understanding the Formation and Evolution of Interstellar Prebiotic Molecules

Duncan V. Mifsud^{1,2*}, Perry A. Hailey¹, Alejandra Traspas Muiña³, Olivier Auriacombe⁴, Nigel J. Mason¹, Sergio Ioppolo³

¹Centre for Astrophysics and Planetary Science, School of Physical Sciences, University of Kent, Canterbury CT2 7NH, United Kingdom

²Institute for Nuclear Research (Atomki), Debrecen H-4026, PO Box 51, Hungary

³School of Electronic Engineering and Computer Science, Queen Mary University of London, London E1 4NS, United Kingdom

⁴Microwave Electronics Laboratory, Department of Microtechnology and Nanoscience, Chalmers University of Technology, Göteborg 41296, Sweden

* Correspondence:

Duncan V. Mifsud
duncanvmifsud@gmail.com

Keywords: terahertz spectroscopy, far-IR spectroscopy, astrochemistry, interstellar chemistry, prebiotic chemistry, review

Abstract

Stellar systems are often formed through the collapse of dense molecular clouds which, in turn, return copious amounts of atomic and molecular material to the interstellar medium. An in-depth understanding of chemical evolution during this cyclic interaction between the stars and the interstellar medium is at the heart of astrochemistry. Systematic chemical composition changes as interstellar clouds evolve from the diffuse stage to dense, quiescent molecular clouds to star-forming regions and proto-planetary disks further enrich the molecular diversity leading to the evolution of ever more complex molecules. In particular, the icy mantles formed on interstellar dust grains and their irradiation are thought to be the origin of many of the observed molecules, including those that are deemed to be ‘prebiotic’; that is those molecules necessary for the origin of life. This review will discuss both observational (e.g., ALMA, SOFIA, Herschel) and laboratory investigations using terahertz and far-IR (THz/F-IR) spectroscopy, as well as centimeter and millimeter spectroscopies, and the role that they play in contributing to our understanding of the formation of prebiotic molecules. Mid-IR spectroscopy has typically been the primary tool used in laboratory studies, particularly those concerned with interstellar ice analogues. However, THz/F-IR spectroscopy offers an additional and complementary approach in that it provides the ability to investigate intermolecular interactions compared to the intramolecular modes available in the mid-IR. THz/F-IR spectroscopy is still somewhat under-utilized, but with the additional capability it brings, its popularity is likely to significantly increase in the near future. This review will discuss the strengths and limitations of such methods, and will also provide some suggestions on future research areas that should be pursued in the coming decade exploiting both space-borne and laboratory facilities.

1 Interstellar Chemistry: From Principles to Practice

The idea of active chemistry occurring within interstellar space is a relatively new one. Historically, interstellar temperatures and pressures were considered to be far too low to permit atoms or ions to come into contact with each other and overcome a reaction activation energy barrier. However, since the first evidence of interstellar molecular species arose in the late 1930s, the scientific community has come to recognize that the interstellar medium is home to a plethora of structurally diverse molecules, with different environments displaying their own characteristic chemistry. A more complete description of the history of astrochemistry may be found in the work of Feldman (2001).




Figure 1 Matter in the interstellar medium is cycled through various phases: the aggregation of matter results in optically transparent diffuse clouds evolving into dense, dark molecular clouds. These clouds are considered to be ‘stellar nurseries’ in which localized gravitational collapse results in the formation of a proto-star, with further accretion of material occurring via magneto-hydrodynamic flows onto the growing star. Within a few million years, the stellar object evolves into a pre-main sequence star surrounded by a proto-planetary disk. Collisions within the disk result in the accretion of dust grains which slowly grow to form planetesimals, thus taking the first steps towards the formation of a planetary system. Once the star’s nuclear fuel has been exhausted, it may follow one of two routes depending on the stellar mass: low-mass stars undergo several mass-loss processes in which molecular material is dispersed back into the interstellar medium leaving behind a remnant white dwarf, whereas high-mass stars instead explode as supernovae with their cores eventually becoming either neutron stars or black holes.

Matter in the interstellar medium is composed of around 99% atomic or molecular hydrogen and 1% carbonaceous or silicate dust grains. This matter is cycled through a number of phases each displaying a characteristic chemistry (Figure 1). The smallest over-densities result in the formation of diffuse clouds, which possess particle densities of 10^2 - 10^3 cm^{-3} and kinetic temperatures of a few hundred K (Smith 1992). Diffuse clouds are relatively transparent to visible light and are continuously exposed to

the interstellar radiation field which results in the dissociation and ionization of molecular material. As such, these clouds are dominated by the presence of atoms and ions, particularly H and H⁺, although a few molecular species do survive (Snow and Bierbaum 2008). Indeed, the first molecules to be detected in interstellar space were detected in diffuse interstellar media via optical spectroscopy (Smith 1992, Feldman 2001, Larsson et al. 2012). With the notable exception of the formation of H₂ (Wakelam et al. 2017), most of the chemistry within diffuse interstellar clouds is mediated by gas-phase reactions (Geppert and Larsson 2013).

Accumulation of gaseous material within diffuse clouds may occur as a result of random perturbations, stellar winds, or more violent phenomena such as supernovae-induced shock waves. As diffuse clouds begin to accumulate matter, the proportion of molecular material (mainly H₂) increases. This is accompanied by a diminishing of the proportion of ionized species present: for example, the dominant form of carbon changes to the atomic form from C⁺ and the number of free electrons also decreases (Snow and Bierbaum 2008). Thus, the accumulation of matter results in an evolution of so-called ‘atomic’ diffuse clouds to ‘molecular’ diffuse clouds.

Further accumulation of material as a result of gravitational attractions results in the formation of a dense interstellar cloud (also known as a dark interstellar cloud). These clouds are, as their name suggests, characterized by higher particle number densities of up to 10⁶ cm⁻³ and are the primary location of stellar formation. The increase in density results in a concomitant increase in cloud opacity. As such, the far- and vacuum-UV components of the interstellar radiation field are attenuated at the edges of the cloud, resulting in an interior with a temperature of 7-20 K (Galli et al. 2002). The higher particle densities of these dark clouds enhance the efficiency of reactions in the gas phase, particularly those between ions and molecules, and contribute to the formation of several new species including complex organic molecules (Bergin and Tafalla 2007, Öberg 2016, Arumainayagam et al. 2019).

At the low temperatures encountered within the interiors of dense interstellar clouds, many species are able to freeze-out onto the surfaces of the carbonaceous and silicate dust grains, thus forming an icy mantle. This icy mantle is typically composed of two zones (Figure 2): a lower polar layer which results from the deposition and synthesis (via hydrogenation reactions) of polar molecules such as H₂O, NH₃, and also CH₄; and an upper apolar layer characterized by the presence of CO and related molecules such as CO₂ and CH₃OH (Larsson et al. 2012, Öberg 2016).





Figure 2 Median structure of icy grain mantles within dense interstellar clouds. Shown in the image, a diagram (not to scale) of a typical icy mantle divided into two zones: a lower polar layer rich in H₂O molecules and hydrogenated heteroatoms, and an upper apolar layer largely consisting of CO and related molecules. More detailed information can be found in the work of Öberg (2016).

These molecular icy grain mantles play host to a rich chemistry mediated by several phenomena and which may result in the formation of complex molecules. For instance, galactic cosmic rays are not attenuated by the boundaries of the dense cloud, and so are able to penetrate through to the cloud core and induce chemistry within the icy layers. Interactions between these cosmic rays and gas-phase H₂ results in the emission of Lyman- α photons which may induce photochemistry. Photochemistry in icy grain mantles may also be induced by lower wavelength photons which enter the cloud via less dense regions. Thermal chemistry may also occur in ices proximal to a heat source, such as a nascent star.



In this section, we review the chemistry which occurs within diffuse and dense interstellar clouds with a particular focus on reactions involving ionic species, and those which result in the formation of complex organic molecules. We should note that, in the literature, the term ‘complex organic molecule’ typically refers to a saturated molecule which may be conceivably formed by geological or biochemical processes. However, for the purposes of this review, we extend this definition to include the unsaturated polyatomic carbon chains which have been detected in astrophysical settings. An understanding of interstellar chemistry is an important prerequisite for a discussion of the experimental and observational techniques used in astrochemistry, and although mentions of detections via terahertz and far-IR (THz/F-IR) spectroscopy and related techniques will be made, further details on such techniques will be given in the subsequent section. Reviews on the chemistry occurring during the various other phases of the astrochemical cycle (Figure 1) are also available elsewhere (van Dishoeck and Blake 1998, Williams and Hartquist 1999, Larsson et al. 2012, Geppert and Larsson 2013, van Dishoeck 2014).

1.1 Diffuse Interstellar Clouds: Overview of Ion Chemistry

As mentioned previously, matter within most diffuse interstellar clouds is primarily composed of atomic and ionic material, with a significantly smaller number of molecules also being present due to the high rates of photon and cosmic ray induced dissociation and ionization processes. Consequentially, the chemistry within these clouds is dominated by gas-phase reactions including those involving or producing ions. At the edges of the diffuse cloud, where stellar UV radiation is more intense, photo-ionization of carbon atoms produces C⁺, while within the diffuse cloud interior cosmic ray driven chemistry is more dominant and allows for the ionization of H and H₂ (Smith 1992). The formation of H₂⁺ in particular is important, as this species is the direct precursor to H₃⁺: 124



For a long time, the formation and presence of H₃⁺ within diffuse clouds were highly controversial due to its suspected high rate of dissociated recombination mediated by electrons emitted from the photo-ionization of carbon atoms (Larson et al. 2000, Larsson et al. 2008). However, its direct observation in diffuse clouds (McCall et al. 1998) combined with the discovery of a new method of recombination (Kokoouline and Greene 2003) confirmed its presence and role within interstellar chemistry: H₃⁺ efficiently donates a proton to other atoms and molecules due to the low proton affinity of H₂. The resultant protonated species is reactive, and is thus capable of triggering various chains of ion-molecule reactions. For instance, the reaction with CO produces HCO⁺, while the reaction with atomic oxygen produces OH⁺:



OH^+ may also be formed as a result of a two-step process starting with the charge-transfer reaction between atomic oxygen and H^+ to yield O^+ , which subsequently reacts with H_2 to furnish OH^+ (Yamamoto 2017). The formation of OH^+ is the first step towards the formation of interstellar gas-phase H_2O . The reaction of OH^+ with one moiety of H_2 yields H_2O^+ , and the reaction with a second moiety produces H_3O^+ (Larsson et al. 2012). The H_3O^+ molecular ion is isoelectronic with NH_3 and so is a stable species which resists further reactions with H_2 (Geppert and Larsson 2013). Instead, it undergoes dissociative recombination with electrons to yield either OH (~60% of outcomes) or H_2O (~25% of outcomes).

The detections of OH^+ , H_2O^+ , and H_3O^+ within diffuse interstellar environments have been made using ground- and space-based telescopes working in the THz/F-IR range, such as the Atacama Pathfinder Experiment (APEX) and the HIFI instrument aboard the Herschel Space Observatory (for more detail, refer to section 2), thus confirming their contribution to chemistry in such settings (Wyrowski et al. 2010, Gerin et al. 2010a, Gupta et al. 2010, Wiesemeyer et al. 2016). A reaction scheme analogous to the formation of H_2O from OH^+ has also been invoked to explain the formation of HCl within diffuse clouds starting from Cl^+ . Such a reaction scheme has been validated by the detection of the relevant intermediate species HCl^+ and H_2Cl^+ by the HIFI instrument (Lis et al. 2010, De Luca et al. 2012).

Reactions leading to the formation of carbon-bearing molecules within diffuse interstellar media are more challenging to explain compared to the oxygen and chlorine chemistry outlined above. Indeed, the formation of CH^+ , one of the earliest known interstellar molecules (Douglas and Herzberg 1941), is still not completely understood and several simulations have underestimated its observed abundance within diffuse clouds (e.g., van Dishoeck and Black 1986, Pan et al. 2004, Godard et al. 2009). It is apparent that the formation of CH^+ cannot rely on the reaction between H_2 and C^+ due to the considerable endothermicity of this reaction (Myers et al. 2015, Valdivia et al. 2017, Moseley et al. 2021, Plašil et al. 2021). In light of this, alternative mechanisms have been suggested to account for its presence within the diffuse interstellar medium. For instance, it has been suggested that the radiative association of C^+ and H_2 to yield CH_2^+ with the concomitant emission of a photon is the first step towards interstellar CH^+ , as the photo-dissociation of the former may lead to the latter (Yamamoto 2017). However, the rate at which radiative association occurs is fairly low (Graff et al. 1983, Barinova and van Hemert 2006) and indeed the primary reaction pathway of CH_2^+ within diffuse interstellar media is the reaction with H_2 which furnishes CH_3^+ . As such, the potential contribution of this reaction network in accounting for interstellar CH^+ abundances is probably low.

Another proposed formation route for CH^+ is the direct reaction between C^+ and H_2 if the endothermicity of the reaction is overcome by some mechanism which causes a localized warming effect which raises the temperature to a few thousand K. Initially, it was thought that this localized warming could be achieved in regions of the diffuse cloud processed by magneto-hydrodynamic shock waves (Elitzur and Watson 1980, Draine and Katz 1986). However, such processes are now thought to be an unlikely source of CH^+ as the radial velocity for this molecule and for CH have been observed to be similar. Among other ideas, gas heating caused by turbulences within the cloud has also been found to be unsatisfactory in accounting for the observed abundances of CH^+ (Godard and Cernicharo 2013), as have reactions between C^{2+} and H_2 in X-ray dominated regions of the diffuse cloud (Plašil et al. 2021). Alfvén waves have also been suggested as a potential source of CH^+ (Federman et al. 1996), however, to the best of the authors' knowledge, this has been neither corroborated nor firmly rejected. As such, the elusive source of CH^+ within diffuse interstellar media remains an unanswered question. There thus exists a strong motivation to address this problem, as the formation of CH^+ may be an important step in the synthesis of significantly more complex carbon-based molecules such as fullerenes and polycyclic aromatic hydrocarbons (PAHs), which are suspected of being the primary

carriers of observed diffuse interstellar bands and which may bear relevance to the synthesis of prebiotic molecules (Taylor and Duley 1997).

With regards to molecular detections in diffuse interstellar clouds, a significant number have been made using methods and techniques which exploit the **submillimeter, millimeter, and centimeter** regions of the electromagnetic spectrum, including THz/F-IR spectroscopy (Snow and McCall 2006, Yamamoto 2017). Indeed, such techniques are the preferred method for detecting polyatomic molecules, as these molecules are difficult to identify using optical spectroscopy (as was done historically) due to the pre-dissociative nature of their excited states. The HIFI instrument aboard the Herschel Space Observatory in particular has proven to be invaluable to the detection and mapping of molecules in diffuse interstellar media, having provided much information on the chemistry of hydrogen-, carbon-, oxygen-, nitrogen-, sulfur-, fluorine-, and chlorine-bearing molecules within diffuse clouds (Sonnentrucker et al. 2010, Langer et al. 2010, Persson et al. 2010, Neufeld et al. 2010a, 194 2010b, 2012a, Gerin et al. 2010a, 2010b, 2012, Monje et al. 2011, 2013, De Luca et al. 2012, Esplugues et al. 2014, Schilke et al. 2014, Indriolo et al. 2015, Jacob et al. 2020).

Other instruments and telescopes have also aided greatly in the elucidation of the chemical complexity of the diffuse interstellar medium. For example, APEX, the Atacama Large Millimeter/Submillimeter Array (ALMA), and the GREAT instrument aboard the Stratospheric Observatory for Infrared Astronomy (SOFIA) have all been used in the detection of sulfur-bearing molecules, including isotopologues (Menten et al. 2011, Neufeld et al. 2012b, 2015, Muller et al. 2017). Perhaps the most exciting of all has been the series of recent ALMA observations of complex molecules. The work of Thiel et al. (2017), for instance, investigated the presence of prebiotic complex organic molecules within diffuse media and detected several species, including CH_3OH , CH_3CN , CH_3CHO , HC_3N , and CHONH_2 . Such results are of immense consequence, as they imply that certain molecular precursors to life either form early on in the astrochemical cycle (Figure 1), or else survive a process of expansion from denser interstellar media.

1.2 Dense (Dark) Interstellar Clouds: Overview of Ion Chemistry

Condensation of diffuse clouds results in the formation of dense (dark) clouds. During this process, the dominant material is converted from atomic and ionic species to molecular ones. For instance, H_2 becomes the dominant form of hydrogen, while the dominant form of carbon changes sequentially from C^+ , to C , and finally to CO (Snow and Bierbaum 2008, Larsson et al. 2012). As such, these structures have also been appropriately dubbed molecular clouds. In their review, Arumainayagam et al. (2019) highlighted the chemical productivity of dense clouds and identified three milieux for astrochemical reactions within them. The first of these is gas-phase chemistry, which includes radical-radical, radical-neutral, and ion-neutral reactions. The second is chemistry occurring on bare carbonaceous or silicate dust grain surfaces, while the third includes all chemical reactions occurring via energetic and non-energetic processing of icy grain mantles. **In this sub-section, we will largely limit ourselves to a brief overview of ion-molecule reactions which occur in the gas-phase within dense interstellar clouds.**

Gas-phase ion-neutral reactions within dense interstellar clouds are favorable, barrierless chemical processes. The source of the ionic species within the cloud is somewhat dependent upon the location of the reaction within the cloud structure. Within the densest regions (surrounding the cloud core), ionization of molecules is caused almost exclusively by penetrating galactic cosmic rays. However, within the less dense regions of the clumpy cloud structure, far- and vacuum-UV components of the interstellar radiation field are the primary cause of molecular ionization via photon-induced electron

loss processes. Furthermore, in the presence of OB stars, irradiation of molecular material by X-rays may also contribute to the formation of ionic species.

Cosmic ray induced ionization of H_2 leads indirectly to the formation of H_3^+ as was described for diffuse interstellar clouds. The low proton affinity of H_2 allows H_3^+ to be an efficient proton donor to other species (Larsson et al. 2012). For example, the reaction with N_2 yields N_2H^+ , which is frequently used as a tracer for N_2 within interstellar space when observing with radio telescopes, since N_2 does not possess a permanent dipole moment. Ionized hydrogen, primarily in the form of H^+ , readily undergoes reactions with other atoms such as nitrogen atoms. The resultant product NH^+ may undergo subsequent hydrogenation reactions with H_2 to sequentially form NH_2^+ , NH_3^+ , and NH_4^+ (Geppert and Larsson 2013). Finally, dissociative recombination with an incident electron efficiently affords gas-phase NH_3 (Öjekull et al. 2004).

Interestingly, the analogous reaction pathway to the formation of gas-phase CH_4 is not thought to be significant within dense clouds. This is because once CH_5^+ is formed, the dominant dissociative recombination mechanism involves the simultaneous loss of two hydrogen atoms, creating CH_3 rather than CH_4 (Semaniak et al. 1998, Viti et al. 2000). The formation of CH_4 and several other carbon-based molecules within the interstellar medium is instead likely to rely on other mechanisms, such as hydrogenation reactions at the surfaces of dust grains (Qasim et al. 2020).

A large number of carbon-based complex organic molecules are known to exist within dense clouds. Until recently, it was thought that these molecules formed within the solid phase in icy grain mantles and were released into the gas phase upon thermal or photo-desorption. However, a growing body of literature has documented the presence of such gas-phase complex organic molecules within starless and pre-stellar cores, where such desorption processes do not take place. Scibelli and Shirley (2020), for instance, have detected CH_3OH and CH_3CHO in several such environments and have suggested that the radicals and molecules responsible for the formation of these species may be formed in the ice phase. The release of energy upon their formation allows them to sublime into the gas phase (a process termed reactive desorption) whereupon they react to form these product molecules.

Another region of dense clouds which may host a rich gas-phase chemistry leading to the formation of complex organic molecules is the edge of the cloud: the photo-dissociation region (PDR). In this region, the cloud interacts with the intense UV components of the interstellar radiation field resulting in the desorption of solid-phase molecules and an increase in the occurrence of gas-phase processes. Using the IRAM 30 m telescope, Cuadrado et al. (2017) were able to perform a millimeter linesurvey towards the edge of the Orion Bar PDR and successfully detected a plethora of complex molecules including carboxylic acids, aldehydes, alkenes, and sulfur-containing molecules. Soma et al. (2018) reported similar results. The production and survival of complex molecules in such an environment is significant from the perspective of astrobiology and, although their exact formation mechanisms are still not completely understood, it is likely that a non-negligible contribution comes from gas-phase reactions.

Of course, other processes leading to the formation of complex organic molecules within dense interstellar clouds are also known. Garrod et al. (2008), for instance, demonstrated that the warm up of icy grain mantles previously processed by galactic cosmic rays under cold conditions yielded various complex organic molecules in abundances which matched those documented observationally, while Balucani et al. (2015) invoked radiative association as a gas-phase process leading to their formation. An interesting mechanism which has been suggested to account for the formation of complex organic molecules is that of non-diffusive surface reactions, which have been experimentally shown to yield

such species in reasonably high abundances (Fedoseev et al. 2015, 2017, Chuang et al. 2016, Ioppolo et al. 2021). 271

The presence and distribution of complex, potentially prebiotic molecules within the interstellar medium have been extensively probed using various ground- and space-based observatories. Perhaps the most productive of these studies, however, have been those which have made use of the ALMA for molecular detections (Belloche et al. 2016, 2019, Garrod et al. 2017, Jørgensen et al. 2018, Calcutt et al. 2019, Willis et al. 2020). The impact of ALMA observations on the detection of interstellar molecules has been massive: first detections of prebiotic molecules which were previously controversial or unconfirmed, such as urea (NH_2CONH_2), have been achieved (Belloche et al. 2019), and more information has been gleaned on interstellar chemical processes such as deuteration (Belloche et al. 2016).

To conclude this sub-section, we provide an overview of two especially fruitful ongoing observational projects which include THz/F-IR, millimeter, and centimeter frequencies in their operational ranges: the GOTHAM and ARKHAM projects. These projects are run using the Robert C. Byrd 100 m Green Bank Telescope in the United States and have provided significant data demonstrating the presence of complex organic molecules within interstellar space, with a particular focus on aromatic species. Although operating in a similar fashion, the projects themselves have somewhat different objectives.

The GOTHAM project is primarily concerned with establishing a chemical inventory of the Taurus Molecular Cloud, one of the nearest dense clouds, by performing high-sensitivity wide-band spectral line surveys. Such work is motivated by the fact that benzonitrile (*c*- $\text{C}_6\text{H}_5\text{CN}$) was recently discovered in this pre-stellar source (McGuire et al. 2018a), contrasting with a working hypothesis which postulates that larger interstellar aromatic molecules are produced in the circumstellar regions of post-AGB stars and subsequently break down into smaller sub-units such as benzene (*c*- C_6H_6) under energetic processing (Tielens 2008): so-called ‘top-down’ chemistry. However, these results suggest that reactions between smaller precursor molecules may also contribute to the formation of larger aromatics within pre-stellar regions through a ‘bottom-up’ chemistry approach.

GOTHAM observations of the Taurus Molecular Cloud, combined with laboratory-generated spectra and computational models, have successfully identified many molecules, some of which had not yet been described in the interstellar medium. The unsaturated molecules detected within this dense cloud may play a pivotal role in the formation of larger aromatic molecules, as well as other species relevant to the origins of life. Early work from GOTHAM identified propargyl cyanide (HCCCH_2CN) and the cyanopolyne HC_{11}N in the Taurus Molecular Cloud (McGuire et al. 2020, Loomis et al. 2021). Cyclic molecules have also been discovered, including 1-cyanocyclopentadiene and 2-cyanocyclopentadiene (McCarthy et al. 2021, Burkhardt et al. 2021a). Other cyano and isocyano species have also been detected (Xue et al. 2020, Lee et al. 2021b).

The ARKHAM project is focused on investigating whether the chemistry observed in the Taurus Molecular Cloud is also found in other pre-stellar and proto-stellar sources so as to assess whether or not the resultant molecules survive the birth and evolution of a proto-star. Results from the project have shown that benzonitrile has been detected in four additional sources: Serpens 1A, Serpens 1B, Serpens 2, and MC27/L1521F (Burkhardt et al. 2021b). Such detections illustrate that aromatic chemistry is favorable and extensive throughout the very earliest stages of star formation. Interestingly, the observed abundances of benzonitrile exceeded those predicted by models, despite these models being well-suited to other large carbon-based molecules such as cyanopolyynes. It is therefore probable that

the formation mechanisms of other aromatic species in pre-stellar sources differ from those of linear molecules and are not yet well understood.

1.3 The Link to Prebiotic Chemistry

Joint observational and laboratory investigations are key to deciphering the exact mechanisms of formation of the various complex organic molecules observed within interstellar and circumstellar media. To date, these studies have largely relied on a ‘bottom-up’ chemical perspective in which large complex molecules are produced from simpler starting molecules undergoing some form of processing (such as photolysis or radiolysis). However, as has been demonstrated by the GOTHAM and ARKHAM projects, the formation of certain molecules (such as aromatic species) is likely to be dependent on some combination of ‘bottom-up’ and ‘top-down’ chemistry (McGuire et al. 2020): that is to say that they are formed both from the joining together of smaller molecules, ions, and radicals, as well as the break-up of larger structures (Figure 3).




Figure 3 Schematic diagram exhibiting models of ‘top-down’ and ‘bottom-up’ chemistry using carbon molecules as an example. In the ‘bottom-up’ perspective, the association of atoms and simple molecules allows for the formation of larger molecules such as CH_4 and CH_3OH . Further additions to the molecular structure may result in the formation of a complex organic molecule, such as an aromatic molecule or a PAH. Two-dimensional extension of the PAH structure produces graphite, which itself may be considered the basis of (sub-)micron sized interstellar carbonaceous dust grains. The ‘top-up’ chemical perspective is essentially the reverse of this, in which larger molecular structures are broken up as a result of energetic processing, thus yielding simpler molecules.

PAHs are perhaps the best class of interstellar molecules to exemplify the contributions of ‘top-down’ and ‘bottom-up’ astrochemistry due to the fact that they are **thought to be** relatively ubiquitous in interstellar media, and also because they are able to undergo many chemical transformations yielding a plethora of molecular structures with various sizes, complexities, and arrangements of the aromatic rings, as well as different chemical characteristics such as the inclusion of heteroatoms and the addition of functional groups (such as amino or carboxylic moieties). Indeed, such chemical transformations are thought to be key to the formation of very small grains and fullerenes which result from the respective agglomeration and destruction of PAHs, as well as serving as a carbon feedstock for the production of molecules relevant to biology (Tielens 2008, Groen et al. 2012).

For instance, when mixed with H₂O and subjected to energetic processing (a scenario which is plausible within icy grain mantles in the dense interstellar medium), PAHs have been shown to incorporate oxygen into their molecular structure and display new functional groups such as alcohols, ethers, and ketones (Cook et al. 2015). The addition of a ketone moiety to the PAH is biologically significant, as this provides the base structure for the vitamin K chemical family. Incorporation of heteroatoms (such as oxygen and nitrogen) into the ring structure of the PAH as a result of the energetic processing of mixed icy grain mantles is also possible (Materese et al. 2015), and the resultant heterocyclic molecules are also of great biological significance with nitrogen-containing heterocycles such as pyrimidine and purine being the building blocks for nucleobases.

Pyrimidine has yet to be formally detected within the interstellar medium, however it possesses many similar physico-chemical characteristics to PAHs and so is expected to exist to some extent within icy grain mantles. The UV irradiation of frozen pyrimidine in the presence of H₂O, CH₃OH, NH₃, and CH₄ has been shown to yield the nucleobases uracil, cytosine, and thymine (Nuevo et al. 2009, 2012, 2014, Bera et al. 2016). Similarly, the irradiation of purine in ices containing NH₃ and H₂O leads to the formation of the nucleobases guanine and adenine (Bera et al. 2015, Materese et al. 2017, 2018). Although the idea of an extended lifetime for these nucleobases within interstellar and near-Earth environments has been called into question (Peeters et al. 2003), if there exists some plausible mechanism by which they survive the stellar evolution phases of the astrochemical cycle depicted in Figure 1, then it is possible that they may have been delivered to the early Earth via meteoritic or cometary impacts, thus providing the starting material for the so-called ‘RNA World Hypothesis’.

The RNA World Hypothesis (Orgel 2004, Bernhardt 2012, Neveu et al. 2013, Higgs and Lehman 2015) hypothesizes that RNA or RNA-like molecules mediated the necessary information processing and metabolic transformations required for life to emerge from the early Earth’s prebiotic environment. Systems based on RNA could act as a precursor to the significantly more complex system of RNA, DNA, and proteins on which current life is based, and it has been recognized that the ribonucleotide co-enzymes now used by many proteins may in fact be molecular fossils from an RNA-based metabolism (White 1976, Raffaelli 2011). The existence of naturally occurring ribozyme catalysts, such as self-splicing introns and the ribonuclease P catalyst (Kruger et al. 1982, Doudna and Cech 2002), along with the fact that ribosomal RNA actually catalyzes the formation of peptide bonds in the ribosome (Lang et al. 2008), has lent some weight to this hypothesis. However, much is still unknown as to the exact mechanisms by which life could arise in the RNA World, and so the hypothesis remains one among many seeking to explain the origins of biology from chemistry.

The ‘top-down’ aspect of PAH astrochemistry has also been considered by previous studies (Merino et al. 2014, Dartois et al. 2020). For instance, Merino et al. (2014) demonstrated experimentally that hydrogen atom interaction with graphene on top of SiC at elevated temperatures triggers an erosion process which results in the ejection of PAH-like flakes. Such results offer an alternative and efficient

synthetic pathway for the formation of gas-phase PAHs in the interstellar medium. In the envelopes of evolved stars, SiC condenses to form grains whereupon thermal annealing by the proximal star results in an elemental segregation and the synthesis of a carbon-rich surface (Frenklach et al. 1989, Merino et al. 2014). Exposure of this surface to hydrogen atoms is likely in this environment, and the elevated temperatures (>1000 K) permit a thermally assisted etching effect to take place, thus resulting in the desorption of PAHs.

The dissociation of PAHs themselves into smaller molecules is also a possible process, and has been invoked to explain the interstellar presence of fullerenes, particularly C_{60} . Experimental and modelling studies have shown that PAHs containing 60-66 carbon atoms may be completely dehydrogenated by UV photons to a graphene-like structure, after which the structure folds into a cage. Photon-induced shrinking of the structure then proceeds via the rate-limiting loss of C_2 units until the C_{60} fullerene structure is attained (Berné and Tielens 2012, Zhen et al. 2014, Berné et al. 2015). The need to further characterize this ‘top-down’ chemistry, particularly as it relates to other smaller carbon-based molecular species which may be relevant to prebiotic chemistry, was highlighted by van Dishoeck (2017).

2 The Use of THz/F-IR Spectroscopy in Astrochemistry

THz/F-IR frequencies cover the region of the electromagnetic spectrum ranging between 0.3-10 THz (10 - 333 cm^{-1}), although the upper boundary is somewhat arbitrary and some sources consider it to be as high as 30 THz (Dhillon et al. 2017). Since, by some popular definitions, the THz/F-IR range begins at a wavelength of 1 mm and proceeds to shorter wavelengths, it is sometimes also referred to as the submillimeter band. The main advantages of performing spectroscopy at such frequencies are the possibility of resolving rotational and ro-vibrational spectral transitions of gaseous molecular species, and the potential to probe long-range interactions between molecules in the solid phase in terms of low-energy intra- and intermolecular modes (Brown and Carrington 2003, Profeta and Scandolo 2011, Townes and Schawlow 2013). Thus, THz/F-IR spectroscopy makes for a unique tool in the detection of gases and ices in interstellar clouds and circumstellar disks against the widespread dust continuum via a number of distinct absorption and emission spectral signatures.

For example, the H_2O molecules (whose energy transitions are well-defined) is an asymmetric rotor with an irregular set of energy levels characterized by the rotational quantum number J and the pseudo-quantum numbers K_a and K_c (which replace the quantum number K used when considering symmetric rotors). These pseudo-quantum numbers represent hypothetical quantum states for the asymmetric rotor were it to assume the geometry of a prolate and oblate rotor, respectively (Cooke and Ohring 2012). Depending on the orientation of the nuclear spins of the constituent hydrogen atoms, the H_2O molecule is either ortho- (if the spins of the hydrogen atoms are parallel) or para- (if the spins are anti-parallel). Each excitation or de-excitation of electrons from one energy level to another results in the propagation of photons of a certain energy (i.e., frequency). Figure 4 shows the energy levels of ortho- and para- H_2O in terms of their pseudo-quantum numbers as detected by the Herschel Space Observatory. In this section, we provide a detailed description of the technical fundamentals of THz/F-IR spectroscopy, as well as an overview of its current applications to astrochemistry.




Figure 4 Energy level diagram of ortho- and para-H₂O showing the transitions observed by the PACS and HIFI instruments aboard the Herschel Space Observatory. Further information may be found in the work of van Dishoeck et al. (2013).

2.1 Spectrometer and Detector Technology

THz/F-IR laboratory spectroscopy measurements may be acquired using several different methods: Fourier-transform IR spectroscopy (Griffiths and de Haseth 1986), spectroscopy based on photo-mixing (Preu et al. 2011) or parametric conversion (Kawase et al. 1996, Kiessling et al. 2013), backward-wave oscillator spectroscopy (Medvedev et al. 2010), chirped-pulse Fourier-transform spectroscopy (Park and Field 2016), time-domain spectroscopy (Auston et al. 1975, van Exter et al. 1989, Mittleman et al. 1998), and cascaded frequency multiplication (Drouin et al. 2005, Li and Yao 2010). These methods are all based on the detection of a continuous wave, broadband source, or pulsed THz/F-IR radiation illuminating gaseous or solid (i.e., ice) molecular samples of a species, and having varying frequency spectral ranges, sensitivities, and performances. For example, chirped-pulse spectrometers have a frequency bandwidth of up to 100 GHz (Park and Field 2016), while backward oscillation and time-domain spectrometers have frequency ranges of up to 2.1 THz and 7.5 THz, respectively (Komandin et al. 2013, Allodi et al. 2014).

THz/F-IR time-domain spectroscopy can combine the generation of a THz/F-IR pulse in a zinc telluride crystal via opto-electronic rectification or from plasma filamentation in air using (in both cases) a femtosecond laser with a delay line and opto-electronic detection method in order to record the time domain trace (Allodi et al. 2013). This kind of spectroscopy has been widely used for gas-phase spectral line characterization in vacuum chambers (Kilcullen et al. 2015), in the study of refractive indices (Giuliano et al. 2019), and in the study of intermolecular transitions in astrophysical ice analogues, where the THz/F-IR signal is reflected by or transmitted through the gas or ice. Although it is able to provide critical information regarding the frequency of the spectral transition and the absorption coefficient, it does have a few disadvantages including slow acquisition times and a fine-tuning requirement (Trofimov and Varentsova 2016). As such, these instruments are appropriate only for laboratory settings, and other detectors are usually found in observatories.

One form of THz/F-IR spectroscopy which has found widespread use within laboratory astrochemistry settings is that based on cascaded frequency multiplications (Drouin et al. 2005, Li and Yao 2010). This technology is based on using a sweep synthesizer as the source of radiation, which is used in a

phase-locked continuous wave mode during measurements and which has a low phase-noise output. A filter (typically an yttrium iron garnet) is swept simultaneously to this synthesizer so as to suppress any frequency spurs. Frequency sextuplers allow for the production of millimeter radiation having a few milliwatts of power, which is then amplified via the use of microwave integrated circuit amplifiers to generate millimeter radiation with a few hundred milliwatts of power. Cascaded doublers or triplers then follow these amplifiers and generate the submillimeter (i.e., THz/F-IR) wavelength radiation.

Incoherent or direct detector arrays are commonly used in THz/F-IR observatories combined with multiplexer readout systems. They require a higher sensitivity than the photon background noise (such as the microwave background) and high pixel counts. The main technologies used are bolometers (exploiting the neutron transmutation doping process or implanted silicon), superconducting transition-edge-sensed bolometers (TESBs), and kinetic inductance detectors (KIDs) (Farrah et al. 2019). Prior to the development of TESBs and KIDs, bolometers operating at temperatures of 300 K or lower were used in observatories. Incoming radiation results in a change in the detector temperature which is read out as a change in its resistivity. Standard bolometers, however, are limited in their total pixel number which has led to the development and preferential use of other technologies, including TESBs and KIDs.

A TESB consists of a superconducting film operating near its superconducting transition temperature (Henderson et al. 2016, Thornton et al. 2016). The signal is seen as a current through a resistive film at a low temperature of around 100 mK (Suzuki et al. 2016). TESBs have high sensitivities which have allowed them to find use in many space-based instruments, as shown in Table 1. KIDs have a similar sensitivity to TESBs but operate below their superconducting transition temperature; below 300 mK (Griffin et al. 2016). The inductance of the detector material is increased due to broken Cooper pairs from incident photons on the superconducting film. In a resonant circuit, the shift in inductance causes a change in the resonant frequency which may be detected (Day et al. 2005).

Coherent systems used in astronomy are based on the heterodyne detection principle: the THz/F-IR high-frequency (RF) signals from molecules in space are down-converted to a lower frequency band typically of a few GHz known as the intermediate frequency (IF). The IF is easily measurable by sensitive back-ends operating at low frequency (such as power meters, spectrum analyzers, digital spectrometers, etc.). The device used for detection and down-conversion is a mixer which requires the input of a local oscillator (LO) signal at the same frequency as the studied RF signal. LO generators need to have narrow linewidths, low noise, high stability, and a broad tunability with sufficient output power to couple to the mixer. In the THz/F-IR domain, multiple chains based on Schottky diodes and quantum cascade lasers (>3 THz) are most commonly used (Valavanis et al. 2019).

There are three main types of mixers used in radio telescopes: superconductor-insulator-superconductor (SIS) diodes (Kojima et al. 2017), hot electron bolometers (HEBs) (Burke et al. 1999), and Schottky diodes (Maestrini et al. 2010). State of the art SIS diodes and HEBs offer the best system noise temperature performances at around 5-10 times the quantum noise limit. These mixers require only a few microwatts of LO power to operate and are very common for instruments with frequency ranges higher than 1 THz and between 1-6 THz for SIS diodes and HEBs, respectively. SIS diode mixers offer wider IF bandwidths and better sensitivities than do HEBs. However, SIS diodes require cooling to 4 K thus necessitating the use of a complex cryostat. On the other hand, Schottky mixers operate at temperatures above 70 K, a frequency range of up to 3 THz, and an IF bandwidth wider than 8 GHz. Their main disadvantages are their lower sensitivity which is between 30-50 times the quantum noise limit, and their milliwatt LO power requirement (Wilson et al. 2008).

The high-resolution spectroscopy possible when using a coherent detector in telescope facilities is achieved by using a digital technique with real-time fast Fourier-transform (FFT) so as to produce high-resolution spectral data. This idea was first proposed by Weinreb et al. (1961) and has been improved upon over the past 60 years in terms of acquisition speed, frequency bandwidth, power efficiency, cost, and size. Nowadays, three main digital systems are used in astronomy to sample the signal waveform at set time intervals: opto-acoustic spectrometers, digital autocorrelation spectrometers (Emrich 1997), and FFT spectrometers (Klein et al. 2012, Price 2016).

Opto-acoustic spectrometers are based on the diffraction of the signal on a Bragg crystal illuminated by a laser beam and detected by charge-coupled devices. This technique was used in the HIFI instrument aboard the Herschel Space Observatory (Siebertz et al. 2007). The main drawback of this spectrometer is its size, which defines its precision and spectral resolution. Digital autocorrelation spectrometers are based on the multiplication of a signal by a delayed version of the same signal using a series of delays. Spectrum measurement is achieved after the application of a Fourier transformation. Such spectrometers can have a bandwidth of a few GHz with a spectral resolution on the scale of one-hundredth of a kHz. Lastly, FFT spectrometers are based on field-programmable gate array (FPGA) chips. The FPGAs are combined with analogue-to-digital converters having a high data sampling rate of a few GHz samples per second. Their numerous spectral channels decompose the incoming RF signal into small sections giving an instantaneous FFT of a few kHz, as seen on the APEX telescope (Klein et al. 2006).

2.2 Telescope Facilities

Since the construction of the first radio telescope in 1937 (Kraus 1988), several telescopes operating in the THz/F-IR range have been successfully built and used by the astronomical community. A principal advantage of performing observations in the THz/F-IR range is the high sensitivity to low-abundance molecules within star-forming regions. Furthermore, observed emission and absorption features can be mapped for all areas within the stellar environment, such as snowlines where ice desorption processes begin to occur.

Progress in the sensitivity of detector technology over the past 50 years has permitted the investigation of interstellar molecular clouds at various wavelengths which has allowed us to improve our understanding of space chemistry. Powerful telescopes have been used with imaging and high spectral resolution instruments combined with large dish antennae. These telescopes have been instrumental in major scientific breakthroughs, such as the discovery of new molecules in the interstellar medium and elucidating the mechanisms for planetary system formation. A summary of the most important telescopes operating in the THz/F-IR range is given in Table 1.

In general, two varieties of observatory exist: ground-based (sometimes also referred to as atmosphere-based) and space-borne telescopes. Ground-based observatories are installed either on the ground at high altitudes or in airborne facilities. Atmospheric transmission has a direct impact on the sensitivity of these observatories at THz/F-IR frequencies since atmospheric molecules (especially H₂O) present strong absorption lines over these frequencies. As such, high atmospheric H₂O content limits atmospheric transmission, thus making observation heavily dependent upon weather conditions and altitude: drier sites to install ground-based telescopes are found at higher altitudes (Figure 5). Brief descriptions of the ground-based ALMA and space-borne Herschel Space Observatory now follow.

ALMA is a ground-based international astronomy facility built over an area of 6596 m² at an altitude of 5000 m located at Llano de Chajnantos in the Atacama Desert, Chile. The 66 high-precision antennae making up the facility are either 12 m (54 of them) or 7 m (12 of them) in diameter. These sizes,

combined with the area of the array, give a spatial resolution of 0.2-0.004 arcseconds (1 arcsecond = 1/3600 degrees). ALMA is composed of ten receivers using Schottky local oscillator sources and SIS diode mixers cooled to 4 K which give up to 16 GHz bandwidth with a spectral resolution of a few kHz. The covered frequency bands range from 31.3-950 GHz in ten windows (Figure 5) with a maximum 8 GHz IF. The lower frequency Bands 1 and 2 are not yet operational. The number of spectral channels available is 4096 per IF. For Bands 3-8, receivers operate in single sideband detection mode acquiring both H and V polarization while Bands 9 and 10 operate in double sideband mode.

Table 1 Examples of the sensitivities achieved by selected observatories operating in the THz/F-IR range.

Observatory	Aperture (m)	Instrument	Frequency Band (GHz)	Detector Technology
IRAM Telescope	30	NIKA2	150, 260	KIDs
CSO (Caltech Submillimeter Observatory)	10.4	MUSIC	142, 212, 272	KIDs
		SHARC II	666, 857	Bolometers
		MAKO	857	KIDS
		Heterodyne Receiver	117-920	SIS
		Z-Spec	190-230	Bolometers
JCMT (James Clerk Maxwell Telescope)	15	SCUBA	353, 666	Bolometers
		SCUBA2	353, 666	TES
		HARP	325-375	SIS
		Nāmakanuī	86, 215-371	SIS
ODIN	1.1	SMR	118-580	Schottky
SMA (Submillimeter Array)	6×6	Band 1-4	180-700	SIS
APEX (Acatama Pathfinder Experiment)	12	ArTeMis	666, 856	Bolometers
		SEPIA	157-738	SIS
		nFLASH	196-507	SIS
		CHAMP	620-850	SIS
		PI230	157-211	SIS
		LABOCA	345	Bolometers
NEOMA Interferometer	10×15	Band 1-4	71-371	SIS
Herschel Space Observatory	3.5	SPIRE	500-1500	Bolometers
		PACS bol.	1427-5000	Bolometers
		PACS phot.	1363-6000	Photoconductors
		HIFI	83-1000	Bolometers
SOFIA (Stratospheric Observatory for Infrared Astronomy)	2.5	HAWC+	1250-6000	TES
		FIFI-LS	1500-5880	Photoconductors
		GREAT	490-4745	HEB
LMT (Large Millimeter Telescope)	50	ToITEC	150, 220, 280	KIDs
		SEQUOIA	85-116	SIS
		AzTEC	272	Bolometers
		RSR	75-110	SIS
ALMA (Atacama Large Millimeter/Sub- millimeter Array)	54×12 12×7	Band 3-10	84-950	SIS
Millimetron	10	Band 1-2	33-116	HEMT
		Band 3-5	211-650	SIS
OST (Origins Space Telescope)	5.9 to 9.1	Imaging	1000-3000	-
		Spectroscopy	1000-3000	-




Figure 5 Zenith atmospheric transmission of H₂O vapor of the ALMA telescope with precipitable H₂O vapors of 0.5 mm simulated from the *Atmospheric Transmission at Microwaves* model described by Pardo et al. (2001). Further information may be found in the work of Maiolino (2008).

Space-based telescopes are not limited by atmospheric attenuations in the way that ground-based observatories are, but do have limitations connected with the size of their antenna aperture. The Herschel Space Observatory was commissioned by the European Space Agency and launched in 2009. The telescope was composed of a 3.5 m diameter dish which was passively cooled by liquid helium at 4 K. Over time, the liquid helium reserves necessary for detector cooling were gradually depleted, and the end of the telescope's operations came about in April 2013.

Three scientific instruments were on board the Herschel Space Observatory: a high-spectral resolution heterodyne **spectrometer** called HIFI (Heterodyne Instrument for the Far-Infrared), and imaging photometer called PACS (Photodetector Camera Array), and a medium resolution grating **spectrometer** called SPIRE (Spectral and Photometric Imaging Receiver). THz/F-IR frequencies were covered by the HIFI instrument from 408-1908 GHz separated in 12 windows using HEB and SIS diode mixer technology. PACS was composed of a camera integral field **spectrometer** combined with an imaging photometer operating between 1.427-6 THz and bolometers at frequencies centered around 1.873 THz, 2.997 THz, and 4.282 THz (Rosenthal et al. 2002). Finally, the SPIRE instrument had three main photometers at 600 GHz, 856 GHz, and 119 GHz, as well as a two-band imaging Fourier-transform **spectrometer** with bolometers operating between 447-1550 GHz (Griffin et al. 2010).

Amongst the key scientific objectives of the Herschel Space Observatory was an investigation of the formation of stellar and planetary systems, and increasing our understanding of the physics and chemistry of the interstellar medium, our Solar System, and extra-galactic galaxies. The HIFI instrument detected more than 100,000 spectral features in a single spectrum (Bergin et al. 2010). Through the Water in Star-Forming Regions with Herschel (WISH) program, many molecules such as H₂O, CO, and O₂ were detected in various astrophysical environments (Goldsmith et al. 2011).

Molecules and radicals including H_2O and OH^- were detected further out and deeper in proto-planetary disks than before from surveys of THz/F-IR spectral lines using PACS (Fedele et al. 2013). Cold water reservoirs were discovered in two extra-solar planet-forming disks (TW Hydrae) through the detection of H_2O gas produced by UV photo-desorption of the ice made by HIFI (Hogerheijde et al. 2011). H_2O vapor was also detected in a pre-stellar core at the early stage of stellar formation using HIFI observations combined with models of the UV photo-desorption processes (Caselli et al. 2012).

Ground-based and space-borne observatories operating in the THz/F-IR range, including those listed in Table 1, have aided astronomers in elucidating the chemistry of interstellar space by allowing them to obtain spectroscopic information of molecular species (colloquially termed ‘molecular fingerprints’). Indeed, the improvement of technology throughout these past decades has allowed for the definitive identification of many new molecular signatures, largely through observation of their rotational emission spectra. Figure 6 shows the comparison of spectra acquired by ALMA Band 10 and the HIFI instrument aboard the Herschel Space Observatory while observing the NGC 6334I region (McGuire et al. 2018b). As can be seen, the number of spectral lines attributed to CO and CH_3OH which are resolved by ALMA Band 10 is significantly higher than that detected by HIFI (Zernickel et al. 2012).

Computer modelling and laboratory measurements of molecular spectra are often needed to assign observed spectral lines to particular molecular species. This is especially true in the case of complex organic molecules such as isopropyl cyanide ($(\text{CH}_3)_2\text{CHCN}$), which was the first branched carbon-chain molecule to be detected and which was found in Sgr B2 using the ALMA Band 3 (Belloche et al. 2014). Later, $\text{CH}_3\text{OCH}_2\text{OH}$ and CH_3Cl were observed for the first time by ALMA Bands 6 and 7 in NGC 6334I, and by ALMA Band 7 in IRAS 16293-2422, respectively (Fayolle et al. 2017, McGuire et al. 2017). However, many molecules and radical species may not be observed in the gaseous phase but may play a central role in the formation of observed gaseous species as a result of their presence and reactions in solid icy grain mantles. The use of laboratory THz/F-IR measurements in astrochemical research, particularly as it pertains to solid-phase chemistry, will be elaborated on in the next sub-section.




Figure 6 Comparison of the spectra acquired by ALMA Band 10 and HIFI while observing the NGC 6334I region. Figure taken from McGuire et al. (2018b) and reproduced with permission from the AAS.

2.3 Laboratory Astrochemistry Using THz/F-IR Spectroscopy

Over 200 molecules have been detected and spectroscopically resolved in interstellar regions within 601 our galaxy, and more than 60 have been observed in extra-galactic molecular clouds (Müller et al.

2005), ranging in complexity from simple diatomics to the more structurally complex fullerenes and PAHs. An overarching aim of laboratory astrochemistry is to be able to understand the physico-chemical processes which occur in these astrophysical environments so as to better comprehend the chemical reactions which lead to the formation of such molecules. Such work is motivated by the knowledge that such reactions can result in the production of various molecules and minerals. For instance, interstellar biomolecules which may form in dense, quiescent molecular clouds may be subsequently incorporated into the stellar and planetary systems which evolved from these clouds and thus play a defining role in the emergence of life within these systems (Watanabe et al. 2002, Jones et al. 2011, Fedoseev et al. 2017, Sandford et al. 2020, Chuang et al. 2020, Ioppolo et al. 2021). The synthesis of smaller, volatile molecules in astrophysical environments is equally important, as such molecules may contribute to the development of planetary and lunar atmospheres or transient exospheres (Milillo et al. 2016, Teolis and Waite 2016). Additionally, condensation processes and grain alterations contribute to the formation of minerals which are significant to the geological and geochemical evolution of planets and moons, including graphite, corundum, moissanite, and forsterite (Frenklach et al. 1989, Hazen et al. 2008, Hazen and Ferry 2010).

The unambiguous identification of such molecules by infrared and radio telescopes (including those which operate in the THz/F-IR range) relies on the availability and completeness of gas- and solid-phase spectroscopic databases at the corresponding frequencies for a great variety of molecules. Such information is obtained by performing classical laboratory measurements (Włodarczak 1995, Smith 2011, Allodi et al. 2013, Cataldo et al. 2013a, Widicus-Weaver 2019, Mifsud et al. 2021). Presently, the two most commonly referenced catalogues are the Cologne Database for Molecular Spectroscopy (Müller et al. 2001, Endres et al. 2016) and the Jet Propulsion Laboratory (Pickett et al. 1998) catalogues, which give the frequency and amplitude of several molecular transitions. Two accessible databases listing vibrational transitions at IR wavelengths are also available: the HITRAN database (Rothman et al. 2009) and the EXOMOL line list.

Broadly speaking, laboratory astrochemistry studies are largely concerned with investigating reactions which occur in either the gas phase or the condensed (ice) phase. Experiments are typically performed using high or ultra-high vacuum chambers. For instance, a novel method of performing gas-phase spectroscopy is to make use of a heterodyne radiometer so as to observe the absorption features of various gaseous species within the chamber, thus functioning in a similar fashion to several telescope facilities. Two laboratory broadband emission THz/F-IR radiometers based on heterodyne detection methods have been developed: the first is based on Schottky-barrier technology combined with a FFT digital spectrometer operating between 80-110 GHz (Wehres et al. 2017), while the second is based on cryo-cooled SIS diode mixer technology operating between 270-290 GHz (Wehres et al. 2018). By using such instruments, the spectroscopic features of pyridine and CH₃CN have been successfully measured and matched with analytical simulations. Similarly, a Schottky-based receiver has been used in combination with a FFT digital spectrometer and vacuum chamber to observe the emission line of N₂O at 355.6 GHz (Parkes et al. 2018).

Laboratory experiments concerned with the solid phase also require the use of high or ultra-high vacuum chambers, wherein interstellar ice analogues may be synthesized on cold substrates via the direct or background deposition of dosed gases or vapors, or the effusive evaporation of refractories. The deposited ices are then processed in such a way as to simulate astrophysical conditions: atom (or free radical) additions represent non-energetic pathways towards molecule formation (Linnartz et al. 2015, Chuang et al. 2016, Potapov et al. 2017, Ioppolo et al. 2021); bombardment with energetic charged particles (i.e., electrons and ions) simulates ice interactions with galactic cosmic rays, the solar wind, or planetary magnetospheric plasmas (Baragiola et al. 1999, Dalton et al. 2013, Ding et al. 2013,

Boamah et al. 2014, Boduch et al. 2015, Fulvio et al. 2019, Ioppolo et al. 2020); irradiation with 649 photons (both ionizing and non-ionizing) simulates extra-terrestrial photochemistry (Bernstein et al. 650 2012, Lo et al. 2014, Öberg 2016, Mullikin et al. 2018); energetic shocks simulate collisions between 651 interstellar or Solar System bodies (Goldman et al. 2010, Martins et al. 2013), and thermal processing 652 simulates the chemical and structural changes induced within interstellar icy grain mantles proximal to 653 nascent stars, or in Solar System bodies as they approach perihelion (Ehrenfreund et al. 1999, 654 Kaňuchová et al. 2017).

The changes in the physico-chemical properties of the ice analogues induced by such processing have 656 been traditionally monitored using mid-IR or UV-VIS spectroscopy (Mason et al. 2006, Boersma et al. 657 2014). This has been advantageous in that such laboratory work is complementary to observational 658 work using space-borne telescopes working within these ranges of the electromagnetic spectrum, such 659 as the Spitzer Space Telescope and the Hubble Space Telescope. Although not yet widely used within 660 laboratory settings, the utility and applicability of THz/F-IR spectroscopy is being increasingly 661 recognized due to its ability to detect lower frequency molecular vibrations, as well as intermolecular 662 interactions and translational vibrations within solid lattices (Bertie 1968, Bertie and Jacobs 1977, 663 Kulesa 2011, McIntosh et al. 2012). The remainder of this sub-section will be devoted to a discussion 664 on the application of THz/F-IR spectroscopy to laboratory studies of astrophysical ice analogues.

THz/F-IR spectra differ depending upon the structural morphology of the ice (e.g., amorphous or 666 crystalline solid phases), as demonstrated by Moore and Hudson (1992, 1994, 1995). For instance, 667 recent studies investigating CO₂ in H₂O and CH₃OH ice matrices at various temperatures have revealed 668 that the measured THz/F-IR spectra are especially sensitive to the degree of segregation within the ice 669 structure. Results have shown that CO₂ roaming during warming of the ice results in a localized 670 accumulation, or segregation. When mixed with crystalline H₂O ice, this CO₂ segregation results in a 671 disruption of the hydrogen bonding network between adjacent H₂O molecule layers, causing the H₂O 672 spectral resonance features to be shifted, distorted, or attenuated (Allodi et al. 2014). On the other hand, 673 CO₂ absorption bands were observed to become narrower and more defined when segregation from 674 CH₃OH ice occurred as a result of warming (McGuire et al. 2016).

Such work clearly demonstrates the sensitivity of THz/F-IR spectroscopy not only to the nature of the 676 chemical species present within the ice, but also to the physical structure of the ice. Given that the 677 nature of the observed spectral absorption features is dependent upon the temperature, degree of 678 crystallinity, segregation, and thermal history of the ice, THz/F-IR spectroscopy appears to be highly 679 suitable for assessing not only the composition and temperature characteristics of astrophysical ices, 680 but also their morphology (Ioppolo et al. 2014). To date, relatively few of the observed molecules have 681 been investigated using THz/F-IR spectroscopy. Amorphous solid H₂O, for instance, presents a 682 spectral line at 1 THz. CH₃OH, CH₃CHO, and (CH₃)₂CO have been found to have a broad feature at 683 around 4 THz, while in HCOOH and CH₃COOH this broad absorption feature is centered at about 7 684 THz (Ioppolo et al. 2014).

Laboratory THz/F-IR spectroscopy has also been used in the analysis of more complex molecular 686 structures such as PAHs. Although these molecules are believed to be ubiquitous in interstellar and 687 circumstellar media (Tielens 2008, Kwok and Zhang 2011), few single structures have been positively 688 identified. Systematic analysis of a series of PAHs, as well as their hydrogenated and alkylated 689 derivatives, has revealed that the THz/F-IR spectra of these molecules display so-called 'Jumping Jack' 690 modes which correspond to the in-plane vibrations around the central molecular core (Cataldo et al. 691 2013b). As such, these spectra provide crucial information related to the number of fused aromatic 692 rings present within the molecule, and are thus valuable in identifying the individual molecular carriers

of PAH diffuse interstellar absorption bands. Other complex molecules, including those which may be 694 of relevance to prebiotic chemistry, have also been studied using THz/F-IR spectroscopy (Widicus- 695 Weaver et al. 2005, Carroll et al. 2010).

Recently, a new laboratory-based method for observing desorption mechanisms occurring within the 697 interstellar medium and star-forming regions has been developed. Studying such desorption has been 698 traditionally performed using mid-IR spectrophotometers and quadrupole mass spectrometers, 699 however this has the drawback of not being able to detect ions and radicals and is difficult to relate to 700 observations made using radio telescopes (Fraser et al. 2002, Collings et al. 2003, Fuchs et al. 2009, 701 Muñoz-Caro et al. 2010). In principle, it is possible to desorb (either via thermal or non-thermal 702 mechanisms) ices formed of pure or layered molecules and observe the resultant gas phase in the 703 THz/F-IR frequency range. One way to do this is to passively observe the emission of those molecules 704 during desorption in an astrochemical chamber equipped with a heterodyne radiometer (Auriacombe 705 et al. 2015, 2016); another is by performing microwave absorption spectroscopy. The region in which 706 desorbing molecules are observed may be directly above the ice, and the relevant spectral transitions 707 may be detected with a HEB (see Fig. 2 in Yocum et al. 2019). Alternatively, a chirped-pulse Fourier- 708 transform **spectrometer** can be used to observe the desorption occurring inside a waveguide (Theulé et 709 al. 2020). These emerging techniques are capable of bringing new insights and possibilities to our 710 understanding of the desorption processes in star-forming regions.

Although the use of THz/F-IR spectroscopy is growing within laboratory-based astrochemical research 712 (driven by developments in the relevant technologies and an increase in the commercial availability of 713 the necessary **spectrometers**), much of the current work takes place at large-scale facilities. A complete 714 and thorough description of the research conducted using THz/F-IR radiation at such facilities would 715 go beyond the scope of this review, and we instead direct the interested reader to other works (e.g., 716 Neil 2014). However, it is worthwhile to take some time to briefly describe the Free Electron Lasers 717 for Infrared Experiments (FELIX) facility located in Nijmegen, the Netherlands, where an ultrahigh- 718 vacuum chamber has been installed as the Laboratory Ice Surface Astrophysics (LISA) end-station 719 thus permitting the characterization of astrophysical ice analogues which have been processed as a 720 result of THz/F-IR irradiation.

The FELIX facility is home to four free electron lasers which provide short-pulsed coherent light 722 covering the microwave to mid-IR regions of the electromagnetic spectrum. The FELIX-1 and FELIX- 723 2 free electron laser beamlines respectively operate over the $\sim 65\text{-}330\text{ cm}^{-1}$ and $\sim 225\text{-}3300\text{ cm}^{-1}$ ranges 724 and are capable of providing bursts of micro-pulses with selective and tunable wavelengths whose 725 duration may be readily controlled. These beamlines are therefore highly suited to providing THz/F- 726 IR (as well as mid-IR) radiation for use in laboratory studies of astrophysical ice analogues, which are 727 prepared in the LISA end-station.

The LISA is an ultrahigh-vacuum chamber with nominal base pressure in the region of a few 10^{-10} 729 mbar containing an elongated rectangular gold-coated deposition substrate which may be cooled down 730 to 15 K using a closed-cycle helium cryostat head with a cooled compressor. The gold-coated substrate 731 is manipulated by means of a xyz-linear translator allowing for the irradiation of different spots of the 732 ice during a single experiment, as well as a rotary stage mostly employed for alignment purposes. Pure 733 gases and gas mixtures are pre-prepared in a dosing line prior to being injected into the main chamber 734 via an all-metal leak valve, wherein background deposition of the gases occurs to form the 735 astrophysical ice analogues. Such background deposition of gases onto the deposition substrate allows 736 for a more uniform ice deposition, and the resultant ice analogue may be used in systematic studies. 737 Ices processed by mid-IR or THz/F-IR irradiation delivered by the FELIX beamlines may be monitored

in two ways: mid-IR reflection-absorption spectroscopy may be used to monitor any physico-chemical effects induced as a result of the ice stimulation, and quadrupole mass spectrometry may be used to monitor species which desorb thermally from the bulk parent ice.

The FELIX-LISA facility has found regular use in effectively characterizing the changes induced in astrophysical ice analogues as a result of stimulation by incident irradiation. For instance, Noble et al. (2020) investigated the effect of resonant irradiation of amorphous solid H₂O using mid-IR radiation supplied by the FELIX-2 beamline. Their results indicated that irradiation with wavelengths corresponding to the mid-IR stretching, bending, and libration modes of amorphous H₂O resulted in a wide-ranging structural rearrangement of the ice into a crystalline-like form due to vibrational relaxation of the intermolecular hydrogen bonding network. Results such as these, combined with the experimental capabilities of large-scale facilities such as FELIX, make future dedicated studies looking into the irradiation of lower frequency THz/F-IR modes of astrophysical ice analogues especially attractive.

As has been demonstrated, the application of THz/F-IR spectroscopy to observational and laboratory astrochemistry has the potential to unravel much information related to the structure and reactivity of molecules in space. To date, however, THz/F-IR experiments at large-scale facilities such as FELIX tasked with investigating the chemistry of interstellar and Solar System ice analogues have not been explored to their fullest potential due to the fact that the dedicated pump-probe experiments required to study such chemistry are not available at the time of writing. The present LISA end-station configuration comprises an FTIR spectrophotometer with an extended spectral range to the far-IR. Such a system may be used to time-resolve transient events within the ice structure (such as diffusion of molecules through different ice layers and segregation events in mixed ices) at millisecond timescales.

To fully benefit from the unique capabilities of a free electron laser (i.e., a wide-range tunability, high peak power, and a controllable repetition rate), THz/F-IR time-domain spectroscopy with single-shot detection techniques should be employed to extend time resolutions to a few tens of picoseconds (Redlich et al. 2003, Perakis et al. 2013, Shalit et al. 2013, Mead et al. 2019). With such a configuration, free electron laser THz/F-IR radiation could be used to ‘pump’ (i.e., inject energy into) inter- and intramolecular vibrations of solid species that, having been excited by the sudden excess of energy, will subsequently rearrange selectively or diffuse within the ice and possibly react with other species. Diffusion and reaction of molecules could be triggered and controlled by operating a free electron laser across its spectral range and could then be monitored by means of a THz/F-IR single-shot spectrometer in the solid phase, while desorbed species could be monitored by implementing a broadband emission THz/F-IR radiometer as described previously. The combination of a free electron laser and new advanced techniques in laboratory astrochemistry thus has the potential to reveal unprecedented details on fundamental phenomena which could play important roles in the formation of complex prebiotic molecules in space, and would thus expand our understanding of the relevant extra-terrestrial physics and chemistry.

Experimental work such as that proposed to be performed at the FELIX-LISA facility on broadband emission THz/F-IR radiometry would also contribute much to our interpretation of observational data. The importance of commissioning new space-borne observational facilities operating in the THz/F-IR was highlighted in the recently published *Pathways to Discovery in Astronomy and Astrophysics for the 2020s* roadmap (National Academies of Sciences, Engineering, and Medicine 2021), particularly in light of the fact that the Space Infrared Telescope for Cosmology and Astrophysics (SPICA) is no longer being considered for launch. Interestingly, the far-IR is one of the spectral ranges where

astronomical data with sub-arcsecond resolution is not currently available: a fact which is somewhat 784 at odds with the potential usefulness of spectroscopic measurements in this range (Linz et al. 2021). A 785 number of space-based observatories, such as the Far-Infrared Spectroscopic Surveyor (FIRSS) and 786 the Terahertz Exploration and Zooming-In for Astrophysics (THEZA) telescopes have also been 787 proposed to fill in this gap (Gurvits et al. 2021, Rigopoulou et al. 2021), and the acquisition of 788 laboratory reference data will doubtlessly aid in further characterizing the molecular composition of 789 the cosmos.

3 Conclusions and Recommendations for Future Work

In summary, we have provided a brief overview of the chemistry occurring in both diffuse and dense 792 (dark) interstellar clouds with a focus on ion-molecule reactions in the gas phase which are relevant to 793 the formation of complex prebiotic molecules, as well as how THz/F-IR techniques have aided in the 794 elucidation of these reactions and how such reactions may relate to the RNA World Hypothesis on the 795 abiogenic origins of life. We have also provided a review on current THz/F-IR **spectrometer** and 796 detector technology in both laboratory and observatory settings, as well as their present and potential 797 future applications to **laboratory solid-phase** astrochemistry. We conclude by emphasizing that further 798 progress in THz/F-IR astrochemistry has the potential to provide great insight to several fundamental 799 astrochemical processes and should therefore be pursued.

4 Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial 802 relationships that could be construed as a conflict of interest.

5 Author Contributions

DVM, PAH, ATM, OA, and SI wrote the manuscript and all authors were responsible for corrections 805 and improvements.

6 Funding

Our research has benefitted from support provided by the Europlanet 2024 RI, which has received 808 funding from the European Union's Horizon 2020 Research Innovation Program under grant 809 agreement No. 871149.

7 Acknowledgments

DVM is the grateful recipient of a University of Kent Vice-Chancellor's Research Scholarship. ATM 812 thanks Queen Mary University of London for doctoral funding. SI acknowledges the Royal Society for 813 financial support.

8 ORCID ID Numbers

Duncan V. Mifsud:	0000-0002-0379-354X
Perry A. Hailey:	0000-0002-8121-9674
Alejandra Traspas Muiña:	0000-0002-4304-2628
Olivier Auriacombe:	0000-0002-5810-8650

Sergio Ioppolo: 0000-0002-2271-1781

Nigel J. Mason: 0000-0002-4468-8324

9 References

- 822 Allodi, M.A., Baragiola, R.A., Baratta, G.A., Barucci, M.A., Blake, G.A., Boduch, P., Brucato, J.R., Contreras, C., Cuyllé,
823 S.H., Fulvio, D., Gudipati, M.S., Ioppolo, S., Kaňuchová, Z., Lignell, A., Linnartz, H., Palumbo, M.E., Raut, U.,
824 Rothard, H., Salama, F., Savchenko, E.V., Sciamma-O'Brien, E., Strazzulla, G. (2013). Complementary and
825 Emerging Techniques for Astrophysical Ices Processed in the Laboratory. *Space Sci. Rev.* 180, 101.
- 826 Allodi, M.A., Ioppolo, S., Kelley, M.J., McGuire, B.A., Blake, G.A. (2014). The structure and dynamics of carbon dioxide
827 and water containing ices investigated via THz and mid-IR spectroscopy. *Phys. Chem. Chem. Phys.* 16, 3442.
- 828 Arumainayagam, C.R., Garrod, R.T., Boyer, M.C., Hay, A.K., Bao, S.T., Campbell, J.S., Wang, J., Nowak, C.M., 829
830 Arumainayagam, M.R., Hodge, P.J. (2019). Extraterrestrial prebiotic molecules: Photochemistry vs. radiation
chemistry of interstellar ices. *Chem. Soc. Rev.* 48, 2293.
- Auriacombe, O., Fraser, H., Ellison, B., Rea, S. (2015). Desorption emission spectroscopy using THz radiometry (THz
DES). *IET Conference Proceedings*, 3.
- Auriacombe, O., Fraser, H., Ellison, B., Ioppolo, S., Rea, S. (2016). Terahertz desorption emission spectroscopy (THz DES)
– ALMA in the lab. Paper presented at the American Astronomical Society Meeting 228, 104.03.
- Auston, D.H. (1975). Picosecond optoelectronic switching and gating in silicon. *Appl. Phys. Lett.* 26, 101.
- Balucani, N., Ceccarelli, C., Taquet, V. (2015). Formation of complex organic molecules in cold objects: The role of gas-
phase reactions. *Mon. Not. R. Astron. Soc.* 449, L16.
- Baragiola, R.A., Atteberry, C.L., Bahr, D.A., Jakas, M.M. (1999). Solid-state ozone synthesis by energetic ions. *Nucl.*
Instrum. Methods Phys. Res. B 157, 233.
- Barinovs, Ģ., van Hemert, M.C. (2006). CH⁺ radiative association. *Astrophys. J.* 636, 923.
- Belloche, A., Garrod, R.T., Müller, H.S.P., Menten, K.M. (2014). Detection of a branched alkyl molecule in the interstellar
medium: iso-propyl cyanide. *Science* 345, 1584.
- Belloche, A., Müller, H.S.P., Garrod, R.T., Menten, K.M. (2016). Exploring molecular complexity with ALMA (EMoCA):
Deuterated complex organic molecules in Sagittarius B2(N2). *Astron. Astrophys.* 587, 91.
- Belloche, A., Garrod, R.T., Müller, H.S.P., Menten, K.M., Medvedev, I., Thomas, J., Kisiel, Z. (2019). Re-exploring
molecular complexity with ALMA (ReMoCA): Interstellar detection of urea. *Astron. Astrophys.* 628, 10.
- Bera, P.P., Nuevo, M., Materese, C.K., Sandford, S.A., Lee, T.J. (2016). Mechanisms for the formation of thymine under
astrophysical conditions and implications for the origin of life. *J. Chem. Phys.* 144, 144308.
- Bera, P.P., Stein, T., Head-Gordon, M., Lee, T.J. (2017). Mechanisms of the formation of adenine, guanine and their
analogues in UV-irradiated mixed NH₃:H₂ ices containing purine. *Astrobiology* 17, 771.
- Bergin, E.A., Tafalla, M. (2007). Cold dark clouds: The initial conditions for star formation. *Annu. Rev. Astron. Astrophys.*
852 45, 339.
- 853 Bergin, E.A., Phillips, T.G., Comito, C., Crockett, N.R., Lis, D.C., Schilke, P., Wang, S., Bell, T.A., Blake, G.A., Bumble, 854
B., Caux, E., Cabrit, S., Ceccarelli, C., Cernicharo, J., Daniel, F., de Graauw, T., Dubernet, M.L., Emprechtinger,
855 M., Encrenaz, P., Falgarone, E., Gerin, M., Giesen, T.F., Goicoechea, J.R., Goldsmith, P.F., Gupta, H., Hartogh, P.,
856 Helmich, F.P., Herbst, E., Joblin, C., Johnstone, D., Kawamura, J.H., Langer, W.D., Latter, W.B., Lord, S.D., Maret,
857 S., Martin, P.G., Melnick, G.J., Menten, K.M., Morris, P., Müller, H.S.P., Murphy, J.A., Neufeld, D.A., Ossenkopf,

- 858 V., Pagani, L., Pearson, J.C., Péroult, M., Plume, R., Roelfsema, P., Qin, S.L., Salez, M., Schlemmer, S., Stutzki, J.,
859 Tielens, A.G.G.M., Trappe, N., van der Tak, F.F.S., Vastel, C., Yorke, H.W., Yu, S., Zmuidzinas, J. (2010). Herschel
860 observations of extra-ordinary sources (HEXOS): The present and future of spectral surveys with Herschel/HIFI.
861 *Astron. Astrophys.* 521, L20.
- 862 Berné, O., Tielens, A.G.G.M. (2012). Formation of buckminsterfullerene (C₆₀) in interstellar space. *Proc. Nat. Acad. Sci.*
863 *USA* 109, 401.
- Berné, O., Montillaud, J., Joblin, C. (2015). Top-down formation of fullerenes in the interstellar medium. *Astron. Astrophys.* 577, A133.
- Bernhardt, H.S. (2012). The RNA world Hypothesis: The worst theory of the early evolution of life (except for all the others). *Biol. Direct* 7, 23.
- Bernstein, M.P., Dworkin, J.P., Sandford, S.A., Cooper, G.W., Allamandola, L.J. (2002). Racemic amino acids from the ultraviolet photolysis of interstellar ice analogues. *Nature* 416, 401.
- Bertie, J.E. (1968). Far-infrared spectra of the ices. *Appl. Spectrosc.* 22, 638.
- Bertie, J.E., Jacobs, S.M. (1977). Far-infrared absorption by ices Ih and Ic at 4.3 K and the powder diffraction pattern of ice Ic. *J. Chem. Phys.* 67, 2445.
- Boamah, M.D., Sullivan, K.K., Shulenberger, K.E., Soe, C.M., Jacob, L.M., Yhee, F.C., Atkinson, K.E., Boyer, M.C., 874
Haines, D.R., Arumainayagam, C.R. (2014). Low-energy electron-induced chemistry of condensed methanol: 875
Implications for the interstellar synthesis of prebiotic molecules. *Faraday Discuss.* 168, 249.
- 876 Boduch, P., Dartois, E., de Barros, A.L.F., da Silveira, E.F., Domaracka, A., Lv, X.Y., Palumbo, M.E., Pilling, S., Rothard,
877 H., Seperuelo Duarte, E., Strazzulla, G. (2015). Radiation effects in astrophysical ices. *J. Phys. Conf. Ser.* 629,
878 012008.
- 879 Boersma, C., Bauschlicher, C.W., Ricca, A., Mattioda, A.L., Cami, J., Peeters, E., Sánchez de Armas, F., Puerta Saborido, 880
G., Hudgins, D.M., Allamandola, L.J. (2014). The NASA Ames PAH IR spectroscopic database version 2.00:
881 Updated content, web site, and on(off)line tools. *Astrophys. J. Suppl.* 211, 8.
- 882 Brown, P.J., Carrington, A. (2003). *Rotational Spectroscopy of Diatomic Molecules*. Cambridge: Cambridge University
883 Press.
- 884 Burke, P.J., Schoelkopf, R.J., Prober, D.E., Skalare, A., Karasik, B.S., Gaidis, M.C., McGrath, W.R., Bumble, B., LeDuc, 885
H.G. (1999). Mixing and noise in diffusion and phonon cooled superconducting hot-electron bolometers. *J. Appl.*
886 *Phys.* 85, 1644.
- 887 Burkhardt, A.M., Lee, K.L.K., Changala, P.B., Shingledecker, C.N., Cooke, I.R., Loomis, R.A., Wei, H., Charnley, S.B., 888
Herbst, E., McCarthy, M.C., McGuire, B.A. (2021a). Discovery of the pure polycyclic aromatic hydrocarbon indene
889 (*c*-C₉H₈) with GOTHAM observations of TMC-1. *ArXiv [Preprint]*. Available at: <https://arxiv.org/abs/2104.15117>
890 (Accessed May 16, 2021).
- 891 Burkhardt, A.M., Loomis, R.A., Shingledecker, C.N., Lee, K.L.K., Remijan, A.J., McCarthy, M.C., McGuire, B.A.
892 (2021b). Ubiquitous aromatic carbon chemistry at the earliest stages of star formation. *Nat. Astron.* 5, 181.
- 893 Calcutt, H., Willis, E.R., Jørgensen, J.K., Bjerkeli, P., Ligterink, N.F.W., Coutens, A., Müller, H.S.P., Garrod, R.T., 894
Wampfler, S.F., Drozdovskaya, M.N. (2019). The ALMA-PILS survey: propyne (CH₃CCH) in IRAS 16293-2422.
895 *Astron. Astrophys.* 631, 137.
- 896 Carroll, P.B., Drouin, B.J., Widicus-Weaver, S.L. (2010). The submillimeter spectrum of glycolaldehyde. *Astrophys. J.*
897 723, 845.

- 898 Casselli, P., Keto, E., Bergin, E.A., Tafalla, M., Aikawa, Y., Douglas, T., Pagani, L., Yıldız, U.A., van der Tak, F.F.S., 899
900 Walmsley, C.M., Codella, C., Nisini, B., Kristensen, L.E., van Dishoeck, E.F. (2012). First detection of water vapor
in a pre-stellar core. *Astrophys. J. Lett.* 759, L37.
- 901 Cataldo, F., García-Hernández, D.A., Manchado, A. (2013a). Far- and mid-infrared spectroscopy of complex organic matter
902 of astrochemical interest: Coal, heavy petroleum fractions and asphaltene. *Mon. Not. R. Astron. Soc.* 429, 3025.
- 903 Cataldo, F., Angelini, G., García-Hernández, D.A., Manchado, A. (2013b). Far-infrared (terahertz) spectroscopy of a series
904 of polycyclic aromatic hydrocarbons and application to structure interpretation of asphaltene and related
905 compounds. *Spectrochim. Acta A* 111, 68.
- 906 Chuang, K.J., Fedoseev, G., Ioppolo, S., van Dishoeck, E.F., Linnartz, H. (2016). H-atom addition and abstraction reactions
907 in mixed CO, H₂CO and CH₃OH ices – an extended view on complex organic molecule formation. *Mon. Not. R.*
908 *Astron. Soc.* 455, 1702.
- 909 Chuang, K.J., Fedoseev, G., Qasim, D., Ioppolo, S., Jäger, C., Henning, T., Palumbo, M.E., van Dishoeck, E.F., Linnartz, 910
911 H. (2020). Formation of complex molecules in translucent clouds: acetaldehyde, vinyl alcohol, ketene, and ethanol
via ‘non-energetic’ processing of C₂H₂ ice. *Astron. Astrophys.* 635, 199.
- 912 Collings, M.P., Dever, J., Fraser, H., McCoustra, M.R.S. (2003). Laboratory studies of the interaction of carbon monoxide
913 with water ice. *Astrophys. Space Sci.* 285, 663.
- 914 Cook, A.M., Ricca, A., Mattioda, A.L., Bouwman, J., Roser, J., Linnartz, H., Bregman, J., Allamandola, L.J. (2015). 915
916 Photochemistry of polycyclic aromatic hydrocarbons in cosmic water ice: The role of PAH ionization and
concentration. *Astrophys. J.* 799, 14.
- 917 Cooke, S.A., Ohring, P. (2012). Decoding pure rotational molecular spectra for asymmetric molecules. *J. Spectrosc.* 2013,
918 698392.
- Cuadrado, S., Goicoechea, J.R., Cernicharo, J., Fuente, A., Pety, J., Tercero, B. (2017). Complex organic molecules in
strongly UV-irradiated gas. *Astron. Astrophys.* 603, 124.
- Dalton, J.B., Cassidy, T., Paranicas, C., Shirley, J.H., Prockter, L.M., Kamp, L.W. (2013). Exogenic controls on sulfuric
acid hydrate production at the surface of Europa. *Planet. Space Sci.* 77, 45.
- Dartois, E., Charon, E., Engrand, C., Pino, T., Sandt, C. (2020). Mechanochemical synthesis of aromatic infrared band
carriers: The top-down chemistry of interstellar carbonaceous dust grain analogues. *Astron. Astrophys.* 637, A82.
- Day, P.K., LeDuc, H.G., Mazin, B.A., Vayonakis, A., Zmuidzinas, J. (2003). A broadband superconducting suitable for
use in large arrays. *Nature* 425, 817.
- De Luca, M., Gupta, H., Neufeld, D., Gerin, M., Teyssier, D., Drouin, B.J., Pearson, J.C., Lis, D.C., Monje, R., Phillips, 928
929 T.G., Goicoechea, J.R., Godard, B., Falgarone, E., Coutens, A., Bell, T.A. (2012). Herschel/HIFI discovery of HCl⁺
in the interstellar medium. *Astrophys. J. Lett.* 751, L37.
- 930 Dhillon, S.S., Vitiello, M.S., Linfield, E.H., Davies, A.G., Hoffmann, M.C., Booske, J., Paoloni, C., Gensch, M., 931
932 Weightman, P., Williams, G.P., Castro-Camus, E., Cumming, D.R.S., Simoens, F., Escorcia-Carranza, I., Grant, J.,
933 Lucyszyn, S., Kuwata-Gonokami, M., Konishi, K., Koch, M., Schuttenmaer, C.A., Cocker, T.L., Huber, R.,
934 Markelz, A.G., Taylor, Z.D., Wallace, V.P., Zeitler, J.A., Sibik, J., Korter, T.M., Ellison, B., Rea, S., Goldsmith, P.,
935 Cooper, K.B., Appleby, R., Pardo, D., Huggard, P.G., Krozer, V., Shams, H., Fice, M., Renaud, C., Seeds, A., Stöhr,
936 A., Naftaly, M., Ridler, N., Clarke, R., Cunningham, J.E., Johnston, M.B. (2017). The 2017 terahertz science and
technology roadmap. *J. Phys. D: Appl. Phys.* 50, 043001.
- Ding, J.J., Boduch, P., Domaracka, A., Guillous, S., Langlinay, T., Lv, X.Y., Palumbo, M.E., Rothard, H., Strazzulla, G.
(2013). Implantation of multiply charged sulfur ions in water ice. *Icarus* 226, 860.
- Doudna, J.A., Cech, T.R. (2002). The chemical repertoire of natural ribozymes. *Nature* 418, 222.

- Douglas, A.E., Herzberg, G. (1941). CH⁺ in interstellar space and in the laboratory. *Astrophys. J.* 94,381.
- Draine, B.T., Katz, N. (1986). Magnetohydrodynamic shocks in diffuse clouds II. Production of CH⁺, OH, CH, and other species. *Astrophys. J.* 310,392.
- Drouin, B.J., Maiwald, F.W., Pearson, J.C. (2005). Application of cascaded frequency multiplication to molecular spectroscopy. *Rev. Sci. Instrum.* 76,093113.
- Ehrenfreund, P., Kerkhof, O., Schutte, W.A., Boogert, A.C.A., Gerakines, P.A., Dartois, E., d'Hendecourt, L., Tielens, A.G.G.M., van Dishoeck, E.F., Whittet, D.C.B. (1999). Laboratory studies of thermally processed H₂O-CH₃OH-CO₂ ice mixtures and their astrophysical implications. *Astron. Astrophys.* 350,240. 946
- Elitzur, M., Watson, W.D. (1980). Interstellar shocks and molecular CH⁺ in diffuse clouds. *Astrophys. J.* 236,172.
- Emrich, A. (1997). "Autocorrelation spectrometers for spaceborne (sub) millimeter astronomy." in *The Far Infrared and Submillimeter Universe*, ed. A. Wilson (Noordwijk), 361.
- Endres, C.P., Schlemmer, S., Schilke, P., Stutzki, J., Müller, H.S.P. (2016). The Cologne Database for Molecular Spectroscopy, CDMS, in the Virtual Atomic and Molecular Data Centre, VAMDC. *J. Mol. Spectrosc.* 327, 95.
- Esplugues, G.B., Viti, S., Goicoechea, J.R., Cernicharo, J. (2014). Modelling the sulphur chemistry evolution in Orion KL. *Astron. Astrophys.* 567,95.
- Farrah, D., Sminth, K.E., Ardila, D., Bradford, C.M., DiPirro, M.J., Ferkinhoff, C., Glenn, J., Goldsmith, P.F., Leisawitz, D.T., Nikola, T., Rangwala, N., Rinehart, S.A., Staguhn, J.G., Zemcov, M., Zmuidzinas, J., Bartlett, J., Carey, S.J., Fischer, W.J., Kamenetzky, J.R., Kartaltepe, J., Lacy, M.D., Lis, D.C., Locke, L.S., Lopez-Rodriguez, E., MacGregor, M., Mills, E., Moseley, S.H., Murphy, E.J., Rhodes, A., Richter, M.J., Rigopoulou, D., Sanders, D.B., Sankrit, R., Savini, G., Smith, J.D., Stierwalt, S. (2019). Review: Far-infrared instrumentation and technological development for the next decade. *J. Astron. Telesc. Instrum. Syst.* 5,020901. 956
- 961 Fayolle, E.C., Öberg, K.I., Jørgensen J.K., Altwegg, K., Calcutt, H., Müller, H.S.P., Rubin, M., der Wiel, M.H.D., Bjerke, P., Bourke, T.L., Coutens, A., van Dishoeck, E.F., Drozdovskaya, M.N., Garrod, R.T., Ligterink, N.F.W., Persson, M.V., Wampfler, S.F., Balsiger, H., Berthelier, J.J., De Keyser, J., Fiethe, B., Fuselier, S.A., Gasc, S., Gombosi, T.I., Sémon, T., Tzou, C.Y. (2017). *Nat. Astron.* 1,703. 962
- 963 Fedele, D., Bruderer, S., van Dishoeck, E.F., Carr, J., Herczeg, G.J., Salyk, C., Evans, N.J., Bouwman, J., Meeus, G., Henning, T., Green, J., Najita, J.R., Gudel, M. (2013). Digit survey of far-infrared lines from protoplanetary disks. *Astron. Astrophys.* 559,77. 964
- 965 Federman, S.R., Rawlings, J.M.C., Taylor, S.D., Williams, D.A. (1996). Synthesis of interstellar CH⁺ without OH. *Mon. Not. R. Astron. Soc.* 279, L41. 966
- 967 Fedoseev, G., Cuppen, H.M., Ioppolo, S., Lamberts, T., Linnartz, H. (2015). Experimental evidence for glycolaldehyde and ethylene glycol formation by surface hydrogenation of CO molecules under dense molecular cloud conditions. *Mon. Not. R. Astron. Soc.* 448, 1288. 970
- 971 Fedoseev, G., Chuang, K.J., Ioppolo, S., Qasim, D., van Dishoeck, E.F., Linnartz, H. (2017). Formation of glycerol through hydrogenation of CO ice under prestellar core conditions. *Astrophys. J.* 842, 52. 972
- 973 Feldman, P.A. (2001). Molecular astronomy from a Canadian perspective: The early years. *Can. J. Phys.* 79, 89. 974
- 975 Fraser, H.J., Collings, M.P., McCoustra, M.R.S. (2002). Laboratory surface astrophysics experiment. *Rev. Sci. Instrum.* 73, 2161. 976
- 977 Frenklach, M., Carmer, C., Feigelson, E. (1989). Silicon carbide and the origin of interstellar carbon grains. *Nature* 339, 196. 978
- 979

- 980 Fuchs, G.W., Cuppen, H.M., Ioppolo, S., Romanzin, C., Bisschop, S.E., Andersson, S., van Dishoeck, E.F., Linnartz, H.
981 (2009). Hydrogenation reactions in interstellar CO ice analogues. *Astron. Astrophys.* 505, 629.
- 982 Fulvio, D., Baratta, G.A., Sivaraman, B., Mason, N.J., da Silveira, E.F., de Barros, A.L.F., Pandoli, O., Strazzulla, G.,
983 Palumbo, M.E. (2018). Ion irradiation of N₂O ices and NO₂:N₂O₄ ice mixtures: first steps to understand the evolution of
984 molecules with the N–O bond in space. *Mon. Not. R. Astron. Soc.* 483, 381.
- 985 Galli, D., Walmsley, M., Gonçalves, J. (2002). The structure and stability of molecular cloud cores in external radiation
986 fields. *Astron. Astrophys.* 394, 275.
- 987 Garozzo, M., Fulvio, D., Kaňuchová, Z., Palumbo, M.E., Strazzulla, G. (2010). The fate of S-bearing species after ion
988 irradiation of interstellar icy grain mantles. *Astron. Astrophys.* 509, 67.
- 989 Garrod, R.T., Wicinus-Weaver, S.L., Herbst, E. (2008). Complex chemistry in star-forming regions: An expanded gas-
990 grain warm-up chemical model. *Astrophys. J.* 682, 283.
- 991 Garrod, R.T., Belloche, A., Müller, H.S.P., Menten, K.M. (2017). Exploring molecular complexity with ALMA (EMoCA):
992 Simulations of branched carbon-chain chemistry in Sgr B2(N). *Astron. Astrophys.* 601, 48.
- 993 Geppert, W.D., Larsson, M. (2013). Experimental investigations into astrophysically relevant ionic reactions. *Chem. Rev.*
994 113, 8872.
- 995 Gerin, M., De Luca, M., Black, J., Goicoechea, J.R., Herbst, E., Neufeld, D.A., Falgarone, E., Godard, B., Pearson, J.C.,
996 Lis, D.C., Phillips, T.G., Bell, T.A., Sonnentrucker, P., Boulanger, F., Cernicharo, J., Coutens, A., Dartois, E., Encrenaz,
997 P., Giesen, T., Goldsmith, P.F., Gupta, H., Gry, C., Hennebelle, P., Hily-Blant, P., Joblin, C., Kazmierczak, M., Kolos, R.,
998 Krelowski, J., Martin-Pintado, J., Monje, R., Mookerjee, B., Perault, M., Persson, C., Plume, R., Rimmer, P.B., Salez, M.,
999 Schmidt, M., Stutzki, J., Teyssier, D., Vastel, C., Yu, S., Contursi, A., Menten, K., Geballe, T., Schlemmer, S., Shipman,
1000 R., Tielens, A.G.G.M., Philipp-May, S., Cros, A., Zmuidzinas, J., Samoska, L.A., Klein, K., Lorenzani, A. (2010a).
1001 Interstellar OH⁺, H₂O⁺ and H₃O⁺ along the sight-line G10.6-0.4. *Astron. Astrophys.* 518, L110.
1002
- 1003 Gerin, M., Kaźmierczak, M., Jastrzebska, M., Falgarone, E., Hily-Blant, P., Godard, B., De Luca, M. (2010b). The tight
1004 correlation of CCH and c-C₃H₂ in diffuse and translucent clouds. *Astron. Astrophys.* 525, 116.
- 1005 Gerin, M., Levrier, F., Falgarone, E., Godard, B., Hennebelle, P., Le Petit, F., De Luca, M., Neufeld, D., Sonnentrucker,
1006 P., Goldsmith, P., Flagey, N., Lis, D.C., Persson, C.M., Black, J.H., Goicoechea, J.R., Menten, K.M. (2012). Hydride
1007 spectroscopy of the diffuse interstellar medium: New clues on the gas fraction in molecular form and cosmic ray ionization
1008 rate in relation to H⁺. *Phil. Trans. R. Soc. A* 370, 5174.
- 1009 Giuliano, B.M., Gavdush, A.A., Müller, B., Zaytsev, K.I., Grassi, T., Ivlev, A.V., Palumbo, M.E., Baratta, G.A., Scirè, C.,
1010 Komandin, G.A., Yurchenko, S.O., Caselli, P. (2019). Broadband spectroscopy of astrophysical ice analogues I: Direct
1011 measurement of the complete refractive index of CO ice using terahertz time-domain spectroscopy. *Astron. Astrophys.* 629,
1012 112.
- 1013 Godard, B., Falgarone, E., Pineau des Forêts (2009). Models of turbulent dissipation regions in the diffuse interstellar
1014 medium. *Astron. Astrophys.* 495, 847.
- 1015 Godard, B., Cernicharo, J. (2013). A complete model of CH⁺ rotational excitation including radiative and chemical pumping
1016 processes. *Astron. Astrophys.* 550, 8.
- 1017 Goldman, N., Reed, E.J., Fried, L.E., Kuo, I.F.W., Maiti, A. (2010). Synthesis of glycine-containing complexes in impacts
1018 of comets on early Earth. *Nat. Chem.* 2, 949.
- 1019 Goldsmith, P.F., Liseau, R., Bell, T.A., Black, J.H., Chen, J.H., Hollenbach, D., Kaufman, M.J., Li, D., Lis, D.C., Melnick,
1020 G., Neufeld, D., Pagani, L., Snell, R., Benz, A.O., Bergin, E.A., Bruderer, S., Caselli, P., Caux, E., Encrenaz, P.,
1021 Falgarone, E., Gerin, M., Goicoechea, J.R., Hjalmarson, Å., Larsson, B., Le Bourlot, J., Le Petit, F., De Luca, M.,
1022 Nagy, Z., Roueff, E., Sandqvist, A., van der Tak, F., van Dishoeck, E.F., Vastel, C., Viti, S., Yıldız, U.A. (2011).
1023 Herschel measurements of molecular oxygen in Orion. *Astrophys. J.* 737, 96.

- 1024 Graff, M.M., Moseley, J.T., Roueff, E. (1983). Resonant and non-resonant processes in the formation of CH⁺ by radiative
1025 association. *Astrophys. J.* 269, 796.
- 1026 Griffin, M.J., Abergel, A., Abreu, A., Ade, P.A.R., André, P., Augeres, J.L., Babbedge, T., Bae, Y., Baillie, T., Baluteau,
1027 J.P., Barlow, M.J., Bendo, G., Benielli, D., Bock, J.J., Bonhomme, P., Brisbin, D., Brockley-Blatt, C., Caldwell, M., Cara,
1028 C., Castro-Rodriguez, N., Cerulli, R., Chaniel, P., Chen, S., Clark, E., Clements, D.L., Clerc, L., Coker, J., Communal, D.,
1029 Conversi, L., Cox, P., Crumb, D., Cunningham, C., Daly, F., Davis, G.R., De Antoni, P., Delderfield, J., Devin, N., Di Giorgio,
1030 A., Didschuns, I., Dohlen, K., Donati, M., Dowell, A., Dowell, C.D., Duband, L., Dumaye, L., Emery, R.J., Ferlet, M.,
1031 Ferrand, D., Fontignie, J., Fox, M., Franceschini, A., Frerking, M., Fulton, T., Garcia, J., Gastaud, R., Gear, W.K., Glenn,
1032 J., Goizel, A., Griffin, D.K., Grundy, T., Guest, S., Guilletmet, L., Hargrave, P.C., Harwit, M., Hastings, P., Hatziminaoglou,
1033 E., Herman, M., Hinde, B., Hristov, V., Huang, M., Imhof, P., Isaak, K.J., Israelsson, U., Ivison, R.J., Jennings, D., Kiernan,
1034 B., King, K.J., Lange, A.E., Latter, W., Laurent, G., Laurent, P., Leeks, S.J., Lellouch, E., Levenson, L., Li, B., Li, J.,
1035 Lilienthal, J., Lim, T., Liu, S.J., Lu, N., Madden, S., Mainetti,
1036 G., Marliani, P., McKay, D., Mercier, K., Molinari, S., Morris, H., Mosely, H., Mulder, J., Mur, M., Naylor, N.A.,
1037 Nguyen, H., O'Halloran, B., Olivier, S., Olofsson, G., Olofsson, H.G., Orfei, R., Page, M.J., Pain, I., Panuzzo, P.,
1038 Papegeorgiou, A., Parks, G., Parr-Burman, P., Pearce, A., Pearson, C., Pérez-Fournon, I., Pinsard, F., Pisano, G.,
1039 Podosek, J., Pohlen, M., Polehampton, E.T., Pouliquen, D., Rigopoulou, D., Rizzo, D., Roseboom, I.G., Roussel,
1040 H., Rowan-Robinson, M., Rownd, B., Saraceno, P., Sauvage, M., Savage, R., Savini, G., Sawyer, E., Scharmberg,
1041 C., Schmitt, D., Schneider, N., Schulz, B., Schwartz, A., Shafer, R., Shupe, D.L., Sibthorpe, B., Sidher, S., Smith,
1042 A., Smith, A.J., Smith, D., Spencer, L., Stobie, B., Sudiwala, R., Sukhatme, K., Surace, C., Stevens, J.A., Swinyard,
1043 B.M., Trichas, M., Tourette, T., Triou, H., Tseng, S., Tucker, C., Turner, A., Vaccari, M., Valtchanov, I., Vigroux,
1044 L., Virique, E., Voellmer, G., Walker, H., Ward, R., Waskett, T., Weilert, M., Wesson, R., White, G.J., Whitehouse,
1045 N., Wilson, C.D., Winter, B., Woodcraft, A.L., Wright, G.S., Xu, C.K., Zavagno, A., Zemcov, M., Zhang, L., Zonca,
1046 E. (2010). The Herschel-SPIRE instrument and its in-flight performance. *Astron. Astrophys.* 518, L3.
- 1047 Griffin, M.J., Baselmans, J., Baryshev, A., Doyle, S., Grim, M., Hargrave, P., Klapwijk, T., Martin-Pintado, J., Monfardini,
1048 A., Neto, A., Steenbeek, H., Walker, I., Wood, K., D'Addabbo, A., Barry, P., Bideaud, A., Blázquez, B., Bueno, J., Calvo,
1049 M., Costa-Kramer, J.L., Ferrari, L., Gómez-Gutiérrez, A., Goupy, J., Llombart, N., Yates, S. (2016). SPACEKIDS: Kinetic
1050 inductance detectors for space applications. *Proc. SPIE 9914, Millimeter, Submillimeter, and Far-Infrared Detectors and*
1051 *Instrumentation for Astronomy VIII*, 991407.
- 1052 Griffiths, P., de Haseth, J. (1986). *Fourier Transform Infrared Spectroscopy*. New York City (NY): John Wiley and Sons.
- 1053 Groen, J., Deamer, D.W., Kros, A., Ehrenfreund, P. (2012). Polycyclic aromatic hydrocarbons as plausible prebiotic
1054 membrane components. *Orig. Life Evol. Biosph.* 42, 295.
- 1055 Gupta, H., Rimmer, P., Pearson, J.C., Yu, S., Herbst, E., Harada, N., Bergin, E.A., Neufeld, D.A., Melnick, G.J., Bachiller,
1056 R., Baechtold, W., Bell, T.A., Blake, G.A., Caux, E., Ceccarelli, C., Cernicharo, J., Chattopadhyay, G., Comito, C., Cabrit,
1057 S., Crockett, N.R., Daniel, F., Falgarone, E., Diez-Gonzalez, M.C., Dubernet, M.L., Erickson, N., Emprechtinger, M.,
1058 Encrenaz, P., Gerin, M., Gill, J.J., Giesen, T.F., Goicoechea, J.R., Goldsmith, P.F., Joblin, C., Johnstone, D., Langer, W.D.,
1059 Larsson, B., Latter, W.B., Lin, R.H., Lis, D.C., Liseau, R., Lord, S.D., Maiwald, F.W.,
1060 Maret, S., Martin, P.G., Martin-Pintado, J., Menten, K.M., Morris, P., Müller, H.S.P., Murphy, J.A., Nordh, L.H.,
1061 Olberg, M., Ossenkopf, V., Pagani, L., Pérou, M., Phillips, T.G., Plume, R., Qin, S.L., Salez, M., Samoska, L.A.,
1062 Schilke, P., Schlecht, E., Schlemmer, S., Szczerba, R., Stutzki, J., Trappe, N., van der Tak, F.F.S., Vastel, C., Wang,
1063 S., Yorke, H.W., Zmuidzinas, J., Boogert, A., Güsten, R., Hartogh, P., Honingh, N., Karpov, A., Kooi, J., Krieg,
1064 J.M., Schieder, R., Zaal, P. (2010). Detection of OH⁺ and H₂O⁺ towards Orion KL. *Astron. Astrophys.* 521, L47.
- 1065 Gurvits, L.I., Paragi, Z., Casasola, V., Conway, J., Davelaar, J., Falcke, H., Fender, R., Frey, S., Fromm, C.M., García Miró,
1066 C., Garrett, M.A., Giroletti, M., Goddi, C., Gómez, J.L., van der Gucht, J., Guirado, J.C., Haiman, Z., Helmich, F.,
1067 Humphreys, E., Impellizzeri, V., Kramer, M., Lindqvist, M., Linz, H., Liuzzo, E., Lobanov, A.P., Mizuno, Y., Rezzolla,
1068 L., Roelofs, F., Ros, E., Rygl, K.L.J., Savolainen, T., Schuster, K., Venturi, T., Wiedner, M.C., Zensus,
1069 J.A. (2021). THEZA: TeraHertz Exploration and Zooming-in for Astrophysics. *Exp. Astron.* DOI: 10.1007/s10686-
1070 021-09714-y
- 1071 Hazen, R.M., Papineau, D., Bleeker, W., Downs, R.T., Ferry, J.M., McCoy, T.J., Sverjensky, D.A., Yang, H. (2008).
1072 Mineral evolution. *Am. Mineral.* 93, 1693.
- 1073 Hazen, R.M., Ferry J.M. (2010). Mineral evolution: Mineralogy in the fourth dimension. *Elements* 6, 9.

- 1074 Henderson, S.W., Stevens, J.R., Amiri, M., Austermann, J., Beall, J.A., Chaudhuri, S., Cho, H.M., Choi, S.K., Cothard,
1075 N.F., Crowley, K.T., Duff, S.M., Fitzgerald, C.P., Gallardo, P.A., Halpern, M., Hasselfield, M., Hilton, G., Ho, S.P.P.,
1076 Hubmayr, J., Irwin, K.D., Koopman, B.J., Li, D., Li, Y., McMahon, J., Nati, F., Niemack, M., Reintsema,
1077 C.D., Salatino, M., Schillaci, A., Schmitt, B.L., Simon, S.M., Staggs, S.T., Vavagiakis, E.M., Ward, J.T. (2016).
1078 Readout of two-kilopixel transition-edge sensor arrays for advanced ACTPol. Proc. SPIE 9914, Millimeter,
1079 Submillimeter, and Far-Infrared Detectors and Instrumentation for Astronomy VIII, 9914.
- 1080 Higgs, P.G., Lehman, N. (2015). The RNA world: Molecular cooperation at the origins of life. *Nat. Rev. Genet.* 16, 7.
- 1081 Hogerheidje, M.R., Bergin, E.A., Brinch, C., Cleaves, L.I., Fogel, J.K.J., Blake, G.A., Dominik, C., Lis, D.C., Melnick, G.,
1082 Neufeld, D., Panić, O., Pearson, J.C., Kristensen, L., Yıldız, U.A., van Dishoeck, E.F. (2011). Detection of the water
1083 reservoir in a forming planetary system. *Science* 334, 338.
- 1084 Indriolo, N., Neufeld, D.A., Gerin, M., Schilke, P., Benz, A.O., Winkel, B., Menten, K.M., Chambers, E.T., Black, J.H.,
1085 Bruderer, S., Falgarone, E., Godard, B., Goicoechea, J.R., Gupta, H., Lis, D.C., Ossenkopf, V., Persson, C.M.,
1086 Sonnentrucker, P., van der Tak, F.F.S., van Dishoeck, E.F., Wolfire, M.G., Wyrowski, F. (2015). HERSCHEL survey of
1087 galactic OH⁺, H₂O⁺, and H₃O⁺: Probing the molecular hydrogen fraction and cosmic-ray ionization rate. *Astrophys. J.* 800,
1088 40.
- 1089 Ioppolo, S., McGuire, B.A., Allodi, M.A., Blake, G.A. (2014). THz and mid-IR spectroscopy of interstellar ice analogues:
1090 methyl and carboxylic acid groups. *Faraday Discuss.* 168, 461.
- 1091 Ioppolo, S., Kaňuchová, Z., James, R.L., Dawes, A., Jones, N.C., Hoffmann, S.V., Mason, N.J., Strazzulla, G. (2020).
1092 Vacuum ultraviolet photoabsorption spectroscopy of space-related ices: 1 keV electron irradiation of nitrogen- and oxygen-
1093 rich ices. *Astron. Astrophys.* 641, 154.
- 1094 Ioppolo, S., Fedoseev, G., Chuang, K.J., Cuppen, H.M., Clements, A.R., Jin, M., Garrod, R.T., Qasim, D., Kofman, V., van
1095 Dishoeck, E.F., Linnartz, H. (2021). A non-energetic mechanism for glycine formation in the interstellar medium. *Nat.*
1096 *Astron.* 5, 197.
- 1097 Jacob, A.M., Menten, K.M., Wyrowski, F., Winkel, B., Neufeld, B.A. (2020). Extending the view of ArH⁺ chemistry in
1098 diffuse clouds. *Astron. Astrophys.* 643, 91.
- 1099 Jones, B.M., Bennett, C.J., Kaiser, R.I. (2011). Mechanistical studies on the production of formamide (H₂NCHO) within
1100 interstellar ice analogs. *Astrophys. J.* 734, 78.
- 1101 Jørgensen, J.K., Müller, H.S.P., Calcutt, H., Coutens, A., Drozdovskaya, M.N., Öberg, K.I., Persson, M.V., Taquet, V., van
1102 Dishoeck, E.F., Wampfler, S.F. (2018). The ALMA-PILS survey: Isotopic composition of oxygen-containing complex
1103 organic molecules toward IRAS 16293-2422B. *Astron. Astrophys.* 620, 170.
- 1104 Kaňuchová, Z., Boduch, P., Domaracka, A., Palumbo, M.E., Rothard, H., Strazzulla, G. (2017). Thermal and energetic
1105 processing of astrophysical ice analogues rich in SO₂. *Astron. Astrophys.* 604, 68.
- 1106 Kawase, K., Sato, M., Taniuchi, T., Ito, H. (1996). Coherent tunable THz-wave generation from LiNbO₃ with monolithic
1107 grating coupler. *Appl. Phys. Lett.* 68, 2483.
- 1108 Kiessling, J., Breunig, I., Schunemann, P.G., Buse, K., Vodopyanov, K.L. (2013). High power and spectral purity
1109 continuous wave photonic THz source tunable from 1 to 4.5 THz for nonlinear molecular spectroscopy. *New J. Phys.* 15,
1110 105014.
- 1111 Kilcullen, P., Hartley, I.D., Jensen, E.T., Reid, M. (2015). Terahertz time domain gas-phase spectroscopy of carbon
1112 monoxide. *J. Infrared Milli. Terahz. Waves* 36, 380.
- 1113 Klein, B., Philipp, S.D., Güsten, R., Krämer, I., Samtleben, D. (2006). A new generation of spectrometers for radio
1114 astronomy: Fast Fourier-transform spectrometer. Proc. SPIE 6275, 62751.
- 1115 Klein, B., Hochgürtel, S., Krämer, I., Bell, A., Meyer, K., Güsten, R. (2012). High-resolution wide-band fast Fourier-
1116 transform spectrometers. *Astron. Astrophys.* 542, L3.

- 1117 Kojima, T., Kroug, M., Uemizu, K., Niizeki, Y., Takahashi, H., Uzawa, Y. (2017). Performance and characterization of a
 1118 wide if SIS-mixer-preamplifier module employing high-j c SIS junctions. *IEEE Trans. Terahertz Sci. Technol.* 7, 694.
 1119
- 1120 Kokoouline, V., Greene, C.H. (2003). Unified theoretical treatment of dissociative recombination of D_{3h} triatomic ions:
 1121 applications to H^+ and D^+ . *Phys. Rev. A* 68, 012703.
- 1122 Komandin, G.A., Chuchupal, S.V., Lebedev, S.P., Goncharov, Y.G., Korolev, A.F., Porodinkov, O.E., Spektor, I.E.,
 1123 Volkov, A.A. (2013). BWO generators for terahertz dielectric measurements. *IEEE Trans. Terahertz Sci. Technol.* 3, 440.
 1124
- 1125 Kraus, J.D. (1988). Grote Reber – Founder of Radio Astronomy. *J. R. Astron. Soc. Can.* 82, 107.
- 1126 Kruger, K., Grabowski, P.J., Zaug, A.J., Sands, J., Gottschling, D.E., Cech, T.R. (1982). Self-splicing RNA: Autoexcision
 1127 and autocyclization of the ribosomal RNA intervening sequence of tetrahymena. *Cell* 31, 147.
- 1128 Kulesa, C. (2011). Terahertz spectroscopy for astronomy: From comets to cosmology. *IEEE Trans. Terahertz Sci. Technol.*
 1129 1, 232.
- 1130 Kwok, S., Zhang, Y. (2011). Mixed aromatic-aliphatic organic nanoparticles as carriers of unidentified infrared emission
 1131 features. *Nature* 479, 80.
- 1132 Lang, K., Erlacher, M., Wilson, D.N., Micura, R., Polacek, N. (2008). The role of 23S ribosomal RNA residue A2451 in
 1133 peptide bond synthesis revealed by atomic mutagenesis. *Chem. Biol.* 15, 485.
- 1134 Langer, W.D., Velusamy, T., Pineda, J.L., Goldsmith, P.F., Li, D., Yorke, H.W. (2010). C^+ detection of warm dark gas in
 1135 diffuse clouds. *Astron. Astrophys.* 521, L17.
- 1136 Larson, Å, Djurić, N., Zong, W., Greene, C.H., Orel, A.E., Al-Khalili, A., Derkatch, A.M., Le Padellec, A., Neau, A.,
 1137 Rosén, S., Shi, W., Vikor, L., Danared, H., af Ugglas, M., Larsson, M., Dunn, G.H. (2000). Resonant ion-pair formation in
 1138 electron collisions with HD^+ and OH^+ . *Phys. Rev. A* 62, 042707.
- 1139 Larsson, M., McCall, B.J., Orel, A. (2008). The dissociative recombination of H_3^+ – a saga coming to an end? *Chem. Phys.*
- 1140 1157
- 1141
- 1142
- 1143
- 1144
- 1145
- 1146
- 1147
- 1148
- 1149
- 1150
- 1151
- 1152
- 1153
- 1154
- 1155
- 1156

Larsson, M., Geppert, W.D., Nymann, G. (2012). Ion chemistry in space. Rep. Prog. Phys. 75, 066901.

- Lee, K.L.K., Loomis, R.A., Burkhardt, A.M., Cooke, I.R., Xue, C., Siebert, M.A., Shingledecker, C.N., Remijan, A., Charnley, S.B., McCarthy, M.C. (2021a). Interstellar detection of 2-cyanocyclopentadiene, C_5H_5CN , a second five-membered ring toward TMC-1. *Astrophys. J. Lett.* 910, L2.
- Lee, K.L.K., Loomis, R.A., Burkhardt, A.M., Cooke, I.R., Xue, C., Siebert, M.A., Shingledecker, C.N., Remijan, A., Charnley, S.B., McCarthy, M.C. (2021b). Discovery of interstellar *trans*-cyanovinylacetylene ($HC\equiv CCH=CHC\equiv N$) and vinylcyanoacetylene ($H_2C=CHC\equiv N$) in GOTHAM observations of TMC-1. *Astrophys. J. Lett.* 908, L11.
- Li, W., Yao, J. (2010). Investigation of Photonically Assisted Microwave Frequency Multiplication Based on External Modulation. *IEEE Trans. Microw. Theory Tech.* 58, 3259.
- Linnartz, H., Ioppolo, S., Fedoseev, G. (2015). Atom addition reactions in interstellar ice analogues. *Int. Rev. Phys. Chem.* 2, 205.

Linz, H., Beuther, H., Gerin, M., Goicoechea, J.R., Helmich, F., Krause, O., Liu, Y., Molinari, S., Ossenkopf-Okada, V., Pineda, J., Sauvage, M., Schinnerer, E., van der Tak, F., Wiedner, M., Amiaux, J., Bhatia, D., Buinhas, L., Durand, G., Förstner, R., Graf, U., Lezius, M. (2021). Bringing high spatial resolution to the far-infrared. *Exp. Astron.* DOI: 10.1007/s10686-021-09719-7

Lis, D.C., Pearson, J.C., Neufeld, D.A., Schilke, P., Müller, H.S.P., Gupta, H., Bell, T.A., Comito, C., Phillips, T.G., Bergin, E.A., Ceccarelli, C., Goldsmith, P.F., Blake, G.A., Bacmann, A., Baudry, A., Benedettini, M., Benz, A., Black, J.,

Lee,
K.L.
.K.,
Cha
ngal
a,
P.B.

,
Loo
mis,
R.A

,,
Bur
khar
dt,
A.
M.,
Xue
, C.,
Cor
dine
r,
M.
A.,
Cha

- 1158 Boogert, A., Bottinelli, S., Cabrit, S., Caselli, P., Castets, A., Caux, E., Cernicharo, J., Codella, C., Coutens, A.,
 1159 Crimier, N., Crockett, N.R., Daniel, F., Demyk, K., Dominic, C., Dubernet, M.L., Emprechtinger, M., Encrenaz, P.,
 1160 Falgarone, E., Fuente, A., Gerin, M., Giesen, T.F., Goicoechea, J.R., Helmich, F., Hennebelle, P., Henning, T.,
 1161 Herbst, E., Hily-Blant, P., Hjalmarsen, Å., Hollenbach, D., Jack, T., Joblin, C., Johnstone, D., Kahane, C., Kama,
 1162 M., Kaufman, M., Klotz, A., Langer, W.D., Larsson, B., Le Boulrot, J., Lefloch, B., Le Petit, F., Li, D., Liseau, R.,
 1163 Lord, S.D., Lorenzani, A., Maret, S., Martin, P.G., Melnick, G.J., Menten, K.M., Morris, P., Murphy, J.A., Nagy,
 1164 Z., Nisini, B., Ossenkopf, V., Pacheco, S., Pagani, L., Parise, B., Péroult, M., Plume, R., Qin, S.L., Roueff, E., Salez,
 1165 M., Sandqvist, A., Saraceno, P., Schlemmer, S., Schuster, K., Snell, R., Stutzki, K., Tielens, A.G.G.M., Trappe, N.,
 1166 van der Tak, F.F.S., van der Wiel, M.H.D., van Dishoeck, E.F., Vastel, C., Viti, S., Wakelam, V., Walters, A., Wang,
 1167 S., Wyrowski, F., Yorke, H.W., Yu, S., Zmuidzinas, J., Delorme, Y., Desbat, J.P., Güsten, R., Krieg, J.M., Delforge,
 1168 B. (2010). Herschel/HIFI discovery of interstellar chloronium (H_2Cl^+). *Astron. Astrophys.* 521, L9.
- 1169 Lo, J.I., Chou, S.L., Peng, Y.C., Lin, M.Y., Lu, H.C., Cheng, B.M. (2014). Photochemistry of solid interstellar molecular
 1170 samples exposed to vacuum-ultraviolet synchrotron radiation. *J. Electron Spectros. Relat. Phenomen.* 196, 173.
- 1171 Loomis, R.A., Burkhardt, A.M., Shingledecker, C.N., Charnley, S.B., Cordiner, M.A., Herbst, E., Kalenskii, S., Lee,
 1172 K.L.K., Willis, E.R., Xue, C., Remijan, A.J., McCarthy, M.C., McGuire, B.A. (2021). An investigation of spectral line
 1173 stacking techniques and application to the detection of HC_{11}N . *Nat. Astron.* 5, 188.
- 1174 Maestrini, A., Thomas, B., Wang, H., Jung, C., Treuttel, J., Jin, Y., Chattopadhyay, G., Mehdi, I., Beaudin, G. (2010).
 1175 Schottky diode-based terahertz frequency multipliers and mixers. *C. R. Phys.* 11, 480.
- 1176 Maiolino, R. (2008). Prospects for AGN studies with ALMA. *New Astron. Rev.* 52, 339.
- 1177 Martins, Z., Price, M.C., Goldman, N., Sephton, M.A., Burchell, M.J. (2013). Shock synthesis of amino acids from
 1178 impacting cometary and icy planet surface analogues. *Nat. Geosci.* 6, 1045.
- 1179 Mason, N.J., Dawes, A., Holtom, P.D., Mukerji, R.J., Davis, M.P., Sivaraman, B., Kaiser, R.I., Hoffmann, S.V., Shaw,
 1180 D.A. (2006). VUV spectroscopy and photo-processing of astrochemical ices: An experimental study. *Faraday*
 1181 *Discuss.* 133, 311.
- 1182 Materese, C.K., Nuevo, M., Sandford, S.A. (2015). N- and O-heterocycles produced from the irradiation of benzene and
 1183 naphthalene in $\text{H}_2\text{O}/\text{NH}_3$ -containing ices. *Astrophys. J.* 800, 116.
- 1184 Materese, C.K., Nuevo, M., Sandford, S.A. (2017). The formation of nucleobases from the ultraviolet photo-irradiation of
 1185 purine in simple astrophysical ice analogues. *Astrobiology* 17, 761.
- 1186 Materese, C.K., Nuevo, M., McDowell, B.L., Buffo, C.E., Sandford, S.A. (2018). The photochemistry of purine in ice
 1187 analogs relevant to dense interstellar clouds. *Astrophys. J.* 864, 44.
- 1188 McCall, B.J., Geballe, T.R., Hinkle, K.H., Oka, T. (1998). Detection of H^+ in the diffuse interstellar medium toward
 1189 Cygnus OB2 No. 12. *Science* 279, 1910.
- 1190 McCarthy, M.C., Lee, K.L.K., Loomis, R.A., Burkhardt, A.M., Shingledecker, C.N., Charnley, S.B., Cordiner, M.A.,
 1191 Herbst, E., Kalenskii, S., Willis, E.R., Xue, C., Remijan, A.J., McGuire, B.A. (2021). Interstellar detection of the highly
 1192 polar five-membered ring cyanocyclopentadiene. *Nat. Astron.* 5, 176.
- 1193 McGuire, B.A., Ioppolo, S., Allodi, M.A., Blake, G.A. (2016). THz time-domain spectroscopy of mixed CO_2 - CH_3OH
 1194 interstellar ice analogues. *Phys. Chem. Chem. Phys.* 18, 20199.
- 1195 McGuire, B.A., Shingledecker, C.N., Willis, E.R., Burkhardt, A.M., El-Abd, S., Motiyenko, R.A., Brogan, C.L., Hunter,
 1196 T.R., Margulès, L., Guillemin, J.C. (2017). ALMA detection of interstellar methoxymethanol ($\text{CH}_3\text{OCH}_2\text{OH}$). *Astrophys.*
 1197 *J. Lett.* 851, 2.
- 1198 McGuire, B.A., Burkhardt, A.M., Kalenskii, S., Shingledecker, C.N., Remijan, A.J., Herbst, E., McCarthy, M.C. (2018a).
 1199 Detection of the aromatic molecular benzonitrile ($c\text{-C}_6\text{H}_5\text{CN}$) in the interstellar medium. *Science* 359, 202.

- 1200 McGuire, B.A., Brogan, C.L., Hunter, T.R., Remijan, A.J., Blake, G.A., Burkhardt, A.M., Carroll, P.B., van Dishoeck,
1201 E.F., Garrod, R.T., Linnartz, H., Shingledecker, C.N., Willis, E.R. (2018b). First results of an ALMA Band 10 spectral line
1202 survey of NGC 6334I: Detections of glycolaldehyde (HC(O)CH₂OH) and a new compact bipolar outflow in HDO and CS.
1203 *Astrophys. J.* 863, L35.
- 1204 McGuire, B.A., Burkhardt, A.M., Loomis, R.A., Shingledecker, C.N., Lee, K.L.K., Charnley, S.B., Cordiner, M.A., Herbst,
1205 E., Kalenskii, S., Momjian, E., Willis, E.R., Xue, C., Remijan, A.J., McCarthy, M.C. (2020). Early science from GOTHAM:
1206 Project overview, methods, and the detection of interstellar propargyl cyanide (HCCCH₂CN) in TMC-
1207 1. *Astrophys. J. Lett.* 900, L10.
- 1208 McGuire, B.A., Loomis, R.A., Burkhardt, A.M., Lee, K.L.K., Shingledecker, C.N., Charnley, S.B., Cooke, I.R., Cordiner,
1209 M.A., Herbst, E., Kalenskii, S., Siebert, M.A., Willis, E.R., Xue, C., Remijan, A.J., McCarthy, M.C. (2021). Detection of
1210 two interstellar polycyclic aromatic hydrocarbons via spectral matched filtering. *Science* 371, 1265.
- 1211 McIntosh, A.I., Yang, B., Goldup, S.M., Watkinson, M., Donnan, R.S. (2012). Terahertz spectroscopy: A powerful new
1212 tool for the chemical sciences? *Chem. Soc. Rev.* 41, 2072.
- 1213 Mead, G., Katayama, I., Takeda, J., Blake, G.A. (2019). An echelon-based single shot optical and terahertz Kerr effect
1214 spectrometer. *Rev. Sci. Instrum.* 90, 053107.
- 1215 Medvedev, I.R., Neese, C.F., Plummer, G.M., Lucia, F.C.D. (2010). Submillimeter spectroscopy for chemical analysis with
1216 absolute specificity. *Opt. Lett.* 35, 1533.
- 1217 Menten, K.M., Wyrowski, F., Belloche, A., Güsten, R., Dedes, L., Müller, H.S.P. (2011). Submillimeter absorption from
1218 SH⁺, a new widespread interstellar radical, ¹³CH⁺ and HCl. *Astron. Astrophys.* 525, 77.
- 1219 Merino, P., Švec, M., Martinez, J.I., Jelinek, P., Lacovig, P., Dalmiglio, M., Lizzit, S., Soukiassian, P., Cernicharo, J.,
1220 Martin-Gago, J.A. (2014). Graphene etching on SiC grains as a path to interstellar polycyclic aromatic hydrocarbons
1221 formation. *Nat. Commun.* 5, 3054.
- 1222 Mifsud, D.V., Kaňuchová, Z., Herczku, P., Ioppolo, S., Juhász, Z., Kovács, S.T.S., Mason, N.J., McCullough, R.W., Sulik,
1223 B. (2021). Sulfur ice astrochemistry: A Review of laboratory studies. *Space Sci. Rev.* 217, 14.
- 1224 Milillo, A., Plainaki, C., De Angelis, E., Mangano, V., Massetti, S., Mura, A., Orsini, S., Rispoli R. (2016). Analytical
1225 model of Europa's O₂ exosphere. *Planet. Space Sci.* 130, 3.
- 1226 Mittleman, D., Jacobsen, R., Neelamani, R., Baranuik, R., Nuss, M. (1998). Gas sensing using terahertz time-domain
1227 spectroscopy. *Appl. Phys. B* 67, 379.
- 1228 Monje, R.R., Emprechtinger, M., Phillips, T.G., Lis, D.C., Goldsmith, P.F., Bergin, E.A., Bell, T.A., Neufeld, D.A.,
1229 Sonnentrucker, P. (2011). Herschel/HIFI observations of hydrogen fluoride toward Sagittarius B2(M). *Astrophys. J. Lett.*
1230 734, L23.
- 1231 Monje, R.R., Lis, D.C., Roueff, E., Gerin, M., De Luca, M., Neufeld, D.A., Godard, B., Phillips, T.G. (2013). Hydrogen
1232 chloride in diffuse interstellar clouds along the line of sight to W31C (G10.6-0.4). *Astrophys. J.* 767, 81.
- 1233 Moore, M.H., Hudson, R.L. (1992). Far-infrared spectral studies of phase changes in water ice induced by proton
1234 irradiation. *Astrophys. J.* 401, 353.
- 1235 Moore, M.H., Hudson, R.L. (1994). Far-infrared spectra of cosmic-type pure and mixed ices. *Astron. Astrophys. Suppl.*
1236 Ser. 103, 45.
- 1237 Moore, M.H., Hudson, R.L. (1995). Far-infrared spectral changes accompanying proton irradiation of solids of
1238 astrochemical interest. *Radiat. Phys. Chem.* 45, 779.
- 1239 Mosely, E.R., Draine, B.T., Tomida, K., Stone, J.M. (2021). Turbulent dissipation, CH⁺ abundance, H₂ line luminosities,
1240 and polarization in the cold neutral medium. *Mon. Not. R. Astron. Soc.* 500, 3290.

- 1241 Müller, H.S.P., Thorwith, S., Roth, D.A., Winnewisser, G. (2001). The Cologne Database for Molecular Spectroscopy,
1242 CDMS. *Astron. Astrophys.* 370, L49.
- 1243 Müller, H.S.P., Schlöder, F., Stutzki, J., Winnewisser, G. (2005). The Cologne Database for Molecular Spectroscopy: A
1244 useful tool for astronomers and spectroscopists. *J. Mol. Struct.* 742, 215.
- 1245 Muller, S., Müller, H.S.P., Black, J.H., Gerin, M., Combes, F., Curran, S., Falgarone, E., Guélin, M., Henkel, C., Martín,
1246 S., Menten, K.M., Roueff, E., Aalto, S., Beelen, A., Wiklind, T., Zwaan, M.A. (2017). Detection of CH⁺, SH⁺, and their
1247 ¹³C- and ³⁴S-isotopologues toward PKS 1830-211. *Astron. Astrophys.* 606, 109.
- 1248 Mullikin, E., van Mulbregt, P., Perea, J., Kasule, M., Huang, J., Buffo, C., Campbell, J., Gates, L., Cumberbatch, H.M.,
1249 Peeler, Z., Schneider, H., Lukens, J., Bao, S.T., Tano-Menka, R., Baniya, S., Cui, K., Thompson, M., Hay, A., Widdup, L.,
1250 Caldwell-Overdier, A., Huang, J., Boyer, M.C., Rajappan, M., Echebiri, G., Arumainayagam, C.R. (2018). Condensed-
1251 phase photochemistry in the absence of radiation. *ACS Earth Space Chem.* 2, 863.
- 1252 Muñoz-Caro, G.M., Jiménez-Escobar, A., Martín-Gago, J.Á., Rogero, C., Atienza, C., Puertas, S., Sobrado, J.M., Torres-
1253 Redondo, J. (2010). New results on thermal and photodesorption of CO ice using the novel InterStellar Astrochemistry
1254 Chamber (ISAC). *Astron. Astrophys.* 522, 108.
- 1255 Myers, A.T., McKee, C.F., Li, P.S. (2015). The CH⁺ abundance in turbulent, diffuse molecular clouds. *Mon. Not. R. Astron.*
1256 *Soc.* 453, 2747.
- 1257 National Academies of Sciences, Engineering, and Medicine (2021). Pathways to discovery in astronomy and astrophysics
1258 for the 2020s. Washington DC: The National Academies Press. DOI: 10.17226/26141.
- 1259 Neil, G.R. (2014). Accelerator sources for THz science: A review. *J. Infrared Millim. Terahertz Waves* 35, 5.
- 1260 Neufeld, D.A., Goicoechea, J.R., Sonnentrucker, P., Black, J.H., Pearson, J., Yu, S., Phillips, T.G., Lis, D.C., De Luca, M.,
1261 Herbst, E., Rimmer, P., Gerin, M., Bell, T.A., Boulanger, F., Cernicharo, J., Coutens, A., Dartois, E., Kaźmierczak, M.,
1262 Encrenaz, P., Falgarone, E., Geballe, T.R., Giesen, T., Godard, B., Goldsmith, P.F., Gry, C., Gupta, H., Hennebelle, P.,
1263 Hily-Blant, P., Joblin, C., Kołos, R., Krelowski, J., Martín-Pintado, J., Menten, K.M., Monje, R., Mookerjea, B., Perault,
1264 M., Persson, C., Plume, R., Salez, M., Schlemmer, S., Schmidt, M., Stutzki, J., Teyssier, D., Vastel, C., Cros, A., Klein, K.,
1265 Lorenzani, A., Philipp, S., Samoska, L.A., Shipman, R., Tielens, A.G.G.M., Szczerba, R., Zmuidzinas, J. (2010a).
1266 Herschel/HIFI observations of interstellar OH⁺ and H₂O⁺ towards W49N: A probe of diffuse clouds with a small molecular
1267 fraction. *Astron. Astrophys.* 521, L10.
- 1268 Neufeld, D.A., Sonnentrucker, P., Phillips, T.G., Lis, D.C., De Luca, M., Goicoechea, J.R., Black, J.H., Gerin, M., Bell,
1269 T.A., Boulanger, F., Cernicharo, J., Coutens, A., Dartois, E., Kaźmierczak, M., Encrenaz, P., Falgarone, E., Geballe, T.R.,
1270 Giesen, T., Godard, B., Goldsmith, P.F., Gry, C., Gupta, H., Hennebelle, P., Herbst, E., Hily-Blant, P., Joblin, C., Kołos, R.,
1271 Krelowski, J., Martín-Pintado, J., Menten, K.M., Monje, R., Mookerjea, B., Pearson, J., Perault, M., Persson, C., Plume,
1272 R., Salez, M., Schlemmer, S., Schmidt, M., Stutzki, J., Teyssier, D., Vastel, C., Yu, S., Cais, P., Caux, E., Liseau, R., Morris,
1273 P., Planesas, P. (2010b). Strong absorption by interstellar hydrogen fluoride: Herschel/HIFI observations of the sigh-line to
1274 G10.6-0.4 (W31C). *Astron. Astrophys.* 518, L108.
- 1275 Neufeld, D.A., Roueff, E., Snell, R.L., Lis, D.C., Benz, A.O., Bruderer, S., Black, J.H., De Luca, M., Gerin, M., Goldsmith,
1276 P.F., Gupta, H., Indriolo, N., Le Boulrot, J., Le Petit, F., Larsson, B., Melnick, G.J., Menten, K.M., Monje, R., Nagy, Z.,
1277 Phillips, T.G., Sandqvist, A., Sonnentrucker, P., van der Tak, F.F.S., Wolfire, M.G. (2012a). Herschel observations of
1278 interstellar chloronium. *Astrophys. J.* 748, 37.
- 1279 Neufeld, D.A., Falgarone, E., Gerin, M., Godard, B., Herbst, E., Pineau des Forêts, G., Vasyunin, A.I., Güsten, R.,
1280 Wiesemeyer, H., Ricken, O. (2012b). Discovery of interstellar mercapto radicals (SH) with the GREAT instrument on
1281 SOFIA. *Astron. Astrophys.* 542, L6.
- 1282 Neufeld, D.A., Godard, B., Gerin, M., Pineau des Forêts, G., Bernier, C., Falgarone, E., Graf, U.U., Güsten, R., Herbst, E.,
1283 Lesaffre, P., Schilke, P., Sonnentrucker, P., Wiesemeyer, H. (2015). Sulfur-bearing molecules in diffuse molecular clouds:
1284 New results from SOFIA/GREAT and the IRAM 30 m telescope. *Astron. Astrophys.* 577, 49.
- 1285 Neveu, M., Kim, H.J., Benner, S.A. (2013). The ‘strong’ RNA world hypothesis: Fifty years old. *Astrobiology* 13, 391.

- 1286 Noble, J.A., Cuppen, H.M., Coussan, S., Redlich, B., Ioppolo, S. (2020). Infrared resonant vibrationally induced re-
1287 structuring of amorphous solid water. *J. Phys. Chem. C* 124, 20864.
- 1288 Nuevo, M., Milam, S.N., Sandford, S.A., Elsila, J.E., Dworkin, J.P. (2009). Formation of uracil from the ultraviolet photo-
1289 irradiation of pyrimidine in pure H₂O ices. *Astrobiology* 9, 683.
- 1290 Nuevo, M., Milam, S.N., Sandford, S.A. (2012). Nucleobases and prebiotic molecules in organic residues produced from
1291 the ultraviolet photo-irradiation of pyrimidine in NH₃ and H₂O+NH₃ ices. *Astrobiology* 12, 295.
- 1292 Nuevo, M., Materese, C.K., Sandford, S.A. (2014). The photochemistry of pyrimidine in realistic astrophysical ices and
1293 the production of nucleobases. *Astrophys. J.* 793, 125.
- 1294 Öberg, K.I. (2016). Photochemistry and astrochemistry: Photochemical pathways to interstellar complex organic molecules.
1295 *Chem Rev.* 116, 9631.
- 1296 Öjekull, J., Andersson, P.U., Någård, M.B., Pettersson, J.B.C., Derkatch, A.M., Neau, A., Rosén, S., Thomas, R., Larsson,
1297 M., Österdahl, F., Semanjak, J., Danared, H., Källberg, A., af Ugglas, M., Marković, N. (2004). Dissociative
1298 recombination of NH⁺ and ND⁺ ions: Storage ring experiments and *ab initio* molecular dynamics. *J. Chem. Phys.*
1299 120, 7391.
- 1300 Orgel, L.E. (2004). Prebiotic chemistry and the origin of the RNA world. *Crit. Rev. Biochem. Mol. Biol.* 39, 99.
- 1301 Pan, K., Federman, S.R., Cunha, K., Smith, V.V., Welty, D.E. (2004). Cloud structure and physical conditions in star-
1302 forming regions from optical observations I. Data and component structure. *Astrophys. J. Suppl. Ser.* 151, 313.
- 1303 Pardo, J.R., Cernicharo, J., Serabyn, E. (2001). Atmospheric Transmission at Microwaves (ATM): An improved model for
1304 mm/sub-mm applications. *IEEE Trans. Antennas Propagation* 49, 1683.
- 1305 Park, G.B., Field, R.W. (2016). Perspective: The first ten years of broadband chirped pulse Fourier-transform microwave
1306 spectroscopy. *J. Chem. Phys.* 144, 200901.
- 1307 Parkes, S., Dunstan, M., Scott, P., Dillon, D., Spark, A., Ellison, B., Auriacombe, O. (2018). A wideband spectrometer in
1308 the Microsemi RTG4 FPGA. Paper presented at Data Systems in Aerospace Conference, DASIA 2018 (Oxford, UK)
- 1309 Peeters, Z., Botta, O., Charnley, S.B., Ruitenkamp, R., Ehrenfreund, P. (2003). The astrobiology of nucleobases. *Astrophys.*
1310 *J.* 593, L129.
- 1311 Perakis, F., Borek, J.A., Hamm, P. (2013). Three-dimensional infrared spectroscopy of isotope-diluted ice I_h. *J. Chem.*
1312 *Phys.* 139, 014501.
- 1313 Persson, C.M., Black, J.H., Cernicharo, J., Goicoechea, J.R., Hassel, G.E., Herbst, E., Gerin, M., De Luca, M., Bell, T.A.,
1314 Coutens, A., Falgarone, E., Goldsmith, P.F., Gupta, H., Kaźmierczak, M., Lis, D.C., Mookerjee, B., Neufeld, D.A., Pearson,
1315 J., Phillips, T.G., Sonnentrucker, P., Stutzki, J., Vastel, C., Yu, S., Boulanger, F., Dartois, E., Encrenaz, P., Geballe, T.R.,
1316 Giesen, T., Godard, B., Gry, C., Hennebelle, P., Hily-Blant, P., Joblin, C., Kołos, R., Krełowski, J., Martín-Pintado, J.,
1317 Menten, K., Monje, R., Perault, M., Plume, R., Salez, M., Schlemmer, S., Schmidt, M., Teyssier, D., Péron, I., Cais, P.,
1318 Gaufre, P., Cros, A., Ravera, L., Morris, P., Lord, S., Planesas, P. (2010). Nitrogen hydrides in interstellar gas. *Astron.*
1319 *Astrophys.* 521, L45.
- 1320 Pickett, H.M., Poynter, I.R.L., Cohen, E.A., Delitsky, M.L., Pearson, J.C., Müller, H.S.P. (1998). Submillimeter,
1321 millimeter, and microwave spectral line catalog. *J. Quant. Spectrosc. Radiat. Transfer* 60, 883.
- 1322 Plašil, R., Rednyk, S., Kovalenko, A., Tran, T.D., Roučka, Š., Dohnal, P., Novotný, O., Glosík, J. (2021). Experimental
1323 study on the CH⁺ formation from doubly charged carbon and molecular hydrogen. *Astrophys. J.* 910, 155.
- 1324 Potapov, A., Jäger, C., Henning, T., Jonusas, M., Krim, L. (2017). The formation of formaldehyde on interstellar
1325 carbonaceous grain analogues by O/H atom addition. *Astrophys. J.* 846, 131.
- 1326

- 1327 Preu, S., Döhler, G.H., Malzer, S., Wang, L.J., Gossard, A.C. (2011). Tunable continuous-wave terahertz photomixer
1328 sources and applications. *J. Appl. Phys.* 109, 061301.
- 1329 Price, D.C. (2016). Spectrometers and polyphase filterbanks in radio astronomy. ArXiv [Preprint]. Available at:
1330 <https://arxiv.org/abs/1607.03579> (Accessed April 24, 2021).
- 1331 Profeta, G., Scandolo, S. (2011). Far-infrared spectrum of ice I_h: A first-principles study. *Phys. Rev. B* 84, 024103.
- 1332 Qasim, D., Fedoseev, G., Chuang, K.-J., He, J., Ioppolo, S., van Dishoeck, E.F., Linnartz, H. (2020). An experimental study
1333 of the surface formation of methane in interstellar molecular clouds. *Nat. Astron.* 4, 781.
- 1334 Raffaelli, N. (2011). “Nicotinamide Coenzyme Synthesis: A Case of Ribonucleotide Emergence or a Byproduct of the RNA
1335 World?” in *Origins of Life: The Primal Self-Organization*, eds. R. Egel, D.H. Lankenau, A.Y. Mulikidjanian (Heidelberg,
1336 Springer), 185-208.
- 1337 Redlich, B., van der Meer, L., Zacharias, H., Meijer, G., von Helden, G. (2003). FEL induced dynamics of small molecules
1338 on surfaces: N₂O on NaCl (1 0 0). *Nucl. Instrum. Methods Phys. Res. A* 507, 556.
- 1339 Rigopoulou, D., Pearson, C., Ellison, B., Wiedner, M., Ossenkopf-Okada, V., Tan, B.K., Garcia-Bernette, I., Gerin, M.,
1340 Yassin, G., Caux, E., Molinari, S., Goicoechea, J.R., Savini, G., Hunt, L.K., Lis, D.C., Goldsmith, P.F., Aalto, S., Magdis,
1341 G., Kramer, C. (2021). The Far-Infrared Spectroscopic Surveyor (FIRSS). *Exp. Astron.* DOI: 10.1007/s10686-021-09716-
1342 w
- 1343 Rosenthal, D., Beeman, J.W., Geis, N., Grözinger, U., Höhle, R., Katterloher, R.O., Kraft, S., Looney, L.W., Poglitsch, A.,
1344 Raab, W., Richter, H. (2002). Stressed Ge:Ga detector arrays for PACS and FIFI LS. Far-IR, sub-mm and mm Detector
1345 Technology Workshop (Monterey, CA).
- 1346 Rothman, L.S., Gordon, I.E., Barbe, A., Benner, D.C., Bernath, P.F., Birk, M., Boudon, V., Brown, L.R., Campargue, A.,
1347 Champion, J.P., Chance, K., Coudert, L.H., Dana, V., Devi, V.M., Fally, S., Flaud, J.M., Gamache, R.R., Goldman,
1348 A., Jacquemart, D., Kleiner, I., Lacorne, N., Lafferty, W.J., Mandin, J.Y., Massie, S.T., Mikhailenko, S.N., Miller,
1349 C.E., Moazzen-Ahmadi, N., Naumenko, O.V., Nikitin, A.V., Orphal, J., Perevalov, V.I., Perrin, A., Predoi-Cross,
1350 A., Rinsland, C.P., Rotger, M., Šimečková, M., Smith, M.A.H., Sung, K., Tashkun, S.A., Tennyson, J., Toth, R.A.,
1351 Vandaele, A.C., Vander Auwera, J. (2009). The HITRAN 2008 molecular spectroscopic database. *J. Quant.*
1352 *Spectrosc. Radiat. Transfer*, 110, 533.
- 1353 Sandford, S.A., Nuevo, M., Bera, P.P., Lee, T.J. (2020). Prebiotic astrochemistry and the formation of molecules of
1354 astrobiological interest in interstellar clouds and protostellar disks. *Chem. Rev.* 120, 4616.
- 1355 Schilke, P., Neufeld, D.A., Müller, H.S.P., Comito, C., Bergin, E.A., Lis, D.C., Gerin, M., Black, J.H., Wolfire, M., Indriolo,
1356 N., Pearson, J.C., Menten, K.M., Winkel, B., Sánchez-Monge, Á., Möller, T., Godard, B., Falgarone, E. (2014). Ubiquitous
1357 argonium (ArH⁺) in the diffuse interstellar medium: A molecular tracer of almost purely atomic gas. *Astron. Astrophys.*
1358 566, 29.
- 1359 Scibelli, S., Shirley, Y. (2020). Prevalence of complex organic molecules in starless and prestellar cores within the Taurus
1360 molecular cloud. *Astrophys. J.* 891, 73.
- 1361 Semaniak, J., Larson, Å., Le Padellec, A., Strömholm, C., Larsson, M., Rosén, S., Peverall, R., Danared, H., Djuric, N.,
1362 Dunn, G.H. (1998). Dissociative recombination and excitation of CH⁺: Absolute cross sections and branching
1363 fractions. *Astrophys. J.* 498, 886.
- 1364 Shalit, A., Perakis, F., Hamm, P. (2013). Two-dimensional infrared spectroscopy of isotope-diluted low density amorphous
1365 ice. *J. Phys. Chem. B* 117, 15512.
- 1366 Siebertz, O., Schmölling, F., Gal, C., Schloeder, F., Hartogh, P., Natale, V., Schieder, R. (2007). The wide-band
1367 spectrometer (WBS) for the HIFI instrument of Herschel. *Proc. ISSTT*, 135.
- 1368 Smith, D. (1992). The ion chemistry of interstellar clouds. *Chem. Rev.* 92, 1473.

- 1369 Smith, I.W.M. (2011). Laboratory astrochemistry: Gas-phase processes. *Annu. Rev. Astron. Astrophys.* 49, 29.
- 1370 Snow, T.P., McCall, B.J. (2006). Diffuse atomic and molecular clouds. *Annu. Rev. Astron. Astrophys.* 44, 367.
- 1371 Snow, T.P., Bierbaum, V.M. (2008). Ion chemistry in the interstellar medium. *Annu. Rev. Anal. Chem.* 1, 229.
- 1372 Soma, T., Sakai, N., Watanabe, Y., Yamamoto, S. (2018). Complex organic molecules in Taurus Molecular Cloud-1.
1373 *Astrophys. J.* 854, 116.
- 1374 Sonnentrucker, P., Neufeld, D.A., Phillips, T.G., Gerin, M., Lis, D.C., De Luca, M., Goicoechea, J.R., Black, J.H., Bell,
1375 T.A., Boulanger, F., Cernicharo, J., Coutens, A., Dartois, E., Kaźmierczak, M., Encrenaz, P., Falgarone, E., Geballe, T.R.,
1376 Giesen, T., Godard, B., Goldsmith, P.F., Gry, C., Gupta, H., Hennebelle, P., Herbst, E., Hily-Blant, P., Joblin, C., Kołos, R.,
1377 Krełowski, J., Martín-Pintado, J., Menten, K.M., Monje, R., Mookerjea, B., Pearson, J., Perault, M., Persson, C.M., Plume,
1378 R., Salez, M., Schlemmer, S., Schmidt, M., Stutzki, J., Teyssier, D., Vastel, C., Yu, S., Caux, E., Güsten, R., Hatch, W.A.,
1379 Klein, T., Mehdi, I., Morris, P., Ward, J.S. (2010). Detection of hydrogen fluoride absorption in diffuse molecular clouds
1380 with Herschel/HIFI: an ubiquitous tracer of molecular gas. *Astron. Astrophys.* 521, L12.
- 1381
- 1382 Suzuki, T., Khosropanah, P., Ridder, M.L., Hijmering, R.A., Gao, J.R., Akamatsu, H., Gottardi, L., van der Kuur, J.,
1383 Jackson, B.D. (2016). Development of ultra-low noise TES bolometer arrays. *J. Low Temp. Phys.* 184, 52.
- 1384 Taylor, S.D., Duley, W.W. (1997). The formation of long carbon chains in diffuse clouds. *Mon. Not. R. Astron. Soc.* 286,
1385 344.
- 1386 Teolis, B.D., Waite, J.H. (2016). Dione and Rhea seasonal exospheres revealed by Cassini CAPS and INMS. *Icarus* 272,
1387 277.
- 1388 Theulé, P., Endres, C., Hermanns, M., Bossa, J.B., Potapov, A. (2020). High-resolution gas-phase spectroscopy of
1389 molecules desorbed from an ice surface: A proof-of-principle study. *ACS Earth Space Chem.* 4, 86.
- 1390 Thiel, V., Belloche, A., Menten, K.M., Garrod, R.T., Müller, H.S.P. (2017). Complex organic molecules in diffuse clouds
1391 along the line of sight to Sagittarius B2. *Astron. Astrophys.* 605, L6.
- 1392 Thornton, R.J., Ade, P.A.R., Aiola, S., Angilè, F.E., Amiri, M., Beall, J.A., Becker, D.T., Cho, H.M., Choi, S.K., Corlies,
1393 P. (2016). The Atacama cosmology telescope: The polarization sensitive ACTPol Instrument. *Astrophys. J. Suppl.*
1394 *Ser.* 227, 21.
- 1395 Tielens, A.G.G.M. (2008). Interstellar polycyclic aromatic hydrocarbon molecules. *Annu. Rev. Astron. Astrophys.* 46, 289.
- 1396 Townes, C., Schawlow, A. (2013). *Microwave Spectroscopy*. Mineola (NY): Dover Publications.
- 1397 Trofimov, V., Varentsova, S.A. (2016). Essential limitations of the standard THz TDS method for substance detection and
1398 identification and a way of overcoming them. *Sensors* 16, 502.
- 1399 Valavanis, A., Auriacombe, O., Rawlings, T., Han, Y., Rea, S., Crook, M., Arena, C., Walker, D., Brooks, D., Yu, G., Li,
1400 L., Davies, A.G., Savini, G., Linfield, E.H., Ellison, B., Saenz, E. (2019). Waveguide-integrated THz quantum-cascade
1401 lasers for atmospheric research satellite payloads. *Proc. IEEE MTT-S IWS*, 1.
- 1402 Valdivia, V., Godard, B., Hennebelle, P., Gerin, M., Lesaffre, P., Le Bourlot, J. (2017). Origin of CH⁺ in diffuse molecular
1403 clouds. *Astron. Astrophys.* 600, 114.
- 1404 van Dishoeck, E.F., Black, J.H. (1986). Comprehensive models of diffuse interstellar clouds: physical conditions and
1405 molecular abundances. *Astrophys. J. Suppl. Ser.* 62, 109.
- 1406 van Dishoeck, E.F., Blake, G.A. (1998). Chemical evolution of star-forming regions. *Annu. Rev. Astron. Astrophys.* 36,
1407 317.

- 1408 van Dishoeck, E.F., Herbst, E., Neufeld, D.A. (2013). Interstellar water chemistry: From laboratory to observations. *Chem.*
1409 *Rev.* 113, 9043.
- 1410 van Dishoeck, E.F. (2014). Astrochemistry of ice, dust, and gas: Introduction and overview. *Faraday Discuss.* 168, 9.
- 1411 van Dishoeck, E.F. (2017). Astrochemistry: Overview and challenges. *Proc. Int. Astron. Union* 13, 3.
- 1412 van Exter, M., Fattinger, C., Grischkowsky, D. (1989). High-brightness terahertz beams characterized with an ultrafast
1413 detector. *Appl. Phys. Lett.* 55, 337.
- 1414 Viti, S., Williams, D.A., O'Neill, P.T. (2000). Hydrocarbons in diffuse and translucent clouds. *Astron. Astrophys.* 354,
1415 1062.
- 1416 Wakelam, V., Bron, E., Cazaux, S., Dulieu, F., Gry, C., Guillard, P., Habart, E., Hornkær, L., Morisset, S., Nyman, G.,
1417 Pirronello, V., Price, S.D., Valdivia, V., Vidali, G., Watanabe, N. (2017). H₂ formation on interstellar dust grains: The
1418 viewpoints of theory, experiments, models and observations. *Mol. Astrophys.* 9, 1.
- 1419 Watanabe, N., Kouchi, A. (2002). Efficient formation of formaldehyde and methanol by the addition of hydrogen atoms to
1420 CO in H₂O-CO ice at 10 K. *Astrophys. J.* 571, 173.
- 1421 Wehres, N., Heyne, B., Lewen, F., Hermanns, M., Schmidt, B., Endres, C., Graf, U.U., Higgins, D.R., Schlemmer, S.
1422 (2017). 100 GHz room-temperature laboratory emission spectrometer. *Proc. Int. Astron. Union* 13, 332
- 1423 Wehres, N., Maßsen, J., Borisov, K., Schmidt, B., Lewen, F., Graf, U.U., Honingh, C.E., Higgins, D.R., Schlemmer, S.
1424 (2018). A laboratory heterodyne emission spectrometer at submillimeter wavelengths. *Phys. Chem. Chem. Phys.* 20, 5530.
1425
- 1426 Weinreb, S. (1961). Digital Radiometer. *Proc. IEEE* 49, 1099.
- 1427 White, H.B. (1976). Coenzymes as fossils of an earlier metabolic state. *J. Mol. Evol.* 7, 101.
- 1428 Widicus-Weaver, S.L., Butler, R.A.H., Drouin, B.J., Petkie, D.T., Dyl, K.A., De Lucia, F.C., Blake, G.A. (2005).
1429 Millimeter-wave and vibrational state assignments for the rotational spectrum of glycolaldehyde. *Astrophys. J. Suppl. Ser.*
1430 158, 188.
- 1431 Widicus-Weaver, S.L. (2019). Millimeter wave and submillimeter wave laboratory spectroscopy in support of observational
1432 astronomy. *Annu. Rev. Astron. Astrophys.* 57, 79.
- 1433 Wiesemeyer, H., Güsten, R., Heyminck, S., Hübers, H.W., Menten, K.M., Neufeld, D.A., Richter, H., Simon, R., Stutzki,
1434 J., Winkel, B., Wyrowski, F. (2016). Far-infrared study of tracers of oxygen chemistry in diffuse clouds. *Astron. Astrophys.*
1435 585, 76.
- 1436 Williams, D.A., Hartquist, T.W. (1999). The chemistry of star-forming regions. *Acc. Chem. Res.* 32, 334.
- 1437 Willis, E.R., Garrod, R.T., Belloche, A., Müller, H.S.P., Barger, C.J., Bonfard, M., Menten, K.M. (2020). Exploring
1438 molecular complexity with ALMA (EMoCA): Complex isocyanides in Sgr B2(N). *Astron. Astrophys.* 636, 29.
- 1439 Wilson, T.L., Rohlfs, K., Hüttemeister, S. (2008). *Tools of Radio Astronomy*. Heidelberg: Springer.
- 1440 Wlodarczak, G. (1995). Rotational spectroscopy and astrochemistry. *J. Mol. Struct.* 347, 131.
- 1441 Wyrowski, F., Menten, K.M., Güsten, R., Belloche, A. (2010). First interstellar detection of OH⁺. *Astron. Astrophys.* 518,
1442 26.
- 1443 Xue, C., Willis, E.R., Loomis, R.A., Lee, K.L.K., Burkhardt, A.M., Shingledecker, C.N., Charnley, S.B., Cordiner, M.A.,
1444 Kalenskii, S., McCarthy, M.C., Herbst, E., Remijan, A.J., McGuire, B.A. (2020). Detection of interstellar HC₄NC and an
1445 investigation of isocyanopolyne chemistry under TMC-1 conditions. *Astrophys. J. Lett.* 900, L9.

- 1446 Yamamoto, S. (2017). Introduction to Astrochemistry: Chemical Evolution from Interstellar Clouds to Star and Planet
1447 Formation. Tokyo: Springer Japan.
- 1448 Yocum, K.M., Smith, H.H., Todd, E.W., Mora, L., Gerakines, P.A., Milam, S.N., Widicus-Weaver, S.L. (2019).
1449 Millimeter/submillimeter spectroscopy detection of desorbed ices: A new technique in laboratory astrochemistry. J.
1450 Phys. Chem. A 123, 8702.
- 1451 Zernickel, A., Schilke, P., Schmiedeke, A., Lis, D.C., Brogan, C.L., Ceccarelli, C., Comito, C., Emprechtinger, M., Hunter,
1452 T.R., Möller, T. (2012). Molecular line survey of the high-mass star-forming region NGC 6334I with Herschel/HIFI
1453 and the submillimeter array. Astron. Astrophys. 546, 87.
- 1454 Zhen, J., Castellanos, P., Paardekooper, D.M., Linnartz, H., Tielens, A.G.G.M. (2014). Laboratory formation of fullerenes
1455 from PAHs: Top-down interstellar chemistry. Astrophys. J. Lett. 797, L30.