



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Growth mormone and prolactin - so much still to learn

Citation for published version:

Le Tissier, P & Grattan, DR 2020, 'Growth mormone and prolactin - so much still to learn', *Journal of Neuroendocrinology*. <https://doi.org/10.1111/jne.12909>

Digital Object Identifier (DOI):

[10.1111/jne.12909](https://doi.org/10.1111/jne.12909)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Neuroendocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Editorial

Growth Hormone And Prolactin- So Much Still To Learn.

Paul R Le Tissier¹ and David R Grattan²

1. Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh UK

2. Centre for Neuroendocrinology, University of Otago, Dunedin, New Zealand

The existence and importance of prolactin and growth hormone (GH) has been known for close to a century^{1,2} and key aspects of their regulation and actions are well understood. This may have led to the assumption by many that there is little still to learn about the regulation, production and actions of these hormones, a view hotly contested by those of us involved in their study. The articles in this special issue, based on research presented at the 2019 FASEB meeting on Growth Hormone/Prolactin Family in Health and Disease, are testament to this. Presentations at the meeting ranged from understanding hypothalamic regulation of these two hormones, their production and secretion from both the pituitary and placenta, the wide range of their physiological functions in a range of tissues, including the brain, and the mechanism and consequences of dysregulation. One thing is for certain – it is no longer appropriate to be thinking about these hormones as having the one-dimensional function implied by their names: “growth” or “lactation”. Rather, these pleiotropic hormones have evolved to contribute to multiple adaptive functions, ranging from metabolism, immune function, reproduction and aging (to only mention a few). The common ancestral origin of the two hormones, as well as their receptors and downstream signal transduction pathways, makes a meeting considering similarities and differences in their regulation and function highly informative. and this is highlighted in the collection of reviews and research articles in this special issue that focuses on some of the neuroendocrine topics covered within the meeting.

The hypothalamic regulation of prolactin by tuberoinfundibular dopamine (TIDA) neurons is well established³ but the regulation of these neurons themselves less clear, as is how this is altered to reduce short-loop feedback of prolactin and ensure high levels of hormone to support lactation. Further understanding of TIDA neuron regulation is described here by Amamri and Broberger, who postulated that γ -aminobutyric acid (GABA) produced by a subset of TIDA neurons may act through GABAB receptors in the arcuate nucleus, with GABAB acting as an autoreceptor⁴. Their results demonstrate a role for GABAB receptors in both pre- and post-synaptic regulation of TIDA neurons, suggesting a key role for GABA in the regulation of prolactin by dopamine.

Recent studies have shown that TIDA neurons remain responsive to prolactin during lactation, discounting a mechanism where a loss of short-loop feedback supports the high level of hormone at this time⁵. Two studies reported here provide insights into the plasticity of TIDA neurons that may underlie their altered function during lactation. Yip et al⁶ elegantly demonstrate a change in somatic spine density that would be expected to result in increased TIDA neuron

excitatory output; it is possible that this supports release of a TIDA neuron neuropeptide, such as met-enkephalin that has been shown to be produced in these neurons in lactation⁷, that would stimulate prolactin release in a feed-forward loop. This possibility is supported by the study of Silva et al.⁸ in this issue who show that reduced dopamine alone does not account for suckling induced rises in prolactin, indicative of a role for a prolactin releasing factor, such as met-enkephalin.

In addition to hypothalamic regulation, peripheral signals also regulate GH and prolactin production and release but the mechanism of many of these are poorly understood. Cell-specific manipulation of receptor expression can provide insights into this, as is demonstrated in the study of Allensworth-James et al⁹ who have shown that the decreased GH secretion following loss of leptin receptors specifically in GH-secreting somatotrophs is mediated by alterations in both gene transcription and translation. Key mediators in the regulation of prolactin have also been identified in fish, where prolactin has an osmoregulatory role¹⁰. Despite the altered role for prolactin, regulatory mechanisms are likely conserved across phyla and the review of Seale et al¹¹ identifies regulators of prolactin gene expression likely to have similar roles in mammals. It is also important to remember that members of the prolactin and GH family are also produced in the placenta, with dramatic effects on physiology in pregnancy; Cattini, et al.¹² review studies of the mechanism linking obesity with reduced placental lactogen production in pregnant women, which may increase the risk of peripartum depression.

An additional feature of both prolactin and GH are their pleiotropic actions, an excellent example of which is the description of the role of prolactin in vascularisation of the retina¹³. This adds to the multiple roles of prolactin, which have recently focused on its importance for regulation of metabolism, which are reviewed here by Lopez-Vichi, et al.¹⁴. Of course, GH has a major role in metabolic regulation and two papers from the Berryman/Kopchick group highlight the effects of GH on adipose tissue in mouse models with either a loss¹⁵ or increase¹⁶ in GH actions on this tissue, which differs between fat depots. A deeper understanding of the intracellular pathways mediating GH and prolactin actions may provide insights into tissue specific effects and the importance of temporal aspects of hormone action. Two papers describing actions of prolactin over differing timescales highlight the necessity of considering the canonical pathways activated by GH and prolactin through Stat5 activation¹⁷ and potential novel pathways mediating rapid action¹⁸.

The diversity of papers presented in this special issue and the novelty of many findings illustrates that study of the prolactin and GH systems is not merely a matter of refining understanding of a well-described system. Exciting advances in all aspects of both hormones are currently being made and will continue for the foreseeable future, from regulation by the brain and periphery, to temporal and differential effects on tissues that are only beginning to be understood.

1. Medeiros A, Siegel Watkins E. Live Longer Better: The Historical Roots of Human Growth Hormone as Anti-Aging Medicine. *Journal of the History of Medicine and Allied Sciences*. 2018;73(3):333-359.
2. Bates RW, Riddle O. THE PREPARATION OF PROLACTIN. *Journal of Pharmacology and Experimental Therapeutics*. 1935;55(3):365.
3. Grattan DR. 60 YEARS OF NEUROENDOCRINOLOGY: The hypothalamo-prolactin axis. *Journal of Endocrinology*. 2015;226(2):T101-T122.
4. Ammari R, Broberger C. Pre- and post-synaptic modulation by GABAB receptors of rat neuroendocrine dopamine neurones. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12881.
5. Romanò N, Yip SH, Hodson DJ, et al. Plasticity of hypothalamic dopamine neurons during lactation results in dissociation of electrical activity and release. *Journal of Neuroscience*. 2013;33(10):4424-4433.
6. Yip SH, Araujo-Lopes R, Szawka RE, et al. Morphological plasticity of the tuberoinfundibular dopaminergic neurones in the rat during the oestrous cycle and lactation. *Journal of Neuroendocrinology*.n/a(n/a):e12884.
7. Yip SH, Romanò N, Gustafson P, et al. Elevated Prolactin during Pregnancy Drives a Phenotypic Switch in Mouse Hypothalamic Dopaminergic Neurons. *Cell Reports*. 2019;26(7):1787-1799.e1785.
8. Silva KSC, Aquino NSS, Gusmao DO, Henriques PC, Reis AM, Szawka RE. Reduced dopaminergic tone during lactation is permissive to the hypothalamic stimulus for suckling-induced prolactin release. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12880.
9. Allensworth-James ML, Odle AK, Lim J, et al. Metabolic signalling to somatotrophs: Transcriptional and post-transcriptional mediators. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12883.
10. Hirano T. The spectrum of prolactin action in teleosts. *Prog Clin Biol Res*. 1986;205:53-74.
11. Seale AP, Malintha GHT, Celino-Brady FT, et al. Transcriptional regulation of prolactin in a euryhaline teleost: characterization of gene promoters through in silico and transcriptome analyses. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12905.
12. Cattini PA, Jin Y, Jarmasz JS, Noorjahan N, Bock ME. Obesity and regulation of human placental lactogen production in pregnancy. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12859.
13. Vázquez-Membrillo M, Siqueiros-Márquez L, Núñez FF, et al. Prolactin stimulates the vascularisation of the retina in newborn mice under hyperoxia conditions. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12858.
14. Lopez-Vicchi F, De Winne C, Brie B, Sorianello E, Ladyman SR, Becu-Villalobos D. Metabolic functions of prolactin: Physiological and pathological aspects. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12888.
15. Young JA, Henry BE, Benencia F, et al. GHR^{-/-} Mice are protected from obesity-related white adipose tissue inflammation. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12854.
16. Duran-Ortiz S, Young JA, Jara A, et al. Differential gene signature in adipose tissue depots of growth hormone transgenic mice. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12893.

17. Gustafson P, Ladyman SR, McFadden S, et al. Prolactin receptor-mediated activation of pSTAT5 in the pregnant mouse brain. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12901.
18. Georgescu T, Ladyman SR, Brown RSE, Grattan DR. Acute effects of prolactin on hypothalamic prolactin receptor expressing neurones in the mouse. *Journal of Neuroendocrinology*. 2020;n/a(n/a):e12908.