



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Heritable variation in telomere length predicts mortality in soay sheep

Citation for published version:

Froy, H, Underwood, S, Dorrens, J, Seeker, L, Watt, K, Wilbourn, R, Pilkington, J, Harrington, L, Pemberton, JM & Nussey, DH 2021, 'Heritable variation in telomere length predicts mortality in soay sheep', *Proceedings of the National Academy of Sciences*, vol. 118, no. 15, e2020563118.
<https://doi.org/10.1073/pnas.2020563118>

Digital Object Identifier (DOI):

[10.1073/pnas.2020563118](https://doi.org/10.1073/pnas.2020563118)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Proceedings of the National Academy of Sciences

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Heritable variation in telomere length predicts mortality in**

2 **Soay sheep**

3 Hannah Froy^{1,2*}, Sarah L. Underwood², Jennifer Dorrens², Luise A. Seeker², Kathryn Watt²,
4 Rachael V. Wilbourn², Jill G. Pilkington², Lea Harrington³, Josephine M. Pemberton², Daniel
5 H. Nussey^{2*}

6 ¹Centre for Biodiversity Dynamics, Institute for Biology, Norwegian University for Science
7 and Technology (NTNU), Trondheim 7014 Norway

8 ²Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh,
9 Edinburgh EH9 3FL UK

10 ³Institute for Research in Immunology and Cancer, Université de Montréal, Montréal H3C
11 3J7 Canada.

12 Corresponding authors:

13 *Hannah Froy; Centre for Biodiversity Dynamics, Institute for Biology, Norwegian University
14 for Science and Technology (NTNU), Trondheim 7014 Norway; Tel: +44 775 3285418;
15 Email: hannah.froy@ntnu.no

16 *Daniel Nussey; Institute of Evolutionary Biology, University of Edinburgh, Ashworth
17 Laboratories, Charlotte Auerbach Road, Edinburgh EH9 3FL UK; Tel: +44 131 6505494;
18 Email: dan.nussey@ed.ac.uk

19 ORCIDs: Hannah Froy (0000-0003-2965-3526), Luise Seeker (0000-0001-6375-6372), Lea
20 Harrington (0000-0002-4977-2744), Josephine Pemberton (0000-0002-0075-1504), Daniel
21 Nussey (0000-0002-9985-0317).

22 **Classification:** Biological sciences – Evolution.

23 **Keywords (3–5):** Survival, longevity, ageing, senescence, biomarker.

24 **Author Contributions**

25 HF & DHN designed the study and wrote the first draft of the manuscript; JMP & JGP
26 conducted fieldwork and managed the long-term study; SLU, JD, LAS, KW & RVW
27 conducted the labwork; HF conducted the analyses; all authors contributed to the revised
28 manuscript.

29 **This PDF file includes:** Main Text; Figures 1 to 3.

30

31 **Abstract**

32 Telomere length (TL) is considered an important biomarker of whole-organism health and
33 ageing. Across humans and other vertebrates, short telomeres are associated with
34 increased subsequent mortality risk, but the processes responsible for this correlation
35 remain uncertain. A key unanswered question is whether TL–mortality associations arise
36 due to positive effects of genes or early-life environment on both an individual’s average
37 lifetime TL and their longevity, or due to more immediate effects of environmental stressors
38 on within-individual TL loss and increased mortality risk. Addressing this question requires
39 longitudinal TL and life-history data across the entire lifetimes of many individuals, which are
40 difficult to obtain for long-lived species like humans. Using longitudinal data and samples
41 collected over nearly two decades, as part of a long-term study of wild Soay sheep, we
42 dissected an observed positive association between TL and subsequent survival using
43 multivariate quantitative genetic models. We found no evidence that telomere attrition was
44 associated with increased mortality risk, suggesting that TL is not an important marker of
45 biological ageing or exposure to environmental stress in our study system. Instead, we find
46 that among-individual differences in average TL are associated with increased lifespan. Our
47 analyses suggest that this correlation between an individual’s average TL and lifespan has a
48 genetic basis. This demonstrates that TL has the potential to evolve under natural
49 conditions, and suggests an important role of genetics underlying the widespread
50 observation that short telomeres predict mortality.

51 **Significance Statement**

52 Telomeres play an important role in ageing and having relatively short telomeres is
53 associated with an increased risk of death in humans and other animals. Telomere length is
54 influenced by both genetic and environmental factors, both of which could potentially drive
55 the positive association with survival. We used lifelong telomere length measurements from
56 a population of wild sheep to disentangle this relationship. For the first time in a natural
57 population, our analyses reveal a genetic correlation between telomere length and longevity
58 but no association between telomere shortening and mortality risk. These findings have
59 important implications for our understanding of telomere dynamics and their role in health
60 and lifespan.

61

62 Main text

63 Introduction

64 Telomeres are repetitive sequences of non-coding DNA found at the terminal ends of linear
65 chromosomes, and they play an important role in maintaining DNA stability and integrity (1-
66 3). Telomeres shorten during cell replication and in response to oxidative stress (4, 5), and
67 cellular senescence and apoptosis is triggered once telomeres reach a critically short
68 threshold (2). The important role of telomeres in cellular senescence has led to telomere
69 shortening being considered as one of nine “hallmarks of ageing”, and average telomere
70 length (TL) as an important biomarker of whole-organism health and biological ageing (6). In
71 humans, relatively short leukocyte telomeres have been linked to a range of age-related
72 diseases such as diabetes, cancer and cardiovascular disease (7-9) and increased
73 subsequent mortality risk (10-12). A recent meta-analysis suggests this pattern may
74 generalise beyond humans: across studies from 20 non-model vertebrate species
75 (predominantly birds) there was an overall positive association between telomere length and
76 subsequent survival (13). Although evidence for a causal role for telomeres in whole-
77 organism ageing and longevity remains weak (14), these findings highlight the potential
78 significance of TL as a biomarker of human and animal health (15, 16) and for our
79 understanding of life-history evolution (17, 18).

80 Studies in humans and other vertebrates have found evidence for consistent differences in
81 TL among individuals over multiple measurements (19, 20). Such repeatable among-
82 individual differences in any trait may result from the trait being under genetic influence, from
83 long-term effects of the early-life environment, and/or environmental conditions that persist
84 across the lifetime. There is good evidence that variation in average TL in blood cells has a
85 genetic basis in humans and other vertebrates, although estimates of the heritability (the
86 proportion of variation attributed to additive genetic effects) of TL are variable (21, 22).
87 Recent studies of wild vertebrates have also revealed considerable variation in adult TL
88 among birth cohorts, suggesting persistent impacts of early-life environment (23, 24). At the
89 same time, there is growing evidence that TL is highly dynamic across an individual’s
90 lifetime, and meta-analyses of human and non-human animal studies show that experience
91 of diverse forms of environmental stress are predictive of shorter TL (25-27). Indeed, some
92 studies using longitudinal TL data have found that telomere shortening over successive
93 measurements rather than telomere length *per se* is predictive of mortality (28-30). Thus, the
94 emerging picture from studies in humans and other vertebrates is that shorter TL generally

95 predicts increased risk of subsequent mortality, and that variation in TL is under the
96 influence of both genetics and environmental stressors.

97 The observation that shorter TL measurements predict increased mortality risk could be
98 underpinned by two, non-mutually exclusive processes operating across the lifetimes of
99 individuals. Firstly, individuals may differ in their average TL across life, and individuals with
100 shorter TL may be shorter lived. This pattern is referred to as the 'selective disappearance'
101 of individuals with shorter telomeres, and it implies that TL reflects constitutive differences
102 among individuals (for example due to genetics or differences in early-life environment)
103 which shape their longevity (31, 32). Secondly, individuals may differ in their pattern of TL
104 change over time, and individuals showing the greatest telomere loss across successive
105 measurements are more likely to die subsequently. This pattern is consistent with the idea
106 that within-individual telomere dynamics reflect recent and cumulative experiences of
107 environmental stress and physiological deterioration that also predict mortality. Neither
108 pattern necessarily implies a causal role for telomeres in driving the mortality risk of an
109 organism, because associations between TL and survival could result from both traits being
110 correlated with underlying, unmeasured variables which causally impact survival (14, 18).
111 Nevertheless, unravelling the contribution of genetics, early-life environment and more
112 immediate telomere shortening to the observed association between TL and survival is
113 essential for our understanding of TL as a biomarker of health and ageing (19).

114 To our knowledge, no study to date has assessed the relative importance of the different
115 processes underlying the relationship between TL and mortality risk across the entire
116 lifespan. To do so demands repeated measurements from across life to characterise among-
117 and within-individual variation in telomere length, a population pedigree or genomic
118 information to separate genetic and environmental sources of variation, and detailed
119 information on individual health and fitness outcomes over the lifetime. Here, we use a
120 multivariate mixed-effects modelling approach to analyse extensive, longitudinal data from a
121 long-term study of wild Soay sheep living on St Kilda, Scotland, to distinguish between
122 possible models of *why* shorter TL predicts increased mortality risk. We find that the
123 observed positive association between TL and mortality in this system is underpinned by
124 selective disappearance of individuals with shorter average TL. Importantly, our results
125 suggest this is largely driven by genetically-based differences in both TL and longevity.

126 **Results**

127 Soay sheep resident to our study area on St Kilda have been individually marked at birth and
128 closely monitored and repeatedly blood sampled across their lifetimes. Here, we measured
129 relative telomere length (RTL) in 3641 samples collected from 1586 individual sheep over a
130 19-year period (see Methods). We found that RTL declined with age in Soay sheep, with a
131 more rapid initial decline between measurements at around 4 and 16 months, followed by a
132 slower linear decline thereafter (Figure 1). The best fitting age function in our models of RTL
133 included a two-level factor for age class (lambs and adults aged ≥ 1 year) and a linear term
134 for age in years, which is equivalent to a segmented regression with a threshold at one year
135 of age (SI Appendix, Tables S1 & S2). There was considerable variation in RTL within age
136 groups (SI Appendix, Fig. S1). There was limited evidence for a sex difference in either
137 average RTL or the rate of change in RTL with age, although the significance of sex in our
138 model depended on model structure (see SI Appendix). The individual repeatability of RTL
139 over the lifespan was 0.214 (95%CI 0.169–0.252, SI Appendix, Table S2). Excluding the
140 variance attributed to qPCR plate and row (which represents measurement error) from the
141 denominator, the repeatability was 0.241 (95%CI 0.204–0.282, SI Appendix, Table S2).
142 Although RTL declined with age on average, we found evidence for consistent differences in
143 RTL among individuals.

144 There was detectable additive genetic variance for relative telomere length across all ages in
145 the population (Figure 2; SI Appendix, Table S3). The heritability of telomere length (the
146 proportion of variance explained by additive genetic effects) was 0.204 (95%CI: 0.158–
147 0.252). The permanent environment effect was bound at zero (<0.001 , 95%CI: <0.001 –
148 0.017), indicating that individual repeatability in telomere length could be almost entirely be
149 attributed to genetic rather than environmental effects. Maternal effects also explained a very
150 small proportion of the variance (<0.001 , 95%CI <0.001 –0.034, estimate bound at zero). The
151 year the sample was collected, qPCR plate and qPCR row each explained 3-4% of the
152 variance in telomere length (year: 0.031, 95%CI 0.014–0.084; qPCR plate: 0.043, 0.027–
153 0.066; qPCR row: 0.038, 0.014–0.172). Excluding the measurement error terms of qPCR
154 plate and row from the total phenotypic variance, the heritability of telomere length was
155 0.233 (95%CI 0.189–0.279). This shows that variation in telomere length has a genetic basis
156 in wild Soay sheep.

157 We went on to estimate the genetic correlation between RTL expressed in lambs, which are
158 still developing at the time of measurement (aged 4 months), and in older individuals, which
159 are sexually mature and have largely completed growth. When estimated with a bivariate
160 model of lamb and adult telomere length, the heritability of telomere length in lambs was
161 estimated to be 0.285 (95%CI 0.206–0.369), and in adults 0.210 (95%CI: 0.156–0.255)

162 (Figure 2; SI Appendix, Table S4). The genetic correlation between lambs and adults was
163 close to 1 (0.916, 95%CI: 0.806–0.996, SI Appendix, Table S4), implying that largely the
164 same or linked genes influenced RTL across age groups. The residual correlation between
165 lamb and adult telomere length was close to zero (0.036, 95% CI: -0.196–0.008, SI
166 Appendix, Table S4). The negligible residual correlation was expected given the lack of
167 permanent environment effect underlying repeatable differences in individual RTL across all
168 ages (SI Appendix, Table S3). Repeatable among-individual differences across ages in RTL
169 were therefore driven predominantly by genetic rather than environmental effects, and a
170 similar set of genes influenced RTL across ages.

171 Using multivariate mixed-effects models, we next tested the strength and direction of
172 correlations among RTL, body weight (both measured in August) and subsequent overwinter
173 survival at different hierarchical levels (among-individual, genetic, among-year, within-
174 individual). Although we were primarily interested in the RTL–survival association, the
175 inclusion of weight in our models allowed us to contrast the magnitude of the RTL–survival
176 association with a well-studied trait which is known to be linked to condition and fitness in
177 our study system (33). As previously documented, August body weight was strongly
178 predictive of improved winter survival prospects ((33-35); Figure 3). Our multivariate models
179 revealed that this association was present at all hierarchical levels (Figure 3; SI Appendix,
180 Tables S5 & S6). Our initial phenotypic model indicated that heavier individuals on average
181 tended to have longer lifespans (Figure 3A; SI Appendix, Tables S5). Developing this model
182 into a pedigree-based quantitative genetic model revealed that this among-individual effect
183 was driven by similar sized contributions from genes and environment (Figure 3B; SI
184 Appendix, Table S6). Overwinter survival probabilities were higher in years where the
185 average body weight was higher (among-year effect), and individuals with relatively low
186 weight compared to their average weight were less likely to survive (residual effect; Figure
187 3). In our phenotypic model, there was little evidence for an association between RTL and
188 body weight at any hierarchical level (Figure 3A; SI Appendix, Table S5). However, the
189 quantitative genetic model revealed a negative genetic correlation (Figure 3B; SI Appendix,
190 Table S6). This was consistent in separate models of lambs and adults, although here
191 credible intervals overlapped zero more widely (SI Appendix, Tables S7-S10).

192 We found a positive association between telomere length and overwinter survival probability
193 at the among-individual level (Figure 3A; SI Appendix, Table S5). This suggests that
194 individuals with longer telomeres on average across their lives tended to have longer
195 lifespans. The 95% credible intervals overlapped zero for both the among-year and the
196 residual covariance between telomere length and overwinter survival (Figure 3A; SI

197 Appendix, Table S5). The absence of an association at the within-individual (or residual)
198 level indicates that if an individual had a relatively short RTL measurement compared to their
199 average in a particular year, this was not associated with an increased risk of mortality.
200 Overall, these findings imply that average telomere length after weaning, rather than more
201 immediate changes relative to an individual's average TL, predicted overwinter survival.

202 The quantitative genetic model revealed a positive association between telomere length and
203 overwinter survival probability at the genetic level, though the credible intervals narrowly
204 overlapped zero (Figure 3B; SI Appendix, Table S6). The covariance between TL and
205 survival was very small (<0.001) at the permanent environment level (SI Appendix, Table
206 S6) and the correlation at this level had wide credible intervals which extensively crossed
207 zero (Figure 3B). Our results were broadly consistent for lambs and adults across separate
208 models, although the credible intervals of the among-individual and genetic effects did
209 narrowly cross zero within some age groups (Figures S2 & S3; SI Appendix, methods &
210 results, Tables S7-10). Overall, our results suggest that the among-individual positive
211 association between telomere length and survival identified in the phenotypic model was
212 driven largely by genetic effects (Figure 3B; SI Appendix, Table S6). In other words, genes
213 conferring longer average telomeres after weaning also tended to be associated with longer
214 lifespan.

215 **Discussion**

216 A growing number of studies across vertebrate species have shown that TL is heritable (21)
217 and that relatively short TL is predictive of increased mortality risk (13). We have dissected
218 the TL–mortality relationship across the natural lifespan of wild Soay sheep to identify the
219 processes that drive this association. We show that selective disappearance of individuals
220 with shorter average TL is the key process, rather than an association between the pattern
221 of TL attrition and mortality. If TL was a marker of biological ageing, or it reflected variation in
222 condition resulting from environmental stress, we would expect telomere shortening to
223 predict mortality, resulting in a positive residual correlation between TL and survival in our
224 models. We found no support for a correlation at the residual level, and only the among-
225 individual correlations between TL and survival were consistently different from zero. This
226 suggests that TL is not a useful marker of biological ageing in our system. Furthermore, it is
227 inconsistent with the hypothesis that TL reflects variation in physiological state or condition
228 resulting from recent or accumulating experience of environmental stress during adulthood,
229 which also influences mortality risk. However, our first TL measurement was taken at four
230 months of age – around the time of weaning for most lambs in our study population – and it

231 is possible that associations between average TL and survival are driven by some
232 combination of initial TL and the rate of telomere attrition prior to first measurement. An
233 individual's average TL could therefore reflect its ability to maintain homeostasis and resist
234 environmental stress during very early development, which in turn could predict its
235 subsequent lifespan. Nonetheless, our results demonstrate that variation in TL is heritable
236 and under directional selection, meaning it has the potential to evolve under natural
237 conditions.

238 Our moderate estimate of the heritability of TL (around 20%, Figure 2) represents the first
239 evidence for an appreciable genetic contribution to variation in TL in a wild mammal. Recent
240 studies in wild badgers and bats have used similar pedigree-based 'animal models' but
241 found both the repeatability and heritability of TL to be negligible (36, 37). Our estimate was
242 lower than in a recent longitudinal study of farmed dairy cattle (32–38%; (38)), which is
243 unsurprising given the expectation of reduced environmental variation in livestock compared
244 to wild systems. It is notable that both studies found negligible permanent environment
245 effects, implying that consistent differences in TL across life were largely the result of genetic
246 rather than early environmental effects in both systems ((38); Figure 2). More broadly, while
247 human studies have tended to find moderate to high heritability of TL, the growing literature
248 in birds presents a much more variable picture (21). Two laboratory studies of birds have
249 estimated the heritability of TL to be >100% (39, 40), whilst some studies in the wild suggest
250 little genetic influence on TL (heritability <5% (41, 42)). The reasons for the variation in
251 heritability estimates across studies remain an important area for future research. As well as
252 variation in the way that genes influence telomere dynamics, differences in the
253 environmental variation experienced by different populations or species, and the degree of
254 error associated with different telomere measurement methodologies are also likely to play a
255 role (21). Our study used a qPCR method to measure TL, which is often described as having
256 greater measurement error than the 'gold standard' terminal restriction fragment (TRF)
257 approach (43). However, we note that the technical repeatability of our qPCR assay is high
258 and compares favorably to those reported for TRF studies (see Methods). Furthermore,
259 most previous studies in humans and birds have focused on specific age classes (e.g.
260 elderly humans or pre-fledging birds). Our data spans the entire lifetimes of individuals, and
261 we were able to demonstrate a very high genetic correlation between TL measured in lambs
262 and adults. This provides clear evidence that the same or linked genes influence TL in early
263 and later life, an assumption that has rarely been tested in any study system (though see
264 (44)).

265 In a previous study, we identified an association between TL and mortality in a much smaller
266 sample of female Soay sheep (45). In that study, we reported a considerably lower
267 repeatability for TL (13%) than in the present study and were only able to detect a TL–
268 mortality relationship among younger females experiencing a high mortality winter. This
269 previous study was focused on a subset of females from four birth cohorts, and thus had
270 relatively few samples from later adulthood (45). Our present study is distinguished by the
271 much larger data set and more complete population coverage, which has allowed us to
272 assess the repeatability of TL and determine that it is the among-individual differences in TL
273 across life that are predictive of mortality. We note that, when splitting our multivariate
274 models by age groups, the among-individual covariance between TL and survival was
275 positive and of a similar magnitude in adults and lambs, though the credible intervals
276 overlapped zero (SI Appendix, Figs. S2 & S3; Table S4). This indicates that the association
277 was not simply due to effects on lamb survival. Although some longitudinal vertebrate
278 studies have found that telomere attrition predicted survival better than average or recent TL
279 (28, 29, 46), others have identified extremely high consistency in individual TL across
280 measurements (19, 20) and long-term associations between early-life TL and adult lifespan
281 (24, 47, 48). Most of these studies involved only one or two measurements of TL per
282 individual, and none used a multivariate mixed-effects modelling approach capable of fully
283 dissecting the contributions of genetic, individual, annual and residual sources to observed
284 TL–survival covariance.

285 Recent studies of elderly human cohorts have identified candidate SNP loci associated with
286 telomerase genes, which are involved in maintenance of telomeres and genomic integrity,
287 that not only predict average leukocyte TL but also subsequent morbidity and mortality (49-
288 51). However, human twin studies also suggest that relatively short TL, independent of
289 genetic factors, predicts mortality (52). In support of a causal role for telomerase genetics in
290 mortality and ageing, studies of laboratory mice without telomerase appear to show early
291 onset of ageing phenotypes, while mice with genetically enlarged telomeres are longer lived
292 (53, 54). Our findings provide the first support for a role for genetics in observed TL–mortality
293 relationships from a non-human system outside the laboratory. Further work is required to
294 determine whether specific genes, such as those involved in telomere maintenance, are
295 implicated in this relationship or whether the observed genetic correlations result from minor
296 effects of many different genes.

297 Our results imply that the observed relationship between TL and mortality is not causal. A
298 causal effect of short telomeres on survival would lead to the expectation that both
299 individuals with low average TL (among-individual covariance) and short TL at measurement

300 relative to their average (residual covariance) should be positively related to survival. But this
301 is not what we observed. Instead, it seems likely that the genes influencing some yet to be
302 determined aspect of an individual's overall frailty (or robustness) have a correlated
303 influence on TL. These genes could be influencing TL determined during early embryonic
304 development and/or the rate of telomere attrition during gestation and neonatal life, as our
305 first TL measurement was not taken until around four months of age. However, comparing
306 the magnitude of the TL–survival correlations with the correlations between body weight and
307 survival highlights that the association between TL and mortality is modest compared to a
308 measure that is more directly relatable to condition and health. It remains to be determined
309 whether our findings in Soay sheep will generalize to other species and systems, but our
310 work highlights the importance of using large data sets that span the entire lifetimes of many
311 individuals, in order to fully understand the drivers of associations between TL, health and
312 mortality risk. Overall, our results provide important insights into the genetics of lifespan in
313 the wild and highlight the importance of long-term, longitudinal studies across different
314 species for our understanding of TL as a biomarker of health and fitness.

315 **Materials and methods**

316 **Study system and sample collection**

317 The Soay sheep (*Ovis aries*) is a primitive breed of domestic sheep that has been living on
318 the remote St Kilda archipelago with minimal human management for the last few millennia
319 (57°49'N, 8°34'W; (33)). Since 1985, the sheep resident within the Village Bay area of the
320 main island in the archipelago, Hirta, have been the subject of an individual-based study
321 (33). Individuals are caught and tagged within a few days of birth in the spring. Ten
322 censuses are conducted on each of three annual field seasons, during spring (March –
323 April), summer (July – August) and autumn (October – November), meaning the timing of
324 individual's disappearance is known with a high degree of accuracy (33). The vast majority
325 of sheep mortality occurs in late winter (85% of adult deaths occur January – April), and daily
326 carcass searches during this period mean that death dates are known to the nearest month
327 for most individuals. Each August, 50–60% of the resident population are caught in
328 temporary corral traps and blood sampled. Blood is collected from each individual into 9 ml
329 lithium heparin Vacuettes and kept in a cool box or fridge from the point of sampling. The
330 blood is processed within 24 hours to separate the plasma and buffy coat fractions. The
331 Vacuette is then spun at 1,008 x g for 10 minutes, the plasma layer drawn off and replaced
332 by the same quantity of 0.9% NaCl solution, gently mixed and spun again at 1,008 x g for 10
333 minutes. The intermediate buffy coat layer, comprising mainly white blood cells, is then

334 drawn off into a 1.5 ml Eppendorf tube and stored at -20 °C until used to assay leukocyte
335 telomere length. All data collection was approved by the UK Home Office and carried out in
336 accordance with the relevant guidelines.

337 **Pedigree reconstruction**

338 Parentage was inferred by genetic methods, except for some maternal links inferred by
339 observation (55). Multi-generational pedigree reconstruction was performed in the R
340 package *Sequoia* (56), using 431 unlinked SNP markers. This likelihood-based approach
341 infers not only parent-offspring relationships, but also siblings and second-degree relatives.
342 In the resulting pedigree, a mother and/or father was assigned to 6082 individuals. After
343 pruning to only those individuals informative to the current analyses (using the R package
344 *MasterBayes* (57)) , the pedigree had a maximum depth of 13 generations and consisted of
345 2411 individuals, of which 2273 were non-founders and a total of 2050 maternities and 2172
346 paternities were assigned.

347 **Sample selection and randomization**

348 A total of 3891 August buffy coat samples from 1647 animals of known age collected
349 between 1998 and 2016 were selected for telomere length measurement. These samples
350 were selected from the total available buffy coat freezer archive (n=6775 from 3315 sheep)
351 based on the following exclusion criteria: samples collected before 1998 (n=1924); samples
352 from individuals of unknown age (n=97); samples collected between 2013 and 2016 from
353 lambs or yearlings that were caught only once (n=454); samples collected from individuals
354 that were only captured once but survived, and were therefore available for sampling in
355 subsequent years (n=409). These criteria were designed to maximise the longitudinality of
356 the dataset while avoiding biasing the dataset against short-lived individuals (i.e. individuals
357 only sampled once because they died). Sample years were then randomly allocated to one
358 of four batches, each comprising 4–5 years (to reduce the number of samples that needed to
359 be removed from the freezer at any one time). Samples were then fully randomised within
360 each batch, assigned a unique identifier from the start of batch 1 to the end of batch 4, and
361 processed from DNA extraction through to qPCR in this order.

362 **DNA extraction**

363 Genomic DNA was extracted from buffy coat on 96 well plates using the Macherey-Nagel
364 Nucleospin 96 Blood kit (Cat# 740665). The samples were extracted on a liquid handling
365 robot (Freedom Evo-2 150; Tecan) using a vacuum manifold. In order to facilitate passage of
366 the sample through the DNA binding plate, the following step was included prior to
367 automated extraction on the robot. 50 µl of buffy coat was mixed with 300 µl RBC lysis

368 solution (Qiagen; Cat# 158902) and incubated at room temperature for 5 minutes, before
369 centrifugation at 12200 rpm for 30 seconds. 250 μ l of the supernatant was discarded and the
370 cell pellet was re-suspended in the residual supernatant, before it was transferred to a 96
371 well MN lysis block (Macherey-Nagel Nucleospin 96 Blood kit; Cat# 740665) and sealed.
372 The liquid handling robot was loaded with the MN lysis block with the seal removed, the
373 silica DNA binding plate, 100% Ethanol, lysis buffer BQ1, wash buffers B5 and BW and
374 finally a master mix containing PBS, proteinase K and RNase A (binding plate, buffers and
375 proteinase K supplied with Macherey-Nagel Nucleospin 96 Blood kit, Cat# 740665; RNase
376 A, Qiagen Cat# 158924; PBS, Sigma Cat# D1408). The extraction protocol followed
377 manufacturer's guidelines for use with a vacuum manifold with the following amendments.
378 The robot added 96 μ l 1X PBS, 25 μ l Proteinase K and 4 μ l RNase A to each sample. Lysis
379 was performed on a shaker for 10 minutes. The vacuum steps for binding and the first two
380 washes were increased to 5 minutes and the final wash to 10 minutes with an additional 10-
381 minute vacuum at the end to dry the membrane. If any lysate/wash failed to pass through the
382 silica membrane after a vacuum step the plate was removed from the robot and centrifuged
383 for 3 minutes at 4000 rpm. For the few samples that failed to pass though after centrifugation
384 the wash/lysate was removed by hand with a pipette. After dry centrifugation (3 minutes,
385 4000 rpm), DNA was eluted in a total of 150 μ l elution buffer BE (Macherey-Nagel
386 Nucleospin 96 Blood kit; Cat# 740665) which was warmed to 60 °C prior to adding it onto
387 the silica membrane. Elution was performed in two steps: first 100 μ l buffer BE followed by
388 centrifugation (3 minutes, 4000 rpm), then 50 μ l buffer BE followed by a final centrifugation
389 (3 minutes, 4000 rpm).

390 *DNA extraction quality control*

391 Following DNA extraction, a strict quality control protocol was implemented to measure DNA
392 concentration, integrity and purity (SI Appendix, Figure S4). Of the 3891 samples selected
393 for analysis, 42 samples were missing or accidentally omitted through human error. The
394 available samples were measured on a Nanodrop ND-1000 9 spectrophotometer (Thermo
395 Scientific, Wilmington DE, USA). Samples yielding < 20 ng/ μ l were rejected, and re-
396 extraction attempted for 491 of these samples. Overall, 87 samples failed to yield sufficient
397 DNA and were excluded from the analyses. Samples yielding \geq 20 ng/ μ l were checked for
398 DNA purity. The acceptable range for absorption for 260/280 nm ratio (a measure of DNA vs
399 protein and RNA contamination) was 1.7–2.0. Samples falling outside of this range were
400 excluded from subsequent analyses (n=66, representing a failure rate of 1.76%). Two thirds
401 of samples were assayed for 260/230 nm ratio (a measure of salt and other impurities), and
402 samples with a ratio <1.8 were excluded from subsequent analyses (n=41, representing a
403 failure rate of 1.73%). Samples of sufficient yield and purity were standardized to 10 ng/ μ l

404 and DNA integrity was assessed by running 200 ng of DNA on a 0.5% agarose gel. Samples
405 were scored for integrity on a scale of 1 to 5 by visual examination of their DNA crowns, and
406 samples scoring 3 to 5 were excluded from further analyses (see (58)). Integrity was initially
407 assayed for all of the first 1667 samples run. Only 16 of these samples were given a DNA
408 integrity score of 2, and 7 failed (failure rate of 0.42%). We randomly tested a quarter of all
409 subsequent samples for DNA integrity, with only 4 failing (representing a failure rate of
410 0.76%). In total, 11 samples were excluded due to poor DNA integrity. Overall, DNA was
411 successfully extracted and passed quality control requirements from 3644 buffy coat
412 samples (SI Appendix, Figure S4).

413 **Telomere length measurement**

414 *Quantitative polymerase chain reaction*

415 Relative leukocyte telomere length (RTL) was measured using real-time quantitative PCR
416 (qPCR; (59)), using protocols we have previously developed and validated using blood
417 samples from sheep and cattle (45, 58). The qPCR method estimates the total amount of
418 telomeric sequence present in a sample relative to the amount of a non-variable copy
419 number reference gene. In this study we used the beta-2-microglobulin (B2M) as our
420 reference gene (45, 58), with primers supplied by Primer Design (Cat# HK-SY-Sh-900,
421 Southampton, UK). For telomeric amplification tel1b (5'-CGG TTT GTT TGG GTT TGG GTT
422 TGG GTT TGG GTT TGG GTT-3') and tel 2b (5'-GGC TTG CCT TAC CCT TAC CCT TAC
423 CCT TAC CCT TAC CCT-3') primers were used. Telomere primers were manufactured,
424 HPLC purified and supplied by Integrated DNA Technologies (IDT, Glasgow, UK). Telomere
425 and reference gene reactions were run in separate wells of the same qPCR plate at a
426 concentration of 300 nM and 900 nM, respectively. Samples were diluted to 0.5 ng/μl with
427 buffer BE (Macherey-Nagel Nucleospin 96 Blood kit; Cat# 740665) immediately prior to
428 qPCR analysis. Each reaction was prepared using 5 μl of LightCycler 480SYBR Green I
429 Master Mix (Cat # 04887352001, Roche, West Sussex, UK) and 1 ng of sample DNA in a
430 total reaction volume of 10 μl. We used 384 well plates which were loaded with sample DNA
431 and master mix using an automated liquid handling robot (Freedom Evo-2 150; Tecan).

432 Each plate included 8 calibrator samples (1 ng/μl) to account for plate to plate variation and
433 two non-template controls (NTC) consisting of nuclease free water. The calibrator sample
434 was extracted from a large quantity of buffy coat prepared from blood supplied from a single
435 domestic sheep (Cat# SHP-BUFCT-LIHP, Sera Laboratories International LTD, West
436 Sussex, UK). We carried out a large number of extractions from this sample, applied the
437 same quality control as above and then pooled the extracts and aliquoted them for
438 subsequent use. A five-point standard curve, consisting of a four-fold serial dilution of the

439 calibrator sample (at concentrations: 20 ng/μl, 5ng/μl, 1.25 ng/μl, 0.3125 ng/μl and 0.078125
440 ng/μl) was included on the plate to provide a visual check that the samples amplified at the
441 correct cycle. Samples, calibrators, standard curve and NTCs were all run in triplicate. All
442 qPCRs were performed using a Roche LC480 instrument using the following reaction
443 protocol; 10 min at 95 °C (enzyme activation), followed by 50 cycles of 15 s at 95 °C
444 (denaturation) and 30 s at 58 °C (primer annealing), then 30 s at 72 °C (signal acquisition).
445 Melting curve protocol was 1 min at 95 °C, followed by 30 s at 58 °C, then 0.11 °C/s
446 increase to 95 °C followed by 10 s at 40 °C (cool down).

447 *Calculation of Relative Telomere Length*

448 We used the LinRegPCR software package (version 2016.0; (60)) to correct our
449 amplification curves for baseline fluorescence, and to calculate well-specific reaction
450 efficiencies and Cq values. A constant fluorescent threshold was set within the window of
451 linearity for each amplicon group, calculated using the average Cq across all plates. The
452 threshold values used were 0.2 and 0.25, and the average PCR efficiency across all plates
453 was 1.876 and 1.881 for the B2M and telomere amplicon groups, respectively. Samples
454 were excluded from further analysis if the coefficient of variation (CV) across triplicate Cq
455 values and triplicate PCR efficiency values for either amplicon was > 5% (n=1; note that for
456 > 95% of our samples the triplicate CV was < 2%), or if at least one of their triplicate
457 reactions had an efficiency that was 5% higher or lower than the mean efficiency across all
458 wells on that plate for the respective amplicon (n=2; > 95% of samples were within 2% of
459 mean plate efficiencies; SI Appendix, Figure S4).

460 RLTL for each sample was calculated, following Pfaffl (61), using average reaction
461 efficiencies for each plate and Cq for each sample determined by LinRegPCR as follows:

$$RTL = \frac{E_{TEL}^{(Cq_{TEL}[Calibrator]-Cq_{TEL}[Sample])}}{E_{B2M}^{(Cq_{B2M}[Calibrator]-Cq_{B2M}[Sample])}}$$

462 Where E_{TEL} and E_{B2M} are the mean reaction efficiencies for the respective amplicon group
463 across all samples on a given plate; $Cq_{TEL}[Calibrator]$ and $Cq_{B2M}[Calibrator]$ are the
464 average Cqs for the relevant amplicon across all calibrator samples on the plate; and
465 $Cq_{TEL}[Sample]$ and $Cq_{B2M}[Sample]$ are the average of the triplicate Cqs for the sample for
466 each amplicon.

467 *Repeatability of telomere length*

468 To assess the repeatability of our qPCR assay, the first qPCR plate (n=48 samples) was run
469 8 times over 4 consecutive days: four times over two days with samples in the same position

470 and then four times over two days with samples in an alternative row within the qPCR plate.
471 We calculated the overall repeatability of RTL as the proportion of variance explained by
472 sample identity over the total variance, in a linear mixed-effects model including only sample
473 identity as a random intercept term (using restricted maximum likelihood estimation in
474 *glmmTMB* v.0.2.3 (62)). The overall repeatability of RTL was 0.866 (95% confidence
475 intervals 0.807–0.908). The repeatability of RTL measured in the same location across the
476 first four qPCR plates was 0.948 (95%CI 0.922–0.965), illustrating how repeatability can be
477 inflated if samples are consistently run in the same location. In a second model which
478 included sample identity, qPCR plate and qPCR row as random intercept terms, the
479 proportion of variance explained by qPCR plate and row was 0.005 and 0.063, respectively.
480 The repeatability of RTL in this model was 0.824 (95%CI 0.748–0.880), or 0.884 (95%CI
481 0.829–0.923) if the plate and row terms were excluded from the total variance (since they
482 represent measurement error). These repeatability estimates compare favourably with other
483 studies of telomere length measured by both qPCR (inter-assay repeatability: 0.85 (48); 0.82
484 (24)) and terminal restriction fragment (TRF repeatability: 0.86 (63)).

485 **Data analysis**

486 All analyses were conducted in the program R version 3.6.1 (64) using the package
487 *MCMCglmm* v.2.29 (65) unless otherwise specified.

488 *Relationship between telomere length, age and sex*

489 Relative telomere length was approximately normally distributed in lambs, adults and overall
490 (Figure S5). We ran a series of linear mixed-effects models to determine the function that
491 best described variation in telomere length with age. We included a two-level factor for age
492 class (lamb: ~4 months of age; adult: ≥ 1 year). To account for age-related variation within
493 the adult age class, we included age in years as a fixed covariate. We tested linear,
494 quadratic and cubic age terms, as well as threshold age functions with a range of
495 breakpoints (2–11 years). We also tested a four-level factor for age class (lamb: ~4 months
496 of age; yearling: ~16 months; adult: 2-6 years; geriatric: >6 years. All models included
497 individual identity and sample year as random intercept terms to account for non-
498 independence among observations. The qPCR plate and row for each sample were also
499 included as crossed random intercept terms to account for variation associated with
500 measurement error. The models were run using maximum likelihood estimation in *glmmTMB*
501 v.0.2.3 (62) and AIC model selection was used to determine the best age function. We
502 selected the model with the fewest parameters within 2 Δ AIC of the model with the lowest
503 AIC value. Once we determined the best fitting age function, we tested for a difference in
504 average telomere length between the sexes by including a two-level factor for sex in our

505 model. To test whether the ageing patterns differed between the sexes, we also tested for
506 interactions between sex and the selected age terms. The significance of additive and
507 interactive effects of sex was assessed using likelihood ratio tests (see SI Appendix). The
508 repeatability of telomere length over the lifespan was estimated as the variance explained by
509 the random effect of individual over the total phenotypic variance (the sum of the random
510 effects variance components plus the residual variance). There were 3641 observations of
511 telomere length from 1586 sheep available for this analysis. Of these individuals, 836 had
512 one RTL measurement available, 271 had two, 281 had three or four, and 198 had five or
513 more measurements.

514 *Heritability of telomere length*

515 A quantitative genetic animal model was used to estimate the additive genetic variance for
516 telomere length in Soay sheep. The model contained a two-level fixed factor for age class
517 (lamb: ~4 months; adult: ≥ 1 year) and a linear covariate for age in years. We also included
518 sex as a two-level fixed factor to account for differences in average telomere length between
519 the sexes. The additive genetic effect was estimated using information on individual
520 relatedness from the population pedigree. We included maternal identity in addition to the
521 additive genetic component to capture similarity among maternal siblings that is not
522 explained by the additive genetic effect (known as the maternal effect). We also included
523 individual identity as a random effect to capture consistent differences in measures from the
524 same individual that are not attributed to genetic effects, influenced for example by where
525 the individual lives or aspects of their early life environment (the permanent environment
526 effect). Year of sample collection, qPCR plate and qPCR row were included as random
527 intercept terms. The heritability of telomere length was calculated as the variance explained
528 by the additive genetic effect over the total phenotypic variance. This model was run for
529 1.1×10^5 iterations, with 1×10^4 burn-in and thinning interval of 50, resulting in 2000 stored
530 samples of the MCMC chain with minimal autocorrelation (< 0.2). Parameter estimates are
531 presented as the posterior mode with 95% highest posterior density (HPD) intervals.
532 Parameter expanded priors were used for all variance components, and inverse-Wishart
533 priors for the residual variance. There were 3632 observations of telomere length from 1582
534 sheep available for this analysis ($n=9$ observations from 4 sheep were excluded because
535 maternal identity was unknown).

536 We also ran a bivariate model to estimate the genetic correlation between telomere length in
537 lambs (~4 months) and adults (≥ 1 year) which treated TL in lambs and adults as separate
538 response variables. We included sex as a two-level fixed factor for both lambs and adults,
539 with age in years included as a fixed covariate for adults only. qPCR plate and row were

540 included as random intercept terms across both models. Maternal identity was included as a
 541 random effect in the lamb model and individual identity in the adult model to estimate
 542 maternal and permanent environment effects, respectively. We estimated the unstructured
 543 variance-covariance matrix for the genetic, year and residual effects, which enabled us to
 544 estimate the correlations across age classes at these different hierarchical levels. There is a
 545 possibility that the lamb residual variance could covary with the adult permanent
 546 environment effect, but we did not attempt to model this covariance because we detected no
 547 permanent environment effect in adults (see Results). The model was run 2.1×10^5 iterations,
 548 with 1×10^4 burn-in and thinning interval of 200, resulting in 1000 stored samples of the
 549 MCMC chain (autocorrelation < 0.2). The genetic correlation was taken from the posterior
 550 correlation of the stored samples with 95% HPD intervals. The bivariate model was run using
 551 3632 measurements from 1582 individuals. Both lamb and adult telomere length
 552 measurements were available for 424 individuals.

553 *Associations between telomere length, August weight and overwinter survival*

554 *Phenotypic model*

555 We used a multivariate mixed-modelling approach to examine the association between
 556 telomere length, weight and overwinter survival at different hierarchical levels. Relative
 557 telomere length, body weight in kg (both measured August year t), and annual survival (to 1
 558 May year $t+1$) were our response variables, modelled as Gaussian, Gaussian and threshold
 559 distributions, respectively (corresponding to identity and probit link functions). Unstructured
 560 variance–covariance matrices were estimated for some of the random effects, allowing us to
 561 estimate the covariance among the three traits at different hierarchical levels. For each of
 562 these random effects, we obtained a posterior distribution for the variance and covariance
 563 between telomere length (“RTL”), weight (“Wt”) and overwinter survival (“Surv”):

$$\begin{bmatrix} \sigma_{RTL}^2 & \sigma_{RTL,Wt} & \sigma_{RTL,Surv} \\ \sigma_{RTL,Wt} & \sigma_{Wt}^2 & \sigma_{Wt,Surv} \\ \sigma_{RTL,Surv} & \sigma_{Wt,Surv} & \sigma_{Surv}^2 \end{bmatrix}$$

564 We estimated the covariance at the among-individual, among-year and residual levels. The
 565 among-individual level captures consistent differences between individuals (e.g. if individuals
 566 that are consistently heavier tended to have longer lifespans). The covariance at the year
 567 level captures associations between average trait values among years (e.g. whether the
 568 survival rate is higher in years when the average body weight is higher). The residual level
 569 reflects covariance that is not captured by the among-individual and among-year levels (i.e.
 570 if an individual has a relatively low body weight compared to its average has lower chances
 571 of surviving).

572 The model was run for 5.3×10^5 iterations, with 3×10^4 burn-in and thinning interval of 500,
573 resulting in 1000 stored samples of the MCMC chain (autocorrelation < 0.1). The residual
574 variance was fixed at 1 for survival (as is the standard recommendation for threshold
575 models), and the latent variables were constrained to be between ± 7 to avoid numerical
576 difficulties as the probabilities approached 0 and 1. In addition to the random effects of
577 identity and year, the telomere length model included random intercept terms for qPCR plate
578 and row, and the weight and survival models included random intercept terms for maternal
579 identity and birth year. We fitted age and sex in slightly different ways across the three
580 models, to reflect our understanding of differences in age-related variation in these three
581 traits. In the telomere model, we included age class (lamb vs adult) as a two-level fixed
582 factor. No other fixed effects were included in the telomere model, since we were interested
583 in how telomere length covaried with the other traits across the lifespan. In the weight model,
584 we included age class (lamb vs adult) plus linear, quadratic and cubic age terms (capturing
585 variation in weight with age during adulthood) and their interactions with sex. In a previous
586 study, we detected a decline in weight across the year prior to death in this system using
587 univariate models of weight (35). We did not include a term of final year alive here, as this
588 effect is captured in the residual covariance between weight and survival in our multivariate
589 model. In the survival model, we included age class (lamb vs adult) plus sex and its
590 interaction with age class, as well as linear and quadratic age terms (capturing variation in
591 survival with age during adulthood). This model was run using 3569 measurements from
592 1574 individuals for which telomere length, August weight and overwinter survival
593 measurements were available. Telomere length and body weight were z-transformed prior to
594 inclusion in the model (mean=0, standard deviation=1). To check the consistency of our
595 results, we additionally ran separate models for lambs aged 4 months and adults aged ≥ 1
596 year (see Supplementary methods and results).

597 *Genetic model*

598 We extended our multivariate analysis to examine the association between telomere length,
599 August body weight and overwinter survival probability at the genetic level (the G matrix). As
600 with the univariate heritability analysis, we used information on individual relatedness from
601 the population pedigree to estimate the additive genetic variances and covariances. In
602 addition to the additive genetic (co)variance, we estimated the covariance at the residual,
603 among-year and among-individual levels (the permanent environment covariance: the
604 covariance among traits at the individual level that is not explained by genetic effects). The
605 model was run for 5.3×10^5 iterations, with 3×10^4 burn-in and thinning interval of 500,
606 resulting in 1000 stored samples of the MCMC chain (autocorrelation < 0.1). Parameter
607 expanded priors were used for all variance components, and inverse-Wishart priors for the

608 residual variances. The residual variance was fixed at 1 for survival (as is the standard
609 recommendation for threshold models), and the latent variables were constrained to be
610 between ± 7 to avoid numerical difficulties as the probabilities approached 0 and 1. Other
611 fixed and random effects were as stated above for the phenotypic models. We additionally
612 ran separate animal models for lambs aged 4 months and adults aged ≥ 1 year to check the
613 consistency of our results (see Supplementary methods and results).

614 **Acknowledgements**

615 We thank the many project members and volunteers who have contributed to the long-term
616 study of Soay sheep on St Kilda over many years; the National Trust for Scotland for
617 permission to work on St Kilda; and QinetiQ and Kilda Cruises for logistical support in the
618 field. We thank Camillo Bérénos and Jisca Huisman for reconstruction of the population
619 pedigree; Lorraine Kerr and Eliane Salvo-Chirnside for laboratory support; Jarrod Hadfield
620 and Craig Walling for statistical advice; and Jon Slate for constructive feedback on the
621 manuscript. The long-term field study has been largely funded by the UK Natural
622 Environment Research Council; SNP genotyping was mostly funded by the European
623 Research Council; and the telomere project was funded by the UK Biotechnology and
624 Biological Sciences Research Council (grant no. BB/L020769/1).

625 **References**

- 626 1. E. H. Blackburn, Structure and function of telomeres. *Nature* **350**, 569-573 (1991).
- 627 2. G. Aubert, P. M. Lansdorp, Telomeres and Aging. *Physiol. Rev.* **88**, 557-579 (2008).
- 628 3. J. W. Shay, W. E. Wright, Telomeres and telomerase: three decades of progress. *Nature*
629 *Reviews Genetics* **20**, 299-309 (2019).
- 630 4. M. Z. Levy, R. C. Allsopp, A. B. Futcher, C. W. Greider, C. B. Harley, Telomere end-replication
631 problem and cell aging. *Journal of Molecular Biology* **225**, 951-960 (1992).
- 632 5. T. von Zglinicki, Oxidative stress shortens telomeres. *Trends in Biochemical Sciences* **27**, 339-
633 344 (2002).
- 634 6. C. López-Otín, M. A. Blasco, L. Partridge, M. Serrano, G. Kroemer, The Hallmarks of Aging.
635 *Cell* **153**, 1194-1217 (2013).
- 636 7. A. Benetos *et al.*, Short telomeres are associated with increased carotid atherosclerosis in
637 hypertensive subjects. *Hypertension* **43**, 182-185 (2004).
- 638 8. M. A. Blasco, Telomeres and human disease: ageing, cancer and beyond. *Nature Reviews*
639 *Genetics* **6**, 611-622 (2005).
- 640 9. J. Zhao, K. Miao, H. Wang, H. Ding, D. W. Wang, Association between telomere length and
641 type 2 diabetes mellitus: a meta-analysis. *PLoS One* **8**, e79993 (2013).
- 642 10. R. M. Cawthon, K. R. Smith, E. O'Brien, A. Sivatchenko, R. A. Kerber, Association between
643 telomere length in blood and mortality in people aged 60 years or older. *The Lancet* **361**,
644 393-395 (2003).
- 645 11. M. Kimura *et al.*, Telomere length and mortality: a study of leukocytes in elderly Danish
646 twins. *American journal of epidemiology* **167**, 799-806 (2008).

- 647 12. J. J. Boonekamp, M. J. Simons, L. Hemerik, S. Verhulst, Telomere length behaves as
648 biomarker of somatic redundancy rather than biological age. *Aging cell* **12**, 330-332 (2013).
- 649 13. R. V. Wilbourn *et al.*, The relationship between telomere length and mortality risk in non-
650 model vertebrate systems: a meta-analysis. *Philosophical Transactions of the Royal Society*
651 *B: Biological Sciences* **373**, 20160447 (2018).
- 652 14. M. J. Simons, Questioning causal involvement of telomeres in aging. *Ageing research reviews*
653 **24**, 191-196 (2015).
- 654 15. A. Aviv, Telomeres and human somatic fitness. *The Journals of Gerontology Series A:*
655 *Biological Sciences and Medical Sciences* **61**, 871-873 (2006).
- 656 16. M. Bateson, Cumulative stress in research animals: Telomere attrition as a biomarker in a
657 welfare context? *BioEssays* **38**, 201-212 (2016).
- 658 17. P. Monaghan, S. E. Ozanne, Somatic growth and telomere dynamics in vertebrates:
659 relationships, mechanisms and consequences. *Philosophical Transactions of the Royal*
660 *Society B: Biological Sciences* **373**, 20160446 (2018).
- 661 18. A. J. Young, The role of telomeres in the mechanisms and evolution of life-history trade-offs
662 and ageing. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**,
663 20160452 (2018).
- 664 19. A. Benetos *et al.*, Tracking and fixed ranking of leukocyte telomere length across the adult
665 life course. *Aging Cell* **12**, 615-621 (2013).
- 666 20. C. Bichet *et al.*, Telomere length is repeatable, shortens with age and reproductive success,
667 and predicts remaining lifespan in a long-lived seabird. *Molecular Ecology* **29**, 429-441
668 (2020).
- 669 21. H. L. Dugdale, D. S. Richardson, Heritability of telomere variation: it is all about the
670 environment! *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**,
671 20160450 (2018).
- 672 22. L. Broer *et al.*, Meta-analysis of telomere length in 19 713 subjects reveals high heritability,
673 stronger maternal inheritance and a paternal age effect. *European journal of human genetics*
674 **21**, 1163-1168 (2013).
- 675 23. L. G. Spurgin *et al.*, Spatio-temporal variation in lifelong telomere dynamics in a long-term
676 ecological study. *Journal of Animal Ecology* **87**, 187-198 (2018).
- 677 24. S. H. van Lieshout *et al.*, Individual variation in early-life telomere length and survival in a
678 wild mammal. *Molecular ecology* **28**, 4152-4165 (2019).
- 679 25. B. S. Oliveira *et al.*, Systematic review of the association between chronic social stress and
680 telomere length: A life course perspective. *Ageing research reviews* **26**, 37-52 (2016).
- 681 26. G. V. Pepper, M. Bateson, D. Nettle, Telomeres as integrative markers of exposure to stress
682 and adversity: a systematic review and meta-analysis. *Royal Society open science* **5**, 180744
683 (2018).
- 684 27. M. Chatelain, S. M. Drobniak, M. Szulkin, The association between stressors and telomeres in
685 non-human vertebrates: a meta-analysis. *Ecology letters* **23**, 381-398 (2020).
- 686 28. J. J. Boonekamp, G. A. Mulder, H. M. Salomons, C. Dijkstra, S. Verhulst, Nestling telomere
687 shortening, but not telomere length, reflects developmental stress and predicts survival in
688 wild birds. *Proceedings of the Royal Society B: Biological Sciences* **281** (2014).
- 689 29. E. M. Wood, A. J. Young, Telomere attrition predicts reduced survival in a wild social bird,
690 but short telomeres do not. *Molecular ecology* **28**, 3669-3680 (2019).
- 691 30. E.L. Barrett, T. A. Burke, M. Hammers, J. Komdeur, D. S. Richardson, Telomere length and
692 dynamics predict mortality in a wild longitudinal study. *Molecular Ecology* **22**, 249-259
693 (2013).
- 694 31. J. W. Vaupel, K. G. Manton, E. Stallard, The Impact of Heterogeneity in Individual Frailty on
695 the Dynamics of Mortality. *Demography* **16**, 439-454 (1979).
- 696 32. D. H. Nussey, T. Coulson, M. Festa-Bianchet, J. M. Gaillard, Measuring senescence in wild
697 animal populations: towards a longitudinal approach. *Functional Ecology* **22**, 393-406 (2008).

- 698 33. T. H. Clutton-Brock, J. M. Pemberton, *Soay sheep: dynamics and selection in an island*
699 *population* (Cambridge University Press, 2004).
- 700 34. J. Milner, S. Albon, A. Illius, J. Pemberton, T. Clutton-Brock, Repeated selection of
701 morphometric traits in the Soay sheep on St Kilda. *Journal of Animal Ecology* **68**, 472-488
702 (1999).
- 703 35. D. H. Nussey *et al.*, Patterns of body mass senescence and selective disappearance differ
704 among three species of free-living ungulates. *Ecology* **92**, 1936-1947 (2011).
- 705 36. S. van Lieshout *et al.*, Estimation of environmental and genetic contributions to telomere
706 length variation in a wild mammal. (2020).
- 707 37. N. M. Foley *et al.*, Drivers of longitudinal telomere dynamics in a long-lived bat species,
708 *Myotis myotis*. *Molecular Ecology* (2020).
- 709 38. L. A. Seeker *et al.*, Bovine telomere dynamics and the association between telomere length
710 and productive lifespan. *Scientific reports* **8**, 1-12 (2018).
- 711 39. M. Asghar, S. Bensch, M. Tarka, B. Hansson, D. Hasselquist, Maternal and genetic factors
712 determine early life telomere length. *Proceedings of the Royal Society B: Biological Sciences*
713 **282** (2015).
- 714 40. E. Atema *et al.*, Heritability of telomere length in the Zebra Finch. *Journal of Ornithology* **156**,
715 1113-1123 (2015).
- 716 41. P. J. J. Becker *et al.*, *Mother-offspring and nest-mate resemblance but no heritability in early-*
717 *life telomere length in white-throated dippers* (2015), vol. 282.
- 718 42. A. M. Sparks *et al.*, Telomere heritability and parental age at conception effects in a wild
719 avian population. (2020).
- 720 43. D. H. Nussey *et al.*, Measuring telomere length and telomere dynamics in evolutionary
721 biology and ecology. *Methods in Ecology and Evolution* **5**, 299-310 (2014).
- 722 44. L. A. Seeker *et al.*, Longitudinal changes in telomere length and associated genetic
723 parameters in dairy cattle analysed using random regression models. *PLoS One* **13**, e0192864
724 (2018).
- 725 45. J. Fairlie *et al.*, Lifelong leukocyte telomere dynamics and survival in a free-living mammal.
726 *Aging cell* **15**, 140-148 (2016).
- 727 46. L. A. Seeker *et al.*, Life-long telomere attrition predicts health and lifespan in a large
728 mammal. *BioRxiv* (2020).
- 729 47. B. J. Heidinger *et al.*, Telomere length in early life predicts lifespan. *Proceedings of the*
730 *National Academy of Sciences of the United States of America* **109**, 1743-1748 (2012).
- 731 48. J. R. Eastwood *et al.*, Early-life telomere length predicts lifespan and lifetime reproductive
732 success in a wild bird. *Molecular ecology* **28**, 1127-1137 (2019).
- 733 49. G. Atzmon *et al.*, Genetic variation in human telomerase is associated with telomere length
734 in Ashkenazi centenarians. *Proceedings of the National Academy of Sciences* **107**, 1710-1717
735 (2010).
- 736 50. M. Soerensen *et al.*, Genetic variation in TERT and TERC and human leukocyte telomere
737 length and longevity: a cross-sectional and longitudinal analysis. *Aging cell* **11**, 223-227
738 (2012).
- 739 51. D. Scarabino, M. Peconi, F. Pelliccia, R. M. Corbo, Analysis of the Association Between TERC
740 and TERT Genetic Variation and Leukocyte Telomere Length and Human Lifespan—A Follow-
741 Up Study. *Genes* **10**, 82 (2019).
- 742 52. S. L. Bakaysa *et al.*, Telomere length predicts survival independent of genetic influences.
743 *Aging cell* **6**, 769-774 (2007).
- 744 53. M. Armanios *et al.*, Short telomeres are sufficient to cause the degenerative defects
745 associated with aging. *The American Journal of Human Genetics* **85**, 823-832 (2009).
- 746 54. M. A. Muñoz-Lorente, A. C. Cano-Martin, M. A. Blasco, Mice with hyper-long telomeres show
747 less metabolic aging and longer lifespans. *Nature communications* **10**, 1-14 (2019).

- 748 55. C. Bérénos, P. A. Ellis, J. G. Pilkington, J. M. Pemberton, Estimating quantitative genetic
749 parameters in wild populations: a comparison of pedigree and genomic approaches.
750 *Molecular ecology* **23**, 3434-3451 (2014).
- 751 56. J. Huisman, Pedigree reconstruction from SNP data: parentage assignment, sibship clustering
752 and beyond. *Molecular ecology resources* **17**, 1009-1024 (2017).
- 753 57. J. Hadfield, D. Richardson, T. Burke, Towards unbiased parentage assignment: combining
754 genetic, behavioural and spatial data in a Bayesian framework. *Molecular Ecology* **15**, 3715-
755 3730 (2006).
- 756 58. L. A. Seeker *et al.*, Method specific calibration corrects for DNA extraction method effects on
757 relative telomere length measurements by quantitative PCR. *PLoS One* **11**, e0164046 (2016).
- 758 59. R. M. Cawthon, Telomere measurement by quantitative PCR. *Nucleic Acids Research* **30**, e47
759 (2002).
- 760 60. J. M. Ruijter *et al.*, Amplification efficiency: linking baseline and bias in the analysis of
761 quantitative PCR data. *Nucleic Acids Research* **37**, e45 (2009).
- 762 61. M. W. Pfaffl, A new mathematical model for relative quantification in real-time RT-PCR.
763 *Nucleic Acids Research* **29**, e45 (2001).
- 764 62. M. E. Brooks *et al.*, glmmTMB balances speed and flexibility among packages for zero-
765 inflated generalized linear mixed modeling. *The R journal* **9**, 378-400 (2017).
- 766 63. O. Vedder, S. Verhulst, E. Zuidersma, S. Bouwhuis, Embryonic growth rate affects telomere
767 attrition: an experiment in a wild bird. *Journal of Experimental Biology* **221**, jeb181586
768 (2018).
- 769 64. R. C. Team (2019) R: A language and environment for statistical computing. (R Foundation
770 for Statistical Computing, Vienna, Austria).
- 771 65. J. D. Hadfield, MCMC methods for multi-response generalized linear mixed models: the
772 MCMCglmm R package. *Journal of Statistical Software* **33**, 1-22 (2010).

773

774 **Figure legends**

775

776 **Figure 1.** Relative telomere length varied with age over the lifespan in Soay sheep (n=3641
777 observations of 1586 individuals). The points show raw data medians and standard errors for
778 each age, with females in black and males in grey (for clarity, n=3 observations of females
779 aged > 13 and n=6 observations of males aged > 8 are grouped with ages 13 and 8,
780 respectively). The black lines show predictions from the best model (SI Appendix, Table S2),
781 grey shading represents 95% credible intervals around those predictions. The rug plot on the
782 inside of the x-axis shows the distribution of observations across the age range. Note that
783 the ageing pattern in the raw data and best-fitting age functions from mixed-effects models
784 are not expected to align perfectly when selective disappearance effects are present.

785

786 **Figure 2.** The proportion of variance in relative telomere length in Soay sheep explained by
787 different variance components. Estimates are based on the mode of the posterior distribution
788 from Bayesian quantitative genetic animal models. Estimates for age class 'all' came from a
789 univariate model incorporating all observations (n=3632 of 1582 individuals) that included
790 sex, age class and age in years as fixed effects (SI Appendix, Table S3). The separate
791 estimates for adults (age ≥ 1 year) and lambs (4 months old) came from a bivariate model
792 that accounted for the covariance between lamb and adult telomere length at the genetic,
793 year and residual levels (SI Appendix, Table S4).

794

795 **Figure 3.** The correlation between relative telomere length (RTL), August body weight and
796 overwinter survival probability at different hierarchical levels in Soay sheep (n=3569
797 observations of 1574 individuals). Correlations were estimated as the mode of the posterior
798 distribution with 95% higher probability density intervals from multivariate Bayesian mixed-
799 effects models. Panel A shows estimates for the among-individual (teal), among-year (black)
800 and residual correlations (grey) from a phenotypic model (SI Appendix, Table S5). Panel B
801 shows estimates for the genetic (teal circles), permanent environment ("PermEnv", teal
802 triangles), among-year (black) and residual correlations (grey) from a quantitative genetic
803 animal model (SI Appendix, Table S6).





