



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Prevalence of child maltreatment in India and its association with gender, urbanisation and policy

Citation for published version:

Fernandes, G, Fernandes, M, Vaidya, N, De Souza, P, Plotnikova, E, Geddes, R, Holla, B, Sharma, E, Benegal, V & Choudhry, V 2021, 'Prevalence of child maltreatment in India and its association with gender, urbanisation and policy: a rapid review and meta-analysis protocol', *BMJ Open*, vol. 11, no. 8, pp. e044983. <https://doi.org/10.1136/bmjopen-2020-044983>

Digital Object Identifier (DOI):

[10.1136/bmjopen-2020-044983](https://doi.org/10.1136/bmjopen-2020-044983)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

BMJ Open

Publisher Rights Statement:

This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made.

General rights


Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



BMJ Open Prevalence of child maltreatment in India and its association with gender, urbanisation and policy: a rapid review and meta-analysis protocol

Gwen Fernandes ¹, Megan Fernandes,² Nilakshi Vaidya,³ Philip De Souza,⁴ Evgeniya Plotnikova,⁵ Rosemary Geddes,⁵ Bharath Holla,³ Eesha Sharma,⁶ Vivek Benegal,³ Vikas Choudhry⁷

To cite: Fernandes G, Fernandes M, Vaidya N, *et al*. Prevalence of child maltreatment in India and its association with gender, urbanisation and policy: a rapid review and meta-analysis protocol. *BMJ Open* 2021;**11**:e044983. doi:10.1136/bmjopen-2020-044983

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online. (<http://dx.doi.org/10.1136/bmjopen-2020-044983>).

Received 28 September 2020
Accepted 26 July 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Gwen Fernandes;
gwen.fernandes@bristol.ac.uk

ABSTRACT

Introduction India is home to 20% of the world's children and yet, little is known on the magnitude and trends of child maltreatment nationwide. The aims of this review are to provide a prevalence of child maltreatment in India with considerations for any effects of gender; urbanisation (eg, urban vs rural) and legislation (Protection of Children from Sexual Offences (POCSO) Act 2012).

Methods and analysis A rapid review will be undertaken of all quantitative peer-reviewed studies on child maltreatment in India between 2005 and 2020. Four electronic databases will be systematically searched: PubMed, EMBASE, Cochrane and PsychInfo. The primary outcomes will include all aspects of child maltreatment: physical abuse, sexual abuse, emotional abuse, emotional neglect and physical neglect. Study participants will be between 0 and 18 years and will have reported maltreatment experiences using validated, reliable tools such as the Adverse Childhood Experiences Questionnaire as well as child self-reports and clinician reports. Study selection will follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines, and the methodological appraisal of the studies will be assessed by the Newcastle-Ottawa Quality assessment scale. A narrative synthesis will be conducted for all included studies. Also, if sufficient data are available, a meta-analysis will be conducted. Effect sizes will be determined from random-effects models stratified by gender, urbanisation and the pre-2012 and post-2012 POCSO Act cut-off. I^2 statistics will be used to assess heterogeneity and identify their potential sources and τ^2 statistics will indicate any between-study variance.

Ethics and dissemination As this is a rapid review, minimal ethical risks are expected. The protocol and level 1 self-audit checklist were submitted and approved by the Usher Research Ethics Group panel in the Usher Institute (School of Medicine and Veterinary Sciences) at the University of Edinburgh (Reference B126255). Findings from this review will be disseminated widely through peer-reviewed publications and in various media, for example, conferences, congresses or symposia.

PROSPERO registration number CRD42019150403.

Strengths and limitations of this study

- This is the first rapid review of the peer-reviewed and published evidence on child maltreatment in India drawing on studies from four electronic databases: PubMed, EMBASE, Cochrane and PsychInfo.
- The primary outcome is any type of prospectively recalled child maltreatment including physical abuse and neglect, emotional abuse and neglect and sexual abuse.
- First study to evaluate the effects of the legislative Act pre-2012 and post-2012 (Protection of Children from Sexual Offences Act 2012).
- The review is limited by the exclusion of the grey literature and non-English peer-reviewed publications due to the lack of resources and funding available to conduct the review.
- The review will include a narrative synthesis of non-governmental reports, media articles and other sources of grey literature to inform the findings.

INTRODUCTION

In the past two decades, significant progress has been made in reducing child deaths. UNICEF estimates that, worldwide, there were 12.6 million children who died prematurely before the age of 5 in 1990 and this has declined to 5.3 million in 2018.¹ Of the children who do survive, 200 million do not reach their full potential due to a combination of family socioeconomic status, geographic location, ethnicity, disability or religious orientation.¹ It is every child's right to develop and thrive, not just survive. Balanced nutrition, good healthcare and consistent care and encouragement in supportive home and school environments and positive neighbourhood environments enable children to lead healthy lives.² The early years of childhood therefore provide the basis of intelligence and learning, personality and self-motivation,

social behaviour and collaboration, which extend into adulthood.³

The WHO defines child maltreatment as ‘the abuse and neglect that occurs to children under 18 years of age and includes all types of physical and/or emotional ill-treatment, sexual abuse, and neglect which results in harm’. The WHO reports that worldwide, child maltreatment is widespread with one in four adults reporting being physically abused as children.^{4,5} Research suggests that children who have experienced such adversity do not receive the right nutrition, care or opportunities to learn and grow as children in more affluent geographical areas or economies.^{6,7} However, these findings are not generalisable to low/middle-income countries (LMICs) like India where millions of children may be exposed to other, specific adversities such as conflict, political instability, community violence and other traumatic life experiences.^{7,8} Research on early childhood adversity and consequences has been limited in India, including a lack of cross-sectional and longitudinal studies.⁹ These adversities such as abuse or neglect, along with associated habits such as underage smoking or drinking, formed in childhood, have been linked to a range of poor physical and mental health outcomes throughout the life course and ultimately greater premature mortality.¹⁰ These adversities and consequent health inequalities can be exacerbated by socioeconomic status and poverty, urbanisation (rural areas or slum areas compared with urban zones) or even caste.¹¹

In her book, *Bitter Chocolate*,¹² Virani discusses the context of these adversities in India—a highly patriarchal culture in which reprimanding, punishing or spanking a child (physically abusive behaviour) is a cultural norm.¹³ The Indian economy is further characterised by external adversities from a rapidly growing economy, stark urban–rural divides, religious strife and geopolitical upheavals.¹⁴ Data from UNICEF and the World Bank¹⁵ show a global perspective of child rights and Indian children are marked as experiencing a difficult situation as categorised

by their rights, access to education and healthcare and exposure to child labour and child marriages (figure 1). It is no surprise that physiological and neurobiological studies have found that adversities such as child maltreatment can lead to changes in neural networks, impaired nervous, endocrine and immune systems development, resulting in physical, mental and emotional dysfunction and chronic physiological damage.^{16–18}

Indian context

In India, home to 19% of the world’s children, it is estimated every second child is exposed to sexual abuse and violence.¹⁹ The Indian National Crimes Records Bureau (NCRB) reports a child is sexually abused every 15 min and 53% of children report abuse by a parent, relative or school teacher.²⁰ The prevalence of child sexual abuse (CSA) in high-income countries is 20% for females and 8% for males²¹ but in India, the estimates vary between 4%–66% for females and 4%–57% for males.^{22–24} These figures are considered underestimates because of limited surveillance of direct childhood adversity such as physical abuse or emotional abuse as well as CSA due to stigma particularly affecting girls; under-reporting of cases by healthcare and police authorities; stigma and taboo associated with being a victim or experiencing abuse, and varying prevalence rates by geography (eg, urban vs rural vs slum settings) as well as varying communities (school-based, hospital-based, trafficking victims, etc).^{24–27}

Poverty, and especially urban poverty in India is prevalent due to high economic migration that has translocated poverty from rural settings to urbanised areas.²⁸ While the proportion of deprived populations in India dropped from 45% to 22% between 1994 and 2012, one in every six urban Indians live in slums.²⁹ Health inequalities that exist as a result of poverty and rapid urbanisations need to be reduced so that child maltreatment, violence against girls and women, aims of the United Nations Sustainable Development Goals can be achieved. Bywaters *et al*³⁰ report strong associations between the




Figure 1 World map showing the rights of a child as measured by UNICEF and World Bank data.¹⁵

socioeconomic status of a family and the risk of children within the household experiencing abuse and neglect. The greater economic hardship in areas like urban slums, the greater the likelihood and severity of child maltreatment. By contrast, we see evidence of the reverse in Deb and Modak's³¹ study where violence is proportional to economic status as well as cultural beliefs and practices. Their data on socioeconomic status found that children from high-income households were almost four times more vulnerable to physical violence compared with children from low-income households. This is attributed to parents within high-income households having high expectations of their children especially with regards to academic achievement and a general undermining of a child's status compared with elder perspectives.³² We see the nuanced effects of economics and urbanisation as children from more deprived backgrounds have parents who remain indifferent to their care and well-being especially for female children.³³ Given these findings, this review will carefully consider socioeconomic status of families, urbanisation metrics and specifically, the types of maltreatment affecting each sex within these settings.

Previous research has demonstrated that experiences of child maltreatment rarely occur in isolation and in fact, have a tendency to cluster.⁹ Community-based studies have reported the prevalence and overlap of childhood sexual, physical and emotional abuse, but in India this has been limited. Choudhry *et al*'s²² systematic review focused on one aspect of child maltreatment: CSA, and was seminal work in detailing the high prevalence rates of CSA in Indian children. However, the authors acknowledged that CSA does not occur in silos and often co-occurs with other forms of child maltreatment in the same child.³⁴

Childhood adversity in India

Over the past 20 years, the phrase 'adverse childhood experiences' or (ACEs) have been studied as forms of maltreatment and household dysfunction affecting children either directly or indirectly. Felitti and colleagues³⁵ found a strong graded relationship between exposure to abuse or dysfunction during childhood and multiple risk factors for mortality in adulthood. ACEs have been associated with non-communicable conditions such as cardiovascular disease, cancer or mental illness.^{7 36–38}

While alternative discourse on ACEs exists,³⁹ there are limitations in the application of this ACEs framework including different types of exposures, lack of consideration for frequency or severity of occurrence and the relevance to population level, rather than individual level, policies. Also, the application of this framework within LMICs like India may be limited given the effects of other adversities such as poverty, community violence and political strife.⁴⁰

Using Felitti's ACEs framework and extending it with additional adversities,⁴⁰ figure 2 depicts different types of adversities each bearing significant ramifications extending from childhood to adulthood and beyond—affecting the family and society as a whole.⁴¹ However, in India, the data on child maltreatment across various different states and settings are available but the data on ACEs are limited. A few recent studies have been able to identify the prevalence of ACEs in specific communities.²² Damodaran and Paul⁹ compiled data on ACEs and specifically, child maltreatment in youth based in Kerala (South India) and reported that in a sample of 600 young people, 91% had reported at least one ACE and over half the population reported three or more ACEs. However, in terms of severity, males reported exposure to more




Figure 2 Adverse childhood experiences (ACEs) including traditional direct ACEs (child maltreatment: physical, sexual and emotional abuse and neglect) and indirect ACEs (Felitti *et al*³⁵) and further extended by researchers since the original study (Houtepen *et al*⁴⁰).




Figure 3 Potential pathways explaining child maltreatment using the socioecological model proposed by Sidebotham and Heron.⁴⁵

severe child maltreatment than females such as physical and sexual abuse.

In India, lower incomes, poor living conditions and living with domestic violence are adversities that can lead to neglect and abuse and often ‘cluster’.⁴² Figure 3 combines the socioecological model of child maltreatment⁴³ and suggests potential pathways (eg, from a macrosystem/societal or exosystem/community level) that may contribute to child maltreatment.^{44 45} Particularly, around the individual level factors—there is a gap in the rates of other types of child maltreatment associated with CSA (eg, emotional abuse or neglect). While the topic of ACEs in India has received limited attention, CSA has received considerable attention from academics, clinicians, policymakers and legislators in recent years.^{22 46} With the high prevalence of all types of child abuse in India, an attempt to amalgamate and compile the findings and present these figures in a systematic, cohesive and coherent manner is pressing.⁵

Effect of gender

Previous research has suggested the nuanced effect of gender on the occurrence of child maltreatment in India.²² In India, gender identification is illegal and yet, almost half of girls ‘wish they were born boys’.⁴⁷ Although

anecdotal evidence and case reports suggest that prevalence of sexual abuse is higher in female children than in male children, based on hospital services or care services utilisation,⁴⁸ some studies shows equal rates of sexual abuse across gender.^{47 49} Deb⁵⁰ suggests that boys may be experiencing and reporting more sexual abuse while girls are less likely to be aware of abuse (knowledge and attitudes) or less likely to report abuse for fear of shame or stigma. Society is more dismissive of male abuse than that of girls as patriarchal tendencies to ‘protect’ include girls, but do not extend to boys.⁴⁹ However, when it comes to neglect with regards to food and education security, girls are known to be disadvantaged.⁵¹ India is the only country where girls have greater rates of under-five mortality compared with boys. Indian girls are breastfed for shorter periods and consume less milk.⁵¹ The reasons for this discrepancy are varied: there is a preference for sons over daughters in India and as an extension, more opportunities with regards to food, healthcare and education are afforded to male children.⁵² However, given the complexities of gender effects in different types of abuse, generalisations cannot be drawn due to the limited number of studies, methodological limitations (eg, small sample sizes, non-generalisability of findings to the wider Indian

diaspora), urbanisation and socioeconomic privilege (eg, remote tribal areas being worse affected than wealthier urban ones⁵³) and reporting bias (eg, particularly if parents reported via food diaries what was consumed).

Effect of urbanisation

The prevalence of child maltreatment by urbanisation (rural, slum or urban areas) has received limited attention in the evidence base. Only two studies^{54 55} included community populations from rural areas as well as urban areas. While other cross-sectional studies have been conducted in urban areas of India,⁵⁶ the child maltreatment prevalence figures (32% of boys and 42% of girls) are higher but not comparable due to differences in sample size; socioeconomic status; access to healthcare and support systems in urban areas compared with rural areas, as well as differing definitions of child maltreatment.

Banerjee⁵⁷ discussed that child maltreatment was more prevalent in metropolitan cities than in rural areas of India and their study found a high frequency of physical abuse among the slum and pavement dwellers although the severity of maltreatment was low. The study highlighted child labour practices (with 78.4% of children between the ages of 4–5 years working part-time jobs) as a result of living in a deprived area of the city. This study also highlighted the fact that often child maltreatment is a product of dire economic distress and within this context; small-scale regional or local studies might help understand the prevalence of child maltreatment in specific areas by urbanisation metrics. Over half the children who were classed as homeless (ie, living in drainage pipes, under tarpaulins, flyovers and those living in the open) reported CSA.⁵⁸ While child maltreatment figures such as CSA may seem high in urban or slum areas, the disadvantage with rural areas may be that abuse may go under the radar. They may be under-reported or disguised within local traditions such as a dowry culture or child marriages, all behaviours rooted in poverty and volatile socioeconomic circumstances.⁵⁹

Effect of Protection of Children from Sexual Offences Act

The Protection of Children from Sexual Offences Act (POCSO Act) was established in 2012 by the Indian government to protect children against offences like sexual abuse. The Act makes it mandatory to report acts of sexual offence and document reporting timelines, prosecution timelines and clearly record all evidence (within a 30-day period). While POSCO has been a relatively recent Act, its effect on number of reported offences has been debated.²² While the gold standard remains as a multi-disciplinary, multi-faceted and holistic team providing psychological and legal support to the child and their families, this level of comprehensive care *under one roof* is yet to exist in India.^{60 61} The impact of this study on policymaking will come from its inclusion of the only legislative act in India that focuses on CSA. Given the overlap between different types of abuse and neglect and their tendency to ‘cluster’, we will determine

any effects of POCSO pre-2012 and post-2012 on all forms of child maltreatment. This has not been evaluated by the literature to date but was highlighted for future work by Choudhry and colleagues.²²

Rationale

There are known gaps in the evidence base with regards to the overlap of different forms of child maltreatment and specifically, the gendered effects of child maltreatment. These include the identification and motivation of perpetrators and what happens after maltreatment has happened—a child’s awareness of their rights and attitudes towards abuse and neglect and reporting of child maltreatment. The circumstances that encourage reporting and details on protective factors and risk factors of child maltreatment have received limited attention in the literature.²²

This review will build on these thematic and methodological gaps in the literature, synthesise the evidence by gender, urbanisations and POCSO effects and provide practice, policy and clinical solutions that may help reduce the occurrence and impact of child maltreatment in India.

Research aims and objectives

The primary research aim is to describe the prevalence of child maltreatment in India. The secondary aim is to determine the effect of the POCSO Act on CSA and on other forms of child maltreatment.

The specific objectives were: to determine the prevalence of child maltreatment (physical abuse, sexual abuse, emotional abuse and physical and emotional neglect); to present rates by gender, urbanisation (rural, slum or urban) and study sample types (eg, community, school-going, high-risk such as trafficking victims, etc); to determine any impact of the POCSO Act (by the date of implementation of the Act, ie, 2012) on rates and to identify implications for future policy, practice and research amendments.

METHODS AND ANALYSIS

Study design

This review and meta-analysis will be carried out in compliance with the Cochrane Handbook for Systematic Reviews of Interventions (version 6.1) and reported in compliance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.^{62–64}

Patient and public involvement

It was not appropriate or possible to involve patients or the public in the design, conduct, reporting or dissemination plans of our research. The research team and advisory/expert panel consisted of clinicians, policymakers and research academics who work in the area of early childhood adversity and were able to advise on search terms used; databases selected and dissemination plans for the work on completion.

Participants

Inclusion criteria

Studies between 1 January 2005 and 1 January 2020 will be included to allow a comparison of an 8-year window pre-2012 and post-2012 POCSO Act. Studies with participants aged between 0 and 18 years will be included in the review; and will have experienced any form of child maltreatment as determined using International Society for the Prevention of Child Abuse and Neglect Child Abuse Screening Tool-Child Version, the Childhood Trauma Questionnaire, Adverse Childhood Experiences International Questionnaires (ACEIQ) or self-reports or clinician reports. ACEIQ or WHO definitions will also be used to determine child maltreatment⁶⁵ in addition to those by Day *et al.*⁶⁶ The primary outcome measure is any type of child maltreatment: sexual, physical or emotional abuse; and physical and emotional neglect. In addition to the definition, the time frame of recall, prospective recall in our case, warrants discussion.

Often, prospective and retrospective measures of child maltreatment do not always identify the same individuals.⁶⁷ Retrospective studies of child maltreatment are often regarded as a shortcut as significant time; finance and resources are not required to undertake prospective studies.⁶⁸ There is a low agreement between prospective and retrospective assessments of child maltreatment that can be explained by factors such as motivation of reporting, measurement features and memory bias. First, motivation can reduce agreement if individuals may gain something by intentionally withholding data about their maltreatment for embarrassment or fear or feeling uncomfortable or upset. Second, child maltreatment assessments are known to have imperfect test-retest reliability⁶⁹ and prospective measures might only account for the more severe of cases while retrospective 'true positive' case.⁶⁷ Finally, memory bias can reduce agreement as a result of impaired brain development,⁷⁰ deficit in memory consolidation of traumatic events or disrupted sleep patterns.⁷¹ Therefore, given the variation, the focus on prospective recall (rather than retrospective) of child maltreatment for systematically reviewing the literature on abuse and neglect in children would reduce variation errors, measurement bias and memory bias.⁶⁷

Exclusion criteria

Studies that are qualitative in nature or mixed-methods (due to the time factor of this review), non-Indian study contexts, non-English studies, retrospective studies of adult CSA survivors, studies with insufficient peer-reviewed published data, media reports, newspaper articles and non-peer-reviewed articles will be excluded from this review. Only quantitative studies published in the English language will be included.

Outcomes

Defining childhood maltreatment

The WHO defines child maltreatment as 'the abuse and neglect that occurs to children under 18 years of age

and includes all types of physical and/or emotional ill-treatment, sexual abuse, neglect which results in harm'. The National Society for the Prevention of Cruelty to Children (NSPCC) defines child maltreatment as harm experienced by a child by an adult or child *that is physical, sexual or emotional but can also involve a lack of love, care and attention.*⁷² The NSPCC also set out clear descriptions of what constitutes child maltreatment and specifically each form of abuse and neglect that are summarised here.

Child physical abuse

Physical abuse happens when a child is hurt or harmed deliberately causing injuries such as bruises, burns, cuts and broken bones. It may also include children being administered medication they do not need or not being given essential medicines for illness. The terms used specifically are kicking, shaking, slapping, throwing, poisoning or drowning. In India, these terms and additional ones such as 'spanking', 'rap' or 'pasting' also have equivalent meanings.

Child sexual abuse

Sexual abuse happens when a child is forced or tricked into sexual activities. It happens when an abused makes physical contact with a child and includes touching, kissing and oral sex or non-contact abuse which includes exposing or flashing, exposing a child to sexual acts or making, viewing or distributing child abuse images or videos.

Child emotional abuse

Emotional abuse includes acts that humiliate a child by constantly criticising them, insulting or shouting at them, blaming and making them a scapegoat or manipulating a child. These behaviours also include not allowing them to have friends or never saying anything kind or positive or congratulatory to a child. It perpetuates the feelings that a child is not loved, is worthless or inadequate.

Child physical and emotional neglect

Neglect is the ongoing failure to meet a child's basic requirements and is often also classed as a form of child abuse. Physical neglect is when a child's basic needs such as food, clothing or shelter are not met or they are not properly supervised or kept safe. Emotional neglect is when a child is not provided with the nurture and stimulation they need to grow and thrive.

While child maltreatment is a global problem, it is certainly difficult to assess and manage in LMICs like India. Specific types of maltreatment such as abandonment, child labour, street-begging and corporal punishment are highly prevalent.⁷³ Therefore, while generic methods may exist to assess child maltreatment, the value of self-reports from children or from paediatricians, child psychologists or social workers and the police should not be dismissed.

Table 1 Search term strategies using keywords and Medical Subject Heading (MeSH) terms

Sexual abuse	Sexual abuse (cont)	Physical abuse	Emotional abuse	Emotional neglect	Additional child maltreatment definitions
Child* sexual abuse	Child abuse survivor	Child hitting	Child worthless	Denial clothing	Child neglect
Child sexual assault	Child sexual offender	Child shaking	Child unloved	Denial shelter	Child psychological assault
Child sexual coercion	Child pornography	Child throwing	Child inadequate	Denial food	Child denial
Child rape	Child molestation	Child poisoning	Child failure	Carer failure	Child pasting
Child prostitution	Child sexual crime	Child burning	Child unloved	Parental failure	Child emotional assault
Child crime victims	Child physical	Child scalding	Child corruption	Emotional harm	Non-accidental injury
Child incest	Child assault	Child drowning	Child unheard	Emotional danger	
Child perpetrator/ child victim 3	Child beating	Child suffocating	Cruelty	Child* neglect	
Child physical violence	Child emo* abuse			Abandonment	
Child mental violence	Child maltreat*			Bereavement	
Child sexual violence	Child sexual coercion			Grief	
Child sexual offence	Child sexual aggression				

Search terms: (“child abuse, sexual”[MeSH Terms] OR (“child”[All Fields] AND “abuse”[All Fields] AND “sexual”[All Fields]) OR “sexual child abuse”[All Fields] OR “physical child abuse” [All Fields] OR “emotional child abuse”[All Fields] OR (“child”[All Fields] AND “sexual”[All Fields] AND “abuse”[All Fields]) OR “child sexual abuse”[All Fields]) AND (“india”[MeSH Terms] OR “india”[All Fields])—expanded from Choudhry *et al*²² and Day *et al*⁶⁶ and updated to include child maltreatment definitions.

*Alternative search terms to include adolescent sexual abuse, adolescent physical assault and so on.

Search strategy

The relevant search terms to be included in the search strategy will be identified during a preliminary search reported in the study protocol using PubMed’s Medical Subject Heading’s (MeSH) browser. The search contained both free-text terms and controlled vocabulary like MeSH terms as well as truncation and Boolean operators. In order to improve the quality of the search strategy and avoid bias or errors, the strategy includes choosing all appropriate terms, both descriptors and synonyms including those that may be specific to the Indian diaspora.⁷⁴ First, the initial search terms were developed with reference to seminal works or major publications of child maltreatment and early childhood adversity in India.^{9 26 47 75} The review then used the MeSH terms and keywords documented (table 1) and were further enhanced by the search strategy from a systematic review conducted on CSA in Indian children.²² These were further validated with input from an expert advisory panel in the field of systematic reviews and ACEs in India (online supplemental appendix 1). Five electronic databases will be systematically searched: PubMed, EMBASE, Cochrane and PsychInfo and Campbell Library. PubMed and EMBASE are the two largest healthcare bibliographic databases containing the vast majority of published health-related literature.^{74 76} Searching Cochrane and

PsychInfo will provide nuanced searches of other systematic reviews of child maltreatment or ACEs from the Asian continent and specifically from India. Systematic reviews will be included in the search to identify potentially relevant studies that may have been excluded by the electronic search strategy. Online supplemental appendix 2a–d contains the search strategies used in the four databases, respectively. These search strategies and advice on specific electronic databases will be cross-checked and verified by an information specialist. The final search strategy used will be published in detail in the future published manuscript of results.

Data management

The selection of studies will be carried out systematically by two people independently, and in accordance with PRISMA guidelines, to improve accuracy and to minimise bias.⁶³ The first reviewer (first author) is an epidemiologist with expertise in early childhood adversities and the impact on health outcomes; the secondary reviewer is a clinical psychologist with expertise on ACEs in India while the third and fourth reviewers, are clinical doctors with expertise in general medicine and paediatric care, validated the selection of studies pre-2012 and post-2012 threshold.



All studies identified from the individual search engines will be imported into a reference manager, EndNote, which will be used for de-duplication. The references will then be extracted into Rayyan for management between the first author and secondary reviewers.⁷⁷ All abstracts and full-text articles will be reviewed independently by the first author and secondary reviewers as a fixed percentage of the work (40%). The details of all articles processed (including duplicates and details of excluded texts at each screening stage) as well as reasons for inclusion and exclusion from the study will be recorded and logged for the first author and secondary reviewers and presented in a PRISMA flowchart. At the title/abstract screening stage, the authors will use Rayyan's coding system (traffic light system) with appropriate tags such as 'wrong outcome of interest' in order to code each study (done independently by the authors on Rayyan's blinded review setting). Once all titles/abstracts are screened, we will unblind all authors and discuss those titles/abstracts to be included, or excluded or further discussed with the expert panel as a result of discrepancies.⁷⁷ All selected full-text articles will be screened by the primary author and secondary reviewers and the reasons for exclusion of these articles will be recorded and presented in the final manuscript.

Data extraction

The following data will be extracted from the included studies: title; author information; year of study (not publication date); study design; recruitment strategy and sample characteristics (school-based, community, hospital based, etc); sample size; urbanisation metrics (urban or rural or slum); response rate; definition of child maltreatment used; timeframe of child maltreatment (in years, eg, past 1 year or past 6 months or half a year); mean age of participants; gender distribution; statistical methodology; confounder data including parental relationships, home ownership, family settings (eg, nuclear, joint, mixed), religion, educational status and literacy rates; prevalence rate of each type of child maltreatment; author's key message or conclusions; ethical approval and any conflicts of interest. An example of a completed data extraction form is presented in [table 2](#).²⁶

Quality assessment

The final selected studies will be appraised for quality by the first author and, to ensure robustness, they will also be quality appraised by a second reviewer. A third reviewer will resolve any conflicts, with support from the expert advisory panel if required.

Study quality (largely cross-sectional and case-control) will be assessed using the Newcastle-Ottawa Quality assessment scale (NOS) adapted for use in cross-sectional study designs as it considers key areas: selection (representativeness of the sample, sample size, non-responders and exposure details); comparability (is confounding considered) and outcome (blinding, recording and statistical test used).^{78 79} While there are a myriad range of

assessment tools available, the NOS has been endorsed by the Cochrane Collaboration to assess quality of research studies.⁷⁶ An example of an NOS assessment of a sample study is presented in online supplemental appendix 3.

Data synthesis

The study will include a narrative synthesis of the findings and attempt a meta-analysis dependant on heterogeneity. A meta-analytic approach will be used including any random-effect pooled prevalence of child maltreatment stratified by gender, urbanisation and pre-POCSO and post-POCSO Act. Heterogeneity will be evaluated using forest plots and measuring the impact of the I^2 statistic ($I^2 > 50\%$ indicates significant heterogeneity) and tested with the χ^2 statistic ($p < 0.05$).⁶² We will also report τ^2 values to indicate any between-study variance. We will use a fixed effects model for meta-analysis, except where we identify statistical heterogeneity in which case, we would use a random effects Poisson model⁸⁰ to determine an 'average prevalence' across the population of all included studies. We will use the 'metaprop' command to perform meta-analyses of proportions of child maltreatment in STATA version 16, which builds on the 'metan' command⁸¹ but is typically used to pool effects such as proportions from cross-sectional studies instead of ORs from case-control studies.

In the case of significant heterogeneity derived from our analysis, we will attempt to create subgroups of the paediatric population to understand any extracted data. This could include different age-bands of participants (eg, child (0–12 years), adolescent (13–18 years) using WHO cut-offs for age bands in paediatric populations. We may also wish to compare prevalence rates of child maltreatment based on self-report measures versus objectively assessed measures as well as compare rates by time (1 year compared with lifetime occurrence or frequency data).

Meta-bias(s)

Research papers were written originally in Hindi or any other Indian dialect, that would otherwise be included in the data capture, will not be used for the purpose of this study.

DISCUSSION

The findings of this study will aim to add to the limited knowledge of child maltreatment in India and specifically, the impact of gender, urbanisation and the POCSO Act on the prevalence and determinants of child maltreatment. There are known gaps in the evidence base with regards to the overlap of different forms of child maltreatment and specifically, the gendered effects of child maltreatment. These include the identification and motivation of perpetrators and what happens after maltreatment has happened—a child's awareness of their rights and attitudes towards abuse, neglect and reporting of child maltreatment. The circumstances that encourage

Table 2 Example of a data extraction form

Title of study	'Prevalence of child abuse in Kerala, India: An ICAST-CH based survey'
Authors	Manoj Therayil Kumar, Nilamadhhab Kar, Sebind Kumar
Year of study	2019
Study design	Cross-sectional survey in Thrissur, Kerala
Recruitment strategy	There are 39 schools in Thrissur with 15 150 students in attendance. Random number generation in excel was used to select half the schools on the list and all students in grades 8–10 who attended school on that day were eligible to take part
Sample setting (school/college, clinical, community)	School setting
Sample size and response rate	6957 participants took part in the survey. No details of response rate were provided however the authors state that no questionnaire was returned completely blank
Child maltreatment definition used (timeframe defined)	All abuse-related questions in the ICAST-CH were scored on a severity scale and categorised into both 1-year prevalence and lifetime prevalence
Classification tool validity/reliability of tool	The International Society for the Prevention of Child Abuse and Neglect (ISPCAN) Child Abuse Screening Tool-Child Home Version (ICAST-CH) was used Developed in unison with the WHO and UNICEF and has good criterion validity and was translated into Malayalam and was piloted for use before being administered to the students. Internal consistency measures using Cronbach's alpha for subscales—exposure to violence, physical abuse, emotional abuse, neglect and sexual abuse
Urban/rural/slum-dwelling participants breakdown	Not mentioned
Mean age (SD)	13.9 years (1.12) for males and 13.7 years (0.99) for females
Gender distribution	2071 males (30.1%) 4810 females (69.9%)
Example: CSA prevalence by gender	30.9% in males for 1-year prevalence 10.8% in females for 1-year prevalence 34.2% in males for lifetime prevalence 14% in females for lifetime prevalence
Statistical methods used	Frequency and counts used to determine prevalence and presented with 95% CIs Cronbach alpha used to determine internal consistency of the Malayalam version of the scale Significance was set at the 0.05 level as a maximum but 0.01 and <0.001 were reported
Confounders measured and/or adjusted for	Measured confounders—gender, age, religion, family type, accommodation, parental accommodation and socioeconomic status
Other confounders discussed but not measured	Literacy rates, cultural strata of participants
Participants experience of: Emotional abuse (rate, %) Physical abuse (rate, %) Sexual abuse (rate, %) Emotional neglect (rate, %) Physical neglect (rate, %)	
Conclusion/comments	The abuse reported was more occasional experiences rather than frequent reporting nonetheless as 89.9% of any abuse in the past 1 year and 91% in the lifetime. First, study in Indian diaspora to use the ICAST-CH tool to estimate household abuse. Did not include external environments such as schools
Conflicts of interest	None declared
Ethical approval	Ethical approval was sought from local authority and educational ministry to conduct the research
Funders	Municipal Corporation of Thrissur city

CSA, child sexual abuse.

reporting, details on protective factors and risk factors of child maltreatment have received limited attention in the literature.²²

Furthermore, the impact of child maltreatment and where they occur (eg, urban areas vs slum areas) is important as these may offer an insight into potential interventional pathways that differ by geography, for example, enforcing child welfare policies in rural areas.²² The impacts of urban and rural poverty, socio-economic status; family income; parental occupation; family structure (nuclear, joint, extended, etc) and other confounder data will be collated during this review to aid in the interpretation of results including subgroup

analysis and meta-analysis. Furthermore, the impact of this study on policymaking will come from its inclusion of the only legislative act in India that focusses on child maltreatment. The POCSO Act was established in 2012 by the Indian government to protect children against offences like sexual abuse. The Act makes it mandatory to report acts of sexual offence and document reporting timelines, prosecution timelines and clearly record all evidence (within a 30-day period). While POSCO has been a relatively recent Act, its effect on the number of reported offences in research studies has yet to be determined.



Implications on policy and research

When the evidence on child maltreatment in India is collated and presented, any policy or practice impacts can be tailor-made based on the country-specific context and findings. In India, the challenge will come from addressing cultural norms and traditional values (eg, gender inequity and reducing child marriages), creating safer environments for children within homes and schools (eg, authorities penalising assault) and scaling up response and support services for children in different settings such as tribal regions, slum areas, rural communities (eg, trauma informed health and social care). This approach has worked in higher-income countries but in LMIC's like India, there are challenges. These include low levels of trauma-informed care for injured children in hospital settings with 91% of healthcare workers not receiving any trauma-informed training.⁸² Other challenges included overburdened judicial and criminal systems; poor child welfare services; gender inequality and illiteracy.²² This study will capture data on health inequities and social determinants that predispose to maltreatment in different geographical areas of India including confounders such as poverty and neglect.

Strengths and limitations

There are some strengths to our approaches. This is the first review undertaken focussing on all aspects of child maltreatment in India where previous review articles and studies have focused on a specific facet of maltreatment, for example, CSA. Second, the inclusion of the POCSO Act makes this review nuanced to comment on any effect of legislation where it has never been formally evaluated.⁸³ This study will include an analysis of child maltreatment studies and corresponding prevalence rates, both pre-2012 and post-2012. With the inclusion of the POCSO 2012 threshold, it is expected that there will be an increased number of cases reporting all forms of child maltreatment. Third, the authors will attempt to meta-analyse the data depending on heterogeneity scores generated. Choudhry *et al's* work²² highlighted the significant heterogeneity of studies conducted to date and this might limit the quality of meta-analytics possible given the discrepancies in samples size. Possible reasons for heterogeneity include varying age ranges of participants, selection bias with sample provenance (eg, recruitment from observation homes vs school children), varying urbanisation metrics (eg, samples including children from all types of socioeconomic backgrounds) and variation in outcome assessment definition (eg, self-report vs validated questionnaire).

There are limitations to this review. First, only papers in English were included due to limited access to funding and resources such as a professional translator. We acknowledge the importance of Anglophone bias in the context of this Indian based study therefore endeavour to ensure future extensions of this research does include non-English papers. Second, excluding grey literature from OPEN Grey, narrative reports from non-governmental organisations or World Bank reports would limit some of the 'unmeasured' exposures and confounders that may increase child maltreatment.^{25 49}

However, the authors will make attempts to collate this additional information so that they may be presented as part of a narrative synthesis on the topic of child maltreatment in India, thus improving the quality, generalisability and relevance of our work.⁸⁴ The exclusion of national statistics (eg, NCRB records) might limit interpretation of any effect of the POCSO Act on conviction rates for child maltreatment. Qualitative studies were also excluded from this review which may not effect prevalence rates of child maltreatment but would allow exploration of non-responses from children (interpretative) or indeed, highlight determinants that have not been identified in quantitative or aggregated reviews.^{85–87}

This review will build on these thematic and methodological gaps in the literature, synthesise the evidence by gender, urbanisations and POCSO effects and provide practice, policy and clinical solutions that may help reduce the occurrence and impact of child maltreatment in India.

Ethics

A level 1 self-audit checklist was approved by an ethics panels at the Usher Institute at the University of Edinburgh. Minimal ethical risks are expected. However, Vergnes *et al*,⁶⁴ propose that for systematic reviews, particularly on sensitive topics authors should consider a minimal ethical assessment review and report these in their research studies. As a result, in this review, ethical approval details from each included study will be collected.

Dissemination

The findings of this review will be published in a peer-reviewed journal, and disseminated at national and international conferences and through social media.

Author affiliations

¹Population Health Sciences, University of Bristol, Bristol, UK

²Department of Surgery, Norfolk and Norwich University Hospital NHS Trust, Norwich, UK

³Department of Child and Adolescent Psychiatry, National Institute of Mental Health and Neuro Sciences, Bangalore, Karnataka, India

⁴Department of Medicine, Norfolk and Norwich University Hospital NHS Trust, Norwich, UK

⁵College of Medicine and Veterinary Medicine, The University of Edinburgh Usher Institute of Population Health Sciences and Informatics, Edinburgh, UK

⁶Child and Adolescent Psychiatry, National Institute of Mental Health and Neuro Sciences, Bangalore, Karnataka, India

⁷Public Health, Sambodhi Research and Communications Pvt Ltd, Noida, Uttar Pradesh, India

Acknowledgements The authors would like to extend their sincere gratitude to Mrs Sarah Herring, Subject Librarian for Population Health Sciences, University of Bristol for her advice and guidance during the protocol development stage.

Contributors GF, RG and EP were responsible for the initial protocol drafting and design. GF, MF, NV, PDS then contributed to preliminary searches used to develop the rationale and background of the study. ES, BH, VC, VB, PDS and MF were on the expert advisory panel for the study to advise on methods design and statistical analysis techniques that could be used. All authors contributed and approved this final manuscript.

Funding GF is funded by the UK Medical Research Council and Alcohol Research UK Grant (MR/L022206/1).

Map disclaimer The inclusion of any map (including the depiction of any boundaries therein), or of any geographic or locational reference, does not imply the expression of any opinion whatsoever on the part of BMJ concerning the legal

status of any country, territory, jurisdiction or area or of its authorities. Any such expression remains solely that of the relevant source and is not endorsed by BMJ. Maps are provided without any warranty of any kind, either express or implied.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Gwen Fernandes <http://orcid.org/0000-0003-0203-7053>

REFERENCES

- 1 WORLD HEALTH ORGANISATION. Children: reducing mortality, 2019. Available: <https://www.who.int/news-room/fact-sheets/detail/children-reducing-mortality> [Accessed 02 Feb 2020].
- 2 GRIFFITHS S, HUNTER DJ. *New perspective in public health*. London: Radcliffe Publishing, 2007.
- 3 REYNOLDS AJ, ROLNICK AJ, TEMPLE JA. *Health and education in early childhood: predictors, interventions and policies*. Cambridge: Cambridge University Press, 2015.
- 4 WORLD HEALTH ORGANISATION. Country cooperation strategy at a glance: India, 2017. Available: https://apps.who.int/iris/bitstream/handle/10665/136895/ccsbrief_ind_en.pdf;jsessionid=90BA4E2E08E57D23C4363B8FBDFC85AC?sequence=1 [Accessed 05 Mar 2020].
- 5 WORLD HEALTH ORGANISATION. Preventing child maltreatment: a guide to taking action and generating evidence, 2006. Available: https://apps.who.int/iris/bitstream/handle/10665/43499/9241594365_eng.pdf;jsessionid=555709D037DBAD5DE822D439283A5F1B?sequence=1 [Accessed 01 Feb 2020].
- 6 Blum RW, Li M, Naranjo-Rivera G, Naranjo-Rivera G. Measuring adverse child experiences among young adolescents globally: relationships with depressive symptoms and violence Perpetration. *J Adolesc Health* 2019;65:86–93.
- 7 Bellis MA, Hughes K, Ford K, *et al*. Life course health consequences and associated annual costs of adverse childhood experiences across Europe and North America: a systematic review and meta-analysis. *Lancet Public Health* 2019;4:e517–28.
- 8 Sharma E, Vaidya N, Iyengar U, *et al*. Consortium on vulnerability to Externalizing disorders and addictions (cVEDA): a developmental cohort study protocol. *BMC Psychiatry* 2020;20:2.
- 9 Damodaran DK, Paul VK. Patterning/clustering of adverse childhood experiences (ACEs): the Indian scenario. *Psychol Stud* 2017;62:75–84.
- 10 Hughes K, Bellis MA, Hardcastle KA, *et al*. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health* 2017;2:e356–66.
- 11 Kowal P, Afshar S. Health and the Indian caste system. *The Lancet* 2015;385:415–6.
- 12 VIRANI P. *Bitter chocolate*. New Delhi: Penguin Books, 2000.
- 13 Nijhara K, Bhatia S, Unnikrishnan B. Corporal punishment in children and its implications on mental health. *Indian J Pediatr* 2018;85:405
- 14 KOCHANNEK SA, HARDGRAVE RL. *India: government and politics in a developing nation*. Boston: Thomson Higher Education, 2008.
- 15 UNICEF. Ending violence against children in Asia, 2014. Available: https://www.unicef.org/publications/files/Ending_Violence_Against_Children_Six_strategies_for_action_EN_9_Oct_2014.pdf [Accessed 24 Jan 2019].
- 16 Anda RF, Felitti VJ, Bremner JD, *et al*. The enduring effects of abuse and related adverse experiences in childhood. *Eur Arch Psychiatry Clin Neurosci* 2006;256:174–86.
- 17 Pechtel P, Pizzagalli DA. Effects of early life stress on cognitive and affective function: an integrated review of human literature. *Psychopharmacology* 2011;214:55–70.
- 18 Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiol Behav* 2012;106:29–39.
- 19 Behere PB, Sathyanarayana Rao TS, Mulmule AN. Sexual abuse in women with special reference to children: barriers, boundaries and beyond. *Indian J Psychiatry* 2013;55:316–9.
- 20 NATIONAL CRIME RECORDS BUREAU. Crime in India – statistics, 2016. Available: <http://ncrb.gov.in/StatPublications/CII/CII2016/Press%20release-Crime%20in%20India%202016.pdf> [Accessed 14 Feb 2019].
- 21 Stoltenborgh M, van Ijzendoorn MH, Euser EM, *et al*. A global perspective on child sexual abuse: meta-analysis of prevalence around the world. *Child Maltreat* 2011;16:79–101.
- 22 Choudhry V, Dayal R, Pillai D, *et al*. Child sexual abuse in India: a systematic review. *PLoS One* 2018;13:e0205086.
- 23 WORLD HEALTH ORGANISATION. Child maltreatment, 2016. Available: <https://www.who.int/news-room/fact-sheets/detail/child-maltreatment> [Accessed 16 Oct 2019].
- 24 Radhika K, Manjula M, Jaisooriya TS. Ethical gaps in conducting research among adult survivors of child sexual abuse: a review. *Indian J Med Ethics* 2018;3:186–92.
- 25 Seth R, Bose V, Qaiyum Y, *et al*. Social determinants of child marriage in rural India. *Ochsner J* 2018;18:390–4.
- 26 Kumar MT, Kar N, Kumar S. Prevalence of child abuse in Kerala, India: an ICAST-CH based survey. *Child Abuse Negl* 2019;89:87–98.
- 27 Hailes HP, Yu R, Danese A, *et al*. Long-Term outcomes of childhood sexual abuse: an umbrella review. *Lancet Psychiatry* 2019;6:830–9.
- 28 Datt G, Ravallion M, Murgai R. Poverty and growth in India over six decades. *Am J Agric Econ* 2020;102:4–27.
- 29 Data WB. Population living in slums. Available: <https://data.worldbank.org/indicator/EN.POP.SLUM.UR.ZS?locations=IN> [Accessed 05 Feb 2021].
- 30 BYWATERS P, BUNTING L, DAVIDSON G. The relationship between poverty, child abuse and neglect: an evidence review, 2016. Available: file:///C:/Users/gf18522/Downloads/bywaters_can_final_report.pdf [Accessed 06 Jun 2019].
- 31 Deb S, Modak S. Prevalence of violence against children in families in Tripura and its relationship with socio-economic factors. *J Inj Violence Res* 2010;2:5–18.
- 32 Bandewar SV, Pitre A, Lingam L. Five years post Nirbhaya: critical insights into the status of response to sexual assault. *Indian J Med Ethics* 2018;3:215–21.
- 33 NATIONAL FAMILY HEALTH SURVEY, 2016. Available: http://rchiips.org/nfhs/factsheet_NFHS-4.shtml [Accessed 16 Jan 2019].
- 34 CARSON DK, FOSTER JM, CHOWDHURY A. Sexual abuse of children and youth in India: an anthropological perspective (the oriental anthropologist). *SAGE* 2014;14:343.
- 35 Felitti VJ, Anda RF, Nordenberg D, *et al*. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The adverse childhood experiences (ACE) study. *Am J Prev Med* 1998;14:245–58.
- 36 Springer KW, Sheridan J, Kuo D, *et al*. Long-term physical and mental health consequences of childhood physical abuse: results from a large population-based sample of men and women. *Child Abuse Negl* 2007;31:517–30.
- 37 Kessler RC, McLaughlin KA, Green JG, *et al*. Childhood adversities and adult psychopathology in the who world mental health surveys. *Br J Psychiatry* 2010;197:378–85.
- 38 Ip P, Wong RS, Li SL, *et al*. Mental health consequences of childhood physical abuse in Chinese populations: a meta-analysis. *Trauma Violence Abuse* 2016;17:571–84.
- 39 Kelly-Irving M, Delpierre C. A critique of the adverse childhood experiences framework in epidemiology and public health: uses and misuses. *Social Policy and Society* 2019;18:445–56.
- 40 Houtepen LC, Heron J, Suderman MJ, *et al*. Adverse childhood experiences in the children of the Avon longitudinal study of parents and children (ALSPAC). *Wellcome Open Res* 2018;3:106.
- 41 Gilbert R, Widom CS, Browne K, *et al*. Burden and consequences of child maltreatment in high-income countries. *Lancet* 2009;373:68–81.
- 42 Kiburi SK, Molebatsi K, Obondo A, *et al*. Adverse childhood experiences among patients with substance use disorders at a referral psychiatric hospital in Kenya. *BMC Psychiatry* 2018;18:197.
- 43 Sidebotham P. An ecological approach to child abuse: a creative use of scientific models in research and practice. *Child Abuse Review* 2001;10:97–112.

- 44 DAHLBERG LL, KRUG EG. Violence—a global public health problem. In: Krug E, Dahlberg LL, Mercy JA, *et al*, eds. *World report on violence and health*. Geneva, Switzerland: World Health Organization, 2002: 1–56.
- 45 Sidebotham P, Heron J, ALSPAC Study Team. Child maltreatment in the “children of the nineties”: a cohort study of risk factors. *Child Abuse Negl* 2006;30:497–522.
- 46 THOMAS REUTERS. India: the worlds most dangerous countries for women in 2018, 2018. Available: <http://poll2018.trust.org/country/?id=india> [Accessed 02 Dec 2019].
- 47 KACKER L, VARDAN S, KUMAR P. Study on child abuse: ministry of women and child development, government of India 2007.
- 48 MACIONNIS J, PLUMMER K. *Sociology: a global introduction*. United Kingdom: Harlow, 2012.
- 49 Subramaniyan VKS, Reddy P, Chandra G, *et al*. Silence of male child sexual abuse in India: qualitative analysis of barriers for seeking psychiatric help in a multidisciplinary unit in a general Hospital. *Indian J Psychiatry* 2017;59:202–7.
- 50 DEB S. *An empirical investigation into child abuse and neglect in India: burden, impact and protective measures*. Pondicherry: Springer, 2018.
- 51 Fledderjohann J, Agrawal S, Vellakkal S, *et al*. Do girls have a nutritional disadvantage compared with boys? statistical models of breastfeeding and food consumption inequalities among Indian siblings. *PLoS One* 2014;9:e107172.
- 52 Aurino E. Do boys eat better than girls in India? Longitudinal evidence on dietary diversity and food consumption disparities among children and adolescents. *Econ Hum Biol* 2017;25:99–111.
- 53 Kshatriya GK, Acharya SK. Gender disparities in the prevalence of undernutrition and the higher risk among the young women of Indian tribes. *PLoS One* 2016;11:e0158308
- 54 Pillai A, Andrews T, Patel V. Violence, psychological distress and the risk of suicidal behaviour in young people in India. *Int J Epidemiol* 2009;38:459–69.
- 55 Balaji M, Andrews T, Andrew G, *et al*. The acceptability, feasibility, and effectiveness of a population-based intervention to promote youth health: an exploratory study in Goa, India. *J Adolesc Health* 2011;48:453–60.
- 56 Jaya J, Hindin MJ. Nonconsensual sexual experiences of adolescents in urban India. *J Adolesc Health* 2007;40:573.e7–573.e14. 573.
- 57 Banerjee SR. Physical abuse of street and slum children of Kolkata. *Indian Pediatr* 2001;38:1163–70.
- 58 SAVE THE CHILDREN. Forgotten voices; the world of urban children in India, 2015. Available: <https://resourcecentre.savethechildren.net/library/forgotten-voices-world-urban-children-india> [Accessed 01 Mar 2020].
- 59 STROCHLIC N. India’s forgotten child brides, 2018. Available: <https://www.nationalgeographic.com/photography/proof/2018/04/child-brides-marriage-shravasti-india-culture/> [Accessed 05 Mar 2020].
- 60 Sathyanarayana Rao TS, Andrade C, Nagpal M. Sexual coercion: time to rise to the challenge. *Indian J Psychiatry* 2013;55:211–3.
- 61 Harbishettar V, Math SB. Violence against women in India: comprehensive care for survivors. *Indian J Med Res* 2014;140:157–9.
- 62 Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- 63 Moher D, Liberati A, Tetzlaff J, *et al*. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 64 Vergnes J-N, Marchal-Sixou C, Nabet C, *et al*. Ethics in systematic reviews. *J Med Ethics* 2010;36:771–4.
- 65 WORLD HEALTH ORGANISATION. Adverse childhood experiences international questionnaire. in adverse childhood experiences international questionnaire (ACE-IQ), 2018. Available: http://www.who.int/violence_injury_prevention/violence/activities/adverse_childhood_experiences [Accessed 01 Feb 2019].
- 66 DAY C, HIBBERT P, CADMAN S. A literature review into children abused and/or neglected prior to custody, 2008. Available: <https://dera.ioe.ac.uk/7997/1/Abused%20prior%20to%20custody.pdf> [Accessed 05 Mar 2020].
- 67 Baldwin JR, Reuben A, Newbury JB, *et al*. Agreement between prospective and retrospective measures of childhood maltreatment: a systematic review and meta-analysis. *JAMA Psychiatry* 2019;76:584–93.
- 68 Teicher MH, Samson JA, Anderson CM, *et al*. The effects of childhood maltreatment on brain structure, function and connectivity. *Nat Rev Neurosci* 2016;17:652–66.
- 69 Colman I, Kingsbury M, Garad Y, *et al*. Consistency in adult reporting of adverse childhood experiences. *Psychol Med* 2016;46:543–9.
- 70 Eichenbaum H. Memory: organization and control. *Annu Rev Psychol* 2017;68:19–45.
- 71 Rasch B, Born J. About sleep’s role in memory. *Physiol Rev* 2013;93:681–766.
- 72 NSPCC. Definitions and signs of child abuse, 2017. Available: <https://learning.nspcc.org.uk/media/1188/definitions-signs-child-abuse.pdf> [Accessed 20 Dec].
- 73 Singhi P, Saini AG, Malhi P. Child maltreatment in India. *Paediatr Int Child Health* 2013;33:292–300.
- 74 SALVADOR-OLIVÁN JA, MARCO-CUENCA G, ARQUERO-AVILÉS R. *Errors in search strategies used in systematic reviews and their effects on information retrieval*. JMLA: Journal of the Medical Library Association, 2019: 107. 210–21.
- 75 Beattie TS, Prakash R, Mazzuca A, *et al*. Prevalence and correlates of psychological distress among 13–14 year old adolescent girls in North Karnataka, South India: a cross-sectional study. *BMC Public Health* 2019;19:48.
- 76 HIGGINS JPT, GREEN S, eds. *Cochrane Handbook for Systematic Reviews of Interventions, Version 5.1.0*. London: The Cochrane Collaboration, 2011.
- 77 Uzzani M, Hammady H, Fedorowicz Z, *et al*. Rayyan—a web and mobile APP for systematic reviews. *Syst Rev* 2016;5:210.
- 78 et al WELLS GA, SHEA B, O’CONNELL. D. The Newcastle-Ottawa scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses, 2014. Available: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp [Accessed 01 Feb 2019].
- 79 Herzog R, Álvarez-Pasquín MJ, Diaz C, *et al*. Are healthcare workers’ intentions to vaccinate related to their knowledge, beliefs and attitudes? A systematic review. *BMC Public Health* 2013;13:154.
- 80 DEEKS JJ, HIGGINS JPT, ALTMAN DG. Chapter 10: Analysing data and undertaking meta-analyses. In: Higgins JPT, Thomas J, Chandler J, *et al*, eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019)*. Cochrane, 2019. www.training.cochrane.org/handbook
- 81 Nyaga VN, Arbyn M, Aerts M. Metaprop: a stata command to perform meta-analysis of binomial data. *Arch Public Health* 2014;72:39.
- 82 Hoysted C, Babi FE, Kassam-Adams N, *et al*. Knowledge and training in paediatric medical traumatic stress and trauma-informed care among emergency medical professionals in low- and middle-income countries. *Eur J Psychotraumatol* 2018;9:1468703
- 83 Seth R, Srivastava RN. Child sexual abuse: management and prevention, and protection of children from sexual offences (POCSO) act. *Indian Pediatr* 2017;54:949–53.
- 84 Paez A. Gray literature: an important resource in systematic reviews. *J Evid Based Med* 2017;10:233–40.
- 85 NOBLIT GW, HARE RD. *Meta-ethnography: synthesizing qualitative studies, vol. 11*. California: Sage Publications, 1988.
- 86 HANNES K, LOCKWOOD, C AND. *Synthesising qualitative research: choosing the right approach*. John Wiley and Sons, 2012.
- 87 Seers K. Qualitative systematic reviews: their importance for our understanding of research relevant to pain. *Br J Pain* 2015;9:36–40.

Appendix 1 - Experts in the field who contributed to the research study design

Member, Designation & Location	Responsibilities
Nilakshi Vaidya, Psychology Counsellor, National Institute of Mental Health and Neurosciences (NIMHANS) (India) – 2nd reviewer	Will be the second reviewer for the systematic review and will review 20% of the article titles, abstract and full texts for inclusion criteria. Will be the second author on any journal paper published.
Megan Fernandes, Core Surgical Trainee with experience working in pediatric medicine and surgery (UK)-3rd reviewer	Will be the third reviewer for the systematic review and will review 20% of the article titles, abstract and full texts for inclusion criteria (published pre-2012). Will be the third author on any journal paper published.
Phillip De Souza, Core Internal Medicine Trainee, experience working in internal medicine (UK) -4th reviewer	Will be the fourth reviewer for the systematic review and will review 20% of the article titles, abstract and full texts for inclusion criteria (published post-2012). Will be the fourth author on any journal paper published.
Vikas Choudhry, Director, Vice President of Public Health, SRC (Uttar Pradesh)	Regular contact through the systematic review process. Will input on search strategies and will specifically resolve any conflicts between myself and the second reviewer. Will be the final author (senior author) on any journal publication.
Bharath Holla, Clinical Psychiatrist and Associate Professor, NIMHANS (India)	PPI group for input on search words and search strategies
Vivek Benegal, Clinical Psychiatrist and cVEDA Principal Investigator, NIMHANS (India)	Clinical psychiatrist and expert in child trauma and informing trauma-based policies in India. Regular contact through the review process and specific input on the search strategies and key words and MESH terms used. Penultimate senior author on any publication.

Appendix 2a. PubMed search strategy

a) Child sexual abuse:

(((((child sexual abuse) OR child sexual coercion) OR child sexual offender)) OR ((child sexual assault) OR child sexual porn*) OR child sexual molest*) OR (((child sexual crime) OR child sexual aggression) OR child sexual offence) OR child sexual violence) OR child rape)) OR ((child rape) OR child incest)) AND India

b) Child emotional abuse:

((((((((((child worthless) OR child unloved) OR child inadequate) OR child failure) OR child corrupt*) OR child unheard) OR child cruelty) OR child emo* AND abuse) OR child maltreat*) AND India)) AND (((((((((((child denial*) OR child denial*) OR child denial shelter*) OR child carer failure) OR child parental failure) OR child emo*) OR child emo* AND danger) OR child neglect) OR child abandonment*) OR child bereavment*) OR child grief) AND India[all])

c) Child physical abuse:

((((((((((child hitting) OR child shaking) OR child throwing) OR child poisoning) OR child burning) OR child scalding) OR child drowning) OR child suffoc*) OR child beating) OR child physical violence) OR child past*) OR child rap) AND India

Child neglect:

d) (((((child neglect) OR child psycho* AND assault) OR child denial) OR child emo* AND assault) OR child malnu*) AND India

Appendix 2b EMBASE search strategy

- a) Child sexual abuse
 1. Prevalence/
 2. Limit 1 to human
 3. ("India" or "india").mp.[mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 4. (child sexual abuse or "child sexual coercion" or "child sexual offender" or "child sexual assault" or "child sexual porn*" or "child sexual molest*" or "child sexual crime" or "child sexual aggression" or "child sexual offence") or "child sexual violence" or "child rape" or "child rape" or "child incest").mp. [mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 5. 1 and 2 and 3 and 4
 6. Limit 5 to yr= "2005 to 2020"

- b) child emotional abuse
 1. Prevalence/
 2. Limit 1 to human
 3. ("India" or "india").mp.[mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 4. (child worthless or "child unloved" or "child inadequate" or "child failure" or "child corrupt" or "child unheard" or "child cruelty" or "child emo*" or "child emo abuse" or "child maltreat" or "child denial" or "child denial shelter" or "child carer failure" or "child parental failure" or "child emo*" or "child emo* and danger" or "child neglect" or "child abandonment*" or "child bereavment*" or "child grief")mp.[mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 5. 1 and 2 and 3 and 4
 6. Limit 5 to yr= "2005 to 2020"

- c) Child physical abuse
 1. Prevalence/
 2. Limit 1 to human
 3. ("India" or "india").mp.[mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 4. (child hitting or "child shaking" or "child throwing" or "child poisoning" or "child burning" or "child scalding" or "child drowning" or "child suffoc*" or "child beating" or "child physical violence" or "child past*" or "child rap")mp.[mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 5. 1 and 2 and 3 and 4
 6. Limit 5 to yr= "2005 to 2020"

- d) Child neglect
1. Prevalence/
 2. Limit 1 to human
 3. ("India" or "india").mp.[mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 4. (child neglect or "child psycho*" or "assault" or "child denial" or "child emo*" or "assault*" or "child malnu")mp.[mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword, floating subheading word]
 5. 1 and 2 and 3 and 4
 6. Limit 5 to yr= "2005 to 2020"

Appendix 2c COCHRANE search strategy

1. India or india
2. (child sexual abuse or child sexual coercion or child sexual offender or child sexual assault or child sexual porn or child sexual molest* or child sexual crime or child sexual aggression or child sexual offence or child sexual violence or child rape or child rape or child incest): ti,ab,kw
3. (child worthless or child unloved or child inadequate or child failure or child corrupt or child unheard or child cruelty or child emo* or child emo abuse or child maltreat or child denial or child denial shelter or child carer failure or child parental failure or child emo* or child emo* and danger or child neglect or child abandonment* or child bereavment* or child grief): ti,ab,kw
4. (child hitting or child shaking or child throwing or child poisoning or child burning or child scalding or child drowning or child suffoc* or child beating or child physical violence or child past* or child rap): ti,ab,kw
5. (child neglect or child psycho* or assault or child denial or child emo* or assault* or child malnu*): ti,ab,kw
6. {#2 or #3 or #4 or #5}
7. #1 and #6

Appendix 2d PsychInfo search strategy

1. Prevalence or epidemiology or frequency.tw.
2. child sexual abuse or child sexual coercion or child sexual offender or child sexual assault or child sexual porn or child sexual molest* or child sexual crime or child sexual aggression or child sexual offence or child sexual violence or child rape or child rape or child incest.tw,sh.
3. child worthless or child unloved or child inadequate or child failure or child corrupt or child unheard or child cruelty or child emo* or child emo abuse or child maltreat or child denial or child denial shelter or child carer failure or child parental failure or child emo* or child emo* and danger or child neglect or child abandonment* or child bereavment* or child grief.tw,sh.
4. child hitting or child shaking or child throwing or child poisoning or child burning or child scalding or child drowning or child suffoc* or child beating or child physical violence or child past* or child rap.tw,sh.
5. child neglect or child psycho* or assault or child denial or child emo* or assault* or child malnu*.tw,sh.
6. India
7. 1 or 2 or 3 or 4 or 5
8. 7 and 6
9. Limit 8 to (English language and yr="2005-2020")

Appendix 3 - Newcastle-Ottawa Scale Quality Assessment

Prevalence of child abuse in Kerala, India: An ICAST-CH based survey

Manoj Therayil Kumar, Nilamadhab Kar, Sebind Kumar

Selection:

1. Representativeness of the sample:
 - a. Truly representative of the average in the target population. * (all subjects or random sampling)
 - b. Somewhat representative of the average in the target group. * (non-random sampling)
 - c. Selected group of users/convenience sample.
 - d. No description of the derivation of the included subjects.

Points: *

2. Sample size:
 - a. Justified and satisfactory (including sample size calculation). *
 - b. Not justified.
 - c. No information provided

Points: -

3. Non-respondents:
 - a. Proportion of target sample recruited attains pre-specified target or basic summary of non-respondent characteristics in sampling frame recorded. *
 - b. Unsatisfactory recruitment rate, no summary data on non-respondents.
 - c. No information provided

Points: *

4. Ascertainment of the exposure (risk factor):
 - a. Vaccine records/vaccine registry/clinic registers/hospital records only. **
 - b. Parental or personal recall and vaccine/hospital records. *
 - c. Parental/personal recall only.

Points: *

Comparability: (Maximum 2 stars)

1. Comparability of subjects in different outcome groups on the basis of design or analysis. Confounding factors controlled.
 - a. Data/ results adjusted for relevant predictors/risk factors/confounders e.g. age, sex, time since vaccination, etc. **
 - b. Data/results not adjusted for all relevant confounders/risk factors/information not provided.

Points: **

Outcome:

1. Assessment of outcome:
 - a. Independent blind assessment using objective validated laboratory methods. **
 - b. Unblinded assessment using objective validated laboratory methods. **
 - c. Used non-standard or non-validated laboratory methods with gold standard. *
 - d. No description/non-standard laboratory methods used.

Points: **

2. Statistical test:
 - a. Statistical test used to analyse the data clearly described, appropriate and measures of association presented including confidence intervals and probability level (ρ value). *
 - b. Statistical test not appropriate, not described or incomplete.

Points: *

Total Points scores: 8-star indicating a good study

Cross-sectional Studies:

Very Good Studies: 9-10 points

Good Studies: 7-8 points

Satisfactory Studies: 5-6 points

Unsatisfactory Studies: 0 to 4 points