



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Efficacy of heat-killed and formalin-killed vaccines against Tilapia tilapinevirus in juvenile Nile tilapia (Oreochromis niloticus)

Citation for published version:

Mai, TT, Kayansamruaj, P, Taengphu, S, Senapin, S, Costa, JZ, Del-Pozo, J, Thompson, KD, Rodkhum, C & Dong, HT 2021, 'Efficacy of heat-killed and formalin-killed vaccines against Tilapia tilapinevirus in juvenile Nile tilapia (Oreochromis niloticus)', *Journal of Fish Diseases*. <https://doi.org/10.1111/jfd.13523>

Digital Object Identifier (DOI):

[10.1111/jfd.13523](https://doi.org/10.1111/jfd.13523)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Fish Diseases

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.










Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



RESEARCH ARTICLE

Efficacy of heat-killed and formalin-killed vaccines against *Tilapia tilapinevirus* in juvenile Nile tilapia (*Oreochromis niloticus*)

Thao Thu Mai^{1,2,3}  | Pattanapon Kayansamruaj⁴  | Suwimon Taengphu^{5,6}  |
 Saengchan Senapin^{5,6}  | Janina Z. Costa⁷  | Jorge del-Pozo⁸  |
 Kim D. Thompson⁷  | Channarong Rodkhum^{1,2}  | Ha Thanh Dong^{9,10} 

¹Center of Excellence in Fish Infectious Diseases (CE FID), Department of Veterinary Microbiology, Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand

²The International Graduate Program of Veterinary Science and Technology (VST), Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand

³Division of Aquacultural Biotechnology, Biotechnology Center of Ho Chi Minh City, Ho Chi Minh, Vietnam

⁴Center of Excellence in Aquatic Animal Health Management, Faculty of Fisheries, Kasetsart University, Bangkok, Thailand

⁵Fish Health Platform, Center of Excellence for Shrimp Molecular Biology and Biotechnology (Centex Shrimp), Faculty of Science, Mahidol University, Bangkok, Thailand

⁶National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency (NSTDA), Pathum Thani, Thailand

⁷Aquaculture Research Group, Moredun Research Institute, Edinburgh, UK

⁸Infection and Immunity Division, Roslin Institute, Edinburgh, UK

⁹Faculty of Science and Technology, Suan Sunandha Rajabhat University, Bangkok, Thailand

¹⁰Department of Food, Agriculture and Bioresources, School of Environment, Resources and Development, Asian Institute of Technology, Pathum Thani, Thailand

Correspondence

Ha Thanh Dong, School of Environment, Resources and Development, Asian Institute of Technology, Pathum Thani, Thailand.
 Email: htdong@ait.ac.th

Channarong Rodkhum, Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand.
 Email: Channarong.R@chula.ac.th

Funding information

International Veterinary Vaccinology Network

Abstract

Tilapia tilapinevirus (also known as tilapia lake virus, TiLV) is considered to be a new threat to the global tilapia industry. The objective of this study was to develop simple cell culture-based heat-killed (HKV) and formalin-killed (FKV) vaccines for the prevention of disease caused by TiLV. The fish were immunized with 100 µl of either HKV or FKV by intraperitoneal injection with each vaccine containing 1.8×10^6 TCID₅₀-inactivated virus. A booster vaccination was carried out at 21-day post-vaccination (dpv) using the same protocol. The fish were then challenged with a lethal dose of TiLV at 28 dpv. The expression of five immune genes (*IgM*, *IgD*, *IgT*, *CD4* and *CD8*) in the head kidney and spleen of experimental fish was assessed at 14 and 21 dpv and again after the booster vaccination at 28 dpv. TiLV-specific IgM responses were measured by ELISA at the same time points. The results showed that both vaccines conferred significant protection, with relative percentage survival of 71.3% and 79.6% for HKV and FKV, respectively. Significant up-regulation of *IgM* and *IgT* was observed in the head kidney of fish vaccinated with HKV at 21 dpv, while *IgM*, *IgD* and *CD4* expression increased in the head kidney of fish receiving FKV at the same time point. After booster vaccination, *IgT* and *CD8* transcripts were significantly

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2021 The Authors. *Journal of Fish Diseases* published by John Wiley & Sons Ltd.

increased in the spleen of fish vaccinated with the HKV, but not with FKV. Both vaccines induced a specific IgM response in both serum and mucus. In summary, this study showed that both HKV and FKV are promising injectable vaccines for the prevention of disease caused by TiLV in Nile tilapia.

KEYWORDS

immune responses, inactivated vaccine, Nile tilapia, protection, tilapia lake virus

1 | INTRODUCTION

Tilapia (*Oreochromis* sp.) is the second most farmed fish species worldwide after carps, reaching 6 million tonnes in 2020 (Fletcher, 2020), equivalent to a value of US\$ 7.9 billion (IMARC, 2020). As the demand for animal protein as a food source increases, tilapia has been considered an important freshwater fish for low- and middle-income countries (LMICs) due to its inexpensive price, high adaptability to various environmental conditions and ease to culture (Prabu et al., 2019). Intensification of tilapia farming systems has occurred as a result of this growing demand. This has led to an increased risk of emerging infectious diseases caused by bacteria, viruses, parasites and fungi (Dong et al., 2015; Mesalhy, 2013). In 2013, a new disease with a suspected viral aetiology emerged in Ecuador (Ferguson et al., 2014) and was named syncytial hepatitis of tilapia (SHT) based on its characteristic histopathological features. Around the same time, a novel RNA virus causing mass mortalities in tilapia was discovered in Israel, termed tilapia lake virus (TiLV) (Eyngor et al., 2014). Subsequent studies, supported with molecular analysis, revealed that the disease episodes in Ecuador and Israel shared the same causative virus, TiLV (Bacharach, Mishra, Briese, Zody, et al., 2016; Del-Pozo et al., 2017). The virus has recently been classified as *Tilapia tilapinevirus*, in the *Tilapinevirus* genus, within the *Amnoonviridae* family (Bacharach, Mishra, Briese, Eldar, et al., 2016). Currently, TiLV has been reported in 16 countries/region worldwide (Jassen et al., 2019; Surachetpong et al., 2020).

Current knowledge indicates that TiLV can infect all stages of fish development, including fertilized eggs, larvae, fry, fingerlings, juveniles and large-size fish (Dong et al., 2017a; Senapin et al., 2018) although fingerlings and juveniles appear to be more vulnerable to infection with the virus (Amal et al., 2018; Dong et al., 2017b; Ferguson et al., 2014; Surachetpong et al., 2017b). Cumulative mortalities of up to 80% have been reported for farmed tilapia in Israel, while in a report from Ecuador the percentage of mortalities appeared to fluctuate from 10%–20% up to 80% depending on the fish strain when tilapia fish were transferred to grow-out cages, with fish dying within 4–7 days of transfer (Eyngor et al., 2014; Ferguson et al., 2014). The mortality levels caused by TiLV infection in Thailand were also variable, ranging from 20% to 90% (Dong et al., 2017b), and experimental infections also tended to result in high levels of mortality (66%–100%) (Behera et al., 2017; Dinh-Hung et al., 2021; Eyngor et al., 2014; Tattiyapong et al., 2017). All of these reports suggest that TiLV is highly virulent and will cause significant mortality loss if introduced to a production site.

Vaccines are an effective way to prevent disease caused by either bacteria or viruses in farmed fish (Evensen, 2016). Currently, the majority of licensed vaccines in aquaculture are inactivated vaccines, which contain either single or combined killed pathogens (Kayansamruaj et al., 2020; Ma et al., 2019), inactivated using either physical (e.g. heat, pH and ultraviolet) or chemical (e.g. formalin, β -propiolactone, glutaraldehyde) processes (Delrue et al., 2012; Lelie et al., 1987). Ideally, when a vaccine is administered, the fish's immune response is stimulated to produce antibodies and an immunologic memory against the pathogen (Secombes & Belmonte, 2016), so that the immune system responds more effectively if the fish should encounter the pathogen at a late date. However, to improve the efficacy of the vaccine, a booster dose(s) is often required in order to obtain high antibody titres against the pathogen (Angelidis, 2006; Bøgwald & Dalmo, 2019; Thu Lan et al., 2021). Inactivated vaccines normally stimulate humoral immune responses, involving helper T cells (CD4+ T cells) and antibody-secreting B cells, secreting IgM, IgD or IgT (Smith et al., 2019). The antibodies combat invading pathogens through a variety of mechanisms, including neutralization, phagocytosis, antibody-dependent cellular cytotoxicity and complement-mediated lysis of pathogens or infected cells (Forthal, 2014). Viral vaccines can also activate cell-mediated immunity, involving cytotoxic T cells (also known as CD8+ T cells), the function of which is to destroy virus-infected cells (Secombes & Belmonte, 2016; Smith et al., 2019; Somamoto et al., 2002; Toda et al., 2011).

Many vaccines traditionally formulated from inactivated bacteria or viruses have been licensed and are commercially available for a variety of fish species, mainly salmon, trout and carp (Ma et al., 2019). The few studies that have been reported relating to the development of a vaccine to prevent TiLV infections in tilapia. The first TiLV vaccine was developed in Israel using strains of TiLV that were attenuated by 17–20 subsequent passages in cell culture. The prototype for these vaccines had relative percentage survival (RPS) values of over 50% (Bacharach & Eldar, 2016). More recently, a cell culture-derived vaccine containing virus inactivated with β -propiolactone and adjuvant Montanide IMS 1312 VG, with a virus titre of 10^8 50% tissue culture infectious dose per millilitre (TCID₅₀ ml⁻¹) was developed in China. The vaccine gave a relatively high level of protection, with the RPS value of 85.7%. This vaccine was able to induce specific IgM, as well as up-regulate a variety of immune genes (Zeng, Wang, Hu, et al., 2021). In another study, a DNA vaccine consisting of a pVAX1 DNA vector containing the sequence for TiLV's segment 8, encoding an immunogenic protein VP20, was used for the primary immunization and a recombinant VP20 (rVP20) protein was used as

a booster vaccine given at 3-week post-vaccination (wpv). This vaccine combination resulted in a RPS value of 72.5%, compared with 50% and 52.5%, respectively, for the DNA vaccine or rVP20 alone (Zeng, Wang, Chen, et al., 2021). In the present study, we investigated whether simple cell culture-based vaccines (water-based with no adjuvant), containing either heat-killed or formalin-killed virus, were able to provoke a specific immune response in vaccinated fish and if the vaccines protected them from TiLV infection.

2 | MATERIALS AND METHODS

2.1 | Fish

Juvenile Nile tilapia (*Oreochromis niloticus*) (body weight, 7.3 ± 1.2 g; length, 5.9 ± 1.1 cm) were obtained from a commercial tilapia hatchery with no previous record of TiLV infection. The fish were placed in 100-L containers at a density of 60 fish per tank at around 28°C and fed with a commercial diet daily at 3% of body weight for 15 days before performing the vaccination trial. Prior to the experiment, five fish were randomly selected to screen for the presence of TiLV using a semi-nested PCR (Taengphu et al., 2020) and bacteria using conventional culture method and found to be negative. Water quality parameters including pH, ammonia and nitrite concentration were monitored every 3 days using a standard Aqua test kit (Sera), and water was changed twice per week. The vaccination study was approved by Kasetsart University Institutional Animal Care and Use Committee (ACKU62-FIS-008).

2.2 | Virus preparation

TiLV strain TH-2018-K was isolated from Nile tilapia during a TiLV outbreak in Thailand in 2018 using E11 cell line following the protocol described previously by Eyngor et al. (2014). The virus was cultured in 75-cm² flasks containing confluent E11 cells and 15 ml of L15 medium at 25°C for 5–7 days or until the cytopathic effect (CPE) of around 80% was obtained in the cell monolayer. The culture supernatant containing the virus was centrifuged at 4500 g for 5 min at 4°C (Eppendorf 5810R) and stored at –80°C. The concentration of the virus was determined by calculating the virus titre as 50% tissue culture infective dose per millilitre (TCID₅₀ ml⁻¹) (Reed & Muench, 1938).

2.3 | Vaccine preparation

TiLV TH-2018-K (1.8×10^7 TCID₅₀ ml⁻¹) was used to prepare both HKV and FKV. Viral inactivation was performed at 60°C for 2, 2.5 and 3 h or with formalin (QReC) at a final concentration of 0.002%, 0.004%, 0.006%, 0.008% and 0.01% for 24 h at 25°C. Viral infectivity was then checked on E11 cells. Successful inactivation of the virus was confirmed by the absence of a CPE after 7 days with all inactivation conditions tested (Table S1). Subsequently, inactivation of the virus was performed at 60°C for 2.5 h for HKV, while incubation

of 0.006% formalin at 25°C for 24 h was used for FKV. The inactivated viral solutions were used as vaccine preparations and were not adjuvanted. These were stored at 4°C until used.

2.4 | Immunization, sampling and challenge test

Before immunization, six fish were chosen randomly from the fish population for blood and mucus sampling. The vaccine study comprised of three groups (HKV, FKV and control). Each group consisted of two 100-L replicate tanks with 25 fish each. Prior to vaccination, fish were anaesthetized using clove oil (100 ppm). Fish in the vaccine groups were immunized with either HKV or FKV by intraperitoneal (IP) injection with 100 µl of vaccine using a 28G × 13 mm needle. Booster immunization was carried out at 21 dpv with the same dose of vaccine (Table 2). Fish in the control group were treated the same, except L15 medium was used in place of the virus solution. Three fish from each tank were randomly collected at 14, 21 and 28 dpv for blood, mucus and tissue sampling (six biological replicates per treatment). Before sampling, fish were anaesthetized with clove oil at 100 ppm. Mucus samples were collected from each fish by placing the fish into a plastic bag containing 1 ml phosphate-buffered saline (PBS, 137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄ and 1.8 mM KH₂PO₄) followed by gentle rubbing for 30 s. These were then centrifuged at 4000 g for 10 min. The mucus supernatant samples were collected and stored at –20°C until used. Blood (~200 µl) was withdrawn from caudal vessel using a 25G × 16 mm needle and allowed to clot for 2 h at 4°C. Serum was collected after centrifugation the blood at 4000 g for 10 min (Thermo Scientific) and then stored at –20°C. Tissues (head kidney and spleen) were collected, immediately placed in Trizol solution (Invitrogen) and kept at –20°C until RNA extraction. For the challenge test, a viral stock of TiLV strain TH-2018-K (1.8×10^7 TCID₅₀ ml⁻¹) was diluted two times with sterile distilled water. Each fish was injected IP with 0.1 ml of the diluted TiLV solution (9×10^5 TCID₅₀ fish⁻¹) at 28 dpv, and mortalities were monitored daily for 21 days. Representative dead fish from each group were subjected for TiLV diagnosis using an in-house RT-qPCR (Taengphu et al., 2021).

2.5 | Immune-related gene expression by RT-qPCR

RNA was extracted using Trizol (Invitrogen) following the protocol recommended by the manufacturer. Genomic DNA contamination was removed using DNase I (Ambion) according to the manufacturer's instructions. After DNase I treatment, RNA samples were re-purified using an equal volume of acid phenol:chloroform (5:1, pH 4.7) (Green & Sambrook, 2019) before checking quality and quantity of extracted RNA with Nanodrop ND-1000 Spectrophotometer (Thermo Scientific). DNA contamination in the treated RNA samples was assessed by performing a qPCR cycling with tilapia elongation factor 1α (*EF-1α*) primers using No-RT master mix (absence of reverse transcriptase enzyme provided in iScript™ Reverse Transcription kit, Bio-Rad). The cDNA synthesis (20 µl reactions) was performed using

TABLE 1 Details of primers used for immune-related gene expression in this study

Gene	Oligo sequences	Annealing temperature (°C)	Product size (bp)	Gene functions	References
<i>EF-1α</i>	F-5'-CTACAGCCAGGCTCGTTTCG-3'	56	139	Elongation factor (housekeeping gene)	Velázquez et al. (2018)
	R-5'-CTTGTACTGGTCTCCAGCA-3'				
IgM	F-5'-GGATGACGAGGAAGCAGACT-3'	53	122	Immunoglobulin M (IgM)	Velázquez et al. (2018)
	R-5'-CATCATCCCTTTGCCACTGG-3'				
IgT	F-5'-TGACCAGAAATGGCGAAGTCTG-3'	53	163	Immunoglobulin T (IgT)	Velázquez et al. (2018)
	R-5'-GTTATAGTACATCTTTAGAAATTACC-3'				
IgD	F-5'-AACACCACCCTGTCCCTGAAT-3'	61	127	Immunoglobulin D (IgD)	Wang et al. (2016)
	R-5'-GGGTGAAAACACACATTCCAAC-3'				
CD4	F-5'-GCTCCAGTGTGACGTGAAA-3'	61	106	Receptor on helper T-cell (CD4 ⁺)	XM_025911776.1, designed in this study
	R-5'-TACAGGTTTGAAGTTGAGCTG-3'				
CD8	F-5'-GCTGGTAGCTCTGGCCTTT-3'	49.5	91	Receptor on cytotoxic T-cell (CD8 ⁺)	XM_0054450353.3 ^a , designed in this study
	R-5'-TGTGATGGTGGGCATCTC-3'				

^aHomolog (98% nucleotide sequence identity) of *Oreochromis aureus* CD8 α (XM_031747820.2).

an iScriptTM Reverse Transcription Supermix (Bio-Rad) containing 100 ng RNA and incubated at 25°C for 5 min for priming, followed by 46°C for 20 min for reverse transcription and then 95°C for 1 min for inactivation of the reverse transcriptase. Immune-related gene expression in the head kidney and spleen was analysed using a quantitative real-time PCR, with specific primers as listed in Table 1 and iTaq Universal SYBR Supermix (Bio-Rad). The 10 μ l reaction consisted of 5.0 μ l 2X Supermix, 0.5 μ l forward and reverse primers (10 μ M each), 1.0 μ l cDNA and 3.0 μ l distilled water. The reaction consisted of an initial activation at 95°C for 2 min, followed by 40 amplification cycles of denaturation at 95°C for 30 s, annealing at the optimal temperature of each primer pair (as shown in Table 1) and extension at 72°C for 30 s. Gene expression data for the immune-related genes of vaccinated and control fish were normalized with that of *EF-1 α* gene amplification using the 2^{- $\Delta\Delta$ C_t} method (Livak & Schmittgen, 2001).

2.6 | Measurement of antibody response by ELISA

Polystyrene 96-well ELISA plates were coated with 0.01% poly-L-lysine solution for 1 h. The plates were then rinsed three times with low salt wash buffer (LSWB, 2 mM Tris; 38 mM NaCl; 0.005% Tween 20, pH 7.3) before the addition of 100 μ l of either heat- or formalin-inactivated TiLV (1.8 \times 10⁷ TCID₅₀ ml⁻¹) overnight at 4°C. The plates were washed three times with LSBW, followed by a blocking step with PBS +1% bovine serum albumin (BSA, Sigma) for 2 h at room temperature (around 28°C). Then, 100 μ l mucus (undiluted) or sera (diluted 1:512 in PBS) was added to each well and incubated overnight at 4°C. The following day, the plates were washed five times with high salt wash buffer (HSWB, 2 mM Tris; 50 M NaCl; 0.01% Tween 20, pH 7.7) and incubated with anti-tilapia IgM (Soonthonsrima et al., 2019) diluted at the ratio 1:200 in PBS +1% BSA for 2 h at around 28°C. The plates were then washed five times with HSWB followed by incubation of goat anti-mouse antibody (Merck) conjugated with HRP (diluted 1:3000 in LSBW +1% BSA) for 1 h at around 28°C. The plates were finally washed five times with HSWB before adding 100 μ l of TMB (Merck) to each well. Colour was developed in the dark for 5–10 min before adding 50 μ l of 2 M H₂SO₄ stop solution (Merck). Optical density was read at wavelength 450 nm using the microplate reader (SpectraMax ID3).

2.7 | Statistical analysis

GraphPad Prism 6 was used to generate the graphs. Kaplan–Meier analysis was performed, and the log-rank test was used to compare the survival curves between vaccinated and control groups. The RPS was calculated using the following equation:

$$RPS = 1 - \left[\frac{\text{average \% mortality of vaccinated fish}}{\text{average \% mortality of unvaccinated fish}} \right] \times 100$$

The differences in relative fold change of immune-related gene expression and specific antibody IgM level were compared using

two-way ANOVA followed by the LSD post hoc test. The differences are considered at different levels of significance $p < .05$, $p < .01$, $p < .001$ and $p < .0001$.

3 | RESULTS

3.1 | Efficacy of vaccine

In the challenge experiment, the first mortality occurred at 3-day post-challenge (dpc) in the non-vaccinated group (control) and at 5 and 7 dpc in the HKV and FKV groups, respectively (Figure 1). Mortalities continued until 13–15 dpc. Moribund fish showed gross signs of TiLV infection including abdominal distension, skin erosion, exophthalmos, fin rot, gill pallor and pale liver (Figure S1). The dead fish from each group were tested positive for TiLV by RT-qPCR. The survival rates were $81.3 \pm 0.0\%$ and $86.3 \pm 0.0\%$ for HKV and FKV groups, respectively, compared with $28.13 \pm 30.9\%$ for the control ($p < .0001$). The survival percentage was analysed using Kaplan–Meier curves with the log-rank test (Figure 1). Average RPS values were 71.3% for the HKV and 79.6% for the FKV vaccine (Table 2).

3.2 | Immune-related gene expression

The relative fold changes of five immune genes (*IgM*, *IgT*, *IgD*, *CD4* and *CD8*) were compared with that of the control group (Figure 2). In the head kidney, a non-significant increase in *IgM* mRNA relative to the control was noted at 14 dpv, which was followed by significant

increase relative to the control at 21 dpv for both HKV and FKV groups (Figure 2a, $p < .05$). A similar trend was observed for *IgT* at 14 dpv for both vaccine groups, which was followed by significantly higher expression levels at 21 dpv for the HKV group only (Figure 2b, $p < .05$). Regarding mRNA levels of *IgD*, there was significant up-regulation of *IgD* in the FKV group only at 21 dpv (Figure 2c, $p < .01$). The *CD4* gene was significantly up-regulated at 14 dpv in the HKV only (Figure 2d, $p < .05$) and at 21 dpv in the FKV ($p < .001$). No statistical difference was observed in *CD8* expression between the vaccinated and control groups at the time point examined (Figure 2e).

In the spleen, non-significant, relative up-regulation of *IgM* expression was noted in both HKV and FKV groups compared with the control at 14 dpv (Figure 2f). There was a slight increase in *IgM* mRNA level relative to the control in the HKV group after booster (28 dpv), which were not significant. Also at 28 dpv, *IgT* expression was over 25 times higher in the HKV group ($p < .05$) and almost 20 times higher in the FKV group (Figure 2g). A slight significant increase in *IgD* expression was seen in the HKV group at 14 dpv (Figure 2h, $p < .05$). No significant increase in *CD4* expression was found at any time point (Figure 2i); meanwhile, an approximately ten-fold increase of *CD8* expression was observed at 28 dpv in the HKV group (Figure 2j, $p < .05$).

3.3 | Detection of antibody IgM against TiLV in serum and mucus

Systemic TiLV-specific antibody IgM (anti-TiLV IgM) levels prevaccination (0 dpv) and at 14, 21 and 28 dpv, as indicated by optical




FIGURE 1 Average per cent survival of heat-killed and formaldehyde-killed vaccinated groups (HKV vs. FKV) compared to the non-vaccinated group (Control) during 21-day post-challenge with TiLV (strain TH-2018-K). Statistical analysis of cumulative survival between both vaccinated groups and the control was analysed using Kaplan–Meier curve with log-rank test ($p < .0001$)

density (OD) at 450 nm, were determined by ELISA (Figure 3a). Before immunization, the average OD value of the fish sera was 0.096 ± 0.009 . The OD readings for HKV, FKV and control groups were 0.254 ± 0.053 , 0.363 ± 0.09 and 0.096 ± 0.015 at 14 dpv, respectively. The OD values showed an increase in antibody levels in both groups of vaccinated fish, but were only statistically different in the FKV group ($p < .01$). A slight decrease was seen in OD readings at 3 wpv in both the HKV and FKV groups relative to the control group (0.249 ± 0.049 , 0.317 ± 0.043 and 0.128 ± 0.017 , respectively). One week after the booster vaccination at 28 dpv, the anti-TiLV IgM levels had increased considerably in both the HKV ($p < .001$) and the FKV ($p < .0001$) groups, reaching the highest values obtained between the different sampling points, compared with that of the non-vaccinated group (average OD readings were 0.438 ± 0.127 , 0.483 ± 0.088 and 0.081 ± 0.01 , respectively) (Figure 3a).

A similar pattern was observed with the mucosal anti-TiLV IgM response (Figure 3b). Before vaccination, the average OD value of fish mucus was 0.068 ± 0.003 . At 14 dpv, the TiLV-specific antibody IgM rose in both of the vaccinated groups, HKV and FKV, compared with the non-vaccinated group (0.251 ± 0.104 , 0.404 ± 0.142 and 0.07 ± 0.005 , respectively), but a significant difference was only noted for the FKV group ($p < .01$). At 3 wpv, the antibody levels were not significantly differ between groups, with OD values of 0.159 ± 0.031 (HKV), 0.290 ± 0.064 (FKV) and 0.083 ± 0.007 (control) being recorded. At 4 wpv (after administering the booster vaccination), a considerable increase in anti-TiLV IgM levels was seen in the mucus of the FKV group ($p < .001$) (0.585 ± 0.145), whereas the increase measured in HKV fish (0.235 ± 0.044) was not statically different to that of the control group (0.107 ± 0.018). No significant changes in average OD readings were seen between the non-vaccinated group and preimmunized fish in either sera or mucus (Figure 3a,b).

4 | DISCUSSION

4.1 | Both simple HKV and FKV were effective in protecting tilapia from TiLV infection

Although many different types of vaccines have been developed for aquaculture in recent years, whole-cell inactivated vaccines remain the major type of vaccine licensed for use by the aquaculture industry (Kayansamruaj et al., 2020; Ma et al., 2019). They are safe, relatively simple to produce and are affordable for farmers, especially for species that are intensively cultured, but low in price like tilapia in LMICs. In this study, we prepared two versions of simple water-based inactivated vaccine (HKV and FKV) for TiLV and assessed the ability of both to protective tilapia against the virus. Both HKV and FKV were able to confer relatively high levels of protection (RPS, 71.3% vs. 79.6%) in vaccinated fish. Differences in methods used to inactivate the virus, vaccine formulation, viral strains, antigen concentration, route of vaccine administration and the population of fish can all contribute to the level of protection obtained from

a vaccine (Table 3). Despite this, vaccination is still considered as a promising strategy to protect tilapia from TiLV infection, although the design of the vaccine should be carefully considered to optimize the level of protection obtained. Other inactivated vaccines have shown relatively high levels of protection in fish. For example, other formalin-killed vaccines resulted in RPS values of 79%, 81.9% and 74% for infectious hematopoietic necrosis virus in rainbow trout (*Oncorhynchus mykiss*) (Tang et al., 2016), *Betanodavirus* in European sea bass (*Dicentrarchus labrax*) (Nuñez-Ortiz et al., 2016) and scale drop disease virus (SDDV) in Asian sea bass (*Lates calcarifer*) (de Groof et al., 2015), respectively. In addition, a heat-killed *Aeromonas hydrophila* vaccine gave 84% protection in rainbow trout (Dehghani et al., 2012). Although the efficacy of these and the current vaccines was not tested against heterologous strains of TiLV, the high level of protection elicited against the homologous strain suggests that autogenous inactivated vaccines may be effective as an emergency vaccine to reduce the risk of production losses in affected tilapia farms.

4.2 | Immunization with HKV or FKV activated both branches of the tilapia's specific immune system

Up-regulation in the expression of *IgM*, *IgD* and *IgT* and *CD4* (genes encoding proteins involved in humoral immunity) and *CD8* (cell-mediated immunity) following immunization with HKV and FKV suggests that the vaccines are able to activate both arms of the specific immune response in Nile tilapia. Protection from these vaccines is, therefore, likely to result from a synergistic effect of humoral (B-cell) and cellular immune (T-cell) responses. This is similar to the recent report by Zeng, Wang, Hu, et al. (2021), showing that β -propiolactone-inactivated TiLV vaccines induced up-regulation of *MHC-I* and *MHC-II/CD4*, which belong to different arms of the immune system.

The increase in *CD4* transcripts at 14 and 21 dpv in fish vaccinated with HKV or FKV may reflect activated naïve *CD4+* cells differentiating into helper T-cell subsets, Th1 and Th2. The Th1 cells produce cytokines that stimulate the expression of anti-viral and inflammatory genes, whereas cytokines secreted by Th2 cells stimulate the differentiation of B cells into plasma cells to produce specific antibody (Secombes & Belmonte, 2016; Secombes & Wang, 2012; Smith et al., 2019). On the other hand, *CD8* transcription was only seen to be significantly up-regulated in the spleen of the HKV group after booster vaccination, indicating that the HKV may stimulate *CD8+* cell activation, which then differentiates into cytotoxic T cells. These cells play a crucial role in cell-mediated immunity (Smith et al., 2019; Somamoto et al., 2002).

In addition to assessing the expression of *IgM* transcripts, this study also examined the expression of two additional immunoglobulins *IgD* and *IgT*. Similar patterns of up-regulation were found in head kidney of fish after the primary immunization, suggesting that all three antibodies may be involved in the protective response elicited by the vaccines. Interestingly, significant increases in mRNA *IgT*

TABLE 2 Details of experimental groups and challenge results

Treatment	Administration route	No of fish challenged	Primary vaccination (TCID ₅₀ fish ⁻¹)		Booster vaccination (TCID ₅₀ fish ⁻¹)		Challenge (TCID ₅₀ fish ⁻¹)		Survival rate (%)	RPS (%)	Significant level (compared to control)
			Day 0	Day 21	Day 21	Day 28	Day 28	Day 28			
Control (L15 media)	IP	16 (× 2 rep.)	0	0	0	0	9 × 10 ⁵	9 × 10 ⁵	28.13 ± 30.9	NA	
HKV	IP	16 (× 2 rep.)	1.8 × 10 ⁶	1.8 × 10 ⁶	1.8 × 10 ⁶	1.8 × 10 ⁶	9 × 10 ⁵	9 × 10 ⁵	81.3 ± 0.0	71.3	p < .0001
FKV	IP	15 (× 2 rep.)	1.8 × 10 ⁶	1.8 × 10 ⁶	1.8 × 10 ⁶	1.8 × 10 ⁶	9 × 10 ⁵	9 × 10 ⁵	86.3 ± 0.0	79.6	p < .0001

Note: FKV, formalin-killed vaccine group; IP, intraperitoneal injection; HKV, heat-killed vaccine group; NA, not applicable; rep, replicate.

levels were seen in the head kidney before booster vaccination and in the spleen after the booster vaccination for both the HKV and FKV groups, suggesting that IgT may be strongly associated with the protective response against TiLV. Unfortunately, the function of IgT in tilapia remains poorly understood. Functional localization studies in other fish species have shown that IgT plays an important role against infectious pathogens on mucosal surfaces, such as skin, gills and gut (Salinas et al., 2021; Smith et al., 2019; Zhang et al., 2011). Nevertheless, further studies are required to gain a better understanding on the role of IgT in tilapia's defence system, especially in response to infection.

Although immune genes were significantly up-regulated in the head kidney after primary immunization, this pattern of expression was not observed in the spleen. This suggests that the head kidney, apart from being a primary lymphoid organ, also act as an important secondary lymphoid organ where specific immune responses to the TiLV vaccine occurred. Studies in other fish have shown that the head kidney, containing blast cells, plasma cells and melanomacrophages, is an important site for antigen presentation and antibody production (Kumar et al., 2016; Soulliere & Dixon, 2017). This might be similar in tilapia. However, it was unexpected to find no significant up-regulation of *IgM*, *IgT*, *IgD* and *CD4* in the head kidney at 7 days after the booster vaccination at 28 dpi. It is possible that the increase in gene expression occurred later than 7 days after the booster vaccination or in other secondary lymphoid organs (not assessed in this study). Therefore, future studies should investigate a longer time course for gene expression to better understand the dynamics of immune gene responses after booster vaccination.

4.3 | HKV and FKV induce both systemic and mucosal IgM

In the present study, HKV and FKV were shown to trigger both systemic and mucosal IgM responses, with similar patterns observed between the two vaccines. The increase in systemic and mucosal IgM in teleost is usually derived from the major lymphoid organs, such as head kidney and spleen (Zapata et al., 2006), but also from the mucosa-associated lymphoid organs located in the skin, gills, gut or nasopharynx (not investigated in this study) (Salinas et al., 2021; Smith et al., 2019). In the present study, up-regulation of IgM expression occurred mainly in the head kidney and to a less extent in the spleen, suggesting head kidney to be one of the main organs for IgM production in response to the TiLV vaccines. Although the pathway of IgM secretion in the mucosal compartment (mucus) is unclear, it is possible that mucosal antibodies are produced locally in the mucosa-associated lymphoid organs and/or by the systemic immune system (Esteban & Cerezuela, 2015; Koppang et al., 2015; Salinas et al., 2011, 2021; Salinas & Parra, 2015). In other research using Asian seabass, monovalent and bivalent bacterial vaccines induced both systemic and mucosal IgM (Thu Lan et al., 2021). Similar kinetics have been reported for IgM secretion in the serum of red hybrid tilapia, infected





FIGURE 2 Fold change in gene expressions between non-vaccinated and vaccinated fish at 14-, 21- and 28-day post-vaccination. Data are presented as the mean \pm SE ($n = 6$). Control, non-vaccinated group; HKV, heat-killed vaccine group; FKV, formalin-killed vaccine group. Asterisks show significant levels between groups. * $p < .05$, ** $p < .01$, *** $p < .001$

FIGURE 3 Optical density (OD) at 540 nm for IgM levels against TiLV in fish sera (diluted 1:512) (a) and mucus (undiluted) (b). Data are presented as the mean \pm SE ($n = 6$). Control, non-vaccinated group; HKV, heat-killed vaccine group; FKV, formalin-killed vaccine group. Asterisks indicate significant levels between groups. * $p < .05$, ** $p < .01$, *** $p < .001$, **** $p < .0001$



IP with TiLV (Tattiyapong et al., 2020). The levels of serum IgM increased significantly in Nile tilapia after immunization with β -propiolactone-inactivated virus (Zeng, Wang, Hu, et al., 2021) or with a recombinant vaccine based on segment 8 of TiLV (Zeng, Wang, Chen, et al., 2021). Mucosal IgM was not investigated in these studies, however. The presence of TiLV-specific IgM in the mucus of vaccinated fish suggests that these vaccines may be able to generate a primary immune response in multiple mucosal organs such as skin and gills, which are crucial sites to prevent the initial invasion of pathogenic agents (Esteban & Cerezuela, 2015; Koppang et al., 2015). The IgM levels produced by FKV were always slightly higher than HKV in both serum and mucus at all sampling points analysed, indicating that FKV induces stronger systemic and mucosal IgM responses than HKV. This could be one of the factors explaining for slightly higher level of protection conferred by FKV.

In this study, increased levels of TiLV-specific IgM after booster vaccination in both serum and mucus indicate successful induction of specific immune memory after first immunization. However, low levels of IgM mRNA detected at 28 dpv did not reflect the IgM levels measured by ELISA at this time point. It was likely that the earlier IgM transcripts had already degraded, while its translated products (antibody) remained. B cells are the major component involved in humoral adaptive immunity. They are activated by specific antigen binding to the B-cell receptors on the cell, followed by presentation of processed antigens to naïve CD4-T cells, which then differentiate into helper T cells. With T cells' help, B cells differentiate into plasma cells and memory B cells. Plasma cells are committed to antibody secretion, whereas memory B cells are responsible for the long-lasting protection from subsequent exposure to the same pathogens (Secombes & Belmonte, 2016; Smith et al., 2019).

TABLE 3 Summary of TILV vaccines and their efficacy

Vaccine type	Description	Country of origin	Dose (TCID ₅₀ fish ⁻¹)	Administration route	Adjuvant	% RPS or survival rate	Reference
Live attenuated	17 and 20 passages (P17 & P20) on cell culture	Israel	1.2 × 10 ⁷ (P17) and 8.9 × 10 ⁶ (P20)	IP	No	>50 ^a	Bacharach and Eldar (2016)
DNA vaccine	Segment 8 (VP20)	China	5 µg	IM	No	50 ^b	Zeng, Wang, Chen, et al. (2021)
Recombinant vaccine			20 µg	IM	M402 (China)	52.5 ^b	
DNA +recombinant vaccine			5 µg of DNA vaccine (prime) + 20 µg of recombinant vaccine (booster)	IM	M402 for booster	72.5 ^b	
Inactivated	β-propiolactone	China	10 ⁵ ; 10 ⁶ ; 10 ⁷	IM	Montanide IMS 1312 VG (Seppic)	32.1 ^a –85.7 ^b	Zeng, Wang, Hu, et al. (2021)
	Heat	Thailand	1.8 × 10 ⁶	IM	No	14.3 ^a –42.9 ^a	
	Formalin		1.8 × 10 ⁶	IP	No	71.3 ^a	This study
				IP	No	79.6 ^a	

Note: IM, intramuscular injection; IP, intraperitoneal injection.

^aRPS, Relative percentage survival.

^bSurvival rate.

Although systemic and mucosal IgM levels were assessed in the study, we were unable to measure levels of other antibodies, that is IgD and IgT by ELISA due to a lack of monoclonal antibodies for these immunoglobulin classes in tilapia. Further studies should investigate the cost of the vaccine for commercial production, the persistence of the immune response in vaccinated fish, duration of protection and efficacy testing these vaccines in a commercial setting.

In conclusion, this study reported on the efficacy of two simple TiLV inactivated vaccines without adjuvant (HKV and FKV) in preventing TiLV infection in Nile tilapia. The vaccines activated both branches of adaptive immunity, triggered expression of three immunoglobulin classes and elicited both systemic and mucosal IgM responses. Most importantly, these vaccines showed relatively high levels of protection against TiLV infection and therefore seem very promising for the prevention of disease associated with TiLV.

ACKNOWLEDGEMENTS

This study was financially supported by the GCRF Networks in Vaccines Research and Development, which was co-funded by the MRC and BBSRC and is supported by the International Veterinary Vaccinology Network. Thao Thu Mai acknowledges the PhD scholarship programme of Chulalongkorn University for ASEAN and non-ASEAN countries and the 90th year anniversary of Chulalongkorn University Fund. The authors thank Ms. Putiya Sangpo and Mr. Dinh Hung Nguyen for their technical assistance.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS


H.T.D., P.K. and K.D.T. involved in conceptualization; T.T.M. and S.T. contributed to investigation; T.T.M., P.K. and H.T.D. made formal analysis; S.S., S.T., J.Z.C., J.D.P., K.D.T., T.T.M. and H.T.D. contributed to methodology; H.T.D. and C.R. supervised the study; T.T.M., H.T.D. and S.S. contributed to writing—original draft; C.R., S.S., P.K., J.Z.C., J.D.P. and K.D.T. contributed to review and editing. All authors have read and agreed to the current version of the manuscript.

Data Availability Statement

The data that support the findings of this study are available on request.

ORCID

Thao Thu Mai  <https://orcid.org/0000-0003-1953-1103>

Pattanapon Kayansamruaj  <https://orcid.org/0000-0002-6187-2772>

Suwimon Taengphu  <https://orcid.org/0000-0002-4834-8693>

Saengchan Senapin  <https://orcid.org/0000-0002-5599-4343>

Janina Z. Costa  <https://orcid.org/0000-0002-2243-0146>

Jorge del-Pozo  <https://orcid.org/0000-0002-9582-1228>

Kim D. Thompson  <https://orcid.org/0000-0003-3550-4842>

Channarong Rodkhum  <https://orcid.org/0000-0002-5711-9294>

Ha Thanh Dong  <https://orcid.org/0000-0002-7190-757X>

REFERENCES

- Amal, M., Koh, C. B., Nurliyana, M., Suhaiba, M., Nor-Amalina, Z., Santha, S., Diyana-Nadhirah, K. P., Yusof, M. T., Ina-Salwany, M. Y., & Zamri-Saad, M. (2018). A case of natural co-infection of tilapia lake virus and *Aeromonas veronii* in a Malaysian red hybrid tilapia (*Oreochromis niloticus* × *O. mossambicus*) farm experiencing high mortality. *Aquaculture*, 485, 12–16. <https://doi.org/10.1016/j.aquaculture.2017.11.019>
- Angelidis, P. (2006). Immersion booster vaccination effect on sea bass (*Dicentrarchus labrax* L.) juveniles. *Journal of Animal Physiology and Animal Nutrition*, 90(1–2), 46–49. <https://doi.org/10.1111/j.1439-0396.2005.00572.x>
- Bacharach, E., & Eldar, A. (2016). Tilapia lake virus vaccines. US Patent Application Publication no.US2016/0354458A1. Retrieved from <https://patents.google.com/patent/US20160354458A1/en>
- Bacharach, E., Mishra, N., Briese, T., Eldar, A., Lipkin, W. I., Kuhn, J. H., & Lipkin, W. I. (2016). ICTV Taxonomic Proposal 2016.016a-dM.A.v2.Tilapinevirus. Create the unassigned genus *Tilapinevirus*. Retrieved from: <https://talk.ictvonline.org/ICTV/proposals/2016.016a-dM.A.v2.Tilapinevirus.pdf>
- Bacharach, E., Mishra, N., Briese, T., Zody, M. C., Kembou Tsoufack, J. E., Zamostiano, R., Berkowitz, A., Ng, J., Nitido, A., Corvelo, A., Toussaint, N. C., Abel Nielsen, S. C., Hornig, M., Del Pozo, J., Bloom, T., Ferguson, H., Eldar, A., & Lipkin, W. I. (2016). Characterization of a novel orthomyxo-like virus causing mass die-offs of Tilapia. *mBio*, 7(2), e00431–e516. <https://doi.org/10.1128/mBio.00431-16>
- Behera, B. K., Pradhan, P. K., Swaminathan, T. R., Sood, N., Paria, P., Das, A., Verma, D. K., Kumar, R., Yadav, M. K., Dev, A. K., Parida, P. K., Das, B. K., Lal, K. K., & Jena, J. K. (2017). Emergence of tilapia lake virus associated with mortalities of farmed Nile Tilapia *Oreochromis niloticus* (Linnaeus 1758) in India. *Aquaculture*, 484, 168–174. <https://doi.org/10.1016/j.aquaculture.2017.11.025>
- Børgwald, J., & Dalmo, R. A. (2019). Review on immersion vaccines for fish: An update 2019. *Microorganisms*, 7(12), 627. <https://doi.org/10.3390/microorganisms7120627>
- de Groof, A. D., Guelen, L., Deijs, M., van der Wal, Y., Miyata, M., Ng, K. S., van Grinsven, L., Simmelink, B., Biermann, Y., Grisez, L., van Lent, J., de Ronde, A., Chang, S. F., Schrier, C., & van der Hoek, L. (2015). A novel virus causes scale drop disease in *Lates calcarifer*. *PLoS Pathogens*, 11(8), e1005074. <https://doi.org/10.1371/journal.ppat.1005074>
- Dehghani, S., Akhlaghi, M., & Dehghani, M. (2012). Efficacy of formalin-killed, heat-killed and lipopolysaccharide vaccines against motile aeromonads infection in rainbow trout (*Oncorhynchus mykiss*). *Global Veterinaria*, 9(4), 409–415. <https://doi.org/10.5829/idosi.gv.2012.9.4.6591>
- Del-Pozo, J., Mishra, N., Kabuusu, R., Cheetham, S., Eldar, A., Bacharach, E., & Ferguson, H. W. (2017). Syncytial hepatitis of tilapia (*Oreochromis niloticus* L.) is associated with Orthomyxovirus-like virions in hepatocytes. *Veterinary Pathology*, 54(1), 164–170. <https://doi.org/10.1177/0300985816658100>
- Delrue, I., Verzele, D., Madder, A., & Nauwynck, H. J. (2012). Inactivated virus vaccines from chemistry to prophylaxis: Merits, risks and challenges. *Expert Review of Vaccines*, 11(6), 695–719. <https://doi.org/10.1586/erv.12.38>
- Dinh-Hung, N., Sangpo, P., Kruangkum, T., Kayansamruaj, P., Rungruangkijkrui, T., Senapin, S., & Dong, H. T. (2021). Dissecting the localization of *Tilapia tilapinevirus* in the brain of the experimentally infected Nile tilapia, *Oreochromis niloticus* (L.). *Journal of Fish Diseases*, 1–12. <https://doi.org/10.1111/jfd.13367>
- Dong, H. T., Ataguba, G. A., Khunrae, P., Rattanojpong, T., & Senapin, S. (2017a). Evidence of TiLV infection in tilapia hatcheries from 2012 to 2017 reveals probable global spread of the disease. *Aquaculture*, 479, 579–583. <https://doi.org/10.1016/j.aquaculture.2017.06.035>

- Dong, H. T., Nguyen, V. V., Le, H. D., Sangsuriya, P., Jitrakorn, S., Saksmerprom, V., Senapin, S., & Rodkhum, C. (2015). Naturally concurrent infections of bacterial and viral pathogens in disease outbreaks in cultured Nile tilapia (*Oreochromis niloticus*) farms. *Aquaculture*, 448, 427–435. <https://doi.org/10.1016/j.aquaculture.2015.06.027>
- Dong, H. T., Siriroob, S., Meemetta, W., Santimanawong, W., Gangnonngiw, W., Pirarat, N., Khunrae, P., Rattanarojpong, T., Vanichviriyakit, R., & Senapin, S. (2017b). Emergence of tilapia lake virus in Thailand and an alternative semi-nested RT-PCR for detection. *Aquaculture*, 476, 111–118. <https://doi.org/10.1016/j.aquaculture.2017.04.019>
- Esteban, M. A., & Cerezuela, R. (2015). Fish mucosal immunity: Skin. In B. H. Beck & E. Peatman (Eds.), *Mucosal health in aquaculture* (pp. 67–93). Academic Press. <https://doi.org/10.1016/B978-0-12-417186-2.00004-2>
- Evensen, O. (2016). Development of fish vaccines: Focusing on methods. In A. Adams (Ed.), *Fish vaccines* (pp. 53–74). Springer. https://doi.org/10.1007/978-3-0348-0980-1_3
- Eyngor, M., Zamostiano, R., Kembou Tsofack, J. E., Berkowitz, A., Bercovier, H., Tinman, S., Lev, M., Hurvitz, A., Galeotti, M., Bacharach, E., & Eldar, A. (2014). Identification of a novel RNA virus lethal to tilapia. *Journal of Clinical Microbiology*, 52(12), 4137–4146. <https://doi.org/10.1128/JCM.00827-14>
- Ferguson, H. W., Kabuuu, R., Beltran, S., Reyes, E., Lince, J. A., & del Pozo, J. (2014). Syncytial hepatitis of farmed tilapia, *Oreochromis niloticus* (L.): A case report. *Journal of Fish Diseases*, 37(6), 583–589. <https://doi.org/10.1111/jfd.12142>
- Fletcher, R. (2020). 2020 tilapia production figures revealed. Retrieved from <https://thefishsite.com/articles/2020-tilapia-production-figures-revealed>
- Forthal, D. N. (2014). Functions of antibodies. *Microbiology Spectrum*, 2(4), 1–17. <https://doi.org/10.1128/microbiolspec.AID-0019-2014>
- Green, M. R., & Sambrook, J. (2019). Removing DNA contamination from RNA samples by treatment with RNase-free DNase I. *Cold Spring Harbor Protocols*, 2019(10), 709–710. <https://doi.org/10.1101/pdb.prot101725>
- IMARC (2020). Tilapia market: Global industry trends, share, size, growth, opportunity and forecast 2021–2026. Retrieved from <https://www.imarcgroup.com/tilapia-market>
- Jassen, M. D., Dong, H. T., & Mohan, C. V. (2019). Tilapia lake virus: A threat to the global tilapia industry? *Review in Aquaculture*, 11(3), 725–739. <https://doi.org/10.1111/raq.12254>
- Kayansamruaj, P., Areechon, N., & Unajak, S. (2020). Development of fish vaccine in Southeast Asia: A challenge for the sustainability of SE Asia aquaculture. *Fish and Shellfish Immunology*, 103, 73–87. <https://doi.org/10.1016/j.fsi.2020.04.031>
- Koppang, E. O., Kvellestad, A., & Fischer, U. (2015). Mucosal health in aquaculture fish mucosal immunity: Gill. In B. H. Beck & E. Peatman (Eds.), *Mucosal health in aquaculture* (pp. 93–133). Academic Press. <https://doi.org/10.1016/B978-0-12-417186-2/00005-4>
- Kumar, R., Joy, K. P., & Singh, S. M. (2016). Morpho-histology of head kidney of female catfish *Heteropneustes fossilis*: Seasonal variations in melano-macrophage centers, melanin contents and effects of lipopolysaccharide and dexamethasone on melanins. *Fish Physiology and Biochemistry*, 42, 1287–1306. <https://doi.org/10.1007/s10695-016-0218-2>
- Lelie, P. N., Reesink, H. W., & Lucas, C. J. (1987). Inactivation of 12 viruses by heating steps applied during manufacture of a hepatitis B vaccine. *Journal of Medical Virology*, 23(3), 297–301. <https://doi.org/10.1002/jmv.1890230313>
- Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods*, 25(4), 402–408. <https://doi.org/10.1006/meth.2001.1262>
- Ma, J., Bruce, T. J., Jones, E. M., & Cain, K. D. (2019). A review of fish vaccine development strategies: Conventional methods and modern biotechnological approaches. *Microorganisms*, 7(11), 569. <https://doi.org/10.3390/microorganisms7110569>
- Mesalhy, S. (2013, July). Review of fish diseases in the Egyptian aquaculture. *Cgiar*. Retrieved from <https://cgispace.cgiar.org/bitstream/handle/10568/34870/EgyptAquacultureSectorDiseaseReview.pdf?sequence=1>
- Núñez-Ortiz, N., Pascoli, F., Picchietti, S., Buonocore, F., Bernini, C., Toson, M., Scapigliati, G., & Toffan, A. (2016). A formalin-inactivated immunogen against viral encephalopathy and retinopathy (VER) disease in European sea bass (*Dicentrarchus labrax*): Immunological and protection effects. *Veterinary Research*, 47(89). <https://doi.org/10.1186/s13567-016-0376-3>
- Prabu, E., Rajagopalsamy, C. B. T., Ahilan, B., Jeevagan, I. J. M. A., & Renuhadevi, M. (2019). Tilapia – An excellent candidate species for world aquaculture: A review. *Annual Research & Review in Biology*, 31(3), 1–14. <https://doi.org/10.9734/arrb/2019/v31i330052>
- Reed, L. J., & Muench, H. (1938). The American. *American Journal of Epidemiology*, 27(3), 493–497. <https://doi.org/10.7723/antiochreview.72.3.0546>
- Salinas, I., Fernández-Montero, Á., Ding, Y., & Sunyer, J. O. (2021). Mucosal immunoglobulins of teleost fish: A decade of advances. *Developmental & Comparative Immunology*, 121, 104079. <https://doi.org/10.1016/j.dci.2021.104079>
- Salinas, I., & Parra, D. (2015). Fish mucosal immunity: Intestine. In B. H. Beck & E. Peatman (Eds.), *Mucosal health in aquaculture* (pp. 135–170). Academic Press. <https://doi.org/10.1016/B978-0-12-417186-2.00006-6>
- Salinas, I., Zhang, Y. A., & Sunyer, J. O. (2011). Mucosal immunoglobulins and B cells of Teleost fish Irene. NIH Public Access. *Developmental & Comparative Immunology*, 35(12), 1346–1365. <https://doi.org/10.1016/j.dci.2011.11.009>
- Secombes, C. J., & Belmonte, R. (2016). Overview of the fish adaptive immune system. In A. Adams (Ed.), *Fish vaccines* (pp. 35–52). Springer. https://doi.org/10.1007/978-3-0348-0980-1_2
- Secombes, C. J., & Wang, T. (2012). The innate and adaptive immune system of fish. In B. Austin (Ed.), *Infectious disease in aquaculture: Prevention and control* (pp. 3–68). Woodhead Publishing. <https://doi.org/10.1533/9780857095732.1.3>
- Senapin, S., Shyam, K. U., Meemetta, W., Rattanarojpong, T., & Dong, H. T. (2018). Inapparent infection cases of tilapia lake virus (TiLV) in farmed tilapia. *Aquaculture*, 487, 51–55. <https://doi.org/10.1016/j.aquaculture.2018.01.007>
- Smith, N. C., Rise, M. L., & Christian, S. L. (2019). A Comparison of the innate and adaptive immune systems in cartilaginous fish, ray-finned fish, and lobe-finned Fish. *Frontiers in Immunology*, 10, 2292. <https://doi.org/10.3389/fimmu.2019.02292>
- Somamoto, T., Nakanishi, T., & Okamoto, N. (2002). Role of specific cell-mediated cytotoxicity in protecting fish from viral infections. *Virology*, 297(1), 120–127. <https://doi.org/10.1006/viro.2002.1486>
- Soonthonsrima, T., Wangman, P., Chaivisuthangkura, P., Pengsuk, C., Sithigorgul, P., & Longyant, S. (2019). Generation of mouse monoclonal antibodies specific to tilapia immunoglobulin using fish immunoglobulin/BSA complex for monitoring of the immune response in Nile tilapia (*Oreochromis niloticus*). *Aquaculture Research*, 50(1), 277–283. <https://doi.org/10.1111/are.13894>
- Soulliere, C., & Dixon, B. (2017). Immune system organs of bony fishes. *Reference Module in Life Sciences*. <https://doi.org/10.1016/B978-0-12-809633-8.12179-X>
- Surachetpong, W., Janetanakit, T., Nonhabenjawan, N., Tattiyapong, P., Sirikanjana, K., & Amonsin, A. (2017). Outbreaks of tilapia lake virus infection. *Emerging Infectious Diseases*, 23(6), 1031–1033. <https://doi.org/10.3201/eid2306.161278>

- Surachetpong, W., Roy, S. R. K., & Nicholson, P. (2020). Tilapia lake virus: The story so far. *Journal of Fish Diseases*, 43(10), 1115–1132. <https://doi.org/10.1111/jfd.13237>
- Taengphu, S., Kayansamruaj, P., Kawato, Y., Deboutteville, J. D., Mohan, C. V., Dong, H. T., & Senapin, S. (2021). Concentration and quantification of *Tilapia tilapinevirus* from water using a simple iron flocculation coupled with probe-based RT-qPCR. *bioRxiv*. <https://doi.org/10.1101/2021.08.10.455809>
- Taengphu, S., Sangsuriya, P., Phiwsaiya, K., Debnath, P. P., Delamare-Deboutteville, J., Mohan, C. V., Dong, H. T., & Senapin, S. (2020). Genetic diversity of tilapia lake virus genome segment 1 from 2011 to 2019 and a newly validated semi-nested RT-PCR method. *Aquaculture*, 526, 735423. <https://doi.org/10.1016/j.aquaculture.2020.735423>
- Tang, L., Kang, H., Duan, K., Guo, M., Lian, G., Wu, Y., Li, Y., Gao, S., Jiang, Y., Yin, J., & Liu, M. (2016). Effects of three types of inactivation agents on the antibody response and immune protection of inactivated IHNV vaccine in rainbow trout. *Viral Immunology*, 29(7), 430–435. <https://doi.org/10.1089/vim.2016.0035>
- Tattiyapong, P., Dachavichitlead, W., & Surachetpong, W. (2017). Experimental infection of tilapia lake virus (TiLV) in Nile tilapia (*Oreochromis niloticus*) and red tilapia (*Oreochromis spp.*). *Veterinary Microbiology*, 207, 170–177. <https://doi.org/10.1016/j.vetmic.2017.06.014>
- Tattiyapong, P., Dechavichitlead, W., Waltzek, T. B., & Surachetpong, W. (2020). Tilapia develop protective immunity including a humoral response following exposure to tilapia lake virus. *Fish and Shellfish Immunology*, 106, 666–674. <https://doi.org/10.1016/j.fsi.2020.08.031>
- Thu Lan, N. G., Salin, K. R., Longyant, S., Senapin, S., & Dong, H. T. (2021). Systemic and mucosal antibody response of freshwater cultured Asian seabass (*Lates calcarifer*) to monovalent and bivalent vaccines against *Streptococcus agalactiae* and *Streptococcus iniae*. *Fish and Shellfish Immunology*, 108, 7–13. <https://doi.org/10.1016/j.fsi.2020.11.014>
- Toda, H., Araki, K., Moritomo, T., & Nakanishi, T. (2011). Perforin-dependent cytotoxic mechanism in killing by CD8 positive T cells in ginbuna crucian carp, *Carassius auratus langsdorfii*. *Development and Comparative Immunology*, 35(1), 88–93. <https://doi.org/10.1016/j.dci.2010.08.010>
- Velázquez, J., Acosta, J., Lugo, J. M., Reyes, E., Herrera, F., González, O., Morales, A., Carpio, Y., & Estrada, M. P. (2018). Discovery of immunoglobulin T in Nile tilapia (*Oreochromis niloticus*): A potential molecular marker to understand mucosal immunity in this species. *Developmental and Comparative Immunology*, 88, 124–136. <https://doi.org/10.1016/j.dci.2018.07.013>
- Wang, B., Wang, P., Wu, Z. H., Lu, Y. S., Wang, Z. L., & Jian, J. C. (2016). Molecular cloning and expression analysis of IgD in Nile tilapia (*Oreochromis niloticus*) in response to *Streptococcus agalactiae* stimulus. *International Journal of Molecular Sciences*, 17(3), 348. <https://doi.org/10.3390/ijms17030348>
- Zapata, A., Diez, B., Cejalvo, T., Gutiérrez-De Frías, C., & Cortés, A. (2006). Ontogeny of the immune system of fish. *Fish and Shellfish Immunology*, 20(2), 126–136. <https://doi.org/10.1016/j.fsi.2004.09.005>
- Zeng, W., Wang, Y., Chen, X., Wang, Q., Bergmann, S. M., Yang, Y., Wang, Y., Li, B. O., Lv, Y., Li, H., & Lan, W. (2021). Potency and efficacy of VP20-based vaccine against tilapia lake virus using different prime-boost vaccination regimens in tilapia. *Aquaculture*, 539, 736654. <https://doi.org/10.1016/j.aquaculture.2021.736654>
- Zeng, W., Wang, Y., Hu, H., Wang, Q., Bergmann, S. M., Wang, Y., Li, B. O., Lv, Y., Li, H., Yin, J., & Li, Y. (2021). Cell culture-derived tilapia lake virus-inactivated vaccine containing montanide adjuvant provides high protection against viral challenge for tilapia. *Vaccines*, 9(2), 86. <https://doi.org/10.3390/vaccines9020086>
- Zhang, Y. A., Salinas, I., & Oriol Sunyer, J. (2011). Recent findings on the structure and function of teleost IgT. *Fish and Shellfish Immunology*, 31(5), 627–634. <https://doi.org/10.1016/j.fsi.2011.03.021>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Mai, T. T., Kayansamruaj, P., Taengphu, S., Senapin, S., Costa, J. Z., del-Pozo, J., Thompson, K. D., Rodkhum, C., & Dong, H. T. (2021). Efficacy of heat-killed and formalin-killed vaccines against *Tilapia tilapinevirus* in juvenile Nile tilapia (*Oreochromis niloticus*). *Journal of Fish Diseases*, 00, 1–13. <https://doi.org/10.1111/jfd.13523>