



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A class of spectral bounds for Max k-Cut

Citation for published version:

Anjos, MF & Neto, J 2020, 'A class of spectral bounds for Max k-Cut', *Discrete Applied Mathematics*, vol. 279, pp. 12-24. <https://doi.org/10.1016/j.dam.2019.10.002>

Digital Object Identifier (DOI):

[10.1016/j.dam.2019.10.002](https://doi.org/10.1016/j.dam.2019.10.002)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Discrete Applied Mathematics

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



A Class of Spectral Bounds for MAX k -CUT[☆]

Miguel F. Anjos^{a,b,*}, José Neto^c

^a *School of Mathematics, University of Edinburgh, James Clerk Maxwell Building, Peter Guthrie Tait Road, Edinburgh EH9 3FD, United Kingdom*

^b *GERAD & Polytechnique Montréal, QC, Canada H3C 3A7*

^c *Samovar, CNRS, Télécom SudParis, Institut Polytechnique de Paris, 9 rue Charles Fourier, 91011 Evry, France*

Abstract

In this paper we introduce a new class of bounds for the maximum k -cut problem on undirected edge-weighted simple graphs. The bounds involve eigenvalues of the weighted adjacency matrix together with geometrical parameters. They generalize previous results on the maximum (2-)cut problem and we demonstrate that they can strictly improve over other eigenvalue bounds from the literature. We also report computational results illustrating the potential impact of the new bounds.

Keywords: Max k -cut, Adjacency matrix eigenvalues, Adjacency matrix eigenvectors

1. Introduction

The partitioning of graphs is an important theme of combinatorial optimization that emerges as a natural modeling of many practical problems from very diverse fields such as VLSI design [4], physical statistics [20], or network
5 planning [14]. Basically, it consists in finding a partition of the node set of a graph that maximizes some objective function and possibly satisfies some side

[☆]This work was supported by a public grant as part of the Investissement d'avenir project, reference ANR-11-LABX-0056-LMH, in a joint call with Gaspard Monge Program for optimization, operations research and their interactions with data sciences.

*Corresponding author

Email addresses: anjos@stanfordalumni.org (Miguel F. Anjos),
Jose.Neto@telecom-sudparis.eu (José Neto)

constraints, e.g., with respect to the cardinality of the partition or the number of nodes in its subsets. In this paper we consider the *maximum k -cut problem* denoted by MAX k -CUT, where k denotes a positive integer. The objective is to find a partition of the node set into k subsets so as to maximize the sum of the weights of the edges having their endpoints in different subsets.

Our contribution is a new family of bounds on the optimal objective value of this problem which generalizes previous results for MAX 2-CUT and can improve over other bounds introduced recently for MAX k -CUT, for any integer $k \geq 2$. We also provide computational results that illustrate the potential impact of these new bounds.

The paper is organized as follows. We establish some notation at the end of this section. In Section 2 we proceed to a literature review. The new bounds are introduced in Section 3 and we prove in Section 4 that it is possible to define perturbations of the weighted adjacency matrix such that these bounds dominate (not strictly) a bound stemming from a classical semidefinite relaxation. We then investigate in Section 5 the efficient computation of distances involved in the expression of the new bounds and also present connections with MAX 2-CUT. The computational results are reported in Section 6, and we conclude in Section 7.

We now close the section with some notation. Given a positive integer n , let $[n]$ stand for the set of integers $\{1, 2, \dots, n\}$. Let $G = (V, E)$ be an undirected simple graph having node set $V = [n]$, edge set E , and let $w \in \mathbb{R}^E$ denote a weight function on the edges. The weighted adjacency matrix, denoted by $W \in \mathbb{R}^{n \times n}$, is a symmetric matrix with entries defined by $W_{ij} = w_{ij}$ if $ij \in E$ and $W_{ij} = 0$ otherwise, for all $(i, j) \in V^2$. Let k denote a positive integer. Given any partition (V_1, V_2, \dots, V_k) of V into k subsets V_1, V_2, \dots, V_k (some of which may be empty), the k -cut defined by this partition is the set $\delta(V_1, V_2, \dots, V_k)$ of edges in E having their endpoints in different subsets of the partition, and the weight of the k -cut is the sum of the weights of the edges it contains. The maximum weight of a k -cut in G is denoted by $mc_k(G, W)$.

Given two disjoint node subsets A, B , let $w[A, B]$ denote the sum of the

weights of the edges having one endpoint in A and the other in B : $w[A, B] = \sum_{(i,j) \in A \times B: ij \in E} w_{ij}$. Similarly, let $w[A]$ represent the sum of the weights of the edges with both endpoints in A : $w[A] = \sum_{\substack{(i,j) \in A^2: \\ ij \in E, i < j}} w_{ij}$. Given a real symmetric matrix $M \in \mathbb{R}^{n \times n}$, let $\lambda_1(M) \leq \lambda_2(M) \leq \dots \leq \lambda_n(M)$ denote the eigenvalues of M in increasing order and let $\nu_1(M), \nu_2(M), \dots, \nu_n(M)$ be the corresponding orthonormal eigenvectors. For the particular case when $M = W$, we shall more simply use λ_i (resp. ν_i) instead of $\lambda_i(W)$ (resp. $\nu_i(W)$) for all $i \in [n]$. For any positive integer q , let $\vec{1}_q$ stand for the q -dimensional all-ones vector. Given any vector $x \in \mathbb{R}^n$, $Diag(x)$ stands for the square diagonal matrix of order n , having x for diagonal. The *Laplacian matrix* is $L = Diag(W\vec{1}_n) - W$. The inner scalar product in \mathbb{R}^n is denoted by $\langle \cdot, \cdot \rangle$, and the Euclidean norm by $\| \cdot \|$. Given $\alpha \in \mathbb{R}$ and a matrix X , the notation αX represents the matrix obtained by multiplying all the entries of X by α . Given two matrices X, Y in $\mathbb{R}^{n \times n}$, $X \bullet Y$ stands for the quantity $\sum_{i=1}^n \sum_{j=1}^n X_{ij} Y_{ij}$.

2. Related work

MAX k -CUT is a notorious \mathcal{NP} -hard problem [23]. In particular, there exists no polynomial time approximation scheme for MAX k -CUT for any $k \geq 2$ unless $\mathcal{P} = \mathcal{NP}$ [2, 23]. Also, there can be no polynomial time approximation algorithm with performance ratio $1 - \frac{1}{34k}$, unless $\mathcal{P} = \mathcal{NP}$ [19]. The challenging task of developing methods for solving this problem has generated many works stemming from the communities in discrete mathematics and operations research. The main developed approaches include heuristics [12], approximation [15] and exact algorithms [1]. In this paper, we are interested in different ways of computing bounds for MAX k -CUT using information about the spectrum of W (possibly *perturbed*, as described later). Therefore the literature review that follows is focused on the works which, to the best of our knowledge, are the most relevant ones in this respect. We point out that all the approximation guarantees mentioned hereafter for randomized algorithms only apply to the restricted version of MAX k -CUT with all the edge weights nonnegative.

The simple randomized algorithm, which consists of assigning each vertex uniformly at random to one of the k subsets, has an approximation guarantee of $(1 - \frac{1}{k})$. For the particular case $k = 2$, i.e., the MAX CUT problem, Goemans and Williamson [17] designed a 0.87856-approximation algorithm based on a semidefinite relaxation of the problem. Their work was subsequently extended to $k \geq 2$ by Frieze and Jerrum [15] making use also of a semidefinite relaxation which can be formulated as follows:

$$(k\text{-SDP}) \left\{ \begin{array}{ll} Z_{kSDP}^* = \max \frac{k-1}{k} \sum_{ij \in E} w_{ij} (1 - X_{ij}) & \\ \text{s.t. } X_{ii} = 1, & \forall i, \quad (1) \\ X_{ij} \geq \frac{-1}{k-1}, & \forall i < j, \quad (2) \\ X \succeq 0, X \in \mathbb{R}^{n \times n}, & \quad (3) \end{array} \right.$$

where the constraint $X \succeq 0$ means that the matrix X is symmetric and positive semidefinite. Note that, for $k = 2$, the inequalities (2) can be removed from the formulation since they are implied by the other constraints. Removing them
70 leads to the SDP relaxation of MAX CUT used by Goemans and Williamson [17], and in fact the randomized algorithm proposed by Frieze and Jerrum coincides with the one by Goemans and Williamson for $k = 2$. De Klerk et al. presented another randomized algorithm for MAX k -CUT based on a semidefinite formulation of the Lovász theta function from [13]. They show their algorithm
75 has the same approximation guarantee as Frieze and Jerrum's method [15] for $k \in \{3, \dots, 10\}$; and a consequence of Conjecture 9.1 in [13] is that this also holds for any $k \geq 3$ (if the conjecture is true). We report in Table 1 the approximation guarantee of Frieze & Jerrum's algorithm for MAX k -CUT, denoted by α_k , for some values of k (these results were proved by Goemans and Williamson
80 for $k = 2, 3$ [17, 18], and De Klerk et al. [13] for $k \geq 3$). From the analysis carried out in [17, 18, 13], it follows that the upper bound on $mc_k(G, W)$ given by the relaxation (k -SDP) satisfies $mc_k(G, W) \geq \alpha_k Z_{kSDP}^*$.

The semidefinite relaxation (k -SDP) may be strengthened with different families of linear inequalities that are valid for the k -cut polytope, i.e., the

Table 1: Approximation guarantees of Frieze and Jerrum’s algorithm for MAX k -CUT in the case when all the edge weights are nonnegative [17, 18, 13]

k	2	3	4	5	6	7	8	9	10
α_k	0.87856	0.836008	0.857487	0.876610	0.891543	0.903259	0.912664	0.920367	0.926788

85 convex hull of all the incidence vectors in \mathbb{R}^E of the k -cuts in G (see, e.g., [3, 24] for $k = 2$ and [10, 11, 26] for $k \geq 3$).

Differently from the works mentioned above which rely on a semidefinite relaxation, another line of research [28, 22] derives upper bounds for MAX k -CUT from the spectrum of the Laplacian or the weighted adjacency matrix. Our
90 work investigates further this last line of research that we now present.

For the particular case when $k = 2$, Mohar and Poljak [21] proved the inequality $mc_2(G, W) \leq \frac{n}{4}\lambda_n(L)$. More recently, van Dam and Sotirov [28] generalized this result for MAX k -CUT:

Theorem 2.1. [28]

$$mc_k(G, W) \leq \frac{n(k-1)}{2k}\lambda_n(L). \quad (4)$$

They also provide several graphs for which this bound is tight together with
95 some comparisons with other bounds stemming from semidefinite relaxations. Also recently, Nikiforov [22] introduced an upper bound for the maximum cardinality of a k -cut in G (i.e., the maximum k -cut problem with $w_e = 1$, for all $e \in E$) that is easily extended to the weighted case and can be expressed as follows.

Theorem 2.2. [22]

$$mc_k(G, W) \leq \frac{k-1}{k} \left(w[V] - \frac{\lambda_1 n}{2} \right). \quad (5)$$

100 As noted in [22], the bounds from Theorems 2.1 and 2.2 are equivalent for regular graphs but different in general. For $k = 2$ (MAX CUT), an upper bound on $mc_2(G, W)$ that is at least as good as (5) was introduced in [5], making use of the eigenvalues and eigenvectors of W . Let d_j denote the distance between

the set of vectors in $\{-1, 1\}^n$ and the linear subspace $\text{lin}(\nu_1, \nu_2, \dots, \nu_j)$ that is
 105 generated by the first j eigenvectors of W . Then, the result can be formulated
 as follows.

Theorem 2.3. [5] *The following inequality holds:*

$$mc_2(G, W) \leq \frac{1}{2}w[V] - \frac{1}{4} \left(\lambda_1 n - \sum_{l \in [n-1]} (\lambda_{l+1} - \lambda_l) d_l^2 \right). \quad (6)$$

In the next section we shall introduce a generalization of the bound (6) for
 MAX k -CUT that is less than or equal to (5) for any integer $k \geq 2$. This is
 obtained by combining ideas from the proofs in [22, 5] leading to Theorems
 110 2.2 and 2.3, and by considering particular perturbations of the entries of the
 matrix W . The new bounds share with (6) the drawback that computing all the
 terms involved in their expression is generally \mathcal{NP} -hard. However, we show that
truncated variants (obtained by removing some terms of the last sum appearing
 in their expression), which are still no greater than (5), can be computed in
 115 polynomial time (see Section 5.2).

3. Spectral bounds for Max k -Cut

For our purposes and with no loss of generality, we assume that the graph
 G is complete (setting zero weights on edges not present in G). In order to
 formulate the new bounds, we now introduce quantities to extend the expression
 of the bound (5) (implying a potential improvement, i.e., decrease of its value)
 by involving almost all of the eigenvalues and eigenvectors of the matrix $\widehat{W} =$
 $W + Q$, where $Q \in \mathbb{R}^n \times \mathbb{R}^n$ stands for a symmetric matrix satisfying some
 conditions to be specified later (the zero matrix is a possible choice for Q). Let
 $\widehat{\lambda}_1 \leq \widehat{\lambda}_2 \leq \dots \leq \widehat{\lambda}_n$ stand for the eigenvalues of the matrix $W + Q$ in increasing
 order, and let $\widehat{\nu}_1, \widehat{\nu}_2, \dots, \widehat{\nu}_n$ be the corresponding unit and pairwise orthogonal
 eigenvectors. Given $r \in \mathbb{R} \setminus \{1\}$, let $\widehat{d}_{j,r}$ denote the distance between the set
 of vectors $\{r, 1\}^n$ and the linear subspace $\text{lin}(\widehat{\nu}_1, \widehat{\nu}_2, \dots, \widehat{\nu}_j)$ that is generated by
 the first j eigenvectors of \widehat{W} :

$$\widehat{d}_{j,r} = \min \{ \|z - y\| : z \in \{r, 1\}^n, y \in \text{lin}(\widehat{\nu}_1, \widehat{\nu}_2, \dots, \widehat{\nu}_j) \}. \quad (7)$$

In the definition of $\widehat{d}_{j,r}$, a vector in $\{r, 1\}^n$ may be interpreted as the incidence vector of a node subset (that may itself be interpreted as a subset of a partition), where the nodes in the subset correspond exactly to the entries with value r (as will be the case in the proof establishing the bound). We now formulate our main general result providing an upper bound on $mc_k(G, W)$.

Theorem 3.1. *For any $r \in \mathbb{R} \setminus \{1\}$ and any symmetric matrix $Q \in \mathbb{R}^n \times \mathbb{R}^n$ satisfying $Q_{ij} \leq 0$, for all $i \neq j$, the following inequality holds.*

$$mc_k(G, W) \leq \frac{1}{2(r-1)^2} (A_r - B_r), \quad (8)$$

with A_r and B_r defined as follows,

$$\begin{cases} A_r = (r^2 + k - 1) \left(2w[V] - \widehat{\lambda}_1 n + 2 \sum_{i=1}^n Q_{ii} \right) + 2(2r + k - 2) \sum_{(i,j) \in [n]^2: i < j} Q_{ij}, \\ B_r = k \sum_{l \in [n-1]} \left(\widehat{\lambda}_{l+1} - \widehat{\lambda}_l \right) \left(\widehat{d}_{l,r} \right)^2. \end{cases} \quad (9)$$

Proof. Let (V_1, V_2, \dots, V_k) denote a partition of V corresponding to an optimal solution of MAX k -CUT.

For all $i \in [k]$, let the vector $y^i \in \{r, 1\}^n$ be defined as follows: $y_l^i = r$ if $l \in V_i$ and 1 otherwise. We have:

$$\begin{aligned} \langle y^i, W y^i \rangle &= 2r^2 w[V_i] + 2 \sum_{j \in [k] \setminus \{i\}} w[V_j] + 2r \sum_{j \in [k] \setminus \{i\}} w[V_i, V_j] + \\ &\quad 2 \sum_{(j,l) \in ([k] \setminus \{i\})^2: j < l} w[V_j, V_l]. \end{aligned} \quad (10)$$

Let us now compute the sum of each term occurring in the right-hand-side of (10) over all $i \in [k]$.

$$\begin{aligned} \sum_{i \in [k]} 2r^2 w[V_i] &= 2r^2 (w[V] - mc_k(G, W)), \\ \sum_{i \in [k]} 2 \sum_{j \in [k] \setminus \{i\}} w[V_j] &= 2(k-1) (w[V] - mc_k(G, W)), \\ \sum_{i \in [k]} 2r \sum_{j \in [k] \setminus \{i\}} w[V_i, V_j] &= 4r mc_k(G, W), \\ \sum_{i \in [k]} 2 \sum_{(j,l) \in ([k] \setminus \{i\})^2: j < l} w[V_j, V_l] &= 2(k-2) mc_k(G, W). \end{aligned}$$

Thus, we deduce

$$\sum_{i \in [k]} \langle y^i, W y^i \rangle = 2mc_k(G, W)(-r^2 + 2r - 1) + 2w[V](r^2 + k - 1). \quad (11)$$

Also, we have for all $i \in [k]$,

$$\langle y^i, Qy^i \rangle = r^2 \sum_{j \in V_i} Q_{jj} + 2r^2 \sum_{(j,l) \in V_i^2: j < l} Q_{jl} + 2r \sum_{j \in [n] \setminus V_i} Q_{ij} + 2 \sum_{\substack{(j,l) \in ([n] \setminus V_i)^2: \\ j < l}} Q_{jl}. \quad (12)$$

Now let us consider the summation over all $i \in [k]$ of the right-hand side in
 125 equation (12). Observe that the coefficient of any

- diagonal entry Q_{jj} is $r^2 + k - 1$,
- non-diagonal entry Q_{jl} is

$$\begin{cases} r^2 + k - 1 & \text{if there exists } q \in [k] \text{ such that } \{j, l\} \subseteq V_q, \text{ and} \\ 2r + k - 2 & \text{otherwise.} \end{cases}$$

Let J_1 stand for set of pairs $(j, l) \in [n]^2$ such that $j < l$ and there exists $q \in [k]$ satisfying $\{j, l\} \subseteq V_q$. Similarly, let J_2 stand for set of pairs $(j, l) \in [n]^2$ such that j and l belong to different subsets of the partition (V_1, V_2, \dots, V_k) and $j < l$. Using the observation given earlier and the fact that Q is symmetric, we deduce:

$$\sum_{i \in [k]} \langle y^i, Qy^i \rangle = (r^2 + k - 1) \left(\sum_{j \in [n]} Q_{jj} + 2 \sum_{(j,l) \in J_1} Q_{jl} \right) + 2(2r + k - 2) \sum_{(j,l) \in J_2} Q_{jl}.$$

Then, using the inequality $r^2 + k - 1 \geq 2r + k - 2$ together with the fact that the non-diagonal coefficients of Q are nonpositive, we obtain:

$$\sum_{i \in [k]} \langle y^i, Qy^i \rangle \leq (r^2 + k - 1) \sum_{j \in [n]} Q_{jj} + 2(2r + k - 2) \sum_{(j,l) \in J_1 \cup J_2} Q_{jl}. \quad (13)$$

We now derive a lower bound on $\langle y^i, \widehat{W}y^i \rangle$, where $\widehat{W} = W + Q$, by making use of the spectrum of \widehat{W} . First, we mention some preliminary properties. Since \widehat{W} is a real symmetric matrix, we may assume that the eigenvectors $\widehat{v}_1, \widehat{v}_2, \dots, \widehat{v}_n$ form an orthonormal basis. Considering the expression of y^i in this basis: $y^i = \sum_{l \in [n]} \alpha_l \widehat{v}_l$ with $\alpha \in \mathbb{R}^n$, we have $\|y^i\|^2 = \sum_{l \in [n]} \alpha_l^2 = n + |V_i|(r^2 - 1)$. Using

this equation, we deduce

$$\begin{aligned}
\langle y^i, \widehat{W}y^i \rangle &= \sum_{l \in [n]} \widehat{\lambda}_l \alpha_l^2 \\
&= \widehat{\lambda}_1 (n + |V_i|(r^2 - 1) - \sum_{l=2}^n \alpha_l^2) + \sum_{l=2}^n \widehat{\lambda}_l \alpha_l^2 \\
&= \widehat{\lambda}_1 (n + |V_i|(r^2 - 1)) + \sum_{l=2}^n (\widehat{\lambda}_l - \widehat{\lambda}_1) \alpha_l^2.
\end{aligned}$$

Note that the quantity $\sum_{l=j}^n \alpha_l^2$ can be interpreted as the distance between the vector y^i and the subspace $\text{lin}(\widehat{\nu}_1, \widehat{\nu}_2, \dots, \widehat{\nu}_{j-1})$. From the definition of the distances defined above, we have $(\widehat{d}_{j-1,r})^2 \leq \sum_{l=j}^n \alpha_l^2$, for all $j \in \{2, 3, \dots, n\}$, thus implying $\alpha_j^2 \geq (\widehat{d}_{j-1,r})^2 - \sum_{l=j+1}^n \alpha_l^2$. By iteratively using the latter inequality for $j = 2, \dots, n$ in the expression of $\langle y^i, \widehat{W}y^i \rangle$ above, we deduce

$$\langle y^i, \widehat{W}y^i \rangle \geq \widehat{\lambda}_1 (n + |V_i|(r^2 - 1)) + \sum_{l \in [n-1]} (\widehat{\lambda}_{l+1} - \widehat{\lambda}_l) (\widehat{d}_{l,r})^2.$$

Summing up these inequalities for all $i \in [k]$ we obtain

$$\sum_{i \in [k]} \langle y^i, \widehat{W}y^i \rangle \geq \widehat{\lambda}_1 n (k + r^2 - 1) + k \left(\sum_{l \in [n-1]} (\widehat{\lambda}_{l+1} - \widehat{\lambda}_l) (\widehat{d}_{l,r})^2 \right). \quad (14)$$

Combining (11), (13) and (14), the result follows. \square

Remark Enforcing the value ‘1’ among the two possible values for the components of the vectors used in the definition of the distances (7) is done only for simplicity of the presentation. We are basically interested in the distance between $\text{lin}(\widehat{\nu}_1, \widehat{\nu}_2, \dots, \widehat{\nu}_j)$ and a set of vectors whose components are restricted to take any of two nonzero values. If we denote by \widehat{d}_{j,r_1,r_2} the distance between $\text{lin}(\widehat{\nu}_1, \widehat{\nu}_2, \dots, \widehat{\nu}_j)$ and the set of vectors $\{r_1, r_2\}^n$ with $(r_1, r_2) \in \mathbb{R}^2$ and $0 \neq r_1 \neq r_2$, then $d_{j,r_1,r_2} = |r_1|d_{j,\frac{r_2}{r_1}}$, for all $j \in [n]$, and the results obtained by using such vectors are equivalent to the ones presented.

Note that all the terms occurring in the expression of B_r (see (9) above) are nonnegative, so that even after removing some or all of the terms involved in the sum, the expression (8) obtained still provides an upper bound on $mc_k(G, W)$. This is relevant with respect to complexity aspects (see Section 5.2).

In view of the bound (8) on $mc_k(G, W)$, one may ask for the best choice for the parameter r and the matrix Q . Firstly with respect to the parameter

r and assuming Q is fixed: if we consider the truncated bound obtained from (8) by removing B_r , it is straightforward to check that the ratio $\frac{r^2+k-1}{(r-1)^2}$ is minimized for $r = 1 - k$. Still assuming Q is fixed, if we now consider the whole
 145 expression of the bound (8), computational experiments show that other values of r may lead to strictly better bounds, depending on the instance. However the improvements that we observed in our experiments by considering other values than $1 - k$ for r in (8) tend to be rather small (by comparison with the choice $r = 1 - k$), and $r = 1 - k$ seems to be a fairly robust choice (more on that in
 150 Section 6).

In the statement of Theorem 3.1, the matrix Q may be interpreted as a perturbation of the coefficients of the matrix W . We shall see later how such perturbations may lead to a value of the bound (8) that is less than or equal to the bound stemming from Frieze and Jerrum's semidefinite relaxation (see
 155 Section 4), and we will also observe that the matrix Q may have an important impact on the value of the bound (Section 6).

Taking $Q = 0$ in Theorem 3.1, we obtain the following upper bound with a simpler expression, where the terms $d_{l,r}$ are defined similarly as $\widehat{d}_{l,r}$ above but using the linear subspace $\text{lin}(\nu_1, \nu_2, \dots, \nu_l)$ instead of $\text{lin}(\widehat{\nu}_1, \widehat{\nu}_2, \dots, \widehat{\nu}_l)$.

Corollary 3.2. *For any $r \in \mathbb{R} \setminus \{1\}$, the following inequality holds.*

$$mc_k(G, W) \leq \frac{1}{2(r-1)^2} \left((r^2 + k - 1)(2w[V] - \lambda_1 n) - k \sum_{l \in [n-1]} (\lambda_{l+1} - \lambda_l) d_{l,r}^2 \right). \tag{15}$$

160 Observe that the upper bound (6) is a particular case of (15), obtained by setting $k = 2$ and $r = 1 - k = -1$.

We conclude this section by mentioning that the approach for proving Theorem 3.1 can also be used to obtain lower bounds on the weight of any k -cut and to generalize results from [6, Section 2.1] for $k = 2$. Let $lc_k(G, W)$ denote the minimum weight of a k -cut in G and let $\bar{d}_{j,r}$ denote the distance between the set of vectors $\{r, 1\}^n$ and the subspace $\text{lin}(\nu_j, \nu_{j+1}, \dots, \nu_n)$ that is generated

by the last $n - j + 1$ eigenvectors of W :

$$\bar{d}_{j,r} = \min \{ \|z - y\| : z \in \{r, 1\}^n, y \in \text{lin}(\nu_j, \nu_{j+1}, \dots, \nu_n) \}. \quad (16)$$

Proposition 3.3.

$$lc_k(G, W) \geq \frac{1}{2(r-1)^2} \left((r^2 + k - 1)(2w[V] - \lambda_n n) + k \sum_{l \in [n-1]} (\lambda_{l+1} - \lambda_l) \bar{d}_{l+1,r}^2 \right). \quad (17)$$

Proof. Similar to that of Theorem 3.1 taking $Q = 0$. Alternatively, apply Theorem 3.1 with $Q = 0$ and the weight matrix $-W$ instead of W , which gives an upper bound on $-lc_k(G, W)$. \square

Corollary 3.2 and Proposition 3.3 lead to the definition of the *spectral bound gap*, which is the difference between the upper and lower spectral bounds:

$$sbgc_k = \frac{1}{2(r-1)^2} \left[(r^2 + k - 1) n (\lambda_n - \lambda_1) - k \sum_{l \in [n-1]} (\lambda_{l+1} - \lambda_l) (\bar{d}_{l+1,r}^2 + d_{l,r}^2) \right].$$

165 4. Improving on the bound Z_{kSDP}^*

In this section we show that by making use of a dual optimal solution of (k -SDP) it is possible to define a matrix Q in Theorem 3.1 such that the bound given by (8) is less than or equal to the bound Z_{kSDP}^* .

First note that the upper bound from (k -SDP) can be expressed as follows: $Z_{kSDP}^* = \frac{k-1}{k} (w[V] + Z_1^*)$, where Z_1^* stands for the optimal objective value of the SDP problem

$$(SDP_P) \left\{ \begin{array}{l} Z_1^* = \max \quad (-\frac{1}{2}W) \bullet X \\ \text{s.t.} \quad X_{ii} = 1, \quad \forall i \in [n], \\ X_{ij} - z_{ij} = -\frac{1}{k-1}, \quad \forall i < j, (i, j) \in [n]^2, \\ X \succeq 0, \text{Diag}(z) \succeq 0, \\ X \in \mathbb{R}^{n \times n}, z \in \mathbb{R}^{\binom{n}{2}}. \end{array} \right.$$

The dual problem of (SDP_P) can be expressed as

$$(SDP_D) \begin{cases} Z_2^* = \min & \sum_{i \in [n]} Y_{ii} - \frac{1}{k-1} \sum_{i < j: (i,j) \in [n]^2} Y_{ij} \\ \text{s.t.} & \mathcal{B}(Y) + W \succeq 0, \\ & Y_{ij} \leq 0, \forall i < j, (i,j) \in [n]^2, \\ & Y \in \mathbb{R}^{n \times n}, Y \text{ symmetric,} \end{cases}$$

where $\mathcal{B}(Y)$ has entries

$$\begin{cases} \mathcal{B}(Y)_{ii} = 2Y_{ii}, \forall i \in [n], \\ \mathcal{B}(Y)_{ij} = Y_{ij}, \forall i \neq j, (i,j) \in [n]^2. \end{cases}$$

One can easily check that strong duality holds and thus $Z_1^* = Z_2^*$. Let Y^* denote
170 an optimal solution of (SDP_D) . Then the optimal objective value of $(k\text{-}SDP)$ can be expressed as: $Z_{kSDP}^* = \frac{k-1}{k} \left(w[V] + \sum_{i \in [n]} Y_{ii}^* - \frac{1}{k-1} \sum_{i < j: (i,j) \in [n]^2} Y_{ij}^* \right)$. Also, an observation to be used later is that necessarily the smallest eigenvalue of the matrix $\mathcal{B}(Y^*) + W$ is zero.

We now prove that particular upper bounds given by (8) are always less than or equal to Z_{kSDP}^* . In the statement of Theorem 3.1, let us take $Q = \mathcal{B}(Y^*)$. Then, in the expression of the spectral bound $A_r - B_r$, we have (using $\lambda_1(\mathcal{B}(Y^*) + W) = 0$):

$$A_r = \frac{1}{(r-1)^2} \left((r^2 + k - 1) \left(w[V] + \sum_{i \in [n]} Y_{ii}^* \right) + (2r + k - 1) \sum_{i < j: (i,j) \in [n]^2} Y_{ij}^* \right).$$

And taking for r the value $1 - k$, we obtain: $A_{1-k} = Z_{kSDP}^*$. Since $B_{1-k} \geq 0$,
175 the next result follows.

Corollary 4.1. *In the statement of Theorem 3.1, taking $r = 1 - k$ and $Q = \mathcal{B}(Y^*)$, the spectral bound (8) is less than or equal to Z_{kSDP}^* .*

The last corollary leads to the next result establishing a domination relation between bounds previously introduced for MAX k -CUT.

180 **Proposition 4.2.** *The best spectral bound \widehat{Z} which can be obtained by the general family described in Theorem 3.1 is always no worse than the bounds given by Z_{kSDP}^* , (4) and (5).*

Proof. By Corollary 4.1, we have $\widehat{Z} \leq Z_{kSDP}^*$.

In the statement of Theorem 3.1, considering the matrix $Q = -\text{Diag}(W\vec{1}_n)$,
 185 setting $r = 1 - k$ and removing from (8) the nonnegative term B_r , we obtain the
 bound (4), and thus \widehat{Z} dominates (4). (van Dam and Sotirov also present in [28]
 a strengthened version of (4) obtained by considering a particular perturbation
 of the diagonal entries of the Laplacian matrix, which leads however to a bound
 that is dominated by (i.e. greater than or equal to) Z_{kSDP}^* . One can easily
 190 define a diagonal matrix Q such that their strengthened bound coincides with
 A_{1-k} in (8).)

The fact that \widehat{Z} dominates (5) follows by considering the zero matrix for Q ,
 $r = 1 - k$, and removing B_r in the expression of the bound given by Theorem
 3.1. □

195 In Section 6 we will see on several instances the relevance of the perturbation
 of W as suggested by Proposition 4.1 and the improvement that may be obtained
 over Z_{kSDP}^* by using the whole expression of the bound (8).

5. On the efficient computation of distances

It has been shown [5, Proposition 4.4] that computing the distances $(d_{j,r})_{j=1}^{n-1}$
 200 is \mathcal{NP} -hard in general. In this section, we shall deal with distances that can be
 computed efficiently. We start by considering the instances of MAX k -CUT such
 that the all-ones vector is an eigenvector of W (Section 5.1). This case leads to
 simple expressions of upper bounds for MAX k -CUT and permits us to identify a
 family of instances for which the spectral bound (8) coincides with $mc_k(G, W)$.
 205 Then, we show in Section 5.2 that, for any fixed positive integer $j \leq n - 1$
 and under some additional conditions, the distance $d_{j,r}$ can be computed in
 polynomial time.

5.1. On the case when $\vec{1}_n$ is an eigenvector of W

In this subsection we specialize Corollary 3.2 for the particular case when
 210 $\vec{1}_n$ is an eigenvector of W . In particular, we obtain simple expressions of upper

bounds on $mc_k(G, W)$ that are lower than or equal to the bounds of Theorems 2.1 and 2.2. We start with an auxiliary result on the minimum squared distance between any vector in $\{1, r\}^n$ and the subspace in \mathbb{R}^n that is orthogonal to $\text{lin}(\vec{1}_n)$ and is denoted by $\text{lin}(\vec{1}_n)^\perp$.

Proposition 5.1. *The following equation holds.*

$$\min \left\{ \|y - z\|^2 : y \in \{1, r\}^n, z \in \text{lin}(\vec{1}_n)^\perp \right\} = \begin{cases} n & \text{if } r \geq 1, \\ nr^2 & \text{if } 0 \leq r < 1, \\ \min\left(\frac{(s+r-1)^2}{n}, \frac{s^2}{n}\right) & \text{otherwise,} \end{cases}$$

215 with $n \equiv s \pmod{1-r}$, $0 \leq s < 1-r$, for the case when $r < 0$.

Proof. Let $p \in \{0, 1, \dots, n\}$ and $\hat{y} \in \{r, 1\}^n$ such that \hat{y} has exactly p entries with value r . Let \hat{d}^2 denote the squared distance between \hat{y} and $\text{lin}(\vec{1}_n)^\perp$, that is, the quantity

$$\hat{d}^2 = \frac{\langle \hat{y}, \vec{1}_n \rangle^2}{n} = \frac{(p(r-1) + n)^2}{n},$$

where the first equation follows from the definition of \hat{d} and the normalization of the vector $\vec{1}_n$. The minimum of \hat{d}^2 is obtained for $p = 0$ if $r \geq 1$, for $p = n$ if $0 < r < 1$, and for $p = \lfloor \frac{n}{1-r} \rfloor$ or $p = \lceil \frac{n}{1-r} \rceil$, otherwise. \square

Using Proposition 5.1 together with the fact that $d_{j,r} \geq d_{j+1,r}$, for all $j \in$
220 $[n-1]$, yields the next result.

Corollary 5.2. *If $\vec{1}_n$ is an eigenvector of W associated with the eigenvalue λ_q , then*

$$mc_k(G, W) \leq \frac{1}{2(r-1)^2} \left((r^2 + k - 1)(2w[V] - \lambda_1 n) - \frac{k}{n} \min((s+r-1)^2, s^2) \sum_{l \in [q-1]} (\lambda_{l+1} - \lambda_l) \right) \quad (18)$$

with $r < 0$ and $n \equiv s \pmod{1-r}$, $0 \leq s < 1-r$.

For the case of complete graphs with unit edge weights, taking $r = 1 - k$ in (18) leads to the following simpler expression.

Corollary 5.3. *If G is a complete graph and all the edge weights are equal to one, then*

$$mc_k(G, W) \leq \frac{1}{2k} \left((k-1)n^2 - \min((s-k)^2, s^2) \right), \quad (19)$$

with $n \equiv s \pmod k$, $0 \leq s < k$.

Proof. The eigenvalues of the adjacency matrix of the complete graph K_n are -1 with multiplicity $n-1$ and $n-1$ with multiplicity 1. The vector $\vec{1}_n$ is an eigenvector associated with the eigenvalue $\lambda_n = n-1$. The result follows from (18) with $q = n$ and $r = 1-k$. \square

Corollary 5.3 gives an infinite class of graphs (complete graphs such that $\min((s-k)^2, s^2) > 0$) where Theorem 3.1 strictly improves on Theorem 2.2. The bound (19) has also the feature of coinciding with the optimal objective value of MAX k -CUT for some cases. For completeness, we give the proof of the next result on the number of edges of Turán graphs, i.e., complete k -partite graphs (k integer, $k \geq 2$), whose partition sets differ in cardinality by at most one. Let $T_{n,k}$ stand for the complete k -partite graph on n vertices with partition sizes equal to $\lfloor \frac{n}{k} \rfloor$ or $\lceil \frac{n}{k} \rceil$, and let $e(T_{n,k})$ denote its number of edges.

Proposition 5.4. *The number of edges of a k -partite Turán graph $T_{n,k}$ is*

$$e(T_{n,k}) = \frac{1}{2k} \left((k-1)n^2 + s^2 - sk \right),$$

with $n \equiv s \pmod k$, $0 \leq s < k$.

Proof. Let $n = qk + s$ with q and s nonnegative integers such that $0 \leq s < k$.

The number of edges is then equal to

$$\begin{aligned}
& \binom{2}{n} - s \binom{2}{q+1} - (k-s) \binom{2}{q} \\
&= \frac{n(n-1)}{2} - s \frac{q(q+1)}{2} - (k-s) \frac{q(q-1)}{2} \\
&= \frac{1}{2} (n(n-1) - q(s(q+1) + (k-s)(q-1))) \\
&= \frac{1}{2} (n(n-1) - q(2s + kq - k)) \\
&= \frac{1}{2} (n(n-1) - \frac{n-s}{k} (2s + (n-s) - k)) \\
&= \frac{1}{2k} (kn(n-1) - (n-s)(s+n-k)) \\
&= \frac{1}{2k} ((k-1)n^2 + s^2 - sk).
\end{aligned}$$

□

Let us recall Turán's theorem [27].

240 **Theorem 5.5.** [27] *If G is an n -vertex K_{k+1} -free graph, then it contains at most $e(T_{n,k})$ edges.*

By Theorem 5.5 and Proposition 5.4, the maximum cardinality of a k -cut in the complete graph K_n is $\frac{1}{2k} ((k-1)n^2 + s^2 - sk)$, with $n \equiv s \pmod{k}$, $0 \leq s < k$. Corollary 5.3 leads to next result.

Proposition 5.6. *Let G be a complete graph with all the edge weights equal to one. If*

$$\begin{cases} n \equiv 0 \pmod{k}, \text{ or} \\ n \equiv \frac{k}{2} \pmod{k} \text{ and } k \text{ is even,} \end{cases}$$

245 *then $mc_k(G, W)$ coincides with the bound (19).*

Proposition 5.6 generalizes Proposition 4.4 in [6] (the latter being obtained by setting $k = 2$ and $r = -1$). Note that for $k = 2$, the bound (19) coincides with $mc_k(G, W)$ for all complete graphs with unit weights, whereas this fails for the bounds of Theorems 2.1 and 2.2 for complete graphs having an odd number
250 of vertices. More generally, for any complete graph with unit edge weights such that $n \equiv \frac{k}{2} \pmod{k}$, k positive and even, the bound (19) coincides with the

optimal objective value $mc_k(G, W) = \frac{(k-1)n^2}{2k} - \frac{k}{8}$ and strictly improves over Z_{kSDP}^* , (4) and (5), which are all equal to $\frac{(k-1)n^2}{2k}$ for this family of instances. This is obvious for (4) and (5), since in this case we have $\lambda_n(L) = n$, $\lambda_1 = -1$.
 255 To see that $Z_{kSDP}^* = \frac{(k-1)n^2}{2k}$, note that a feasible solution of (k -SDP) is given by $X = \frac{1}{n-1}(nI_n - J_n)$, $z = 0$, and a feasible solution of (SDP_D) is given by $Y = \frac{1}{2}I_n$, where I_n stands for the identity matrix of order n , and J_n for the all-ones matrix with order n . Since these two solutions have the same objective value $\frac{(k-1)n^2}{2k}$, the result follows. In fact, this family of instances of MAX k -CUT
 260 also illustrates the fact that the gap between (19) and Z_{kSDP}^* (or equivalently (4) or (5) for this family of instances) can be arbitrarily large with increasing values of k .

5.2. Polynomial-time computable distances

Computing all the distances $(\widehat{d}_{j,r})_{j=1}^{n-1}$ involved in the expression of the
 265 bounds (8) is difficult in general. Even for $k = 2$, computing the single distance $d_{n-1,-1}$ is NP-hard in general [5]. However, given a fixed positive integer $p \leq n-1$ and assuming all the eigenvalues and eigenvectors are given and rational, it has been shown that computing the restricted set of distances $(\widehat{d}_{j,-1})_{j=1}^p$ can be done in polynomial time [7]. We now show that this result can be ex-
 270 tended to the computation of the distances $(\widehat{d}_{j,r})_{j=1}^p$ for any $r \in \mathbb{R} \setminus \{1\}$.

Lemma 5.7. *Let $r \in \mathbb{R} \setminus \{1\}$. The computation of the distance $\widehat{d}_{j,r}$ is equivalent to an unconstrained quadratic program of the form $\min_{z \in \{-1,1\}^{n+1}} z^T \overline{Q} z$, where \overline{Q} has rank at most $j+2$ and no positive diagonal entries.*

Proof. The problem of determining the squared distance $(\widehat{d}_{j,r})^2$ can be formulated as follows:

$$(P1) \begin{cases} \min & y^T V V^T y \\ & y \in \{r, 1\}^n, \end{cases}$$

where V stands for the $n \times (n-j)$ matrix having as columns the eigenvectors $\widehat{v}_{j+1}, \widehat{v}_{j+2}, \dots, \widehat{v}_n$. Using the affine transformation

$$z = \frac{2}{1-r} y - \frac{1+r}{1-r} \vec{1}_n,$$

problem (P1) can be reformulated as follows:

$$(P2) \begin{cases} \min & \frac{(1-r)^2}{4} z^T V V^T z + \frac{1-r^2}{2} (\vec{1}_n)^T V V^T z + \frac{(1+r)^2}{4} (\vec{1}_n^T V V^T \vec{1}_n) \\ & z \in \{-1, 1\}^n. \end{cases}$$

Using the fact that for any $z \in \{-1, 1\}^n$ we have :

$$z^T V V^T z = \sum_{j+1}^n (z^T \nu_i)^2 = n - \sum_{i=1}^j (z^T \nu_i)^2,$$

problem (P2) is equivalent to

$$(P3) \begin{cases} \min & z^T Q z + 2b^T z + c \\ & z \in \{-1, 1\}^n, \end{cases}$$

where $Q = -\frac{(1-r)^2}{4} \bar{V}_j \bar{V}_j^T$, \bar{V}_j stands for the $n \times j$ matrix whose columns correspond to the j first eigenvectors $\hat{\nu}_1, \hat{\nu}_2, \dots, \hat{\nu}_j$, $b = \frac{1-r^2}{4} V V^T \vec{1}_n$, and $c = \frac{n}{4}(1-r)^2 + \frac{1}{4}(1+r)^2 (\vec{1}_n^T V V^T \vec{1}_n)$. (P3) can be reformulated as

$$(P4) \begin{cases} \min & \bar{z}^T \bar{Q} \bar{z} \\ & \bar{z} \in \{-1, 1\}^{n+1}, \end{cases}$$

where \bar{Q} is an $(n+1) \times (n+1)$ matrix with rows and columns indexed on $\{0, 1, \dots, n\}$, and entries defined as follows.

$$\bar{Q}_{il} = \begin{cases} 0 & \text{if } i = l = 0, \\ b_{i+l} & \text{if } \{i, l\} \cap \{0\} \neq \emptyset \text{ and } i + l \neq 0, \\ Q_{il} & \text{otherwise.} \end{cases}$$

(Given an optimal solution \bar{z}^* for (P4), an optimal solution for (P3) is given by $z_i^* = \bar{z}_0^* \bar{z}_i^*$, for all $i \in \{1, 2, \dots, n\}$.)

Note that the matrix \bar{Q} has rank at most $j+2$. This can be seen as follows. By definition Q has rank at most j . Adding to Q a row corresponding to vector b , the resulting $(n+1) \times n$ matrix Q' has rank at most $j+1$. Adding to Q' the column vector $(0, b)^T$, we obtain \bar{Q} and its rank is at most $j+2$. \square

The next result directly follows from Lemma 5.7 above and Theorem 2 from [7]. It extends a polynomiality result about the complexity of computing the distance $\hat{d}_{j,-1}$ for a fixed positive integer $j \leq n-1$ (see Corollary 2.7 in [6]).

Theorem 5.8. *Let $r \in \mathbb{R} \setminus \{1\}$. For a fixed positive integer $j \leq n - 1$, assuming the eigenvalues and eigenvectors of the matrix \bar{Q} as mentioned in Proposition 5.7 are given and rational, the distance $\widehat{d}_{j,r}$ can be computed in polynomial time.*

6. Computational experiments

In this section, we report computational results to illustrate the quality of the new bounds in comparison to other spectral bounds and the Frieze & Jerrum bound. We do not concern ourselves with the computational effort to obtain the distances involved in the new bounds, which we compute using a straightforward enumeration procedure. Developing efficient algorithms for determining them and dealing with large instances is a challenging matter for future research work. In the present study, we consider graphs having up to 30 nodes, and with the practical setting described hereafter, the computation of a single bound involving the distances (denoted Sp in what follows) for a fixed value of the parameter r takes about 45 minutes, whereas it is negligible (less than one second) for the other bounds (denoted FJ , vDS and N hereafter).

6.1. Practical setting

All the computational experiments were performed on a laptop using a processor Intel Core i7-2640M CPU @ 2.80GHz x 4, 7.7 Gio RAM. Our implementation is in C, and the SDPs were solved using CSDP [9]. The graphs used in our experiments are as follows, where d stands for a real value in $[0, 1]$:

- $gka1b$, $gka2b$, $gka6a$, $gka7a$: these are four instances of unconstrained binary quadratic programs of the form: $\max_{x \in \{0,1\}^n} x^T Q x$, taken from [16], where Q is a symmetric matrix of order $n \leq 30$. In our experiments, the matrix Q is re-interpreted as the weighted adjacency matrix of a graph of order n , ignoring the diagonal coefficients. The off-diagonal coefficients of Q are integers in $[0, 100]$ for $gka1b$, $gka2b$ and integers in $[-100, 100]$ for $gka6a$, $gka7a$.

- 310 • C_n : the cycle with n nodes.
- K_n : the complete graph with n nodes.
- $P_i(n, d)$: planar graphs of order n , with density parameter $d \in [0, 1]$ (so that the number of edges is about $3(n - 2)d$, $i = 1, 2, \dots, 8$).
- $R_i(n, p)$: random graphs with order n and density parameter d (so that
315 the number of edges is about $\frac{n(n-1)}{2}d$, $i = 1, 2, \dots, 12$).

Except for the four instances from [16], all the instances were generated using *rudu* [25]. (We indicate in Appendix A the input data to generate the instances different from the ones taken from [16].)

In our experiments, we consider both the cases of unit and non-unit edge
320 weights. For the instances from [16], setting the nonzero entries of Q to value one is indicated by the the notation (*unit*) next to the name of the instance. For the case of non-unit edge weights, except for the four instances from [16] (for which we use the original weights), these are uniformly and randomly generated in $[-100, 100]$ using *rudu* [25].

325 The computational results are reported using the following notation:

- FJ : upper bound from (k -SDP).
- vDS : upper bound from [28] (Theorem 2.1).
- N : upper bound from [22] (Theorem 2.2).
- Sp : upper bound from Corollary 3.2 with the value for r that is mentioned
330 in parentheses.
- $FJ + Sp$: upper bound from Theorem 3.1 with the value for r that is mentioned in parentheses and the matrix $Q = \mathcal{B}(Y^*)$ as described in Section 4.
- For Sp and $FJ + Sp$, we also report in additional columns the best upper
335 bound from testing all the values $\{-k + 0.5q: q = 0, 1, \dots, 4\}$ for r and

selecting the value r_{best} that gives the lowest bound (reported in parentheses).

6.2. Computational results

The results for instances of MAX 3-CUT (resp. MAX 4-CUT and MAX 5-
340 CUT) are reported in Table 2 (resp. 4, 6) for the case of unit weights and in
Table 3 (resp. 5, 7) for non-unit weights (see the previous section).

If we first consider the bounds not involving semidefinite programming, i.e.
 vDS , N and Sp , the best results were obtained with Sp (for $k = 3, 4, 5$) with the
exception of two instances from [16] for the case of unit weights. In particular
345 for the case of non-unit weights the gaps are significant between Sp on the one
hand and vDS and N on the other hand. The instance K_{30} with unit weights
for MAX 4-CUT (see Table 4) illustrates our discussion following Proposition
5.6 on cases when Sp coincides with $mc_k(G, W)$ and strictly improves over FJ ,
 vDS and N .

350 Considering now the bounds using (k -SDP) (i.e. FJ and $FJ + Sp$), they
clearly (non strictly) dominate the other bounds. The improvements over FJ
obtained with $FJ + Sp$ seem modest for the case of unit weights but tend to be
more important for the case of non-unit weights.

Finally, concerning the question raised in Section 3 about the best value of
355 the parameter r and its impact on the spectral bound from Theorem 3.1, our
experiments suggest that the best values may be close to $1 - k$, but possibly
different. This is illustrated, for example, by the values of Sp on the planar
instances from Table 3. The results also show that small modifications of this
parameter may lead to substantial improvements of the bound, in particular
360 for the case of non-unit weights and without perturbations of the weighted
adjacency matrix.

Table 2: Computational results on upper bounds for MAX 3-CUT and unit weights

Instance	$ V $	$ E $	FJ	vDS	N	S_p ($r = -2$)	$FJ + S_p$ ($r = -2$)	S_p ($r = r_{best}$)	$FJ + S_p$ ($r = r_{best}$)
<i>gka1b (unit)</i>	20	187	133.33	133.33	138.00	137.25	133.17	137.25 (-2.0)	133.17 (-2.0)
<i>gka2b (unit)</i>	30	429	300.00	300.00	306.00	304.82	300.00	304.82 (-2.0)	300.00 (-2.0)
<i>gka6a (unit)</i>	30	174	156.05	197.81	171.08	162.25	154.81	162.25 (-2.0)	154.81 (-2.0)
<i>gka7a (unit)</i>	30	211	180.69	214.56	195.26	186.87	178.97	186.87 (-2.0)	178.97 (-2.0)
C_{30}	30	30	30.00	40.00	40.00	38.75	30.00	38.17 (-2.5)	30.00 (-2.0)
K_{30}	30	435	300.00	300.00	300.00	300.00	300.00	300.00 (-2.0)	300.00 (-2.0)
$P_1(30, .7)$	30	58	57.00	116.70	72.71	65.50	56.90	65.50 (-2.0)	56.90 (-2.0)
$P_2(30, .7)$	30	58	56.34	136.83	75.05	67.06	56.23	67.06 (-2.0)	56.23 (-2.0)
$P_3(30, .9)$	30	75	70.06	143.74	81.62	76.27	69.73	76.27 (-2.0)	69.73 (-2.0)
$P_4(30, .9)$	30	75	70.29	171.67	82.84	77.03	70.09	77.03 (-2.0)	70.09 (-2.0)
$R_1(30, .25)$	30	109	104.82	138.91	119.87	112.04	104.05	112.04 (-2.0)	104.05 (-2.0)
$R_2(30, .25)$	30	109	103.95	155.33	117.09	110.07	103.16	110.07 (-2.0)	103.16 (-2.0)
$R_3(30, .5)$	30	218	187.87	218.32	199.63	192.62	186.26	192.62 (-2.0)	186.26 (-2.0)
$R_4(30, .5)$	30	218	185.84	222.24	194.68	188.89	184.24	188.89 (-2.0)	184.24 (-2.0)
$R_5(30, .8)$	30	348	270.25	285.63	280.02	275.02	268.99	275.02 (-2.0)	268.99 (-2.0)
$R_6(30, .8)$	30	348	270.26	291.34	281.34	275.60	268.98	275.60 (-2.0)	268.98 (-2.0)

Table 3: Computational results on upper bounds for MAX 3-CUT and non-unit weights

Instance	$ V $	$ E $	FJ	vDS	N	Sp ($r = -2$)	$FJ + Sp$ ($r = -2$)	Sp ($r = r_{best}$)	$FJ + Sp$ ($r = r_{best}$)
<i>gka1b</i>	20	187	7547.21	8451.14	7921.31	7698.05	7502.65	7698.05 (-2.0)	7502.65 (-2.0)
<i>gka2b</i>	30	429	16605.75	18564.99	17148.43	16791.01	16521.28	16791.01 (-2.0)	16521.28 (-2.0)
<i>gka6a</i>	30	174	2454.64	4470.14	3631.95	2972.53	2365.88	2972.53 (-2.0)	2365.88 (-2.0)
<i>gka7a</i>	30	211	2671.19	6728.62	3960.69	3166.80	2545.51	3166.80 (-2.0)	2545.51 (-2.0)
C_{30}	30	30	1122.00	2530.93	1638.77	1496.45	1122.00	1492.34 (-2.5)	1122.00 (-2.0)
K_{30}	30	435	4289.89	7774.93	5314.43	4565.10	4091.73	4565.10 (-2.0)	4091.73 (-2.0)
$P_5(30, .7)$	30	58	1373.12	4079.19	2541.64	2002.70	1354.35	1969.65 (-1.5)	1353.04 (-1.5)
$P_6(30, .7)$	30	58	1103.17	3565.42	2187.73	1665.76	1083.47	1652.84 (-1.5)	1079.88 (-1.5)
$P_7(30, .9)$	30	75	824.25	3470.83	1747.35	1302.61	791.01	1280.31 (-1.5)	791.01 (-2.0)
$P_8(30, .9)$	30	75	1659.94	4287.78	3429.51	2620.02	1645.23	2545.28 (-1.5)	1644.77 (-1.5)
$R_7(30, .25)$	30	109	2316.90	5051.98	3421.89	2883.97	2281.90	2883.97 (-2.0)	2281.90 (-2.0)
$R_8(30, .25)$	30	109	2286.42	5540.40	3726.62	2947.49	2221.64	2947.49 (-2.0)	2221.64 (-2.0)
$R_9(30, .5)$	30	218	2186.70	4476.53	3062.93	2524.49	2047.29	2524.49 (-2.0)	2047.29 (-2.0)
$R_{10}(30, .5)$	30	218	3112.21	6667.96	4360.54	3717.39	3013.75	3717.39 (-2.0)	3013.75 (-2.0)
$R_{11}(30, .8)$	30	348	4312.47	6578.78	5196.61	4629.66	4171.72	4629.66 (-2.0)	4171.72 (-2.0)
$R_{12}(30, .8)$	30	348	4085.90	7113.67	5390.40	4519.50	3931.45	4519.50 (-2.0)	3931.45 (-2.0)

Table 4: Computational results on upper bounds for MAX 4-CUT and unit weights

Instance	$ V $	$ E $	FJ	vDS	N	Sp ($r = -3$)	$FJ + Sp$ ($r = -3$)	Sp ($r = r_{best}$)	$FJ + Sp$ ($r = r_{best}$)
<i>gka1b (unit)</i>	20	187	150.00	150.00	155.25	154.67	150.00	154.67 (-3.0)	150.00 (-3.0)
<i>gka2b (unit)</i>	30	429	337.50	337.50	344.25	343.07	337.00	343.07 (-3.0)	337.00 (-3.0)
<i>gka6a (unit)</i>	30	174	168.90	222.54	192.46	180.36	168.40	180.36 (-3.0)	168.40 (-3.0)
<i>gka7a (unit)</i>	30	211	198.96	241.37	219.67	208.57	197.71	208.57 (-3.0)	197.71 (-3.0)
C_{30}	30	30	30.00	45.00	45.00	40.95	30.00	40.95 (-3.0)	30.00 (-3.0)
K_{30}	30	435	337.50	337.50	337.50	337.00	337.00	337.00 (-3.0)	337.00 (-3.0)
$P_1(30, .7)$	30	58	58.00	131.29	81.80	73.68	58.00	73.23 (-2.5)	58.00 (-3.0)
$P_2(30, .7)$	30	58	58.00	153.94	84.44	76.93	58.00	75.94 (-2.5)	58.00 (-3.0)
$P_3(30, .9)$	30	75	75.00	161.70	91.82	85.49	75.00	85.49 (-3.0)	75.00 (-3.0)
$P_4(30, .9)$	30	75	75.00	193.13	93.19	86.36	75.00	86.13 (-2.5)	75.00 (-3.0)
$R_1(30, .25)$	30	109	109.00	156.28	134.86	124.21	109.00	124.21 (-3.0)	109.00 (-3.0)
$R_2(30, .25)$	30	109	108.98	174.75	131.73	122.35	108.98	122.35 (-3.0)	108.98 (-3.0)
$R_3(30, .5)$	30	218	205.72	245.61	224.58	215.03	204.83	215.03 (-3.0)	204.83 (-3.0)
$R_4(30, .5)$	30	218	204.92	250.02	219.02	211.42	204.26	211.42 (-3.0)	204.26 (-3.0)
$R_5(30, .8)$	30	348	300.77	321.33	315.02	307.25	299.46	307.25 (-3.0)	299.46 (-3.0)
$R_6(30, .8)$	30	348	301.18	327.75	316.51	308.68	299.98	308.68 (-3.0)	299.98 (-3.0)

Table 5: Computational results on upper bounds for MAX 4-CUT and non-unit weights

Instance	$ V $	$ E $	FJ	vDS	N	Sp ($r = -3$)	$FJ + Sp$ ($r = -3$)	Sp ($r = r_{best}$)	$FJ + Sp$ ($r = r_{best}$)
<i>gka1b</i>	20	187	8332.67	9507.53	8911.47	8652.01	8309.91	8652.01 (-3.0)	8309.91 (-3.0)
<i>gka2b</i>	30	429	18409.31	20885.61	19291.98	18763.12	18336.08	18763.12 (-3.0)	18336.08 (-3.0)
<i>gka6a</i>	30	174	2514.45	5028.91	4085.94	3271.60	2419.39	3221.83 (-2.5)	2413.12 (-2.5)
<i>gka7a</i>	30	211	2735.57	7569.69	4455.78	3580.26	2653.06	3515.72 (-2.5)	2618.38 (-2.0)
C_{30}	30	30	1122.00	2847.29	1843.61	1603.17	1122.00	1603.17 (-3.0)	1122.00 (-3.0)
K_{30}	30	435	4435.21	8746.79	5978.73	4956.97	4256.00	4956.97 (-3.0)	4251.25 (-2.5)
$P_5(30, .7)$	30	58	1389.23	4589.09	2859.34	2323.69	1377.79	2246.66 (-2.0)	1373.35 (-2.0)
$P_6(30, .7)$	30	58	1108.82	4011.10	2461.20	1894.45	1097.06	1840.77 (-2.5)	1090.45 (-2.0)
$P_7(30, .9)$	30	75	852.91	3904.68	1965.77	1540.86	830.20	1445.59 (-2.0)	827.43 (-2.0)
$P_8(30, .9)$	30	75	1671.66	4823.75	3858.20	3013.76	1665.97	2921.77 (-2.0)	1662.89 (-2.0)
$R_7(30, .25)$	30	109	2351.97	5683.48	3849.62	3188.21	2332.51	3188.21 (-3.0)	2331.72 (-2.5)
$R_8(30, .25)$	30	109	2330.44	6232.94	4192.45	3396.54	2283.29	3255.73 (-2.5)	2270.30 (-2.0)
$R_9(30, .5)$	30	218	2247.85	5036.09	3445.79	2705.54	2168.96	2705.54 (-3.0)	2140.88 (-2.0)
$R_{10}(30, .5)$	30	218	3203.67	7501.45	4905.60	4106.77	3120.57	4070.72 (-2.5)	3118.76 (-2.5)
$R_{11}(30, .8)$	30	348	4428.04	7401.12	5846.18	5111.59	4314.15	5111.59 (-3.0)	4308.57 (-2.5)
$R_{12}(30, .8)$	30	348	4172.81	8002.88	6064.20	4954.34	4044.38	4948.71 (-2.5)	4016.76 (-2.0)

Table 6: Computational results on upper bounds for MAX 5-CUT and unit weights

Instance	$ V $	$ E $	FJ	vDS	N	Sp ($r = -4$)	$FJ + Sp$ ($r = -4$)	Sp ($r = r_{best}$)	$FJ + Sp$ ($r = r_{best}$)
<i>gka1b (unit)</i>	20	187	160.00	160.00	165.60	165.12	160.00	165.12 (-4.0)	160.00 (-4.0)
<i>gka2b (unit)</i>	30	429	360.00	360.00	367.20	366.48	360.00	366.48 (-4.0)	360.00 (-4.0)
<i>gka6a (unit)</i>	30	174	173.79	237.37	205.29	192.65	173.75	191.94 (-3.5)	173.75 (-4.0)
<i>gka7a (unit)</i>	30	211	207.56	257.47	234.31	223.14	207.00	222.62 (-3.5)	207.00 (-4.0)
C_{30}	30	30	30.00	48.00	48.00	43.28	30.00	43.15 (-3.5)	30.00 (-4.0)
K_{30}	30	435	360.00	360.00	360.00	360.00	360.00	360.00 (-4.0)	360.00 (-4.0)
$P_1(30, .7)$	30	58	58.00	140.04	87.25	79.25	58.00	78.46 (-3.0)	58.00 (-4.0)
$P_2(30, .7)$	30	58	58.00	164.20	90.07	82.57	58.00	82.09 (-3.0)	58.00 (-4.0)
$P_3(30, .9)$	30	75	75.00	172.48	97.94	91.74	75.00	91.26 (-3.5)	75.00 (-4.0)
$P_4(30, .9)$	30	75	75.00	206.00	99.41	92.55	75.00	92.04 (-3.5)	75.00 (-4.0)
$R_1(30, .25)$	30	109	109.00	166.70	143.85	133.11	109.00	132.16 (-3.5)	109.00 (-4.0)
$R_2(30, .25)$	30	109	109.00	186.40	140.51	130.64	109.00	130.09 (-3.5)	109.00 (-4.0)
$R_3(30, .5)$	30	218	213.68	261.98	239.56	228.81	213.36	228.79 (-3.5)	213.36 (-4.0)
$R_4(30, .5)$	30	218	213.69	266.69	233.62	224.64	213.13	224.64 (-4.0)	213.13 (-4.0)
$R_5(30, .8)$	30	348	318.42	342.75	336.02	327.59	317.11	327.59 (-4.0)	317.11 (-4.0)
$R_6(30, .8)$	30	348	318.53	349.60	337.61	328.98	317.53	328.98 (-4.0)	317.53 (-4.0)

7. Conclusion

In this paper we introduced a new class of bounds for the MAX k -CUT problem involving the spectrum of the (possibly perturbed) weighted adjacency matrix. We exhibited a family of instances for which the new bounds are tight. We showed that truncated variants of the bounds can be computed in polynomial time. Computational experiments show that the new bounds compare well with other spectral bounds from the literature. We proved that particular perturbations of the weighted adjacency matrix could be used so that the bound obtained using our results dominates (non strictly) the bound from the Frieze & Jerrum semidefinite relaxation. Future research will look at developing efficient methods for computing the distances involved in the expression of the

Table 7: Computational results on upper bounds for MAX 5-CUT and non-unit weights

Instance	$ V $	$ E $	FJ	vDS	N	Sp ($r = -4$)	$FJ + Sp$ ($r = -4$)	Sp ($r = r_{best}$)	$FJ + Sp$ ($r = r_{best}$)
<i>gka1b</i>	20	187	8708.12	10141.37	9505.57	9215.10	8699.70	9215.10 (-4.0)	8699.70 (-4.0)
<i>gka2b</i>	30	429	19373.01	22277.99	20578.12	19984.58	19314.46	19984.58 (-4.0)	19314.46 (-4.0)
<i>gka6a</i>	30	174	2532.91	5364.17	4358.33	3607.97	2478.77	3408.51 (-3.0)	2452.83 (-3.0)
<i>gka7a</i>	30	211	2755.71	8074.34	4752.83	3840.87	2705.64	3732.70 (-3.0)	2687.73 (-3.0)
C_{30}	30	30	1122.00	3037.11	1966.52	1674.95	1122.00	1664.52 (-3.5)	1122.00 (-4.0)
K_{30}	30	435	4482.41	9329.91	6377.31	5297.62	4361.68	5150.83 (-3.5)	4315.81 (-3.0)
$P_5(30, .7)$	30	58	1395.68	4895.03	3049.96	2535.54	1387.09	2428.06 (-3.0)	1383.64 (-3.0)
$P_6(30, .7)$	30	58	1111.26	4278.51	2625.28	2110.94	1103.19	1957.86 (-3.0)	1099.47 (-3.0)
$P_7(30, .9)$	30	75	864.80	4164.99	2096.82	1673.72	845.99	1598.45 (-3.0)	845.32 (-3.5)
$P_8(30, .9)$	30	75	1674.19	5145.34	4115.42	3352.09	1670.80	3124.17 (-3.0)	1669.49 (-3.0)
$R_7(30, .25)$	30	109	2358.40	6062.38	4106.26	3396.25	2345.38	3330.54 (-3.5)	2342.37 (-3.0)
$R_8(30, .25)$	30	109	2341.76	6648.47	4471.95	3719.89	2317.87	3546.93 (-3.0)	2307.28 (-3.0)
$R_9(30, .5)$	30	218	2262.05	5371.83	3675.51	2961.17	2219.45	2807.62 (-3.0)	2199.66 (-3.0)
$R_{10}(30, .5)$	30	218	3231.45	8001.55	5232.64	4457.79	3176.37	4315.86 (-3.0)	3165.59 (-3.0)
$R_{11}(30, .8)$	30	348	4450.17	7894.53	6235.93	5379.06	4399.91	5363.86 (-3.5)	4376.85 (-3.0)
$R_{12}(30, .8)$	30	348	4200.29	8536.41	6468.48	5316.22	4118.67	5175.52 (-3.5)	4084.17 (-3.0)

new bounds.

References

- 375 [1] Anjos M.F., Ghaddar B., Hupp L., Liers F., Wiegele A.: Solving k-Way Graph Partitioning Problems to Optimality: The Impact of Semidefinite Relaxations and the Bundle Method. In Facets of Combinatorial Optimization, Jünger M., Reinelt G. (Eds) . Springer, Berlin (2013) 355-386
- 380 [2] Arora, S., Lund, C., Motwani, R., Sudan, M., Szegedy, M.: Proof verification and hardness of approximation problems. Proceedings of the 33rd

IEEE Symposium on Foundations of Computer Science, IEEE Computer Society Press (1992) 14-23

- 385 [3] Barahona, F., Mahjoub, A.R.: On the cut polytope. *Mathematical Programming* 36 (1986) 157-173.
- [4] Barahona, F., Grötschel, M., Jünger, M., Reinelt, G.: An application of combinatorial optimization to statistical physics and circuit layout design. *Operations Research* 36:3 (1988) 493-513
- 390 [5] Ben-Ameur, W., Neto, J.: Spectral bounds for the maximum cut problem. *Networks* 52 (2008) 8-13
- [6] Ben-Ameur, W., Neto, J.: Spectral bounds for unconstrained $(-1, 1)$ -quadratic optimization problems. *European Journal of Operational Research* 207 (2010) 15-24
- 395 [7] Ben-Ameur, W., Neto, J.: A polynomial-time recursive algorithm for some unconstrained quadratic optimization problems. *Discrete Applied Mathematics* 159 (2011) 1689-1698
- [8] Ben-Ameur, W., Mahjoub, A.R., Neto, J.: The Maximum Cut Problem. In *Paradigms of Combinatorial Optimization*, V.T. Paschos (Ed.). John Wiley & Sons (2014) 131-172
- 400 [9] Borchers, B.: CSDP, a C Library for Semidefinite Programming. *Optimization Methods and Software* 11:1 (1999) 613-623
- [10] Chopra, S., Rao, M.R.: The partition problem. *Mathematical Programming* 59:1-3 (1993) 87-115
- 405 [11] Chopra, S., Rao, M.R.: Facets of the k -partition polytope. *Discrete Applied Mathematics* 61:1 (1995) 27-48
- [12] Coja-Oghlan, A., Moore, C., Sanwalani, V.: Max k -cut and approximating the chromatic number of random graphs. *Random Structures and Algorithms* 28:3 (2006) 289-322

- [13] de Klerk, E., Pasechnik, D.V., Warners, J.P.: On approximate graph
410 colouring and max- k -cut algorithms based on the θ -function. *Journal of Combinatorial Optimization* 8:3 (2004) 267-294
- [14] Eisenblätter, A.: The semidefinite relaxation of the k -partition polytope is strong. In Cook W.J., Schulz A.S. (Eds) *Integer Programming and Combinatorial Optimization. IPCO 2002. Lecture Notes in Computer Science*,
415 vol 2337. Springer, Berlin (2002) 273-290
- [15] Frieze, A., Jerrum, M.: Improved approximation algorithms for max k -cut and max bisection. *Algorithmica* 18:1 (1997) 67-81
- [16] Glover, F., Kochenberger, G., Alidaee, B.: Adaptive memory tabu search for binary quadratic programs. *Management Science* 44:3 (1998) 336-345
- [17] Goemans, M.X., Williamson, D.P.: Improved approximation algorithms
420 for maximum cut and satisfiability problems using semidefinite programming. *J. ACM* 42:6 (1995) 1115–1145
- [18] Goemans, M.X., Williamson, D.P.: Approximation algorithms for max 3-cut and other problems via semidefinite programming. *Journal of Computer and System Sciences* 68:2 (2004) 442-470
425
- [19] Kann, V., Khanna, S., Lagergren, J., Panconesi, A.: On the hardness of approximating max k -cut and its dual. *Chicago Journal of Theoretical Computer Science* 2 (1997) 1-18
- [20] Liers, F., Jünger, M., Reinelt, G., Rinaldi, G.: Computing exact ground
430 states of hard Ising spin glass problems by branch-and-cut. In *New optimization algorithms in physics*. Wiley (2004) 47-68
- [21] Mohar, B., Poljak, S.: Eigenvalues and the max-cut problem. *Czechoslovak Mathematical Journal* 40:115 (1990) 343-352
- [22] Nikiforov, V.: Max k -cut and the smallest eigenvalue. *Linear Algebra and its Applications* 504 (2016) 462-467
435

- [23] Papadimitriou, C.H., Yannakakis, M.: Optimization, approximation, and complexity classes. *Journal of Computer and System Sciences* 43:3 (1991) 425-440
- [24] Rendl, F., Rinaldi, G., Wiegele, A.: Solving max-cut to optimality by intersecting semidefinite and polyhedral relaxations. *Mathematical Programming* 121 (2008) 307-335
- [25] Rinaldi, G. Rudy.
<https://www-user.tu-chemnitz.de/~helmberg/rudy.tar.gz> (1996)
- [26] Rodrigues De Sousa, V.J., Anjos, M.F., Le Digabel, S.: Computational study of valid inequalities for the maximum k-cut problem. *Annals of Operations Research* 265:1 (2018) 5-27
- [27] Turán, P.: On an extremal problem in graph theory. *Matematikai és Fizikai Lapok* 48 (1941) 436-452
- [28] van Dam, E.R., Sotirov, R.: New bounds for the max- k -cut and chromatic number of a graph. *Linear Algebra and its Applications* 488 (2016) 216-234

Appendix A. Input data to generate the test graphs with “*rudy*”

Instance	Command line
C_{30}	<code>rudy -circuit 30</code>
$C_{30} (W)$	<code>rudy -circuit 30 -random -100 100 4001</code>
K_{30}	<code>rudy -clique 30 -random 1 1 1000</code>
$K_{30} (W)$	<code>rudy -clique 30 -random -100 100 5001</code>
$P_1(30, .7)$	<code>rudy -planar 30 70 1001</code>
$P_2(30, .7)$	<code>rudy -planar 30 70 2001</code>
$P_3(30, .9)$	<code>rudy -planar 30 90 3001</code>
$P_4(30, .9)$	<code>rudy -planar 30 90 4001</code>
$P_5(30, .7)$	<code>rudy -planar 30 70 1001 -random -100 100 1001</code>
$P_6(30, .7)$	<code>rudy -planar 30 70 2001 -random -100 100 2001</code>
$P_7(30, .9)$	<code>rudy -planar 30 90 3001 -random -100 100 3001</code>
$P_8(30, .9)$	<code>rudy -planar 30 90 4001 -random -100 100 4001</code>
$R_1(30, .25)$	<code>rudy -rnd_graph 30 25 1001</code>
$R_2(30, .25)$	<code>rudy -rnd_graph 30 25 2001</code>
$R_3(30, .5)$	<code>rudy -rnd_graph 30 50 3001</code>
$R_4(30, .5)$	<code>rudy -rnd_graph 30 50 4001</code>
$R_5(30, .8)$	<code>rudy -rnd_graph 30 80 5001</code>
$R_6(30, .8)$	<code>rudy -rnd_graph 30 80 6001</code>
$R_7(30, .25)$	<code>rudy -rnd_graph 30 25 1001 -random -100 100 1001</code>
$R_8(30, .25)$	<code>rudy -rnd_graph 30 25 2001 -random -100 100 2001</code>
$R_9(30, .5)$	<code>rudy -rnd_graph 30 50 3001 -random -100 100 3001</code>
$R_{10}(30, .5)$	<code>rudy -rnd_graph 30 50 4001 -random -100 100 4001</code>
$R_{11}(30, .8)$	<code>rudy -rnd_graph 30 80 5001 -random -100 100 5001</code>
$R_{12}(30, .8)$	<code>rudy -rnd_graph 30 80 6001 -random -100 100 6001</code>