

# Edinburgh Research Explorer

# European Stroke Organisation expedited recommendation for the use of short-term dual antiplatelet therapy early after minor stroke and high-risk TIA

# Citation for published version:

Dawson, J, Merwick, Á, Webb, A, Dennis, M, Ferrari, J & Fonseca, AC 2021, 'European Stroke Organisation expedited recommendation for the use of short-term dual antiplatelet therapy early after minor stroke and high-risk TIA', *European Stroke Journal*, vol. 6, no. 2, pp. VI-VI. https://doi.org/10.1177/23969873211027006

### Digital Object Identifier (DOI):

10.1177/23969873211027006

#### Link:

Link to publication record in Edinburgh Research Explorer

#### **Document Version:**

Publisher's PDF, also known as Version of record

## Published In:

European Stroke Journal

#### **General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Édinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



# EUROPEAN STROKE JOURNAL

European Stroke Organisation expedited recommendation for the use of short-term dual antiplatelet therapy early after minor stroke and high-risk TIA

European Stroke Journal 2021, Vol. 6(2) VI © European Stroke Organisation 2021



Article reuse guidelines: sagepub.com/journals-permissions DOI: 10.1177/23969873211027006 journals.sagepub.com/home/eso



Jesse Dawson<sup>1</sup>, Áine Merwick<sup>2</sup>, Alastair Webb<sup>3</sup>, Martin Dennis<sup>4</sup>, Julia Ferrari<sup>5</sup> and Ana Catarina Fonseca<sup>6</sup>; for the European Stroke Organisation

#### **Abstract**

Prevention of early recurrent ischaemic stroke remains a priority in people with TIA or ischaemic stroke. A number of trials have recently been completed assessing the efficacy of short-term dual antiplatelet therapy (DAPT) versus single antiplatelet therapy early after minor or moderate stroke or high-risk TIA. We present an Expedited Recommendation for use of dual antiplatelet therapy early after ischaemic stroke and TIA on behalf of the ESO Guideline Board. We make a strong recommendation based on high quality of evidence for use of 21-days of dual antiplatelet therapy with aspirin and clopidogrel in people with a non-cardioembolic minor ischaemic stroke or high-risk TIA in the past 24 hours. We make a weak recommendation based on moderate quality evidence for 30-days of dual antiplatelet therapy with aspirin and ticagrelor in people with non-cardioembolic mild to moderate ischaemic stroke or high-risk TIA in the past 24 hours.

# Keywords

Stroke, TIA, antiplatelets

Date received: 9 February 2021; accepted: 16 February 2021

#### Corresponding author:

Jesse Dawson. College of Medical Veterinary and Life Sciences, University of Glasgow, University Avenue, Glasgow G12 8QQ, UK. Email: Jesse.dawson@glasgow.ac.uk

<sup>&</sup>lt;sup>1</sup>College of Medical Veterinary and Life Sciences, University of Glasgow, Glasgow, UK

<sup>&</sup>lt;sup>2</sup>Department of Neurology, Cork University Hospital, Cork, Ireland <sup>3</sup>Wolfson Centre for Prevention of Stroke and Dementia, Department of Clinical Neurosciences, John Radcliffe Hospital, University of Oxford, Oxford, UK

 $<sup>^{4}\</sup>mathsf{Centre}$  for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

<sup>&</sup>lt;sup>5</sup>Department of Neurology, St. John's of God Hospital, Vienna Austria <sup>6</sup>Department of Neurosciences (Neurology), Hospital de Santa Maria, University of Lisbon, Lisboa, Portugal