



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

New methods for the synthesis of spirocyclic cephalosporin analogues

Citation for published version:

Zhao, AX, Horsfall, LE & Hulme, AN 2021, 'New methods for the synthesis of spirocyclic cephalosporin analogues', *Molecules*, vol. 26, no. 19, 6035. <https://doi.org/10.3390/molecules26196035>

Digital Object Identifier (DOI):

[10.3390/molecules26196035](https://doi.org/10.3390/molecules26196035)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Molecules

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Communication

New Methods for the Synthesis of Spirocyclic Cephalosporin Analogues

Alan X. Zhao ¹, Louise E. Horsfall ²  and Alison N. Hulme ^{1,*} 

¹ EaStChem School of Chemistry, The University of Edinburgh, Joseph Black Building, David Brewster Road, Edinburgh EH9 3FJ, UK; s1117565@sms.ed.ac.uk

² Institute of Quantitative Biology, Biochemistry, and Biotechnology, School of Biological Science, The University of Edinburgh, Roger Land Building, Alexander Crum Brown Road, Edinburgh EH9 3FF, UK; louise.horsfall@ed.ac.uk

* Correspondence: Alison.Hulme@ed.ac.uk

Abstract: Spiro compounds provide attractive targets in drug discovery due to their inherent three-dimensional structures, which enhance protein interactions, aid solubility and facilitate molecular modelling. However, synthetic methodology for the spiro-functionalisation of important classes of penicillin and cephalosporin β -lactam antibiotics is comparatively limited. We report a novel method for the generation of spiro-cephalosporin compounds through a Michael-type addition to the dihydrothiazine ring. Coupling of a range of catechols is achieved under mildly basic conditions (K_2CO_3 , DMF), giving the stereoselective formation of spiro-cephalosporins (d.r. 14:1 to 8:1) in moderate to good yields (28–65%).

Keywords: spiro-cyclisation; spiro-cephalosporin; Michael-type reaction



Citation: Zhao, A.X.; Horsfall, L.E.; Hulme, A.N. New Methods for the Synthesis of Spirocyclic Cephalosporin Analogues. *Molecules* **2021**, *26*, 6035. <https://doi.org/10.3390/molecules26196035>

Academic Editor: Grzegorz Mlostof

Received: 20 August 2021

Accepted: 29 September 2021

Published: 5 October 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Spiro compounds (or spirocycles) are rigid scaffolds that consist of at least two rings fused through a single atom, known as the spiro atom. The structure of spirocycles was first discovered by Von Baeyer in the late 1890s [1]. Spiro compounds are an important class of molecule, and spirocycles are found in a number of different natural products with pronounced biological activities. Griseofulvin, for example, is a spirocyclic natural product generated by the fungus *Penicillium griseofulvum* that has been used clinically to treat dermatophytosis (ringworm) [2].

In recent years, spirocycles have emerged as attractive synthetic targets in drug discovery. Growing efforts have been made to employ spirocyclic scaffolds as core or periphery structures in drug synthesis, and many show promising biological activities as cholesterol absorption inhibitors [3–5], anti-Parkinsonian agents [6–8], anti-Alzheimer's agents [9–11], antiviral agents [12–15], antibacterial agents [16–18], β -lactamase inhibitors [19] and anti-cancer agents [20–23]. The inherent three-dimensional nature of spiro compounds provides better protein interactions as compared to their planar (hetero)aromatic counterparts [24]. Moreover, spirocyclic scaffolds have relatively rigid structures and possess a limited number of well-defined conformations, which is an advantage when using in silico docking to optimise interactions between drugs and protein targets computationally [25]. The inherent sp^3 nature of spirocycles, means they are also generally more water soluble than the corresponding (hetero)aromatic systems [26], and the introduction of a spirocycle can be used to modulate the water solubility of compounds of interest [24,27].

Nevertheless, spirocycles are considered a synthetic challenge, mainly due to the difficulty in synthesising quaternary carbon centres, as well as controlling the stereochemistry [28,29]. Multiple synthetic strategies have been employed to synthesise spirocycles including alkylation methods, cycloaddition approaches, rearrangement approaches, ring closure methods and radical cyclisation approaches [28–31].

Among the many possible spirocyclic structures, spirocyclic β -lactams have drawn increasing attention, not only because they exhibit desirable antibacterial properties [32–34], but also because they show other interesting activities as enzyme inhibitors [3,19,35,36] and antivirals [37]. Multiple reaction sites can be utilised to modify the penicillin- or cephalosporin-based β -lactam antibiotics with spirocyclic moieties without altering their β -lactam core structures. Figure 1 shows that the spirocycles can be either fused directly onto the β -lactam ring (site a) or attached to the thiazolidine or dihydrothiazine ring (sites b and c).





Figure 1. Modification sites for attaching spirocyclic moieties.

A number of different methods to achieve spiro-modification at site a have been reported including: (i) 1,3-dipolar cycloaddition approaches [38,39]; (ii) a phosphane-catalysed [3+2] cycloaddition approach [40]; and (iii) a rhodium-catalysed cyclopropanation approach [19]. However, methodologies for spiro-modifications at site b and c are rather limited, and mostly rely on 1,3-dipolar cycloaddition reactions [41]. In this work, we report a novel synthetic strategy to synthesise spirocyclic cephalosporin analogues at site c through a Michael-type reaction.

2. Results and Discussion

To achieve spiro-modification at site c of cephalosporin, catechols were used. Catechols are commonly found in natural products, insecticides and pharmaceuticals. They have well-documented antioxidant properties, acting as structural units in many bronchodilator, adrenergic, antioxidant, anti-Parkinsonian and anti-hypertensive drugs [42]. Catechols are important building blocks in organic synthesis and are frequently used as nucleophiles in substitution reactions due to the nucleophilicity of the catechol monoanion; catechols have also been used as nucleophilic catalysts for peptide bond formation [43]. Furthermore, catechols can undergo Michael addition to α,β -unsaturated carbonyl compounds to generate six-membered heterocyclic compounds under mild basic conditions. Cabbidu et al. reported a method to synthesise benzodioxane **3** by reacting pyrocatechol **1** with methyl 4-bromocrotonate **2** in the presence of K_2CO_3 (Scheme 1) [44]. The reaction is thought to proceed via nucleophilic substitution of the bromine, followed by Michael addition to yield the desired product **3**.




Scheme 1. Generation of benzodioxane **3** via a Michael addition reaction.

Inspired by the work by Cabbidu et al., we decided to take a similar approach, utilising the α,β -unsaturated carbonyl moiety within the cephalosporin structure to react with catechols for the generation of spiro-cephalosporin compounds through Michael addition reactions.

2.1. Reaction Scope for Generating Spiro-Cephalosporins

The spiro-cyclisation reaction was first conducted by reacting commercially available cephalosporin derivative **4** with pyrocatechol **1** in the presence of K_2CO_3 [45], under microwave irradiation at 50 °C to accelerate the reaction, as shown in Scheme 2.



Scheme 2. Generation of benzodioxane **3** via Michael addition reaction.

Spiro-cephalosporin **5** was isolated from the reaction mixture by column chromatography as a single diastereomer in moderate yield (40%). Stereochemical assignment was achieved using the NOESY spectrum of **5** (Figure 2), which shows the interaction of the ring junction proton H_a with methylene proton H_b , and the second methylene proton H_c with the axial proton H_d on the benzodioxane ring. An nOe interaction is also observed between the equatorial proton H_e on the benzodioxane ring and H_f , indicating these two protons are on the same face. Therefore, the C-2' and C-2 stereocentres in product **5** are determined as *S* and *R* configurations, as shown in Figure 2A (2D view) and B (3D view).


















Figure 2. Structure of spiro-cephalosporin **5** in (A) 2-D and (B) 3-D; (C) the NOESY spectrum.

To expand further the scope of the spiro-cyclisation reaction, a number of different catechols were screened under the same reaction conditions (Table 1).

Table 1. Reaction scope for the synthesis of spiro-cephalosporins ^a.

Entry	Catechol	Major Product	Isolated Yield (%)	dr ^b
1			59	— ^c
2			62	14:1
3			54	9:1
4			65	12:1
5			51	— ^c
6			— ^d	— ^d
7			28	8:1

^a Reagents and conditions: cephalosporin 4, K₂CO₃, DMF, μW, 50 °C, 50 min. ^b dr values determined by high performance liquid chromatography (HPLC). ^c Only one diastereomer was found. ^d No product was isolated by column chromatography.

The spiro-cyclisation reactions with dihydroxy coumarins (Table 1, entries 1–3) and flavonoids (Table 1, entries 4–6) were explored. Reactions were monitored by HPLC, and it was found that diastereomers were formed in the reactions of coumarins 7 and 8, and flavonoid 9 with diastereomeric ratios of 14:1, 9:1, and 12:1, respectively. For entry 3, the two diastereomers were successfully isolated and the characteristic NMR peaks were assigned to allow NOESY analysis to identify their stereochemistry. Figure 3A shows the NOESY spectra for the major diastereomer 15, in which nOe interactions are observed between H_a and H_b , H_c and H_d , and H_e and H_f . Therefore, the configurations of the newly generated chiral centres at the C-2' and C-2 positions were, again, determined to be *S* and *R*, respectively. In contrast, no interaction is observed between H_e and H_f for the minor diastereomer (Figure 3B), suggesting these two protons are not on the same face and, thus the minor diastereomer has a (2'*S*,2*S*) configuration. For the other reactions, only the major diastereomer was isolated by column chromatography in each case and the stereochemistry was found to be the same as for the major diastereomer 15, i.e., a (2'*S*,2*R*) configuration.






Figure 3. NOESY spectra of (A) the major diastereomer 15 and (B) the minor diastereomer.

The reaction with flavonoid 10 which contains three adjacent hydroxyl groups was also studied (Table 1, entry 5). The reaction was found to be regioselective, as spiro-product 17 was the predominant product (see Section 2.3). Again, the stereochemistry at the C-2' and C-2 positions of 17 was determined to be *S* and *R* configuration by NOESY NMR.

We also investigated the regioselectivity of spiro-cycloaddition when a flavonoid contains multiple hydroxyl groups. Quercetin 11 was reacted with cephalosporin 4 in the presence of K_2CO_3 (Table 1, entry 6). However, the reaction was unsuccessful, and no spiro-product 18 was isolated from the reaction mixture. From the LC-MS analysis, a mass of 753 Da was detected, which corresponds to the mass of an alkylated product. The most acidic phenol in quercetin is known to be the 7-OH on the flavonol core, thus the monoanion is most likely to form here. Reaction with cephalosporin 4 (see Section 2.3) gives an adduct which cannot proceed to spirocyclisation, resulting in the formation of an acyclic product. However, this putative alkylation product could not be isolated by column chromatography on silica gel, suggesting facile decomposition under mild acidic conditions.

Finally, we tested the reaction with the polyphenol, ellagic acid 12 (Table 1, entry 7). The reaction proceeded successfully; two diastereomers were observed in the reaction

Table 2. Cont.

Entry	Starting Material	Product	Isolated Yield (%)
6	17		63
7	19		77

2.3. Proposed Mechanism

For the 6,7-dihydroxycoumarins (**6–8**), the 7-OH ($pK_a = 7.5$) is reported to be much more acidic than the 6-OH ($pK_a = 9.0$) [46]. This difference in acidity can be explained by resonance stabilisation as shown in Figure 4A. Similarly, for flavonoids (**9** and **10**), the 7-OH ($pK_a = 7.7$) is more acidic than the 6-OH ($pK_a = 9.0$) due to resonance stabilisation [47], and, hence, can be deprotonated more easily under basic conditions (Figure 4B). In addition, for 5,6,7-trihydroxyflavonoid **10**, the 5-OH is found to be least acidic ($pK_a = 11.4$) due to the formation of an intramolecular hydrogen bond with the adjacent carbonyl group (Figure 4B) [47,48]. Thus spiro-cyclisation can be initiated by the 7-OH and is completed by the adjacent 6-OH to give the observed regioselectivity in compound **17** shown in Table 1, entry 5. In contrast, the two hydroxyl groups in pyrocatechol **1** have a much higher pK_a value ($pK_a = 9.3$) [49] than the 7-OH in the dihydroxy coumarins and flavonoids, which cannot be easily deprotonated under mild basic conditions, resulting in a lower reaction yield (40%).




Figure 4. Resonance structures of (A) coumarin **6** and (B) flavonoid **10** under mild basic conditions.

A mechanism for the generation of the spiro-cephalosporin product **30** is therefore proposed as shown in Figure 5. Under mildly basic conditions, the more acidic 7-OH in catechol **27** is readily deprotonated to form the reactive phenolate ion **28**. The phenolate ion **28** is then alkylated by cephalosporin **4** in the presence of NaI to form intermediate **29**. The experimental pK_a of the phenol in mono-alkylated catechols is consistently reported to be several log units higher [50,51] than the most acidic phenol in their non-alkylated precursors, hence a second phenolate alkylation reaction is not favoured under these conditions. Rather, phenol **29** undergoes an intramolecular Michael addition to form the

spiro-cephalosporin **30**, with initial attack on the convex face of the cephalosporin core, followed by protonation on the face opposite the newly-introduced, bulky catechol unit to set the observed (2′S,2R) stereochemistry.




Figure 5. Proposed spiro-cyclisation mechanism for the formation of spiro-cephalosporin **30**.

3. Materials and Methods

3.1. General Information

All reagents were obtained from commercial suppliers and were used without further purification. Microwave reactions were performed using Biotage Initiator+. Flash chromatography was carried out using Merck Kieselgel 60 (Merck 9385) under positive pressure. ^1H and ^{13}C NMR spectra were obtained on a Bruker AVA 500 instrument, using TMS as a reference and residual solvent as an internal standard. The data are presented as follows: chemical shift (in ppm on the δ scale relative to $\delta\text{TMS} = 0$), integration, multiplicity, coupling constant and interpretation. ^1H and ^{13}C spectra for all new compounds are presented in the SI file and raw data in the accompanying data deposit. Electrospray ionisation (ESI) mass spectra were obtained on a Bruker microTOF II instrument. Analytical Reverse Phase HPLC (Analytical RP-HPLC) was conducted on a Waters[®] 600 (100 μL) system using a 717plus autosampler and 996 PDA detector (190 to 800 nm) equipped with a Phenomenex[®] SphereClone ODS(2), 5 μm , 100×4.6 mm column for HPLC (method 1) or with a Phenomenex Luna C18(2), 5 μm , 250×4.6 mm column for HPLC (method 2). A binary solvent system was used A = water (0.1% TFA), B = MeCN (0.1% TFA) at a flow rate of $1.00\text{ mL}\cdot\text{min}^{-1}$; and the column was maintained at $30 \pm 1\text{ }^\circ\text{C}$. The HPLC (method 1) was a linear gradient from 0 min (95A:5B) to 10 min (5A:95B), isocratic from 10 min to 12 min (5A:95B), before recovery of the initial conditions over 3 min and equilibration over 5 min, giving a total run time of 20 min. The HPLC (method 2) was a linear gradient from 0 min (95A:5B) to 30 min (5A:95B), isocratic from 30 min to 35 min (5A:95B), before recovery of the initial conditions over 5 min and equilibration over 10 min, giving a total run time of 50 min.

3.2. Typical Procedure for the Spiro-Cyclisation Reaction

Pyrocatechol **1** (30.0 mg, 0.27 mmol), Cephamester **4** (0.13 g, 0.27 mmol), NaI (4.08 mg, 0.03 mmol) and K_2CO_3 (0.11 g, 0.82 mmol) were dissolved in DMF (3 mL). The reaction mixture was stirred at $50\text{ }^\circ\text{C}$ under microwave irradiation for 50 min. The reaction mixture was concentrated in vacuo then diluted with EtOAc (20 mL). The solution was washed with water (20 mL), Na_2SO_3 (20 mL, 5% aq) and brine (20 mL, sat aq), then dried ($MgSO_4$) and concentrated in vacuo. The crude product was purified with column chromatography (hexane:EtOAc, 3:1) to give the desired product **5** [45].

3.3. Characterisation of Spiro-Cephalosporin Products

(4-Methoxyphenyl)methyl (2'*S*,2*R*,6*R*,7*R*)-8-oxo-7-(2-phenylacetamido)-5-thia-1-azaspiro[1'*A'*-benzodioxine-3,2'-bicyclo[4.2.0]octane]-2-carboxylate (**5**): 61.2 mg (40%); R_t (method 1) = 14.52 min; R_f (hexane:EtOAc, 3:1) = 0.16; mp 157 – 160 °C; IR 2920 (NH), 1770 (C=O), 1734 (C=O), 1683 (C=O); $^1\text{H NMR } \delta$ (500 MHz, DMSO- d_6) 9.21 (1H, d, $J = 8.5$ Hz, NH), 7.34 – 7.22 (7H, m, ArH), 6.97 – 6.83 (5H, m, ArH), 6.64 (1H, dd, $J = 7.9, 1.7$ Hz, ArH), 5.42 (1H, dd, $J = 8.5, 4.3$ Hz, NHCH), 5.29 (1H, dd, $J = 4.3$ Hz, SCHN), 5.11 (1H, d, $J = 11.8$ Hz, $\text{OCH}_A\text{H}_B\text{Ar}$), 5.08 (1H, d, $J = 11.8$ Hz, $\text{OCH}_A\text{H}_B\text{Ar}$), 4.76 (1H, d, $J = 12.1$ Hz, ArOCH_CH_D), 4.68 (1H, s, NCHCOO), 4.20 (1H, d, $J = 12.1$ Hz, ArOCH_CH_D), 3.76 (3H, s, CH_3), 3.57 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 3.53 (1H, d, $J = 13.7$ Hz, SCH_GH_H), 3.52 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 2.93 (1H, d, $J = 13.7$ Hz, SCH_GH_H). $^{13}\text{C NMR } \delta$ (126 MHz, DMSO- d_6) 171.04 (C), 166.98 (C), 165.57 (C), 159.87 (C), 142.13 (C), 141.36 (C), 136.13 (C), 130.96 (2 × CH), 129.46 (2 × CH), 128.73 (2 × CH), 127.32 (C), 127.00 (CH), 122.77 (CH), 122.39 (CH), 117.84 (CH), 117.64 (CH), 114.23 (2 × CH), 70.81 (C), 67.26 (CH₂), 65.86 (CH₂), 59.44 (CH), 55.62 (CH₃), 55.58 (CH), 52.76 (CH), 42.28 (CH₂), 26.91 (CH₂); m/z (ESI+, MeCN) 583 ([M+Na]⁺, 5%), 578 ([M+NH₄]⁺, 100), 561 ([M+H]⁺, 30); HRMS (ESI+, MeCN) [M+H]⁺ found 561.1687, C₃₀H₂₉N₂O₇S requires 561.1690.

(4-Methoxyphenyl)methyl (2'*S*,2*R*,6*R*,7*R*)-7',8-dioxo-7-(2-phenylacetamido)-3',7'-dihydro-5-thia-1-azaspiro[1'*A'*-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylate (**13**): 210 mg (59%); R_t (method 1) = 12.39 min; R_f (EtOAc:DCM, 1:5) = 0.34; mp 100–103 °C; IR 3315 (NH), 1726 (C=O), 1681 (C=O), 1512 (C=C); $^1\text{H NMR } \delta$ (601 MHz, DMSO- d_6) 9.21 (1H, d, $J = 8.5$ Hz, NH), 7.79 (1H, d, $J = 9.5$ Hz, COCH=CH), 7.32 – 7.22 (7H, m, ArH), 7.08 (1H, s, ArH), 6.96 (1H, s, ArH), 6.86 (2H, d, $J = 8.7$ Hz, ArH), 6.36 (1H, d, $J = 9.5$ Hz, COCH=CH), 5.42 (1H, dd, $J = 8.5, 4.2$ Hz, NHCH), 5.30 (1H, d, $J = 4.2$ Hz, SCHN), 5.13 (1H, d, $J = 11.7$ Hz, $\text{OCH}_A\text{H}_B\text{Ar}$), 5.03 (1H, d, $J = 11.7$ Hz, $\text{OCH}_A\text{H}_B\text{Ar}$), 4.88 (1H, d, $J = 12.3$ Hz, ArOCH_CH_D), 4.67 (1H, s, NCHCOO), 4.32 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 3.73 (3H, s, CH_3), 3.56 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 3.52 (1H, d, $J = 13.8$ Hz, SCH_GH_H), 3.51 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 2.99 (1H, d, $J = 13.8$ Hz, SCH_GH_H); $^{13}\text{C NMR } \delta$ (151 MHz, DMSO- d_6) 171.05 (C), 166.84 (C), 165.61 (C), 160.51 (C), 159.82 (C), 149.27 (C), 145.33 (C), 144.12 (CH), 138.13 (C), 136.11 (C), 131.05 (2 × CH), 129.47 (2 × CH), 128.74 (2 × CH), 127.10 (CH), 127.02 (C), 115.89 (CH), 114.50 (CH), 114.21 (2 × CH), 113.96 (C), 105.07 (CH), 70.90 (C), 67.38 (CH₂), 66.34 (CH₂), 59.53 (CH), 55.66 (CH₃), 55.56 (CH), 52.60 (CH), 42.31 (CH₂), 26.94 (CH₂); m/z (ESI+, MeCN) 629 ([M+H]⁺, 100%); HRMS (ESI+, MeCN) [M+H]⁺ found 629.1597, C₃₃H₂₉N₂O₉S requires 629.1588.

(4-Methoxyphenyl)methyl (2'*S*,2*R*,6*R*,7*R*)-7',8-dioxo-7-(2-phenylacetamido)-3',7',8',9',10',11'-hexahydro-1',4',6'-trioxo-5-thia-1-azaspiro[bicyclo[4.2.0]octane-3,2'-tetraphene]-2-carboxylate (**14**): 420 mg (62%); R_t (method 1) = 13.63 min; R_f (DCM:EtOAc, 8:1) = 0.25; mp 120 – 123 °C; IR 2934 (NH), 1626 (C=O), 1701 (C=O), 1578 (C=O), 1506 (C=C); $^1\text{H NMR } \delta$ (500 MHz, DMSO- d_6) 9.21 (1H, d, $J = 8.5$ Hz, NH), 7.34 – 7.22 (5H, m, ArH), 7.16 (2H, d, $J = 8.7$ Hz, ArH), 7.07 (1H, s, C_{5'}H), 6.88 (1H, s, C_{12'}H), 6.81 (2H, d, $J = 8.7$ Hz, ArH), 5.44 (1H, dd, $J = 8.5, 4.2$ Hz, NHCH), 5.32 (1H, d, $J = 4.2$ Hz, SCHN), 5.19 (1H, d, $J = 12.1$ Hz, $\text{OCH}_A\text{H}_B\text{Ar}$), 5.04 (1H, d, $J = 12.1$ Hz, $\text{OCH}_A\text{H}_B\text{Ar}$), 4.86 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 4.74 (1H, s, NCHCOO), 4.32 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 3.71 (3H, s, CH_3), 3.57 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 3.56 (1H, d, $J = 13.8$ Hz, SCH_GH_H), 3.52 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 3.01 (1H, d, $J = 13.8$ Hz, SCH_GH_H), 2.60 – 2.54 (1H, m, C_{11'}H), 2.42 – 2.37 (3H, m, C_{8'}H₂, C_{11'}H), 1.72 – 1.71 (4H, m, C_{9'}H₂C_{10'}H₂); $^{13}\text{C NMR } \delta$ (126 MHz, DMSO- d_6) 171.05 (C), 166.97 (C), 165.64 (C), 161.11 (C), 159.60 (C), 147.31 (C), 147.01 (C), 144.07 (C), 138.14 (C), 136.11 (C), 130.17 (2 × CH), 129.47 (2 × CH), 128.74 (2 × CH), 127.27 (C), 127.02 (CH), 121.32 (C), 115.15 (C), 113.94 (2 × CH), 111.58 (CH), 104.83 (CH), 70.98 (C), 67.12 (CH₂), 66.32 (CH₂), 59.50 (CH), 55.73 (CH), 55.46 (CH₃), 52.66 (CH), 42.32 (CH₂), 26.96 (CH₂), 25.13 (CH₂), 24.11 (CH₂), 21.57 (CH₂), 21.26 (CH₂); m/z (ESI+, MeCN) 705 ([M+Na]⁺, 63%), 683 ([M+H]⁺, 94); HRMS (ESI+, MeCN) [M+H]⁺ found 683.2087, C₃₇H₃₅N₂O₉S requires 683.2058. Minor diastereomer: R_t (method 1) = 13.34 min; $^1\text{H NMR } \delta$ (500 MHz, DMSO- d_6)

diagnostic peaks: 9.12 (1H, d, $J = 8.2$ Hz, NH), 5.41 (1H, dd, $J = 8.2, 4.3$ Hz, NHCH), 5.19 (1H, d, $J = 4.3$ Hz, SCHN), 4.63 (1H, s, NCHCO), 3.77 (3H, s, CH₃).

(4-Methoxyphenyl)methyl (2'*S*,2*R*,6*R*,7*R*)-7'-8-dioxo-9'-phenyl-7-(2-phenylacetamido)-3'-7'-dihydro-5-thia-1-azaspiro[1',4'-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylate (**15**): 290 mg (54%); R_t (method 1) = 15.47 min; R_f (DCM:EtOAc, 6:1) = 0.51; mp 165 – 168 °C; IR 3120 (NH), 1762 (C=O), 1701 (C=O), 1514 (C=C); ¹H NMR δ (500 MHz, DMSO-*d*₆) 9.20 (1H, d, $J = 8.5$ Hz, NH), 7.60 – 7.58 (3H, m, ArH), 7.46 – 7.44 (2H, m, ArH), 7.33 – 7.24 (6H, m, ArH), 7.07 (2H, d, $J = 8.6$ Hz, ArH), 6.75 (1H, s, ArH), 6.67 (2H, $J = 8.6$ Hz, ArH), 6.32 (1H, s, =CH), 5.40 (1H, dd, $J = 8.5, 4.2$ Hz, NHCH), 5.27 (1H, d, $J = 4.2$ Hz, SCHN), 5.05 (1H, d, $J = 12.1$ Hz, OCH_AH_BAr), 4.97 (1H, d, $J = 12.1$ Hz, OCH_AH_BAr), 4.84 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 4.72 (1H, s, NCHCOO), 4.41 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 3.70 (3H, s, CH₃), 3.57 (1H, d, $J = 14.1$ Hz, COCH_EH_FAr), 3.51 (1H, d, $J = 14.1$ Hz, COCH_EH_FAr), 3.41 (1H, d, $J = 13.9$ Hz, SCH_GH_H), 2.94 (1H, d, $J = 13.9$ Hz, SCH_GH_H); ¹³C NMR δ (126 MHz, DMSO-*d*₆) 171.07 (C), 166.63 (C), 165.58 (C), 160.24 (C), 159.64 (C), 154.89 (C), 149.44 (C), 145.74 (C), 138.26 (C), 136.08 (C), 135.19 (C), 130.32 (CH), 130.19 (2 × CH), 129.46 (4 × CH), 128.73 (2 × CH), 128.70 (2 × CH), 127.16 (C), 127.01 (CH), 114.04 (3 × CH), 113.82 (C), 113.45 (CH), 105.87 (CH), 71.26 (C), 67.28 (CH₂), 66.41 (CH₂), 59.53 (CH), 55.63 (CH), 55.58 (CH₃), 52.84 (CH), 42.30 (CH₂), 26.78 (CH₂); m/z (ESI+, MeCN) 727 ([M+Na]⁺, 68%), 722 ([M+NH₄]⁺, 14), 705 ([M+H]⁺, 100); HRMS (ESI+, MeCN) [M+H]⁺ found 705.1896, C₃₉H₃₃N₂O₉S requires 705.1901. Minor diastereomer: R_t (method 1) = 15.24 min; ¹H NMR δ (500 MHz, DMSO-*d*₆) diagnostic peaks: 9.10 (1H, d, $J = 8.1$ Hz, NH), 5.41 (1H, dd, $J = 8.1, 4.3$ Hz, NHCH), 5.19 (2H, s, OCH₂Ar), 5.15 (1H, d, $J = 4.3$ Hz, SCHN), 4.54 (1H, s, NCHCO), 4.35 (1H, d, $J = 12.2$ Hz, ArOCH_AH_B), 4.18 (1H, d, $J = 12.2$ Hz, ArOCH_AH_B), 3.77 (3H, s, CH₃), 3.58 (1H, $J = 14.0$ Hz, COCH_CH_DAr), 3.52 (1H, $J = 14.0$ Hz, COCH_CH_DAr), 3.29 (1H, d, $J = 13.4$ Hz, SCH_EH_F), 2.94 (1H, d, $J = 13.4$ Hz, SCH_EH_F).

(4-Methoxyphenyl)methyl (2'*S*,2*R*,6*R*,7*R*)-8,9'-dioxo-7'-phenyl-7-(2-phenylacetamido)-3'-9'-dihydro-5-thia-1-azaspiro[1',4'-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylate (**16**): 67.3 mg (65%); R_t (method 1) = 15.52 min; R_f (DCM:EtOAc, 5:1) = 0.42; mp; 150 – 152 °C; IR 2933 (NH), 1770 (C=O), 1734 (C=O), 1624 (C=O), 1508 (C=C); ¹H NMR δ (500 MHz, DMSO-*d*₆) 9.20 (1H, d, $J = 8.5$ Hz, NH), 8.08 (2H, dd, $J = 7.8, 1.6$ Hz, ArH), 7.64 – 7.57 (3H, m, ArH), 7.47 (1H, d, $J = 1.6$ Hz, ArH), 7.33 – 7.28 (5H, m, ArH), 7.26 – 7.22 (3H, m, ArH), 6.99 (1H, s, =CH), 6.85 (2H, d, $J = 8.7$ Hz, ArH), 5.43 (1H, dd, $J = 8.5, 4.2$ Hz, NHCH), 5.33 (1H, d, $J = 4.2$ Hz, SCHN), 5.15 (1H, d, $J = 11.7$ Hz, OCH_AH_BAr), 5.07 (1H, d, $J = 11.7$ Hz, OCH_AH_BAr), 4.92 (1H, $J = 12.2$ Hz, ArOCH_CH_D), 4.72 (1H, s, NCHCOO), 4.46 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 3.72 (3H, s, CH₃), 3.58 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 3.56 (1H, d, $J = 13.8$ Hz, SCH_GH_H), 3.53 (1H, d, $J = 14.1$ Hz, COCH_EH_F), 3.02 (1H, d, $J = 13.8$ Hz, SCH_GH_H); ¹³C NMR δ (126 MHz, DMSO-*d*₆) 176.53 (C), 171.06 (C), 166.74 (C), 165.57 (C), 162.83 (C), 159.80 (C), 151.54 (C), 147.47 (C), 139.79 (C), 136.10 (C), 132.20 (CH), 131.65 (C), 130.94 (2 × CH), 129.59 (2 × CH), 129.47 (2 × CH), 128.75 (2 × CH), 127.09 (C), 127.02 (CH), 126.68 (2 × CH), 119.13 (C), 114.15 (2 × CH), 111.99 (CH), 106.58 (CH), 106.45 (CH), 71.30 (C), 67.58 (CH₂), 66.36 (CH₂), 59.58 (CH), 55.67 (CH), 55.44 (CH₃), 52.75 (CH), 42.33 (CH₂), 26.84 (CH₂); m/z (ESI+, MeCN) 705 ([M+H]⁺, 100%); HRMS (ESI+, MeCN) [M+H]⁺ found 705.1925, C₃₉H₃₃N₂O₉S requires 705.1901. Minor diastereomer: R_t (method 1) = 15.74 min; ¹H NMR δ (500 MHz, DMSO-*d*₆) diagnostic peaks: 9.13 (1H, d, $J = 8.5$ Hz, NH), 5.75 (1H, dd, $J = 8.4, 4.8$ Hz, NHCH), 5.15 (1H, d, $J = 4.2$ Hz, SCHN), 4.71 (1H, s, NCHCOO), 3.73 (3H, s, CH₃).

(4-Methoxyphenyl)methyl (2'*S*,2*R*,6*R*,7*R*)-10'-hydroxy-8,9'-dioxo-7'-phenyl-7-(2-phenylacetamido)-3',8'-dihydro-5-thia-1-azaspiro[1',4'-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylate (**17**): 40.3 mg (51%); R_t (method 1) = 15.27 min; R_f (DCM:EtOAc, 8:1) = 0.38; mp 113 – 115 °C; IR 2954 (NH), 1747 (C=O), 1734 (C=O), 1653 (C=O), 1516 (C=C); ¹H NMR δ (500 MHz, DMSO-*d*₆) 12.91 (1H, s, OH), 9.20 (1H, d, $J = 8.5$ Hz, NH), 8.11 (2H, d, $J = 7.1$ Hz, ArH), 7.66 – 7.59 (3H, m, ArH), 7.34 – 7.28 (4H, m, ArH), 7.25 (1H, m, ArH), 7.15 (2H, d, $J = 8.6$ Hz, ArH), 7.07 (1H, s, =CH), 6.94 (1H, s, ArH), 6.74 (2H, d, $J = 8.6$ Hz, ArH), 5.43

(1H, d, $J = 8.5, 4.2$ Hz, NHCH), 5.34 (1H, d, $J = 4.2$ Hz, SCHN), 5.11 (1H, d, $J = 11.8$ Hz, OCH_AH_BAr), 5.03 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 4.93 (1H, d, $J = 11.8$ Hz, OCH_AH_BAr), 4.69 (1H, s, NHCOO), 4.34 (1H, d, $J = 12.2$ Hz, ArOCH_CH_D), 3.65 – 3.63 (4H, m, SCH_EH_F, CH₃), 3.58 (1H, d, $J = 14.1$ Hz, COCH_GH_H), 3.53 (1H, d, $J = 14.1$ Hz, COCH_GH_H), 3.07 (1H, d, $J = 13.8$ Hz, SCH_EH_F); ¹³C NMR δ (126 MHz, DMSO-*d*₆) 183.04 (C), 171.05 (C), 166.81 (C), 165.64 (C), 164.25 (C), 159.66 (C), 150.46 (C), 148.87 (C), 148.36 (C), 136.10 (C), 132.73 (CH), 131.08 (C), 130.39 (2 × CH), 129.67 (2 × CH), 129.47 (2 × CH), 128.75 (2 × CH), 127.02 (CH), 126.94 (2 × CH), 126.85 (C), 126.05 (C), 114.03 (2 × CH), 106.12 (C), 105.00 (CH), 95.61 (CH), 70.44 (C), 67.73 (CH₂), 66.78 (CH₂), 59.53 (CH), 55.79 (CH), 55.38 (CH₃), 52.32 (CH), 42.34 (CH₂), 27.11 (CH₂); *m/z* (ESI+, MeCN) 721 ([M+H]⁺, 100%); HRMS (ESI+, MeCN) [M+H]⁺ found 721.1846, C₃₉H₃₃N₂O₁₀S requires 721.1850.

(4-Methoxyphenyl)methyl (2′*S*,2*R*,6*R*,7*R*)-14′,15′-dihydroxy-8,10′,17′-trioxo-7-(2-phenylacetamido)-3′,6′,11′,18′-tetraoxa-5-thia-1-azaspiro-bicyclo[4.2.0]octane-3,4′-pentacyclo-icosane-1′,2′,8′,12′,13′,15′-hexaene-2-carboxylate (**19**): 17.5 mg (28%); R_t (method 2) = 27.61 min; mp 131 – 133 °C; IR 2978 (NH), 1734 (C=O), 1716 (C=O), 1541 (C=O); ¹H NMR δ (500 MHz, DMSO-*d*₆) 9.23 (1H, d, $J = 8.5$ Hz, NH), 7.47 (1H, s, ArH), 7.35 – 7.21 (8H, m, ArH), 6.84 (2H, d, $J = 8.5$ Hz, ArH), 5.54 (1H, dd, $J = 8.5, 4.2$ Hz, NHCH), 5.34 (1H, d, $J = 4.2$ Hz, SCHN), 5.20 (1H, d, $J = 11.8$ Hz, OCH_AH_BAr), 5.06 (1H, d, $J = 11.8$ Hz, OCH_AH_BAr), 5.01 (1H, d, $J = 12.1$ Hz, ArOH_CH_D), 4.77 (1H, s, NCHCO), 4.59 (1H, d, $J = 12.1$ Hz, ArOH_CH_D), 3.73 (3H, s, CH₃), 3.60 – 3.51 (3H, m, SCH_EH_F, COCH₂), 3.07 (1H, d, $J = 13.8$ Hz, SCH_EH_F); ¹³C NMR δ (126 MHz, DMSO-*d*₆) 171.11 (C), 166.67 (C), 165.64 (C), 159.82 (2 × C), 158.91 (C), 158.86 (C), 149.42 (C), 142.78 (C), 137.22 (C), 136.85 (C), 136.10 (2 × C), 135.55 (C), 130.94 (2 × CH), 129.49 (2 × CH), 128.77 (2 × CH), 127.17 (C), 127.04 (CH), 114.11 (2 × CH), 113.76 (C), 113.28 (CH), 112.00 (C), 111.24 (C), 110.83 (CH), 71.79 (C), 67.54 (CH₂), 66.48 (CH₂), 59.65 (CH), 55.72 (CH), 55.42 (CH₃), 52.76 (CH), 42.35 (CH₂), 26.74 (CH₂); *m/z* (ESI-, MeCN) 751 ([M-H]⁻, 100%); HRMS (ESI-, MeCN) [M-H]⁻ found 751.1264, C₃₈H₂₇N₂O₁₃S requires 751.1228. Minor diastereomer: R_t (method 2) = 27.92 min; ¹H NMR δ (500 MHz, DMSO-*d*₆) diagnostic peaks: 9.24 (1H, d, $J = 7.9$ Hz, NH), 5.42 (1H, dd, $J = 7.9, 4.3$ Hz, NHCH), 5.21 (1H, d, $J = 4.3$ Hz, SCHN), 4.74 (1H, s, NCHCO), 3.77 (3H, s, CH₃).

3.4. Typical Procedure for PMB Deprotection

Anisole (0.15 mL, 1.34 mmol) was added to a stirred solution of spiro-cephalosporin **5** (30.0 mg, 0.05 mmol) in DCM (4 mL) and was cooled to 0 °C. TFA (0.21 mL, 2.68 mmol) was added dropwise to the reaction mixture and the mixture was stirred at 0 °C for 1 h, then at rt for 3 h. The reaction mixture was concentrated in vacuo and the crude product was purified by column chromatography (DCM:MeOH:AcOH, 96:4:1) to give the desired product **20** [45].

3.5. Characterisation of PMB Deprotected Products

(2′*S*,2*R*,6*R*,7*R*)-8-Oxo-7-(2-phenylacetamido)-5-thia-1-azaspiro[1′,4′-benzodioxine-3,2′-bicyclo[4.2.0]octane]-2-carboxylic acid (**20**): 16.6 mg (71%); R_t (method 1) = 12.90 min; R_f (DCM:MeOH:AcOH, 96:4:1) = 0.25; mp 206 – 208 °C; IR (neat, cm⁻¹) 3400 – 3061 (OH), 2887 (NH), 1747 (C=O), 1625 (C=O); ¹H NMR δ (500 MHz, DMSO-*d*₆) 9.18 (1H, d, $J = 8.5$ Hz, NH), 7.35 – 7.26 (4H, m, ArH), 7.26 – 7.22 (1H, m, ArH), 7.01 – 6.95 (1H, m, ArH), 6.91 – 6.88 (2H, m, ArH), 6.86 – 6.81 (1H, m, ArH), 5.39 (1H, dd, $J = 8.5, 4.2$ Hz, NHCH), 5.29 (1H, d, $J = 4.2$ Hz, SCHN), 4.73 (1H, d, $J = 12.0$ Hz, ArOH_AH_B), 4.51 (1H, s, NCHCOO), 4.20 (1H, d, $J = 12.0$ Hz, ArOH_AH_B), 3.65 (1H, d, $J = 13.6$ Hz, SCH_CH_D), 3.57 (1H, d, $J = 14.0$ Hz, COCH_EH_F), 3.52 (1H, d, $J = 14.0$ Hz, COCH_EH_F), 2.86 (1H, d, $J = 13.6$ Hz, SCH_CH_D); ¹³C NMR δ (126 MHz, DMSO-*d*₆) 171.03 (C), 168.41 (C), 165.48 (C), 142.36 (C), 141.79 (C), 136.20 (C), 129.49 (2 × CH), 128.73 (2 × CH), 126.99 (CH), 122.76 (CH), 122.18 (CH), 117.87 (CH), 117.66 (CH), 70.84 (C), 66.12 (CH₂), 59.22 (CH), 55.40 (CH), 53.40 (CH), 42.28 (CH₂), 26.96 (CH₂); *m/z* (ESI+, MeCN) 463 ([M+Na]⁺, 29%), 458 ([M+NH₄]⁺, 15), 441 ([M+H]⁺, 100); HRMS (ESI+, MeCN) [M+H]⁺ found 441.1104, C₂₂H₂₁N₂O₆S requires 441.1115.

(2'*S*,2*R*,6*R*,7*R*)-7',8-Dioxo-7-(2-phenylacetamido)-3',7'-dihydro-5-thia-1-azaspiro[1',4'-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylic acid (**21**): 37.0 mg (93%); R_t (method 1) = 12.03 min; R_f (DCM:MeOH:AcOH, 92:8:1) = 0.40; mp 210 – 212 °C; IR 3304 (NH), 1718 (C=O), 1560 (C=O), 1500 (C=C); $^1\text{H NMR } \delta$ (500 MHz, DMSO- d_6) 9.15 (1H, d, J = 8.5 Hz, NH), 7.94 (1H, d, J = 9.6 Hz, COCH=CH), 7.35 – 7.27 (4H, m, ArH), 7.26 – 7.20 (2H, m, ArH), 7.08 (1H, s, ArH), 6.34 (1H, d, J = 9.6 Hz, COCH=CH), 5.34 (2H, m, NHCH, SCHN), 4.78 (1H, d, J = 12.0 Hz, OCH_AH_B), 4.40 (1H, br s, NCHCOO), 4.38 (1H, d, J = 12.0 Hz, OCH_AH_B), 3.76 (1H, br d, J = 13.6 Hz, SCH_CH_D), 3.57 (1H d, J = 14.1 Hz, COCH_EH_F), 3.52 (1H d, J = 14.1 Hz, COCH_EH_F), 2.82 (1H, br d, J = 13.6 Hz, SCH_CH_D); $^{13}\text{C NMR } \delta$ (126 MHz, DMSO- d_6) 171.02 (C), 168.00 (C), 165.24 (C), 160.63 (C), 149.22 (C), 145.81 (C), 144.29 (CH), 138.84 (C), 136.23 (C), 129.49 (2 × CH), 128.73 (2 × CH), 126.98 (CH), 115.93 (CH), 114.44 (CH), 113.99 (C), 104.94 (CH), 71.11 (C), 66.75 (CH₂), 59.10 (CH), 55.43 (CH), 54.44 (CH), 42.29 (CH₂), 26.94 (CH₂); m/z (ESI+, MeCN) 531 ([M+Na]⁺, 100%), 509 ([M+H]⁺, 33); HRMS (ESI+, MeCN) [M+H]⁺ found 509.1038, C₂₅H₂₁N₂O₈S requires 509.1013.

(2'*S*,2*R*,6*R*,7*R*)-7',8-Dioxo-7-(2-phenylacetamido)-3',7',8',9',10',11'-hexahydro-1',4',6'-trioxo-5-thia-1-azaspiro[bicyclo[4.2.0]octane-3,2'-tetraphene]-2-carboxylic acid (**22**): 60.4 mg (74%); R_t (method 1) = 12.94 min; R_f (DCM:MeOH:AcOH, 95:5:1) = 0.50; mp 218 – 220 °C; IR 3400 – 3200 (OH), 2934 (NH), 1751 (C=O), 1676 (C=O); $^1\text{H NMR } \delta$ (500 MHz, DMSO- d_6) 9.19 (1H, J = 8.5 Hz, NH), 7.32 – 7.25 (5H, m, ArH), 7.13 (1H, s, ArH), 7.08 (1H, s, ArH), 5.40 (1H, dd, J = 8.5, 4.1 Hz, NHCH), 5.30 (1H, d, J = 4.1 Hz, SCHN), 4.81 (1H, d, J = 12.1 Hz, ArOCH_AH_B), 4.54 (1H, s, NCHCOO), 4.36 (1H, d, J = 12.1 Hz, ArOCH_AH_B), 3.66 (1H, d, J = 13.6 Hz, SCH_CH_D), 3.57 (1H, d, J = 14.1 Hz, COCH_EH_F), 3.52 (1H, d, J = 14.1 Hz, COCH_EH_F), 2.92 (1H, d, J = 13.6 Hz, SCH_CH_D), 2.76 – 2.64 (2H, m, C_{11'}H), 2.42 – 2.39 (2H, m, C_{11'}H), 1.77 – 1.69 (4H, m, C₉H₂C₁₀H₂). $^{13}\text{C NMR } \delta$ (126 MHz, DMSO- d_6) 171.03 (C), 168.27 (C), 165.53 (C), 161.18 (C), 147.36 (C), 147.01 (C), 144.36 (C), 138.52 (C), 136.18 (C), 129.49 (2 × CH), 128.74 (2 × CH), 127.00 (CH), 121.48 (C), 115.10 (C), 111.65 (CH), 104.85 (CH), 71.05 (C), 66.52 (CH₂), 59.31 (CH), 55.48 (CH), 53.37 (CH), 42.31 (CH₂), 26.90 (CH₂), 25.18 (CH₂), 24.13 (CH₂), 21.60 (CH₂), 21.31 (CH₂). m/z (ESI+, MeCN) 585 ([M+Na]⁺, 23%), 580 ([M+NH₄]⁺, 35), 563 ([M+H]⁺, 100); HRMS (ESI+, MeCN) [M+H]⁺ found 563.1476, C₂₉H₂₇N₂O₈S requires 563.1483.

(2'*S*,2*R*,6*R*,7*R*)-7',8-Dioxo-9'-phenyl-7-(2-phenylacetamido)-3',7'-dihydro-5-thia-1-azaspiro[1',4'-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylic acid (**23**): 150 mg (74%); R_t (method 1) = 13.35 min; R_f (DCM:MeOH:AcOH, 90:10:1) = 0.38; mp 182 – 184 °C; IR 3300 – 2954 (OH), 2924 (NH), 1724 (C=O), 1626 (C=O), 1506 (C=C); $^1\text{H NMR } \delta$ (500 MHz, DMSO- d_6) 13.52 (1H, s, OH), 9.18 (1H, d, J = 8.5 Hz, NH), 7.59 (3H, m, ArH), 7.51 (2H, m, ArH), 7.33 – 7.27 (4H, m, ArH), 7.25 – 7.22 (2H, m, ArH), 6.75 (1H, s, ArH), 6.32 (1H, s, =CH), 5.41 (1H, dd, J = 8.5, 4.2 Hz, NHCH), 5.26 (1H, d, J = 4.2 Hz, SCHN), 4.88 (1H, d, J = 12.2 Hz, ArOCH_AH_B), 4.54 (1H, s, NCHCOO), 4.39 (1H, d, J = 12.2 Hz, ArOCH_AH_B), 3.57 (1H, d, J = 14.1 Hz, COCH_EH_F), 3.55 (1H, d, J = 13.8 Hz, SCH_CH_D), 3.52 (1H, d, J = 14.1 Hz, COCH_EH_F), 2.93 (1H, d, J = 13.8 Hz, SCH_CH_D); $^{13}\text{C NMR } \delta$ (126 MHz, DMSO- d_6) 171.02 (C), 168.26 (C), 165.63 (C), 160.27 (C), 154.80 (C), 149.44 (C), 145.79 (C), 138.39 (C), 136.15 (C), 135.24 (C), 130.27 (CH), 129.49 (2 × CH), 129.46 (2 × CH), 128.79 (2 × CH), 128.74 (2 × CH), 127.01 (CH), 114.17 (CH), 113.74 (C), 113.43 (CH), 105.75 (CH), 70.85 (C), 66.63 (CH₂), 59.39 (CH), 55.53 (CH), 52.88 (CH), 42.32 (CH₂), 26.93 (CH₂); m/z (ESI+, MeCN) 585 ([M+H]⁺, 100); HRMS (ESI+, MeCN) [M+H]⁺ found 585.1315, C₃₁H₂₅N₂O₈S requires 585.1326.

(2'*S*,2*R*,6*R*,7*R*)-8,9'-Dioxo-7'-phenyl-7-(2-phenylacetamido)-3',9'-dihydro-5-thia-1-azaspiro[1',4'-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylic acid (**24**): 61.9 mg (97%); R_t (method 1) = 13.47 min; R_f (DCM:MeOH:AcOH, 90:10:1) = 0.42; mp 177 – 178 °C; IR 3430 – 3120 (OH), 2924 (NH), 1751 (C=O), 1622 (C=O); $^1\text{H NMR } \delta$ (500 MHz, DMSO- d_6) 13.63 (1H, s, OH), 9.18 (1H, d, J = 8.5 Hz, NH), 8.09 (2H, dd, J = 8.0, 1.6 Hz, ArH), 7.61 (3H, m, ArH), 7.52 (1H, s, ArH), 7.36 (1H, s, ArH), 7.34 – 7.27 (4H, m, ArH), 7.25 (1H, m, ArH), 6.96 (1H, s, =CH), 5.42 (1H, dd, J = 8.5, 4.2 Hz, NHCH), 5.31 (1H, d, J = 4.2 Hz, SCHN), 4.96 (1H, d, J = 12.2 Hz, ArOCH_AH_B), 4.59 (1H, s, NCHCOO), 4.43 (1H, d, J = 12.2 Hz, ArOCH_AH_B), 3.65 (1H,

d, $J = 13.7$ Hz, $SCH_C H_D$), 3.58 (1H, d, $J = 14.1$ Hz, $COCH_E H_F Ar$), 3.53 (1H, d, $J = 14.1$ Hz, $COCH_E H_F Ar$), 3.00 (1H, d, $J = 13.7$ Hz, $SCH_C H_D$); ^{13}C NMR δ (126 MHz, DMSO- d_6) 176.56 (C), 171.04 (C), 168.27 (C), 165.60 (C), 162.95 (C), 151.60 (C), 147.71 (C), 140.14 (C), 136.15 (C), 132.21 (CH), 131.70 (C), 129.59 (2 \times CH), 129.50 (2 \times CH), 128.75 (2 \times CH), 127.02 (CH), 126.74 (2 \times CH), 119.10 (C), 111.64 (CH), 106.64 (CH), 106.46 (CH), 71.13 (C), 66.60 (CH₂), 59.45 (CH), 55.51 (CH), 52.94 (CH), 42.34 (CH₂), 27.02 (CH₂); m/z (ESI+, MeCN) 585 ([M+H]⁺, 100%); HRMS (ESI+, MeCN) [M+H]⁺ found 585.1353, C₃₁H₂₅N₂O₈S requires 585.1326.

(2'*S*,2*R*,6*R*,7*R*)-10'-Hydroxy-8,9'-dioxo-7'-phenyl-7-(2-phenylacetamido)-3',9'-dihydro-5-thia-1-azaspiro[1',4'-dioxino-3,2'-bicyclo[4.2.0]octane]-2-carboxylic acid (**25**): 60.0 mg (63%); R_t (method 1) = 13.73 min; R_f (DCM:MeOH:AcOH, 95:5:1) = 0.47; mp 171 – 173 °C; IR 3500 – 3400 (OH), 3000 (NH), 1757 (C=O), 1609 (C=O), 1500 (C=C); 1H NMR δ (500 MHz, DMSO- d_6) 13.61 (1H, brs, COOH), 12.94 (1H, s, ArOH), 9.18 (1H, d, $J = 8.5$ Hz, NH), 8.11 (2H, d, $J = 7.0$ Hz, ArH), 7.66 – 7.57 (3H, m, ArH), 7.34 – 7.28 (4H, m, ArH), 7.25 (1H, m, ArH), 7.03 (1H, s, =CH), 6.97 (1H, s, ArH), 5.42 (1H, dd, $J = 8.5, 4.1$ Hz, NHCH), 5.31 (1H, d, $J = 4.1$ Hz, SCHN), 4.97 (1H, d, $J = 12.1$ Hz, ArOCH_AH_B), 4.56 (1H, s, NHCOO), 4.40 (1H, d, $J = 12.1$ Hz, ArOCH_AH_B), 3.69 (1H, d, $J = 13.7$ Hz, $SCH_C H_D$), 3.58 (1H, d, $J = 14.0$ Hz, $COCH_E H_F$), 3.53 (1H, d, $J = 14.0$ Hz, $COCH_E H_F$), 2.98 (1H, d, $J = 13.7$ Hz, $SCH_C H_D$); ^{13}C NMR δ (126 MHz, DMSO- d_6) 183.15 (C), 171.02 (C), 168.18 (C), 165.56 (C), 164.36 (C), 150.51 (C), 149.03 (C), 148.68 (C), 136.14 (C), 132.70 (CH), 131.14 (C), 129.65 (2 \times CH), 129.48 (2 \times CH), 128.74 (2 \times CH), 127.01 (3 \times CH), 126.26 (C), 106.06 (C), 105.05 (CH), 95.64 (CH), 70.53 (C), 66.84 (CH₂), 59.35 (CH), 55.55 (CH), 52.88 (CH), 42.33 (CH₂), 26.98 (CH₂); m/z (ESI-, MeCN) 599 ([M-H]⁻, 100%); HRMS (ESI-, MeCN) [M-H]⁻ found 599.1110, C₃₁H₂₃N₂O₉S requires 599.1119.

(2'*S*,2*R*,6*R*,7*R*)-14',15'-Dihydroxy-8,10',17'-trioxo-7-(2-phenylacetamido)-3',6',11',18'-tetraoxa-5-thia-1-azaspiro-bicyclo[4.2.0]octane-3,4'-pentacyclo-icosane-1',2',8',12',13',15'-hexaene-2-carboxylic acid (**26**): 12.3 mg (77%); R_t (method 1) = 12.42 min; R_f (DCM:MeOH:AcOH, 86:14:1) = 0.23; mp 184 – 185 °C; IR 3486 – 3070 (OH), 2974 (NH), 1737 (C=O), 1614 (C=O); 1H NMR δ (600 MHz, DMSO- d_6) 9.21 (1H, d, $J = 8.5$ Hz, NH), 7.46 (1H, s, ArH), 7.42 (1H, s, ArH), 7.34 – 7.30 (4H, m, ArH), 7.26 – 7.25 (1H, m, ArH), 5.42 (1H, dd, $J = 8.5, 4.2$ Hz, NHCH), 5.34 (1H, d, $J = 4.1$ Hz, SCHN), 5.06 (1H, d, $J = 12.1$ Hz, ArOCH_AH_B), 4.61 (1H, s, NCHCOO), 4.54 (1H, d, $J = 12.1$ Hz, ArOCH_AH_B), 3.72 (1H, d, $J = 13.5$ Hz, $SCH_C H_D$), 3.59 (1H, d, $J = 14.1$ Hz, $COCH_E H_F$), 3.54 (1H, d, $J = 14.1$ Hz, $COCH_E H_F$), 3.03 (1H, d, $J = 13.5$ Hz, $SCH_C H_D$); ^{13}C NMR δ (151 MHz, DMSO- d_6) 171.06 (C), 168.18 (C), 165.56 (C), 159.02 (C), 158.88 (C), 149.41 (C), 143.19 (C), 137.30 (C), 136.81 (C), 136.17 (C), 135.78 (C), 132.07 (C), 129.51 (2 \times CH), 128.76 (2 \times CH), 127.02 (CH), 120.41 (C), 113.70 (C), 113.02 (CH), 112.01 (C), 111.16 (C), 110.86 (CH), 71.71 (C), 66.78 (CH₂), 59.42 (CH), 55.53 (CH), 53.20 (CH), 42.36 (CH₂), 26.97 (CH₂); m/z (ESI-, MeCN) 631 ([M-H]⁻, 100%); HRMS (ESI-, MeCN) [M-H]⁻ found 631.0064, C₃₀H₁₉N₂O₁₂S requires 631.0653.

4. Conclusions

Novel methodology for the synthesis of spiro-cephalosporins through a Michael-type reaction with the dihydrothiazine ring under mild basic conditions has been developed. A range of catechols were screened, and the spiro-cyclisation reactions were found to be highly diastereoselective (d.r. 14:1 to 8:1) with moderate to good yields (28–65%). The spirocyclisation products were transformed to more drug-like motifs through simple deprotection of the PMB-ester found in the cephalosporin precursor. Although the focus of this work has been the fusion of antioxidants, such as coumarins and flavonoids, with the cephalosporin core, we anticipate that spiro-cyclisation could be achieved with a range of other interesting catechol species, such as dopamine, apomorphine, catechin and caffeic acid. Given the recent rise in interest in the synthesis of spirocyclic motifs in medicinal chemistry [52], we anticipate that this new approach to spirocyclisation might be implemented across other target structures.

Supplementary Materials: The following are available online at <https://www.mdpi.com/article/10.3390/molecules26196035/s1>, Figures S1–S14: ^1H and ^{13}C NMR spectra for compounds 5, and 13–26.

Author Contributions: The work was conceived by A.X.Z., L.E.H. and A.N.H., conducted by A.X.Z., and supervised by L.E.H. and A.N.H. The manuscript was drafted by A.X.Z., and reviewed and edited by L.E.H. and A.N.H. All authors have read and agreed to the published version of the manuscript.

Funding: We thank the EPSRC (Grant Refs: EP/M506515/1 and EP/N509644/1) and IBioIC (Innovation Centre Programme funded by SFC, SE and HIE) for funding (studentship to AZ).

Data Availability Statement: Data is contained within the article or Supplementary Materials.

Acknowledgments: We thank Faye Cruickshank of the SIRCAMS facility at the University of Edinburgh for assistance with and Juraj Bella for assistance with NMR experiments.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Samples of the compounds are not available from the authors.

References

1. Baeyer, A. Systematik und Nomenclatur Bicyclischer Kohlenwasserstoffe. *Ber. Dtsch. Chem. Ges.* **1900**, *33*, 3771–3775. [[CrossRef](#)]
2. Petersen, A.B.; Rønne, M.H.; Larsen, T.O.; Clausen, M.H. The Chemistry of Griseofulvin. *Chem. Rev.* **2014**, *114*, 12088–12107. [[CrossRef](#)]
3. Dugar, S.; Clader, J.W.; Chan, T.-M.; Davis, H. Substituted 2-azaspiro [5.3] nonan-1-ones as potent cholesterol absorption inhibitors: Defining a binding conformation for SCH 48461. *J. Med. Chem.* **1995**, *38*, 4875–4877. [[CrossRef](#)]
4. Kambara, T.; Tomioka, K. Formal Asymmetric Synthesis of a Cholesterol Absorption Inhibitor Bearing a 2-Azaspiro [3.5] nonan-1-one Moiety. *J. Org. Chem.* **1999**, *64*, 9282–9285. [[CrossRef](#)]
5. Howell, K.; Devita, R.J.; Garcia-Calvo, M.; Meurer, R.D.; Lisnock, J.; Bull, H.G.; McMasters, D.R.; McCann, M.E.; Mills, S.G. Spiroimidazolidinone NPC1L1 inhibitors. Part 2: Structure-activity studies and in vivo efficacy. *Bioorganic Med. Chem. Lett.* **2010**, *20*, 6929–6932. [[CrossRef](#)]
6. Khalil, E.M.; Ojala, W.H.; Pradhan, A.; Nair, V.D.; Gleason, W.B.; Mishra, R.K.; Johnson, R.L. Design, Synthesis, and Dopamine Receptor Modulating Activity of Spiro Bicyclic Peptidomimetics of Prolyl-L-leucyl-glycinamide. *J. Med. Chem.* **1999**, *42*, 628–637. [[CrossRef](#)] [[PubMed](#)]
7. Macias, A.; Ramallal, A.M.; Alonso, E.; del Pozo, C.; González, J. Synthesis of Enantiopure Pyrrolidine-Derived Peptidomimetics and Oligo- β -peptides via Nucleophilic Ring-Opening of β -Lactams†. *J. Org. Chem.* **2006**, *71*, 7721–7730. [[CrossRef](#)]
8. Bhagwanth, S.; Mishra, S.; Daya, R.; Mah, J.; Mishra, R.K.; Johnson, R.L. Transformation of Pro-Leu-Gly-NH₂ Peptidomimetic Positive Allosteric Modulators of the Dopamine D₂ Receptor into Negative Modulators. *ACS Chem. Neurosci.* **2012**, *3*, 274–284. [[CrossRef](#)] [[PubMed](#)]
9. Yuan, J.; Venkatraman, S.; Zheng, Y.; McKeever, B.M.; Dillard, L.W.; Singh, S.B. Structure-Based Design of β -Site APP Cleaving Enzyme 1 (BACE1) Inhibitors for the Treatment of Alzheimer’s Disease. *J. Med. Chem.* **2013**, *56*, 4156–4180. [[CrossRef](#)] [[PubMed](#)]
10. Thomas, A.A.; Hunt, K.W.; Newhouse, B.; Watts, R.J.; Liu, X.; Vigers, G.; Smith, D.; Rhodes, S.P.; Brown, K.D.; Otten, J.N.; et al. 8-Tetrahydropyran-2-yl Chromans: Highly Selective Beta-Site Amyloid Precursor Protein Cleaving Enzyme 1 (BACE1) Inhibitors. *J. Med. Chem.* **2014**, *57*, 10112–10129. [[CrossRef](#)]
11. Dineen, T.A.; Chen, K.; Cheng, A.C.; Derakhchan, K.; Epstein, O.; Esmay, J.; Hickman, D.; Kreiman, C.; Marx, I.E.; Wahl, R.C.; et al. Inhibitors of β -site amyloid precursor protein cleaving enzyme (BACE1): Identification of (S)-7-(2-fluoropyridin-3-yl)-3-((3-methyloxetan-3-yl)ethynyl)-5'-H-spiro[chromeno[2,3-b]pyridine-5,4'-oxazol]-2'-amine (AMG-8718). *J. Med. Chem.* **2014**, *57*, 9811–9831. [[CrossRef](#)]
12. Neelamkavil, S.F.; Agrawal, S.; Bara, T.; Bennett, C.; Bhat, S.; Biswas, D.; Brockunier, L.; Buist, N.; Burnette, D.; Cartwright, M.; et al. Discovery of MK-8831, a novel spiro-proline macrocycle as a pan-genotypic HCV-NS3/4a protease inhibitor. *ACS Med. Chem. Lett.* **2016**, *7*, 111–116. [[CrossRef](#)]
13. Link, J.O.; Taylor, J.G.; Xu, L.; Mitchell, M.; Guo, H.; Liu, H.; Kato, D.; Kirschberg, T.; Sun, J.; Squires, N.; et al. Discovery of Ledipasvir (GS-5885): A Potent, Once-Daily Oral NS5A Inhibitor for the Treatment of Hepatitis C Virus Infection. *J. Med. Chem.* **2014**, *57*, 2033–2046. [[CrossRef](#)]
14. Ye, N.; Chen, H.; Wold, E.A.; Shi, P.-Y.; Zhou, J. Therapeutic Potential of Spirooxindoles as Antiviral Agents. *ACS Infect. Dis.* **2016**, *2*, 382–392. [[CrossRef](#)]
15. Das, P.; Boone, S.; Mitra, D.; Turner, L.; Tandon, R.; Raucher, D.; Hamme, A.T. Synthesis and biological evaluation of fluoro-substituted spiro-isoxazolines as potential anti-viral and anti-cancer agents. *RSC Adv.* **2020**, *10*, 30223–30237. [[CrossRef](#)]
16. Guardia, A.; Baiget, J.; Cacho, M.; Pérez, A.; Ortega-Guerra, M.; Nxumalo, W.; Khanye, S.D.; Rullas, J.; Ortega, F.; Jiménez, E.; et al. Easy-To-Synthesize Spirocyclic Compounds Possess Remarkable in vivo Activity against *Mycobacterium tuberculosis*. *J. Med. Chem.* **2018**, *61*, 11327–11340. [[CrossRef](#)]

17. Blondiaux, N.; Moune, M.; Desroses, M.; Frita, R.; Flipo, M.; Mathys, V.; Soetaert, K.; Kiass, M.; Delorme, V.; Djaout, K.; et al. Reversion of antibiotic resistance in *Mycobacterium tuberculosis* by spiroisoxazoline SMART-420. *Science* **2017**, *355*, 1206–1211. [[CrossRef](#)] [[PubMed](#)]
18. Urzúa, A.; Echeverría, J.; Rezende, M.C.; Wilkens, M. Antibacterial properties of 3 *H*-spiro[1-benzofuran-2,1'-cyclohexane] derivatives from *Heliotropium filifolium*. *Molecules* **2008**, *13*, 2385–2393. [[CrossRef](#)] [[PubMed](#)]
19. Sandanayaka, V.P.; Prashad, A.S.; Yang, Y.; Williamson, R.T.; Lin, Y.I.; Mansour, T.S. Spirocyclopropyl β -lactams as mechanism-based inhibitors of serine β -lactamases. Synthesis by rhodium-catalyzed cyclopropanation of 6-diazopenicillanate sulfone. *J. Med. Chem.* **2003**, *46*, 2569–2571. [[CrossRef](#)]
20. Mani, K.S.; Kaminsky, W.; Rajendran, S.P. A facile atom economic one pot multicomponent synthesis of bioactive spiro-indenoquinoxaline pyrrolizines as potent antioxidants and anti-cancer agents. *New J. Chem.* **2018**, *42*, 301–310. [[CrossRef](#)]
21. Islam, M.S.; Al-Majid, A.M.; El-Senduny, F.F.; Badria, F.A.; Rahman, A.F.M.M.; Barakat, A.; Elshaier, Y.A.M.M. Synthesis, Anticancer Activity, and Molecular Modeling of New Halogenated Spiro [pyrrolidine-thiazolo-oxindoles] Derivatives. *Appl. Sci.* **2020**, *10*, 2170. [[CrossRef](#)]
22. Hati, S.; Tripathy, S.; Dutta, P.K.; Agarwal, R.; Srinivasan, R.; Singh, A.; Singh, S.; Sen, S. Spiro[pyrrolidine-3,3-oxindole] as potent anti-breast cancer compounds: Their design, synthesis, biological evaluation and cellular target identification. *Sci. Rep.* **2016**, *6*, 32213. [[CrossRef](#)] [[PubMed](#)]
23. Najim, N.; Bathich, Y.; Zain, M.M.; Hamzah, A.S.; Shaameri, Z. Evaluation of the Bioactivity of Novel Spiroisoxazoline Type Compounds against Normal and Cancer Cell Lines. *Molecules* **2010**, *15*, 9340–9353. [[CrossRef](#)] [[PubMed](#)]
24. Zheng, Y.-J.; Tice, C.M. The utilization of spirocyclic scaffolds in novel drug discovery. *Expert Opin. Drug Discov.* **2016**, *11*, 831–834. [[CrossRef](#)]
25. Shennan, B.D.A.; Smith, P.W.; Ogura, Y.; Dixon, D.J. A modular and divergent approach to spirocyclic pyrrolidines. *Chem. Sci.* **2020**, *11*, 10354–10360. [[CrossRef](#)]
26. Lovering, F.; Bikker, J.; Humblet, C. Escape from Flatland: Increasing Saturation as an Approach to Improving Clinical Success. *J. Med. Chem.* **2009**, *52*, 6752–6756. [[CrossRef](#)]
27. Zheng, Y.; Tice, C.M.; Singh, S.B. The use of spirocyclic scaffolds in drug discovery. *Bioorganic Med. Chem. Lett.* **2014**, *24*, 3673–3682. [[CrossRef](#)]
28. Rios, R. Enantioselective methodologies for the synthesis of spiro compounds. *Chem. Soc. Rev.* **2012**, *41*, 1060–1074. [[CrossRef](#)]
29. Ding, A.; Meazza, M.; Guo, H.; Yang, J.W.; Rios, R. New development in the enantioselective synthesis of spiro compounds. *Chem. Soc. Rev.* **2018**, *47*, 5946–5996. [[CrossRef](#)] [[PubMed](#)]
30. Franz, A.K.; Hanhan, N.V.; Ball-Jones, N.R. Asymmetric Catalysis for the Synthesis of Spirocyclic Compounds. *ACS Catal.* **2013**, *3*, 540–553. [[CrossRef](#)]
31. Pawlowski, R.; Skorka, P.; Stodulski, M. Radical-Mediated Non-Deardomative Strategies in Construction of Spiro Compounds. *Adv. Synth. Catal.* **2020**, *362*, 4462–4486. [[CrossRef](#)]
32. Bycroft, B.W.; Shute, R.E.; Begley, M.J. Novel β -lactamase inhibitory and antibacterial 6-spiro- epoxy penicillins. *J. Chem. Soc. Chem. Commun.* **1988**, 274–276. [[CrossRef](#)]
33. Turos, E.; Long, T.E.; Heldreth, B.; Leslie, J.M.; Reddy, G.S.K.; Wang, Y.; Coates, C.; Konaklieva, M.; Dickey, S.; Lim, D.V.; et al. *N*-Thiolated β -lactams: A new family of anti-Bacillus agents. *Bioorganic Med. Chem. Lett.* **2006**, *16*, 2084–2090. [[CrossRef](#)]
34. Sheehan, J.C.; Chacko, E.; Lo, Y.S.; Ponzi, D.R.; Sato, E. Some new spiro penicillins. *J. Org. Chem.* **1978**, *43*, 4856–4859. [[CrossRef](#)]
35. Benfatti, F.; Cardillo, G.; Gentilucci, L.; Tolomelli, A. Synthesis and biological evaluation of unprecedented classes of spiro- β -lactams and azido- β -lactams as acyl-CoA:cholesterol acyltransferase inhibitors. *Bioorganic Med. Chem. Lett.* **2007**, *17*, 1946–1950. [[CrossRef](#)] [[PubMed](#)]
36. Narender Reddy, A.V.; Fiakpui, C.Y.; Czajkowski, D.P.; Spevak, P.; Kaleta, J.; Micetich, R.G.; Maiti, S.N. Synthesis and human leukocyte elastase inhibitory activity of novel 2-spiro (2',2'-diphenylcyclopropane) cephalosporin sulfones. *Chem. Heterocycl. Compd.* **1998**, *34*, 1289–1295. [[CrossRef](#)]
37. Skiles, J.W.; McNeil, D. Spiro indolinone beta-lactams, inhibitors of pliovirus and rhinovirus 3C-proteinases. *Tetrahedron Lett.* **1990**, *31*, 7277–7280. [[CrossRef](#)]
38. Santos, B.S.; Nunes, S.C.C.; Pais, A.A.C.C.; Pinho e Melo, T.M.V.D. Chiral spiro- β -lactams from 6-diazopenicillanates. *Tetrahedron* **2012**, *68*, 3729–3737. [[CrossRef](#)]
39. Hirai, K.; Iwano, Y.; Saito, T.; Hiraoka, T.; Kishida, Y. Functionalisation of C₆₍₇₎ of penicillins and cephalosporins via 1,3-dipolar intermediate. *Tetrahedron Lett.* **1976**, *17*, 1303–1306. [[CrossRef](#)]
40. Santos, B.S.; Pino e Melo, T.M.V.D. Synthesis of chiral spirocyclopentenyl- β -lactams through phosphane-catalyzed [3+2] annulation of allenates with 6-alkylidenepenicillanates. *Eur. J. Org. Chem.* **2013**, 3901–3909. [[CrossRef](#)]
41. Pitlik, J. Cycloaddition and related reactions of cephalosporin antibiotics. *Bioorganic Med. Chem.* **1995**, *3*, 1157–1181. [[CrossRef](#)]
42. Yang, D.-P.; Ji, H.-F.; Tang, G.-Y.; Ren, W.; Zhang, H.-Y. How Many Drugs Are Catecholics. *Molecules* **2007**, *12*, 878–884. [[CrossRef](#)]
43. Ivanova, G.; Bratovanova, E.; Petkov, D. Catechol as a nucleophilic catalyst of peptide bond formation. *J. Pept. Sci.* **2002**, *8*, 8–12. [[CrossRef](#)]
44. Cabiddu, S.; Floris, C.; Melis, S.; Sotgiu, F.; Cerioni, G. Heterocyclic compound studies. II. Synthesis of 1,4-benzodioxin, 1,4-benzoxathiin, 1,4-benzoxazine and 1,4-benzothiazine derivatives. *J. Heterocycl. Chem.* **1986**, *23*, 1815–1820. [[CrossRef](#)]

45. van Berkel, S.S.; Brem, J.; Rydzik, A.; Salimraj, R.; Cain, R.; Verma, A.; Owens, R.J.; Fishwick, C.W.G.; Spencer, J.; Schofield, C.J. Assay Platform for Clinically Relevant Metallo- β -lactamases. *J. Med. Chem.* **2013**, *56*, 6945–6953. [[CrossRef](#)] [[PubMed](#)]
46. Nowak, P.M.; Sagan, F.; Mitoraj, M.P. Origin of Remarkably Different Acidity of Hydroxycoumarins—Joint Experimental and Theoretical Studies. *J. Phys. Chem. B* **2017**, *121*, 4554–4561. [[CrossRef](#)] [[PubMed](#)]
47. Musialik, M.; Kuzmicz, R.; Pawłowski, T.S.; Litwinienko, G. Acidity of Hydroxyl Groups: An Overlooked Influence on Antiradical Properties of Flavonoids. *J. Org. Chem.* **2009**, *74*, 2699–2709. [[CrossRef](#)]
48. McNab, H.; Ferreira, E.S.B.; Hulme, A.N.; Quye, A. Negative ion ESI-MS analysis of natural yellow dye flavonoids—An isotopic labelling study. *Int. J. Mass Spectrom.* **2009**, *284*, 57–65. [[CrossRef](#)]
49. Schweigert, N.; Zehnder, A.J.B.; Eggen, R.I.L. Chemical properties of catechols and their molecular modes of toxic action in cells, from microorganisms to mammals. *Environ. Microbiol.* **2001**, *3*, 81–91. [[CrossRef](#)]
50. Dhumrongvaraporn, S. Synthesis of mono- and dialkyl catechol ethers. Master' Thesis, Rochester Institute of Technology, Rochester, NY, USA; p. 1983.
51. Zagorevskii, V.A.; Kirsanova, Z.D.; Persianova, I.V.; Zykov, D.A. Alkylation and some physicochemical characteristics of 6,7- and 7,8-dihydroxycoumarins. *Chem. Heterocycl. Compd.* **1974**, *10*, 1023–1028. [[CrossRef](#)]
52. Hiesinger, K.; Dar'in, D.; Proschak, E.; Krasavin, M. Spirocyclic Scaffolds in Medicinal Chemistry. *J. Med. Chem.* **2021**, *64*, 150–183. [[CrossRef](#)] [[PubMed](#)]