



TITLE:

ON SEMIDEFINITE LINEAR FRACTIONAL OPTIMIZATION PROBLEMS (Study on Nonlinear Analysis and Convex Analysis)

AUTHOR(S):

KIM, MOON HEE; KIM, GWI SOO; LEE, GUE MYUNG

CITATION:

KIM, MOON HEE ...[et al]. ON SEMIDEFINITE LINEAR FRACTIONAL OPTIMIZATION PROBLEMS (Study on Nonlinear Analysis and Convex Analysis). 数理解析研究所講究録 2021, 2190: 95-97

ISSUE DATE:

2021-07

URL:

<http://hdl.handle.net/2433/265660>

RIGHT:

ON SEMIDEFINITE LINEAR FRACTIONAL OPTIMIZATION PROBLEMS

MOON HEE KIM, GWI SOO KIM, AND GUE MYUNG LEE

ABSTRACT. Recently, semidefinite optimization problems have been intensively studied since many optimization problems can be changed into the problems and the problems are very tractable ([5]). In this paper, we review the duality results for semidefinite linear fractional optimization problems in the paper ([3] On duality theorems for semidefinite linear fractional optimization problems, Journal of Nonlinear and Convex Analysis, volume 20, 2019, 1907-1912).

1. INTRODUCTION

B. D. Craven [1, 2] considered the following linear fractional program:

$$(LF) \quad \text{Maximize } \frac{c^T x + \alpha}{d^T x + \beta} \quad \text{subject to } x \geq 0, Ax = b,$$

where $c, d \in \mathbb{R}^n$, $\alpha, \beta \in \mathbb{R}$, $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$ are given, and formulated the dual problem for (LF) as follows:

$$(LD) \quad \text{Minimize } \gamma \quad \text{subject to } A^T s + d\gamma \geq c, \quad \beta\gamma - b^T s \geq \alpha,$$

and then studied the duality theorems which holds between (LF) and (LD).

Recently, semidefinite optimization problems have been intensively studied since many optimization problem can be changed into the problems and the problems are very tractable ([5]). In the paper [3], we considered a semidefinite linear fractional optimization problem (SLP) and formulated its dual problem (SLD), and then obtained the duality theorems which hold between (SLF) and (SLD). By using the B. D. Craven's approaches in [1, 2], we can prove the duality theorems which hold between (SLF) and (SLD). But we proved the duality theorems by a direct methods in the paper [3]. In this paper, we review the duality results for semidefinite linear fractional optimization problems in the paper [3].

Date: January 10, 2020.

2010 Mathematics Subject Classification. 90C25, 90C30, 90C46.

Key words and phrases. semidefinite linear fractional optimization problem, weak duality theorem, strong duality theorem, converse duality theorem .

We considered the semidefinite linear fractional optimization problem in the paper [3] :

$$\begin{aligned}
 \text{(SLF)} \quad & \text{Maximize} \quad \frac{\text{Tr}(CX) + \alpha}{\text{Tr}(DX) + \beta} \\
 & \text{subject to} \quad X \succeq 0, \text{Tr}(A_i X) = b_i, i = 1, \dots, m
 \end{aligned}$$

where C, D are given symmetric $n \times n$ matrices, α, β are given real numbers, $A_i, i = 1, \dots, m$ are given symmetric $n \times n$ matrices and $b_i, i = 1, \dots, m$ are given real numbers.

$\text{Tr}A$ is the trace of $n \times n$ matrix A . For a symmetric $n \times n$ matrix A , $A \succeq 0$ means that A is positive semidefinite and $A \succ 0$ means that A is positive definite. Let S^n be the space of $n \times n$ symmetric matrices. Then $\text{Tr}(\cdot)$ is the inner product on S^n and S^n is the finite-dimensional Banach space with the norm $\|\cdot\|$ induced by the inner product $\text{Tr}(\cdot)$ ([4]).

Moreover we formulated the dual problem for (SLF) in the paper [3] as follows;

$$\begin{aligned}
 \text{(SLD)} \quad & \text{Minimize} \quad \gamma \\
 & \text{subject to} \quad \sum_{i=1}^m y_i A_i + \gamma D \succeq C \\
 & \quad \quad \quad \beta \gamma - b^T y \geq \alpha.
 \end{aligned}$$

2. DUALITY THEOREMS

Let $\Delta = \{X \in S^n \mid X \succeq 0, \text{Tr}(A_i X) = b_i, i = 1, \dots, m\}$ and $F = \{(\gamma, y) \in \mathbb{R} \times \mathbb{R}^m \mid \sum_{i=1}^m y_i A_i + \gamma D \succeq C, \beta \gamma - b^T y \geq \alpha\}$.

Now we state the duality theorems in the paper [3] .

Theorem 2.1. [3] **(Weak duality)** Assume that $[X \in \Delta] \Rightarrow [\text{Tr}(DX) + \beta > 0]$. Let X be feasible (SLF) and let (γ, y) be feasible (SLD). Then

$$\frac{\text{Tr}(CX) + \alpha}{\text{Tr}(DX) + \beta} \leq \gamma.$$

Theorem 2.2. [3] **(Strong duality)** Assume that $[X \in \Delta] \Rightarrow [\text{Tr}(DX) + \beta > 0]$ and that $[S - \sum_{i=1}^m \mu_i A_i = 0] \Rightarrow [S = 0, \mu = 0]$. If \bar{X} is a solution of (SLF), then there exists $(\bar{\gamma}, \bar{y}) \in F$ such that $(\bar{\gamma}, \bar{y})$ is a solution of (SLD) and

$$\frac{\text{Tr}(CX) + \alpha}{\text{Tr}(DX) + \beta} = \bar{\gamma}.$$

Theorem 2.3. [3] **(Converse duality)** Assume that $[X \in \Delta] \Rightarrow [\text{Tr}(DX) + \beta > 0]$ and that Δ is bounded. Further assume that there exist $y_0 \in \mathbb{R}^m$ and $\gamma^0 \in \mathbb{R}$ such

that $\sum_{i=1}^m y_0^i A_i + \gamma^0 D - C \succ 0$ and $\beta \gamma^0 - b^T y > \alpha$. If $(\bar{\gamma}, \bar{y}) \in F$ is a solution of (SLD) then there exists a solution \bar{X} of (SLF) such that

$$\bar{\gamma} = \frac{\text{Tr}(C\bar{X}) + \alpha}{\text{Tr}(D\bar{X}) + \beta}.$$

REFERENCES

- [1] B. D. Craven, “*Fractional Programming*”, Heldermann Verlag, Berlin, 1988.
- [2] B. D. Craven and B. Mond, *The dual of a fractional linear program*, Journal of Mathematical Analysis and Applications. Math. Anal. Appl., 42 (1973), 507-512
- [3] M. H. Kim, G. W. Kim and G. M. Lee, *On duality theorems for semidefinite linear fractional optimization problems*, Journal of Nonlinear and Convex Analysis, 20(2019), 1907-1912.
- [4] Etienne de Klerk, “*Aspects of Semidefinite Programming: Interior Point Algorithms and Selected Applications*”, Kluwer Academic Publishers, 2002.
- [5] L. Vandenberghe and S. Boyd, *Semidefinite programming*, SIAM Review, 38 (1996), 49-95.

(M. H. Kim) COLLEGE OF GENERAL EDUCATION, TONGMYONG UNIVERSITY, BUSAN 48520, KOREA.

Email address: mooni@tu.ac.kr

(G. S. Kim) DEPARTMENT OF APPLIED MATHEMATICS, PUKYONG NATIONAL UNIVERSITY, BUSAN 48513, KOREA.

Email address: gwisoo1103@hanmail.net

(G. M. Lee) CORRESPONDING AUTHOR, DEPARTMENT OF APPLIED MATHEMATICS, PUKYONG NATIONAL UNIVERSITY, BUSAN 48513, KOREA

Email address: gmlee@pknu.ac.kr