



TITLE:

Maass forms on  $GL(2)$  over division quaternion algebras of discriminant  $p$   
(Analytic, geometric and  $p$ -adic aspects of automorphic forms and  $L$ -functions)

AUTHOR(S):

Narita, Hiro-aki; Pitale, Ameya; Wagh, Siddhesh

---

CITATION:

Narita, Hiro-aki ...[et al]. Maass forms on  $GL(2)$  over division quaternion algebras of discriminant  $p$  (Analytic, geometric and  $p$ -adic aspects of automorphic forms and  $L$ -functions). 数理解析研究所講義録 2021, 2197: 118-123

ISSUE DATE:

2021-08

URL:

<http://hdl.handle.net/2433/265781>

RIGHT:

# Maass forms on $GL(2)$ over division quaternion algebras of discriminant $p$

Hiroaki Narita, Ameya Pitale, Siddhesh Wagh

## 1 Introduction

In this talk, we will present a construction of Maass forms, that violate the Ramanujan conjecture, on 5-dimensional hyperbolic spaces. To provide some context, let us remind the reader of another famous example of modular forms that violate the Ramanujan conjecture – the Saito-Kurokawa lifts.

**Saito-Kurokawa lifts:** Let  $f \in S_{2k-2}(SL_2(\mathbb{Z}))$ , with  $k$  even, and let  $h \in S_{k-1/2}^+(\Gamma_0(4))$  be the corresponding cusp form in the Kohnen plus space. Let  $\{c(n)\}$  be the Fourier coefficients of  $h$ . For  $T$  half integral, positive definite, symmetric  $2 \times 2$  matrix, define

$$A(T) := \sum_{d|\gcd(T)} c\left(\frac{\det(2T)}{d^2}\right) d^{k-1}.$$

**1.1 Theorem.** *With  $A(T)$  as above, the function  $F_f(Z) = \sum_T A(T) \exp(2\pi i \operatorname{Tr}(TZ))$  is a Siegel cusp form of weight  $k$  with respect to  $Sp_4(\mathbb{Z})$ .*

Let us list some of the properties of the Saito-Kurokawa lifts (see [2] for details).

1. Explicit formula for Fourier coefficients.
2. The map  $f \mapsto F_f$  is linear and injective.
3. Relation between  $L$ -functions

$$L(s, F_f, \text{spin}) = \zeta(s - k + 1) \zeta(s - k + 2) L(s, f).$$

4. The map  $f \mapsto F_f$  preserves Hecke eigenforms.
5. If  $F_f$  is a Hecke eigenform, then let  $\pi_F = \otimes_p \pi_p$  be the irreducible cuspidal automorphic representation of  $GSp_4(\mathbb{A})$  generated by  $F_f$ . Then, for every  $p < \infty$ , the local representation  $\pi_p$  is not tempered, i.e.  $F_f$  violates the generalized Ramanujan conjecture.
6. Characterization of lifts as the Maass space: For  $T = \begin{bmatrix} m & r/2 \\ r/2 & n \end{bmatrix}$ , write  $A(T) = A(m, r, n)$ . Then a Siegel cusp form  $F$  with Fourier coefficients  $A(T)$  is a Saito-Kurokawa lift if and only if we have

$$A(m, r, n) = \sum_{d|\gcd(m, r, n)} d^{k-1} A\left(\frac{mn}{d^2}, \frac{r}{d}, 1\right).$$

2 MAASS FORMS ON 5-DIMENSIONAL HYPERBOLIC SPACE

2 Maass forms on 5-dimensional hyperbolic space

Let  $B$  be a definite division quaternion algebra over  $\mathbb{Q}$ . Let us make the assumption that the discriminant of  $B$  is a prime number  $p$ .

Let  $G$  be the algebraic group such that  $G(\mathbb{Q}) = \text{GL}_2(B)$ . Then  $G(\mathbb{R}) = \text{GL}_2(\mathbb{H})$ , where  $\mathbb{H}$  is the Hamiltonian quaternions. We have the Iwasawa decomposition:  $\text{GL}_2(\mathbb{H}) = ZNAK$ , where  $Z$  is the center, and  $K$  is the maximal compact, and

$$N = \{n(x) = \begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} : x \in \mathbb{H}\}, A = \{a_y = \begin{bmatrix} \sqrt{y} & \\ & \sqrt{y}^{-1} \end{bmatrix} : y \in \mathbb{R}^+\}.$$

We have

$$G/ZK \simeq \{ \begin{bmatrix} y & x \\ & 1 \end{bmatrix} : x \in \mathbb{H}, y \in \mathbb{R}^+ \},$$

a realization of the 5-dimensional hyperbolic space  $\mathbb{H}_5$ . For a discrete subgroup  $\Gamma \subset \text{GL}_2(\mathbb{H})$  and  $r \in \mathbb{C}$  we denote by  $\mathcal{M}(\Gamma, r)$  the space of smooth functions  $F$  on  $\text{GL}_2(\mathbb{H})$  satisfying the following conditions:

1.  $\Omega \cdot F = -\frac{1}{2}(\frac{r^2}{4} + 1)F$ , where  $\Omega$  is the Casimir operator,
2. for any  $(z, \gamma, g, k) \in Z \times \Gamma \times G \times K$ , we have  $F(z\gamma gk) = F(g)$ ,
3.  $F$  is of moderate growth.

We will take  $\Gamma = \text{GL}_2(\mathcal{O})$ , where  $\mathcal{O}$  is any maximal order in  $B$ . Let  $\mathcal{O}'$  be the dual of  $\mathcal{O}$  with respect to trace map on  $B$ . For  $F \in \mathcal{M}(\text{GL}_2(\mathcal{O}), r)$ , we have the Fourier expansion

$$F(n(x)a_y) = u(y) + \sum_{\beta \in \mathcal{O}' \setminus \{0\}} A(\beta)y^2 K_{\sqrt{-1}r}(2\pi|\beta|y)e^{2\pi\sqrt{-1}\text{tr}(\beta x)}$$

3 The Maass lift

Let  $f \in S(\Gamma_0(p), \frac{r^2+1}{4})$  be an Atkin Lehner eigenfunction with eigenvalue  $\epsilon \in \{-1, 1\}$ . Let  $\{c(n) : n \in \mathbb{Z} - \{0\}\}$  be the Fourier coefficients of  $f$ . Let us define the primitive elements of  $\mathcal{O}'$  by

$$\mathcal{O}'_{\text{prim}} := \{ \beta \in \mathcal{O}' : \frac{1}{n}\beta \notin \mathcal{O}' \text{ for all positive integers } n \}.$$

Write  $\beta \in \mathcal{O}'$  as

$$\beta = p^u n \beta_0, \quad u \geq 0, n > 0, p \nmid n \text{ and } \beta_0 \in \mathcal{O}'_{\text{prim}}.$$

Set

$$\delta = \begin{cases} 0 & \text{if } \beta_0 \notin \mathcal{O}; \\ 1 & \text{if } \beta_0 \in \mathcal{O}. \end{cases}$$

Define

$$A_f(\beta) := |\beta| \sum_{t=0}^{2u+\delta} \sum_{d|n} c\left(\frac{-|\beta|^2}{p^{t-1}d^2}\right)(-\epsilon)^t. \tag{1}$$

The main theorem is the following.

4 BORCHERDS THETA LIFTS

**3.1 Theorem.** Let  $f \in S(\Gamma_0(p), \frac{r^2+1}{4})$  be an Atkin Lehner eigenfunction with eigenvalue  $\epsilon \in \{-1, 1\}$  with Fourier coefficients  $\{c(n)\}$ . For  $\beta \in \mathcal{O}'$ , define  $A_f(\beta)$  as above. Then the function  $F_{f, \mathcal{O}}$  on  $\text{GL}_2(\mathbb{H})$  with Fourier coefficients  $A_f(\beta)$  is a cusp form in  $\mathcal{M}(\text{GL}_2(\mathcal{O}), r)$ .

One way to prove the automorphy is to use the converse theorem due to Maass.

**3.2 Theorem. (Maass [3])**  $F$  given by the Fourier expansion is in  $\mathcal{M}(\Gamma_{\mathcal{O}}, r)$  if and only if a family of twisted Dirichlet series are “nice”.

Here,  $\Gamma_{\mathcal{O}} = \langle \begin{bmatrix} 1 & \beta \\ & 1 \end{bmatrix}, \begin{bmatrix} & 1 \\ -1 & \end{bmatrix} : \beta \in \mathcal{O} \rangle$ . Unfortunately, we have  $\text{GL}_2(\mathcal{O}) = \Gamma_{\mathcal{O}}$  if and only if  $p = 2, 3, 5$ . We have used the Maass converse theorem to prove automorphy for  $p = 2$  in joint paper with Muto-Narita [4]. For general  $p$ , the strategy is to use Borcherds theta lifts.

4 Borcherds Theta lifts

In a nutshell, the idea for the theta lift is given by

$$\Phi(n(x)a_y) \sim \int_{\text{SL}_2(\mathbb{Z}) \backslash \mathfrak{h}} f(\tau) \Theta(\tau, n(x)a_y) d\tau.$$

To execute the strategy we have to do the following two things.

1. Replace  $f$  by a vector valued modular form with respect to  $\text{SL}_2(\mathbb{Z})$ .
2. Define the theta kernel.

Let us first define the vector valued modular forms. Define the discriminant form  $D = \mathcal{O}'/\mathcal{O} \simeq (\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/p\mathbb{Z})$ . The group algebra  $\mathbb{C}[D]$  is a  $\mathbb{C}$ -vector space generated by the formal basis vectors  $\{e_{\mu} : \mu \in D\}$  with product defined by  $e_{\mu}e_{\mu'} = e_{\mu+\mu'}$ . Let  $\text{SL}_2(\mathbb{Z})$  act on  $\mathbb{C}[D]$  via the representation  $\rho_D$  as follows:

$$\rho_D\left(\begin{bmatrix} 1 & 1 \\ & 1 \end{bmatrix}\right)e_{\mu} = e(|\mu|^2)e_{\mu}, \rho_D\left(\begin{bmatrix} & -1 \\ 1 & \end{bmatrix}\right)e_{\mu} = -\frac{1}{p} \sum_{\mu' \in D} e(-(\mu, \mu'))e_{\mu'}.$$

Here  $e(x) = \exp(2\pi ix)$ . Now, given  $f \in S(\Gamma_0(p), \frac{r^2+1}{4})$ , define  $\mathcal{L}_D(f) : \mathfrak{h} \rightarrow \mathbb{C}[D]$  by

$$(\mathcal{L}_D(f))(\tau) = \sum_{\Gamma_0(p) \backslash \text{SL}_2(\mathbb{Z})} f(M\langle\tau\rangle) \rho_D(M)^{-1}(e_0).$$

The main result is

**4.1 Proposition.** Let  $f \in S(\Gamma_0(p), \frac{r^2+1}{4})$  be an Atkin Lehner eigenfunction with eigenvalue  $\epsilon \in \{-1, 1\}$  with Fourier coefficients  $\{c(n)\}$ .

1. For all  $\gamma \in \text{SL}_2(\mathbb{Z})$ , we have

$$\mathcal{L}_D(f)|\gamma = \rho_D(\gamma)\mathcal{L}_D(f).$$

4 BORCHERDS THETA LIFTS

2. Write  $\mathcal{L}_D(f) = \sum_{\mu \in D} f_\mu e_\mu$ . Let  $c_\mu(n)$  be the Fourier coefficients of  $f_\mu$ . Then we have

$$c_\mu(n) = \begin{cases} c(n) - \epsilon c(np) & \text{if } \mu = 0; \\ -\epsilon c(n) & \text{if } \mu \neq 0, n \equiv |\mu|^2 \pmod{p}; \\ 0 & \text{otherwise.} \end{cases}$$

Next, let us define the theta kernel. Let  $(\mathcal{O}, |\cdot|^2) \simeq (\mathbb{Z}^4, A_0)$ . Set  $L := [\mathbb{Z}, \mathcal{O}, \mathbb{Z}]^t \simeq (\mathbb{Z}^6, A)$  with  $A = \begin{bmatrix} & & & & & 1 \\ & & & & & -A_0 \\ & & & & & \\ & & & & & \\ & & & & & \\ & & & & & \\ 1 & & & & & \end{bmatrix}$ . Let  $V = (\mathbb{R}^6, Q_A) = L \otimes \mathbb{R} \simeq \mathbb{R}^{1,5}$ . We have that the connected component of  $\text{SO}(V) \simeq \text{SO}(1, 5)$  is isomorphic to  $\text{GL}_2(\mathbb{H})/Z$ . Let  $\mathcal{D}$  be the Grassmanian of positive oriented lines in the quadratic space  $V$ . We can identify the 5-dimensional hyperbolic space  $\mathbb{H}_5$  with the connected component  $\mathcal{D}^+$  via

$$\begin{aligned} \mathbb{H}_5 \ni (x, y) &\mapsto \nu(x, y) := \frac{1}{\sqrt{2}} {}^t(y + y^{-1}Q_{A_0}(x), -y^{-1}x, y^{-1}) \\ &\mapsto \mathbb{R} \cdot \nu(x, y) \in \mathcal{D}^+. \end{aligned}$$

Every  $\nu := \nu(x, y)$  defines an isometry

$$\iota_\nu : V \rightarrow \mathbb{R} \cdot \nu \oplus (\nu^\perp, Q_{A_0}|_{\nu^\perp}) \simeq \mathbb{R}^{1,5}, \quad \lambda \rightarrow (\lambda_\nu, \lambda_{\nu^\perp}).$$

Let  $p : \mathbb{R}^6 \rightarrow \mathbb{R}$  be the polynomial given by  $p(x_1, \dots, x_6) = -2^{-2}x_1^2$ . For  $\tau = u + iv \in \mathfrak{h}$ ,  $(x, y) \in \mathbb{H}_5$ , define the theta function

$$\Theta_L(\tau, \nu(x, y), p) := \sum_{\mu \in D} \left( \sum_{\lambda \in L + \mu} \left( \exp\left(\frac{-\Delta}{8\pi v}\right)(p) \right) (\iota_\nu(\lambda)) e(Q_A(\lambda_\nu)\tau + Q_A(\lambda_{\nu^\perp})\bar{\tau}) \right) e_\mu.$$

Here,  $\Delta$  is the Laplacian on  $\mathbb{R}^{1,5}$ .

**4.2 Proposition. (Borcherds [1])** For  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \text{SL}_2(\mathbb{Z})$ , we have

$$\Theta_L\left(\frac{a\tau + b}{c\tau + d}, \nu(x, y), p\right) = |c\tau + d|^5 \rho_D\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) \Theta_L(\tau, \nu(x, y), p).$$

For  $(x, y) \in \mathbb{H}_5$ , define

$$\Phi_{f, \mathcal{O}}(\nu(x, y)) := \int_{\text{SL}_2(\mathbb{Z}) \backslash \mathfrak{h}} \langle \mathcal{L}_D(f)(\tau), \overline{\Theta_L(\tau, \nu(x, y), p)} \rangle v^{\frac{5}{2}} \frac{dudv}{v^2}.$$

**4.3 Proposition.** For every  $\gamma \in \text{GL}_2(\mathcal{O})$ , we have

$$\Phi_{f, \mathcal{O}}(\gamma\nu(x, y)) = \Phi_{f, \mathcal{O}}(\nu(x, y)).$$

*Proof.*  $\Theta_L$  is invariant under a subgroup of  $\text{GL}_2(\mathcal{O})$  that fixes  $\mathcal{O}'/\mathcal{O}$ . Action of  $\text{GL}_2(\mathcal{O})$  preserves norms on  $\mathcal{O}'/\mathcal{O}$ , and Fourier coefficients of  $f_\mu, \mu \in \mathcal{O}'/\mathcal{O}$  only depend on  $|\mu|^2$ . ■

5 MAASS SPACE

Borchers gives explicit formula for the Fourier coefficients of  $\Phi_{f,\mathcal{O}}(\nu(x, y))$ . We compute this to show that the Fourier coefficients of  $\Phi_{f,\mathcal{O}}(\nu(x, y))$  are exactly  $A_f(\beta)$  defined in (1). Hence, we obtain

$$\begin{aligned} \Phi_{f,\mathcal{O}}(\nu(x, y)) &= \sum_{\beta \in \mathcal{O} \setminus \{0\}} A_f(\beta) y^2 K_{\sqrt{-1}r}(2\pi|\beta|y) e^{2\pi\sqrt{-1}\text{tr}(\beta x)} \\ &= F_{f,\mathcal{O}}(n(x)a_y), \end{aligned}$$

which shows that  $F_{f,\mathcal{O}} \in \mathcal{M}(\text{GL}_2(\mathcal{O}), r)$ . Cuspidality follows from the observation that the Fourier expansion of  $\Phi_{f,\mathcal{O}}$  at a different cusp corresponds to the Fourier expansion of the Borchers lift for a shift of  $\mathcal{O}$ . This completes the proof of Theorem 3.1.

If  $f$  is a non-zero even Hecke eigenform, then  $c(-1) \neq 0$ . Hence  $A_f(1) \neq 0$ , and we get non-vanishing of  $F_{f,\mathcal{O}}$ . To show that  $F_{f,\mathcal{O}}$  is non-zero for a general  $f$ , we use the fact that the space of Maass forms  $f$  for a fixed  $p$  and  $r$  is finite dimensional. In addition, we need to show that  $f \rightarrow F_{f,\mathcal{O}}$  is Hecke equivariant.

If a prime  $\ell \neq p$ , then  $B \otimes \mathbb{Q}_\ell =: B_\ell \simeq M_2(\mathbb{Q}_\ell)$  and  $\text{GL}_2(B_\ell) \simeq \text{GL}_4(\mathbb{Q}_\ell)$ . Hence, we can use the well-known Hecke theory for  $\text{GL}_4$  and show that if  $f$  is a Hecke eigenform, then  $F_{f,\mathcal{O}}$  is also a Hecke eigenform.

Now, let  $F_{f,\mathcal{O}}$  be a Hecke eigenform. Suppose  $\pi_{F,\mathcal{O}} = \otimes \pi_\ell$  is the irreducible cuspidal automorphic representation of  $\text{GL}_2(B_\mathbb{A})$  corresponding to  $F_{f,\mathcal{O}}$ . Let  $\sigma_f = \otimes \sigma_\ell$  be the irreducible cuspidal automorphic representation of  $\text{GL}_2(\mathbb{A})$  associated to  $f$ . Then, for  $\ell \neq p$ , the local representation  $\pi_\ell$  is the spherical component of the induced representation  $\text{Ind}_{P_{2,2}(\mathbb{Q}_\ell)}^{\text{GL}_4(\mathbb{Q}_\ell)}(|\det|^{-1/2}\sigma_\ell \times |\det|^{1/2}\sigma_\ell)$ . We have

$$L(s, \pi_{F,\mathcal{O}}) = L(s + 1/2, \sigma_f)L(s - 1/2, \sigma_f),$$

i.e.  $F_{f,\mathcal{O}}$  does not satisfy the generalized Ramanujan conjecture. Note that the strong multiplicity one theorem for  $\text{GL}_2(B_\mathbb{A})$  implies that, if  $\mathcal{O}_1$  and  $\mathcal{O}_2$  are two maximal orders in  $B$ , then  $\pi_{F,\mathcal{O}_1} = \pi_{F,\mathcal{O}_2}$ . Hence,  $F_{f,\mathcal{O}_1}$  and  $F_{f,\mathcal{O}_2}$  give two vectors in the same representation.

5 Maass space

Let us finish with the description of Maass space in the case  $p = 2$ . Let the Maass space  $\mathcal{M}^*(\text{GL}_2(\mathcal{O}), r)$  denote the subspace of cusp forms  $F$  in  $\mathcal{M}(\text{GL}_2(\mathcal{O}), r)$  with Fourier coefficients  $A(\beta)$  satisfying the following.

1. If  $\beta = \varpi_2^n n \beta_0$ , then  $A(\beta)$  depends only on  $K := |\beta|^2, u$  and  $n$ . We write  $A(\beta)$  as  $A(K, u, n)$ .
2.  $A(K, u, n)$  satisfy the recurrence relation
  - $A(K, u, n) = (-3\epsilon/\sqrt{2})A(K/2, u - 1, n) - A(K/4, u - 2, n)$  for some  $\epsilon \in \{-1, 1\}$ .
  - $A(K, u, n) = \sum_{d|n} dA(K/d^2, u, 1)$ .

**5.1 Theorem. (Wagh [5])**  $F \in \mathcal{M}^*(\text{GL}_2(\mathcal{O}), r)$  if and only if  $F = F_f$  for some  $f \in S(\Gamma_0(2), \frac{r^2+1}{4})$ .

We plan to extend this theorem to  $p > 2$  in the future.

## REFERENCES

## References

- [1] BORCHERDS, R. E.: *Automorphic forms with singularities on Grassmanians*. Invent. Math., **132** (1998), 491–562.
- [2] EICHLER, M., ZAGIER, D.: *The Theory of Jacobi Forms* Progress in Mathematics, 55. Birkhauser, Boston (1985)
- [3] MAASS, H.: *Automorphe Funktionen von mehreren Veränderlichen und Dirchletsche Reihen*. Abh. Math. Sem. Univ. Hamburg, **16**, no. 3-4, (1949) 72–100.
- [4] MUTO, M., NARITA, H., PITALE, A.: *Lifting to  $GL(2)$  over a division quaternion algebra and an explicit construction of CAP representations*. Nagoya Math. J., **222**, issue 01, (2016) 137–185.
- [5] WAGH, S.: *Maass space for lifts to  $GL_2$  over a division quaternion algebra* Journal of Number Theory, **28**, (2020) 230–261

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, TOKYO, JAPAN

*Email address:* [hnarita@waseda.jp](mailto:hnarita@waseda.jp)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK, 73019, USA

*Email address:* [apitale@ou.edu](mailto:apitale@ou.edu)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, USA

*Email address:* [wagsiddhesh@gmail.com](mailto:wagsiddhesh@gmail.com)