

SHORT COMMUNICATION

First record of the gray-legged tinamou, *Crypturellus duidae*, and other poor-soil specialist birds from peatlands in the Putumayo River basin, Loreto, Peru

Francisco A. VÁSQUEZ-ARÉVALO^{1*}, Ricardo ZÁRATE-GÓMEZ², Jacob B. SOCOLAR³, Juan DÍAZ-ALVÁN⁴, Pedro E. PÉREZ-PEÑA⁵

¹ Universidad Nacional de la Amazonía Peruana (UNAP). Facultad de Ciencias Biológicas. Pebas 5ta Cuadra, Iquitos, Maynas, Loreto, Peru

² Instituto de Investigaciones de la Amazonía Peruana (IIAP), Programa de Investigación en Cambio Climático Desarrollo Territorial y Ambiente, Av. A. Quiñones km 2.5 (Iquitos), San Juan Bautista, Maynas, Loreto, Peru

³ Norwegian University of Life Sciences, Universitetstunet 3, 1430 Ås, Norway

⁴ Universidad Científica del Perú (UCP), Av. A. Quiñones km 2.5 (Iquitos), San Juan Bautista, Maynas, Loreto, Peru

⁵ Instituto de Investigaciones de la Amazonía Peruana (IIAP), Programa de Investigación en Biodiversidad Amazónica, Av. A. Quiñones km 2.5 (Iquitos), San Juan Bautista, Maynas, Loreto, Peru

* Corresponding author: fvasquezare@gmail.com; <https://orcid.org/0000-0002-8177-0757>

ABSTRACT

Crypturellus duidae (Tinamidae) is a poor-soil specialist with isolated populations in Amazonia, and is considered restricted to white-sand forest habitats. We report the first record of *C. duidae* in a peatland forest in northern Peru, in the Putumayo River basin. Our record extends the known distribution of *C. duidae* between two disjoint areas of occurrence in Peru and Colombia, and shows its presence in peatland forest, another forest type on nutrient-poor soils. Additionally, we report the presence of other poor-soil specialist bird species that were previously registered in peatlands. Together with the new record of *C. duidae*, these bird records provide evidence of the diversity of poor-soil specialists in peatland forests.

KEYWORDS: *Mauritia flexuosa*, palm swamps, white-sand forests, Amazonia

Primer registro de la perdiz de pata gris, *Crypturellus duidae*, y otras aves especialistas de suelos pobres en turberas en la cuenca del Río Putumayo, Loreto, Perú

RESUMEN

Crypturellus duidae (Tinamidae) es considerada un ave especialista de suelos pobres, con poblaciones aisladas y restringidas a bosques sobre arena blanca en Amazonía. En este trabajo reportamos el primer registro de *C. duidae* en un bosque de turbera al norte de Perú, en la cuenca del Río Putumayo. Nuestro registro extiende la distribución conocida de *C. duidae* entre dos áreas de ocurrencia disjunta entre Perú y Colombia, y muestra su presencia en bosques de turbera, otro tipo de bosque sobre suelos pobres en nutrientes. Adicionalmente reportamos la presencia de otras aves especialistas de suelos pobres anteriormente registradas en turberas. En conjunto con el nuevo registro de *C. duidae*, esos registros de aves proporcionan evidencia de la diversidad de especialistas de suelos pobres en bosques de turberas.

PALABRAS-CLAVE: *Mauritia flexuosa*, pantanos de palmeras, bosque sobre arena blanca, Amazonía

In the Loreto region, in the northern Peruvian Amazon, white-sand forests are a distinctive collection of forest types characterized by high stem densities (roughly 1000 stems > 10 cm DBH per hectare), a canopy as low as 5 m (García-Villacorta *et al.* 2003), and a distinctive floristic assemblage dominated by specialist trees (Zárate *et al.* 2015). The avifauna is also distinctive and includes a variety of specialist species such as *Poliophtila clementsi* Whitney and Álvarez, 2005,

Neopelma chrysocephalum Pelzelin, 1868, *Zimmerius villarejoi* Álvarez and Whitney, 2001, among others (Álvarez *et al.* 2012; Álvarez *et al.* 2013; Borges *et al.* 2016; Díaz-Alván *et al.* 2017; Socolar *et al.* 2018).

The gray-legged tinamou, *Crypturellus duidae* J.T. Zimmer, 1938 (Tinamidae) is a poorly known species distributed disjunctly on nutrient-poor white-sand forests across northern South America (Schulenberg *et al.* 2010; BirdLife

CITE AS: Vásquez-Arévalo, F.A.; Zárate-Gómez, R.; Socolar, J.B.; Díaz-Alván, J.; Pérez-Peña, P.E. . 2020. First record of the gray-legged tinamou, *Crypturellus duidae*, and other poor-soil specialist birds from peatlands in the Putumayo River basin, Loreto, Peru. *Acta Amazonica* 50: 155-158.

International 2017; Borges *et al.* 2017). In Peru, the species is known exclusively from white-sand forests in the Nanay and middle Tigre river basins (Álvarez and Whitney 2003; Díaz-Alván *et al.* 2017). The species is apparently absent from extensive white-sand forests along the Morona River and south of the Amazonas River (O’Shea *et al.* 2015; Schmitt *et al.* 2017).

Recent work in Loreto has documented extensive peatland forests in swampy river floodplains (Draper *et al.* 2014). Forests that receive floodwaters are frequently dominated by the palm *Mauritia flexuosa* L.f., while domed ombrotrophic peatland swamps, that receive water and nutrients only from precipitation, and not through connections with water currents, such as minerotrophic peatlands (Lähteenoja *et al.* 2009), resemble white-sand forests in several important ways, including severe nutrient limitation, stunted physiognomy, and low epiphyte loads (Draper *et al.* 2014). Moreover, while these forests are severely depauperate in tree species, the few that are present are often shared with white-sand forests and include white-sand specialists (Zárate *et al.* 2013; Draper *et al.* 2018). The avifauna of ombrotrophic peatlands also shows clear white-sand affinities. For example, recent records indicate that peatlands along the lower Tigre River, in Loreto-Peru, harbor white-sand specialists (Díaz-Alván *et al.* 2017).

Here, we report a record of *Crypturellus duidae* from a peatland forest in the Putumayo River basin in Loreto, Peru. In addition to being the first record of *C. duidae* from a peatland forest, this is the first record of the species in Peru far from white sand and its known occurrence areas in the upper Nanay River basin in Peru and in the Chiribiquete region of Colombia (Álvarez *et al.* 2003).

From 14 August to 11 September 2017 we surveyed bird communities in peatlands and *terra firme* forests dominated by the palms *Mauritia flexuosa* and *Oenocarpus bataua* Mart., respectively, along the upper Putumayo River, Loreto, Peru. In the vicinity of our sampling site, we assessed habitat in 20 x 50 m tree plots (Stohlgren *et al.* 1995; Judd *et al.* 1999). On 1 September 2017, near the town of Santa Rita, on the Yubinetto River, we recorded the distinctive song of *Crypturellus duidae* from a peatland dominated by *M. flexuosa* (1°00’02.12”S, 74°19’39.35”W; Figure 1). The song was recorded at 07:25 a.m. using an LG H502 smartphone and the RecForce II application for Android system. The voice-recording is diagnostic for *C. duidae* and is available in the Xeno Canto library (XC429170, www.xeno-canto.org). In addition to the voice-recorded individual, we detected two further individuals vocalizing nearby.

Habitat at the site consisted of palms with scattered dicot trees growing on waterlogged substrate. The canopy height varied roughly between 12 to 19 m, with emergent trees and palms as tall as 25 m. Stem densities (≥ 10 cm DBH) varied between 78 and 96 stems per 0.1 ha, of which 17 to 23 stems




Figure 1. A – Location of the record of the gray-legged tinamou, *Crypturellus duidae* in the Yubinetto River drainage, Putumayo, Loreto, Peru. The inset shows the global range of *C. duidae* (based on BirdLife International). B – Image of the habitat in which the gray-legged tinamou was recorded. This figure is in color in the electronic version.

involved the dominant palm *M. flexuosa*. Other common tree species included *Oxandra euneura* Diels, *Euterpe precatoria* Mart., *Cynometra spruceana* Benth., *Virola pavonis* (A. DC.) A.C. Sm., among others.

Apart from *Crypturellus duidae*, we recorded in the upper Putumayo study area a number of other bird species with white-sand forest affinity in peatlands, *terra firme* forests, and non-peatland floodplain forest (Table 1). Our records represent a new habitat and an extension of the known distribution range of *C. duidae* in Peru. Amazonian peatlands are geologically ephemeral formations, with radiocarbon dates for the onset of peat accumulation generally of a few thousand years ago (Lähteenoja *et al.* 2012; Roucoux *et al.* 2013). Thus, the occurrence of *C. duidae* on isolated peatlands in northern Loreto, coupled with the presence of a variety of poor-soil specialist species in the surrounding *terra firme*, suggests that *C. duidae* might maintain low-density populations across a vast forest area between known populations near Iquitos, Peru and in Chiribiquete, Colombia.

Table 1. Poor-soil specialist birds (Álvarez *et al.* 2013; Borges *et al.* 2016) registered in the Putumayo River basin in Loreto, Peru. Habitat codes: F = floodplain forest; P = peatland; T = terra firme forest. Previous peatland record = Species was previously registered in peatland forests according to Díaz-Alván *et al.* (2017).

Species	Family	Common name	Habitat	Previous peatland record
<i>Nyctibius bracteatus</i>	Nyctibiidae	Rufous Potoo	F	
<i>Trogon rufus</i>	Trogonidae	Black-throated Trogon	P, T	X
<i>Galbula dea</i>	Galbulidae	Paradise Jacamar	T	X
<i>Northarchus ordii</i>	Bucconidae	Brown-banded Puffbird	T	X
<i>Megascictus margaritatus</i>	Thamnophilidae	Pearly Antshrike	T	X
<i>Hypocnemis hypoxantha</i>	Thamnophilidae	Yellow-browed Antbird	F, T	
<i>Sclerurus rufularis</i>	Furnariidae	Short-billed Leaf-tosser	T	X
<i>Deconychura longicauda</i>	Furnariidae	Long-tailed Woodcreeper	T	
<i>Lepidocolaptes duidae</i>	Furnariidae	Duida Woodcreeper	T	X
<i>Lophotriccus galeatus</i>	Tyrannidae	Helmeted Pygmy-Tyrant	T	X
<i>Conopias parvus</i>	Tyrannidae	Yellow-throated Flycatcher	F, T	X
<i>Ramphotrigon ruficauda</i>	Tyrannidae	Rufous-tailed Flatbill	F, P, T	X
<i>Attila citriniventris</i>	Tyrannidae	Citron-bellied Attila	F, P, T	X
<i>Heterocercus aurantiivertex</i>	Pipridae	Orange-crowned Manakin	F, P, T	X
<i>Dixiphia pipra</i>	Pipridae	White-crowned Manakin	T	X

While peatland pole forest (a type of ombrotrophic peatland different in physiognomy and composition of plant species than peatland dominated by *M. flexuosa*; Draper *et al.* 2014, 2017) is known to harbor a variety of white-sand specialist birds (Díaz-Alván *et al.* 2017), our records of *C. duidae* and other poor-soil specialist species recorded on peatlands are the first from peatlands dominated by *M. flexuosa* (peatland palm swamp) (Draper *et al.* 2014, 2017). Because *M. flexuosa* palm swamps dominate a vast area of northern Peru (Draper *et al.* 2014), these records suggest that some of these species might be more widely distributed than previously thought.

The mechanisms governing habitat selection for *C. duidae* and other white-sand specialists are not well known. It is perhaps noteworthy that the vast majority of *C. duidae* records from the Nanay basin are from white-sand forests on waterlogged substrates with a deep layer of organic humus (“varillal humedo”; J. Socolar, personal observation), which may affect the foraging or nesting ecology of *C. duidae*. On white-sand, *C. duidae* is often associated with dense understory stands of the palm *Euterpe catinga* Wallace, which do not exist at the Putumayo site, though its congener *E. precatória* is relatively common. It is also possible that *C. duidae* is a dietary specialist on fruit or invertebrate resources that are restricted to moist poor-soil forests.

As in Peru, in other countries, *C. duidae* is considered an indicator species of white-sand forest (Hurd *et al.* 1995; Borges *et al.* 2016, 2017). However, there is a previous record of *C. duidae* within the Chiribiquete National Park, in a flooded forest with a structure similar to forests that grow on poor soils (Álvarez *et al.* 2003). Although it is known that there are peatlands in the Guayana Highland landscape (including the Serranía de Chiribiquete) (Zinck and Huber 2011), it is not known whether the record of *C. duidae* was related

to a peatland forest. The presence of species with isolated populations, such as *C. duidae* and other poor-soil specialist birds in the upper Putumayo region suggests the existence of previously unrecognized poor-soil bird assemblages across northernmost Peru, which possibly afford connectivity between populations on the Guianan shield and populations near Iquitos (Socolar *et al.* 2018).

ACKNOWLEDGMENTS

We thank the Instituto de Investigaciones de la Amazonía Peruana for funding our fieldwork and the Proyecto Especial Binacional Desarrollo Integral de la Cuenca del Río Putumayo for support. We thank the entire field team involved in the inventory: Geancarlo Coello, Claudia Ramos, Guillisa Flores, Ian Medina, Jorge Pizarro, Amilcar Ortiz, Victor Vásquez and Kember Mejía. We thank members of the Mashunta, Santa Rita, and Nueva Jerusalén indigenous communities for support in the field. We thank Luis Torres and Marcos Ríos for sharing information about peatlands in Peru, and Surecht Ruiz for assistance in preparing the map in Figure 1. Finally, we thank Claudia Keller, Sergio Borges and two anonymous reviewers for their comments on the manuscript.

REFERENCES

- Álvarez, A.J.; Whitney, B.M. 2003. New distributional records of birds from white-sand forests of the northern Peruvian Amazon, with implications for biogeography of northern South America. *Condor*, 105: 552–566.
- Álvarez, M.; Umaña, A.M.; Mejía, G.D.; Cajiao, J.; von Hildebrand, P.; Gast, F. 2003. Aves del Parque Nacional Natural Serranía de Chiribiquete, Amazonia-Provincia de la Guyana, Colombia. *Biota Colombiana*, 4: 49-63.
- Álvarez, A.J.; Díaz, A.J.; Shany, N. 2012. Avifauna de la Reserva Nacional Allpahuayo Mishana, Loreto. Peru. *Cotinga*, 34: 61-84.

- Álvarez, A.J.; Metz, M.R.; Fine, P.V.A. 2013. Habitat specialization by birds in western Amazonian white-sand forests. *Biotropica*, 45: 365–372.
- Borges, S.H.; Cornelius, C.; Ribas, C.; Almeida, R.; Guilherme, E.; Aleixo, A.; *et al.* 2016. What is the avifauna of Amazonian white-sand vegetation? *Bird Conservation International*, 26: 192–204.
- Borges, S.H.; Whittaker, A.; Almeida, R.A.; Cornelius, C.; Santos-Jr, M.A.; Moreira, M. 2017. Bird records in the northwestern and central portions of the Amazon Basin highlight the needs for inventories and long-term monitoring in the region. *Revista Brasileira de Ornitologia*, 25: 206–220.
- BirdLife International. 2017. Species factsheet: *Crypturellus duidae* (<http://datazone.birdlife.org/species/factsheet/22678213>). Accessed on 16 Jan 2020.
- Díaz-Alván, J.; Socolar, J.B.; Álvarez, A.J. 2017. The Avifauna of the Río Tigre Basin, Northern Peru. *Ornitología Neotropical*, 28: 11 – 21.
- Draper, F.C.; Roucoux, K.H.; Lawson, I.T.; Mitchard, E.T.A.; Honorio, C.E.N.; Lähenteenoja, O.; Torres, M.L.; Valderrama, S.E.; Zárate, R.; Baker, T.R. 2014. The distribution and amount of carbon in the largest peatland complex in Amazonia. *Environmental Research Letters*, 9: 1–12.
- Draper, F.C.; Honorio, C.E.N.; Roucoux, K.H.; Lawson, I.T.; Pitman, N.C.A.; Fine, P.V.A.; *et al.* 2018. Peatland forests are the least diverse tree communities documented in Amazonia, but contribute to high regional beta-diversity. *Ecography*, 41: 1256 – 1269.
- García-Villacorta, R.; Ahuite-Reátegui, M.; Olortegui-Zumaeta, M. 2003. Clasificación de bosques sobre arena blanca de la Zona Reservada Allpahuayo-Mishana. *Folia Amazonica*, 14: 17–33.
- Hurd, F.G.S.; Telleria, J.L.; Díaz, M. 1995. Observaciones sobre la composición, ecología, y zoogeografía de la avifauna de la Sierra de Chiribiquete, Caquetá, Colombia. *Caldasia*, 17: 481–500.
- Judd, W.; Campbell, C.; Kellogg, E.; Stevens, P. 1999. *Plant Systematics. A Phylogenetic Approach*. Sinauer Associates, Sunderland, Massachusetts. xvi + 464p.
- Lähenteenoja, O.; Ruokolainen, K.; Schulman, L.; Alvarez, J. 2009. Amazonian floodplains harbour minerotrophic and ombrotrophic peatlands. *Catena*, 79: 140–145.
- Lähenteenoja, O.; Reátegui, Y.R.; Räsänen, M.; Torres, D.D.C.; Oinonen, M.; Page, P. 2012. The large Amazonian peatland carbon sink in the subsiding Pastaza-Marañón foreland basin, Peru. *Global Change Biology*, 18: 164–178.
- O’Shea, B.J.; Stotz, D.F.; Saboya, C.P.; Ruelas, I.E. 2015. Birds. In: Pitman, N.A.; Vriesendorp, C.; *et al.* (Ed.). *Peru: Tapiche Blanco. Rapid Biological and Social Inventories Report 27*. The Field Museum, Chicago, p.305–320.
- Roucoux, K.H.; Lawson, I.T.; Jones, T.D.; Baker, T.R.; Honorio, C.E.N.; Gosling, W.D.; Lähenteenoja, O. 2013. Vegetation development in an Amazonian peatland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 374: 242–255.
- Schmitt, F.; Sané, R.; Thibault, M.; Vásquez, G. 2017. New locality for White-masked Antbird *Pithys castaneus* and other avian range extensions for dpto. Loreto, Peru. *Cotinga*, 39: 2–11.
- Schulenberg, T.S.; Stotz, D.F.; Lane, D.F.; O’Neill, J.P.; Parker III, T.A. 2010. *Birds of Peru: Revised and updated edition*. Princeton Univ. Press, Princeton, 664p.
- Socolar, J.B.; Díaz-Alván, J.; Saboya, D. P.; Pormara, L.Y.; O’Shea, B.J.; Cubas, S.P.; *et al.* 2018. Noteworthy bird records from northeastern Peru reveal connectivity and isolation in the western Amazonian avifauna. *The Wilson Journal of Ornithology* 130: 94–111.
- Stohlgren, T.; Falkner, M.; Schell, L. 1995. Modified-Whittaker Nested Vegetation Sampling Method. *Vegetatio*, 117: 113–121.
- Zárate, R.; Mori, T.; Maco, J.T. 2013. Estructura y composición florística de las comunidades vegetales del ámbito de la carretera Iquitos-Nauta, Loreto-Peru. *Folia Amazonica*, 22: 77–89.
- Zárate, R.; Mori, T.; Ramirez, F.; Davila, H.; Gallardo, P.; Cohello, G. 2015. Lista actualizada y clave para la identificación de 219 especies arbóreas de los bosques sobre arena blanca de la Reserva Nacional Allpahuayo Mishana, Loreto, Peru. *Acta Amazonica*, 45:133–156.
- Zinck, J. A.; Huber, O. 2011. *Peatlands of the Western Guayana Highlands, Venezuela. Properties and Paleogeographic Significance of Peats*. 1st ed. Springer, Heidelberg, Dordrecht, London, New York, 280p.

RECEIVED: 26/04/2019

ACCEPTED: 16/01/2020

ASSOCIATE EDITOR: Sérgio H. Borges



This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.