



Publication Year	2015
Acceptance in OA @INAF	2020-04-23T14:12:26Z
Title	Photometric study of HD 15555 in the beta Pictoris Association
Authors	MESSINA, Sergio; Millward, Mervyn; Bradstreet, David H.
DOI	10.1016/j.newast.2014.12.007
Handle	http://hdl.handle.net/20.500.12386/24211
Journal	NEW ASTRONOMY
Number	37

Photometric study of HD 155555C in the β Pictoris Association

Sergio Messina ^{a,*}, Mervyn Millward ^b, David H. Bradstreet ^c

^a*INAF- Catania Astrophysical Observatory, via S.Sofia, 78 I-95123 Catania, Italy*

^b*York Creek Observatory, Georgetown, Tasmania*

^c*Eastern University, St. Davids, PA, USA*

Abstract

We are carrying out a series of photometric monitoring to measure the rotation periods of members in the young β Pictoris Association, as part of the RACE-OC project (Rotation and ACTivity Evolution in Open Clusters). In this paper, we present the results for HD 155555C which is believed to be physically associated to the spectroscopic binary V824 Ara (HD155555) and thus constituting a triple system. We collected B, V, and R-band photometric data timeseries and discovered from periodogram analysis the rotation period $P = 4.43$ d. Combined with stellar radius and projected rotational velocity, we find this star almost equator-on with an inclination $i \simeq 90^\circ$. The rotational properties of HD155555C fit well into the period distribution of other β Pic members, giving further support to the suggested membership to the association and to its physical association to V824 Ara. A comparison with Pre-Main-Sequence isochrones from various models allows us to estimate an age of 20 ± 15 Myr for this triple system.

Key words: Stars: activity - Stars: low-mass - Stars: rotation - Stars: starspots - Stars: pre main sequence: individual: HD155555C

1 Introduction

We are carrying out a systematic investigation aimed at determining the rotation periods and the photospheric activity levels of all confirmed and candidate

* Corresponding author.

Email addresses: sergio.messina@oact.inaf.it (Sergio Messina),
mervyn.millward@yorkcreek.net (Mervyn Millward), dbradstr@eastern.edu
(David H. Bradstreet).

members of the young β Pictoris stellar Association (Zuckerman et al. 2001). This investigation is part of the RACE-OC project (Rotation and Activity Evolution in Open Clusters; Messina [2007]). In an earlier study by Messina et al. [2010], we focused on only confirmed members and derived the rotation periods for a total of 27 of them either from the literature or from our analysis of photometric timeseries in the ASAS (All Sky Automated Survey; Pojmanski [1997]) and SuperWASP (Wide Angle Search for Planets; Butters et al. [2010]) archives. However, a few members have remained with unknown or uncertain periods for a number of reasons: they were not in the archives; the low-photometric precision in combination with low variability level prevented us from rotation period detection; they were in spatially unresolved multiple systems. Meanwhile, new candidate members have been proposed, e.g., by Lepine et al. [2009], Kiss et al. [2011], Shkolnik et al. [2012], Schlieder et al. [2012], Malo et al. [2014]. In order to have a complete rotational characterization of the association, we have undertaken a number of collaborations. In the present paper, we present the results of the monitoring project carried out at the York Creek Observatory on HD 155555C, one of the first proposed members to this Association.

We focus our investigation on the pre-main-sequence triple system HD 155555 consisting of a SB2 spectroscopic binary, HD 155555AB, and a nearby single dM component, HD 155555C. HD 155555AB has been identified as member of the young stellar Association β Pictoris by Zuckerman et al. [2001]. The nearby dM star has been suggested to be physically connected to the SB2 by Pasquini et al. [1989] and Pasquini et al. [1991], and therefore to be a member of the same Association. However, no parallax of the latter is measured, and membership to this stellar system as well as to the Association are only inferred, though significantly supported by a number of evidences. Our rotational investigation aims to add additional information to get a better characterization of this dM component.

For instance, this system, like other multiple systems, see, e.g., Messina et al. [2014], is particularly interesting since a few parameters are equal, like age, initial chemical composition, and mass for some systems. These commonalities allow us to explore the dependence of their rotational evolution on other parameters, such as initial angular velocity and environment properties (presence of disc or nearby stellar/planetary companions), shedding light on the causes of dispersion observed in the rotation period distributions of all young clusters/associations.

2 Literature information

HD 155555 is a triple stellar system in the southern constellation of Ara at a distance $d = 31.4 \pm 0.5$ pc (van Leeuwen [2007]). This system consists of a

double-line spectroscopic binary HD 155555AB and of a fainter M dwarf companion HD 155555C at an angular distance of about $33''$, which corresponds to about 1040 AU.

The spectroscopic binary HD 155555AB was discovered by Bennet et al. [1967] to be composed of G5IV + K0IV stars that rotate with an orbital period $P = 1.687$ d. First evidence of photometric variability of this SB2 was found by Eggen [1978] after comparing his photometry collected in 1978 ($V = 6.83$, $B-V = 0.835$, $U-B = 0.29$, $R = 6.46$, and $R-I = 0.325$ mag) with the earlier photometry collected by Stoy [1963] ($V = 6.67$ and $B-V = 0.80$ mag), even though the latter also included the dM companion, owing to the large aperture of $44''$ used to collect the photoelectric photometry. Subsequent long-term photometric monitoring by Cutispoto (see, e.g., Cutispoto 1998 and references therein) revealed robust evidence of the existence of short- and long-term photometric variability. HD 155555AB has the variable's name V824 Ara in the Combined General Catalogue of Variable Stars [Kholopov 1998]. The most recent determination of orbital and physical properties, and first magnetic maps of both components were provided by the spectro-polarimetric study of Dunstone et al. [1998]. Most recent surface temperature maps have been obtained with an improved Doppler imaging technique by Kristovics et al. [2013]. The membership of HD155555AB to the β Pictoris moving group was first suggested by Zuckerman et al. [2001] based on distance, UVW velocity components, high $v \sin i$, and L_X/L_{bol} ratio. Also the photometric variability observed by Eggen (1978) and Cutispoto (1998) are in agreement with the youth of the system.

The present investigation is focused on the fainter HD 155555C component (also named LDS 587B) which is a M3.5 dwarf (Pasquini et al. 1991). Eggen [1978] reported the following magnitudes and colors: $V = 12.82$, $B-V = 1.54$, $U-B = 1.05$, $R = 11.40$, and $R-I = 1.205$ mag, noticing both UV and IR flux excess with respect to standard colors of main sequence stars. This component is a strong X-ray source (EXO 171224-6653.9), which was a serendipitous discovery by Barstow et al. [1987] during EXOSAT observations of HD 155555AB (EXO 171217-6653.8). Pasquini et al. [1989] identified this X-ray source with HD 155555C, and subsequent spectroscopic investigation (Pasquini et al. 1991) showed high levels of magnetic activity, the HeI, NaI D2 and D1 lines all being in emission. The $H\alpha$ line was also found to be in emission and variable from night to night with the equivalent widths ranging from 6.2 to 9.7 Å. Pasquini et al. [1991] could measure only an upper value ($\leq 3 \text{ km s}^{-1}$) of the projected rotational velocity. Their study concluded that HD155555 AB and HD155555 C are physically associated and are likely PMS stars. They noticed that radial velocities are similar ($V_r = 2.74 \text{ km s}^{-1}$ for HD 155555AB compared to $V_r = 3.5 \text{ km s}^{-1}$ for HD 155555C. Torres et al. [2006] also find that the UVW velocity components and radial velocity of HD155555C are similar to those of HD155555AB, suggesting their physical association.

The physical association is also supported by proper motion studies by Martin et al. [1995], who find the angular separation changed from $33''$ (about




Fig. 1. Light curves of HD155555C phased with the rotation period in three filters. Blue bullets denote observations collected after HJD 2456761 to better show the light curve evolution. The rectangle denotes a possible flare event.

in 1929) to $32.6''$ in 1994, suggesting similar proper motions. Similarly, the Washington Visual Double Star Catalog [Mason et al. 2001] reports a small variation from $33''$ in 1920 to $34''$ in 2000. In the literature, two other measurements of projected rotational velocity are available from Torres et al. [2006] $v \sin i = 6.0 \pm 1.2 \text{ km s}^{-1}$ and from Weise et al. [2010] $v \sin i = 8.8 \pm 0.9 \text{ km s}^{-1}$.

This system has been searched for possible planetary companions using the NACO adaptive optics system at the Very Large Telescope (VLT) [Masciadri et al. 2005], and the MMT telescope [Biller et al. 2007]. It was included in the SIM PlanetQuest Key Project [Tanner et al. 2007], and more recently in the Gemini/NICI planet-finding campaign [Biller et al. 2013]. None of these investigations have so far detected the presence of any additional companion.

3 Photometric observations

HD 155555C was observed in 2014, from Feb 28 to May 10, for a total of 21 nights. Observations were collected at the York Creek Observatory ($41^\circ 06' 06''\text{S}$; $146^\circ 50' 33''\text{E}$, Georgetown, Tasmania) using a f/10 25cm Takahashi Mewlon reflector, equipped with a QSI 683ws-8 camera, and BVR standard Johnson-Cousins filters. The telescope has a FoV of $24.5' \times 18.5'$, and the plate scale is $0.44''/\text{pixels}$. On each night observations were collected from 3 to 6 consecu-




Fig. 2. Correlations among magnitude variations in the three filters. Solid lines are weighted linear fits, whereas labels indicates the linear correlation coefficient (r) and the fit slope with its uncertainty.




Fig. 3. *left panel*: Lomb-Scargle periodogram of HD15555C (solid line) with indication of the power peak corresponding to the rotation period $P = 4.43$ d. The dotted red line indicates the window spectral function, whereas the horizontal dashed line the power level corresponding to a $FAP = 0.01$. *right panel*: Clean periodogram.

tive hours, using 120-s integrations in each filter. A total of 360 frames in B, 359 in V, and 400 in R filter were obtained (see Table 1).

After bias subtraction and flat fielding, we extracted the magnitude timeseries for a total of 30 stars. These series were cleaned by applying a 3σ threshold to remove outliers. The differential photometry was performed using the technique of ensemble photometry (Gilliland et al. 1988; Bailer-Jones et al. 2001; Everett et al. 2001). The differential magnitude of the target star is computed with respect to the average magnitude of a large number of non-variable ref-




Fig. 4. Spectral Energy Distribution of HD155555C.

erence stars. In this averaging process, the uncertainties of the ensemble star magnitudes due to statistical fluctuations as well as short-term small incoherent variations will cancel each other. The uncertainty in the magnitude of the artificial comparison will therefore be smaller than the uncertainty on the magnitude of a single star and this, in turn, will produce less noisy light curves.

We started with 30 sufficiently bright stars, which were isolated, distributed all over the frame but not found to be close to the CCD edges, and common to all the frames. Having selected the ensemble, we computed (i) the magnitude of the artificial comparison star by averaging the instrumental magnitudes of ensemble stars, (ii) time-series differential magnitudes for each star of the ensemble and (iii) mean, median and the standard deviation (σ) associated with each time series. While constructing the final ensemble, we excluded those ensemble stars whose standard deviation was larger than the threshold σ value that we fixed to 0.010 mag. After a number of iterations, we were left with six stars whose standard deviations were smaller than 0.010 mag. In Table 2 we list the 2MASS designation [Cutri et al. 2003] comparison stars, the nearby optical sources from USNO catalogue [Monet et al. 2013], the coordinates and the J-band magnitude. These stars were used to build the ensemble comparison star and to obtain the differential magnitude time series of HD 155555C in B, V, and R filters.

In Fig. 1, we plot the B, V, and R light curves phased with the $P = 4.43$ d

Table 1
Log of observations.

Observation	# frames		
	Date	B	V
28/02/2014	8	10	8
02/03/2014	10	10	43
19/03/2014	4	4	34
24/03/2014	3	3	3
25/03/2014	19	16	17
03/04/2014	3	3	3
04/04/2014	5	6	5
05/04/2014	20	24	24
06/04/2014	22	20	14
12/04/2014	22	22	19
14/04/2014	13	15	15
24/04/2014	32	32	29
27/04/2014	17	17	18
30/04/2014	33	33	33
03/05/2014	24	26	26
04/05/2014	20	18	17
05/05/2014	18	18	17
06/05/2014	27	26	27
07/05/2014	25	24	19
08/05/2014	10	10	9
10/05/2014	25	22	20

rotation period (see Sect. 5). The average photometric uncertainties associated to each magnitude are 0.014, 0.007, and 0.005 mag in the B, V, and R filters, respectively.

4 Analysis

As shown in Fig. 2, the photometric timeseries show that HD 155555C varies in all B, V, and R bands. The magnitude variations are strongly correlated to

Table 2
Comparison stars.

2MASS	USNO-B1.0	RA	DEC	J mag
2MASS J17174644-6656455	09064-02437	17 17 46.44	-66 56 45.60	8.696
2MASS J17174439-6657287	0230-0693233	17 17 44.37	-66 57 28.84	11.285
2MASS J17175417-6657550	0230-0693339	17 17 54.17	-66 57 55.06	12.085
2MASS J17172719-6700468	0229-0763917	17 17 27.19	-67 00 46.86	11.457
2MASS J17172092-6701280	0229-0763856	17 17 20.93	-67 01 28.08	9.411
2MASS J17170984-6700057	0229-0763745	17 17 09.84	-67 00 05.77	12.383

each other. The Pearson linear correlation coefficients are all positive, $r_{BV} = 0.85$, $r_{VR} = 0.92$, and $r_{BR} = 0.89$, and all with a significance level $> 99.9\%$. Weighted linear fits to the magnitude variations (solid lines in Fig. 2) give the following slopes $\Delta B/\Delta V = 0.76 \pm 0.11$, $\Delta V/\Delta R = 1.12 \pm 0.06$, and $\Delta B/\Delta R = 1.23 \pm 0.17$. This is the behavior typically exhibited, according to Messina [2008], by *color-correlated* and *reddening stars*, whose color variations arise from cool spots. Such temperature inhomogeneities generally give rise to light curves whose amplitudes (peak-to-peak) increase towards shorter wavelengths, that is the B light curve has amplitude larger than the V light curve that is larger than the R light curve. In our specific case, we note that whereas V (whose amplitude is $\Delta V = 0.07$ mag) and R (whose amplitude is $\Delta R = 0.05$ mag) curves follow this trend, the B-band light curve has amplitude comparable, if not slightly smaller ($\Delta B = 0.06$ mag), than V ($\Delta B/\Delta V < 1$). Unfortunately, our B-band observations could not cover the rotation phases around the lightcurve maximum (phases 0.15–0.35). Therefore, the true amplitude of variation in the B band may be larger than measured and certainly further observations will address this aspect. The V and R light curves, for which we have the longest timeseries, exhibit evidence of a significant variation, the amplitude has decreased and the phase of light maximum migrated by $\Delta\phi = 0.2$, as likely consequence of the active regions growth and decay (ARGD), which are responsible for the observed variability.

5 Rotation period search

We used the Lomb-Scargle [Scargle 1982] and Clean [Roberts et al. 1987] periodogram analyses to search for significant periodicities in the HD155555C magnitude timeseries related to stellar rotation period. The results of our analysis are plotted in Fig. 3. In the left panel we plot the normalized Lomb-Scargle periodogram (solid line) with overplotted the spectral window function related to the data sampling (red dotted line). The horizontal dashed line indicates

the power level corresponding to a False Alarm Probability (FAP) of 1%, that is the probability that a power peak of that height simply arises from Gaussian noise in the data. The FAP was estimated using a Monte-Carlo method, i.e., by generating 1000 artificial light curves obtained from the real one, keeping the date but scrambling the magnitude values. In the right panel we plot the Clean periodogram where the power peak arising from the light rotational modulation dominates, whereas all secondary peaks, arising from the aliasing, are effectively removed. Our analysis finds that the stellar rotation period is $P = 4.43 \pm 0.05$ d. The uncertainty of the rotation period is computed following the prescription of Lamm et al. [2004].

6 SED analysis

We used the available optical, near-IR, and IR photometry to build the observed spectral energy distribution (SED). The UBVRI magnitudes from Eggen [1978], JHK magnitudes from 2MASS [Cutri et al. 2003], and W1-W4 magnitudes from WISE [Cutri et al. 2013] are listed in Table 3. The SED was fitted with a grid of theoretical spectra from the NextGen Model [Allard et al. 2012] and the best fit is obtained with a model of $T_{\text{eff}} = 3300 \pm 100$ K, $\log g = 4.0 \pm 0.3$ dex and metallicity $[\text{Fe}/\text{H}] = 0.0$ (see Fig.4). We note that no evidence of IR excess was found.

7 Target parameters

We can use effective temperature and luminosity to derive the stellar radius and the inclination of the rotation axis. We assume that HD155555AB and HD155555C are bound and, therefore, they have the same distance $d = 31.4$ pc as measured by Hipparcos for HD155555AB (Zuckerman et al. 2004). Using the brightest observed magnitude $V = 12.71$ mag (Zuckerman et al. 2001), and the bolometric correction $BC_V = -2.03$ mag from Pecaut & Mamajek [2013], we infer the bolometric magnitude $M_{\text{bol}} = 8.19 \pm 0.05$ mag, the luminosity $L = 0.042 \pm 0.005 L_{\odot}$, and the stellar radius $R = 0.65 \pm 0.05 R_{\odot}$. From these values and the stellar rotation period we infer the inclination of the stellar rotation axis $i \simeq 90^{\circ}$ (adopting $\langle v \sin i \rangle = 7.6 \text{ km s}^{-1}$).

In Fig. 5, we compare T_{eff} and luminosity of all components of HD 155555 with a set of isochrones taken from Baraffe et al. [1998], Siess et al. [2000], and D’Antona & Mazzitelli [1997], for solar metallicity to infer the age of each component and see if their coevalness hypothesis is sustainable. The physical parameters of HD 155555A and B are taken from Strassmeier & Rice [2000], with the only difference that luminosities are computed using the BC from Pecaut

Table 3

Photometry from the literature.

U	B	V	R	I	J	H	K	W1	W2	W3	W4
(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)	(mag)
15.41	14.36	12.82	11.40	10.19	8.542	7.916	7.629	7.528	7.358	7.204	6.940

& Mamajek [2013]. We see that the D’Antona & Mazzitelli [1997] model produces the smallest age dispersion among all components with an estimated age of 15 ± 7 Myr for the system and a mass $M=0.25 M_{\odot}$ for HD 155555C. The Baraffe et al. [1998] and Siess et al. [2000] models produce results that emphasize a larger dispersion between the age of the SB2 components and the age of HD155555C. According to these models HD155555AB has an age $A_{\text{Baraffe}} = 30\pm 10$ Myr and $A_{\text{Siess}} = 24\pm 8$ Myr that are about a factor 2 older than the age of $A_{\text{Baraffe}} = 13\pm 7$ Myr and $A_{\text{Siess}} = 12\pm 4$ Myr of HD155555C. Despite these systematic differences between the Siess et al. [2000] and Baraffe et al. [1998] models and the D’Antona & Mazzitelli [1997] model, the uncertainties on the age are quite large and our results are consistent with the age range reported by Zuckermann et al. [2001] who found the system components to be fitted by a set of isochrones in the age range from 5 to 30 Myr. Thus, the coevalness hypothesis is still sustainable with an age of 20 ± 15 Myr. Moreover, such an age is also consistent with the age of 21 ± 9 Myr that Mentuch et al. [2008] estimate for the β Pic Association and based on the Li depletion. We note that the low Li content ($EW_{\text{Li}} < 20 \text{ m}\text{\AA}$) measured by Martin et al. [1995] in the component C is consistent with the values measured on other Association members of similar spectral type. On the contrary, the hotter components A and B are in the un-depleted part of the Li distribution, which prevents us from using the Li content to constrain their ages.

8 Spot model

To show that the presence of cool spots is the most typical explanation of the light modulation observed in a low-mass star as HD155555C we performed a spot modeling of the light curves. We modeled the observed multi-band light curves using Binary Maker V 3.0 [Bradstreet & Steelman 2004] to infer some information on the geometrical and physical properties of spots on the photosphere of HD 155555C. Binary Maker V 3.0 models are almost identical to those generated by Wilson-Devinney program [Wilson & Devinney 1971] and uses Roche equipotentials to create star surfaces. In our modeling the second component is essentially “turned off” (i.e., assigned a near zero mass and luminosity) in order to model a single rotating star. The gravity-darkening coefficient has been assumed $\nu = 0.25$ [Kopal 1959], and limb-darkening coefficients




Fig. 5. HR diagrams. Dashed lines are isochrones whereas the blue solid line is the evolutionary track for a mass $M=0.25 M_{\odot}$. Models are from Baraffe et al. [1998] (top panel), Siess et al. [2000] (middle panel), and D’Antona & Mazzitelli [1997] (bottom panel).

from Claret et al. [2012] were also adopted. We adopted $T_{\text{eff}} = 3300 \text{ K}$, and inclination $i = 90^{\circ}$ as input parameters. To find a suitable spot configuration we took into account that the light curve shape is asymmetric, the decreasing flux branch is shorter than the increasing flux branch. This suggests the existence of two major spot groups separated in longitude. Unfortunately, we had no observations to be fit around the light curve maximum.

Thanks to the dependence of the light curve amplitude on the photometric band wavelength, we could constrain the spot temperature contrast and better determine the area of the spots responsible for the flux rotational modulation. The temperature contrast between spots and surrounding photosphere was




Fig. 6. Spot model of HD155555C. Observed normalized flux (small bullets) in the B, V, and R filters versus rotation phase with overplotted the fit model (solid line).




Fig. 7. Rotation period distribution of β Pic members from Messina et al. [2011]. Squares denote the components of the HD 155555 system.

found $T_{spot}/T_{phot} = 0.85$, which is a value generally measured in dM3 spotted stars (see, e.g., Berdyugina 2005). The only free parameters in our model remained the spot areas and the spot longitudes. The spots latitude cannot be constrained by photometry alone, especially in the case of an equator-on star as in our present case. We found a satisfactory fit of V and R light curves using two spots separated in longitude by about $140 \pm 30^\circ$ and with radius of about $r = 25 \pm 5^\circ$ each, which corresponds to about 17% of the whole photosphere. Such a value refers to that component of spots unevenly distributed in longitude and, then, represents a lower value for the effective percentage of spotted surface. In Fig. 6, we plot the normalized flux in the B, V, and R filters with overplotted the fits from our spot modeling. The B model light curve is to some extent poorer with respect to the other fits, especially at the light minimum, where the observed light level is brighter than modeled.

9 Conclusion

Our investigation allowed us to measure the rotation period $P = 4.43\text{d}$ of the dM component HD 155555C in the triple system HD155555. Comparing all components luminosities and effective temperatures with PMS models, we find that the hypothesis that all three components are coeval is sustainable with an age of 20 ± 15 Myr, which is consistent with literature values and with the age of the β Pic Association. Then, the difference of rotation periods can be explained considering the different nature of the components. The SB2 components that rotate with $P = 1.687\text{d}$ have already reached a tidal synchronization, their orbital and rotation period being equal (Strassmeier et al. 2000; Cutispoto 1998). The single dM is still contracting and spinning up its rotation rate while approaching the zero-age main-sequence. The rotation periods of all components fit well into the period distribution of other β Pic members (see Fig. 7). From the spot modeling we infer a high level of photospheric activity, with spots having a temperature contrast of 0.85 with respect to the unspotted photosphere and a covering fraction larger than about 17% of the whole photosphere. This level is consistent with the high activity level at the higher atmospheric levels reported in the literature.

Acknowledgments

The Authors thank the anonymous Referees for their helpful comments that allowed to improve the quality of the manuscript. The extensive use of the SIMBAD and ADS databases, operated by the CDS center, (Strasbourg, France), is also gratefully acknowledged.

References

- [Allard et al. 2012] Allard, F., Homeier, D., & Freytag, B. 2012, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 370, 2765
- [1998] Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P. H. 1998, *A&A*, 337, 403
- [1987] Barstow, M.A. 1987, *MNRAS*, 228, 251
- [Bailer-Jones et al. 2001] Bailer-Jones C. A. L., Mundt R., 2001, *A&A*, 367, 218
- [1967] Bennett, N.W.W, Evans, D.S., & Laing, J.D. 1967, *MNRAS*, 137, 107
- [2005] Berdyugina, S. V. 2005, *Living Reviews in Solar Physics*, vol. 2, no. 8
- [Biller et al. 2007] Biller, B.A., Close, L.M., Masciadri, E. 2007, *ApJS*, 173, 143
- [Biller et al. 2013] Biller, B.A., Liu, M.C., Wahhaj, Z. 2013, *ApJ*, 777, 160
- [Bradstreet & Steelman 2004] Bradstreet, D. H., & Steelman, D.P. 2004, “Binary Maker 3: Light Curve Synthesis Program”, (Contact Software, Norristown, PA)
- [2010] Butters, O. W. and West, R. G. and Anderson, D. R. et al. 2010, *A&A*, 520, 10
- [2012] Claret, A., Hauschildt, P. H., & Witte, S. 2012, *A&A*, 546, 12
- [1998] Cutispoto, G. 1998, *A&AS*, 131, 321
- [Cutri et al. 2003] Cutri, R. M., Skrutskie, M. F., van Dyk, S. et al. 2003, *2MASS All-Sky Catalog of Point Sources, VizieR On-line Data Catalog: II/246*
- [Cutri et al. 2013] Cutri, R. M., Wright, E. L., Conrow, T. et al. 2013, *AllWISE Data Release, VizieR On-line Data Catalog: II/328*
- [1997] D’Antona, F., & Mazzitelli, I. 1997, *MmSAI*, 68, 807
- [2009] da Silva, L. Torres, C.A.O., de la Rez, R., et al. 2009, *A&A*, 508, 833
- [2004] de La Reza, R. & Pinzon, G. 2004, *AJ*, 128, 1812
- [1998] Dunstone, N.J., Hussain, G.A.J., Collier Cameron, A., et al. 1998, *MNRAS*, 387, 1525
- [1978] Eggen, O.J. 1978, *IBVS*, 1426, 1
- [Everett et al. 2001] Everett M. E., Howell S. B., 2001, *PASP*, 113, 1428
- [Gilliland et al. 1988] Gilliland R. L. & Brown T. M., 1988, *PASP*, 100, 754
- [Kholopov 1998] Kholopov, P.N., Samus, N.N., Frolov, M.S., et al. 1998, *The Combined General Catalogue of Variable Stars, 4.1 Edition*.
- [2011] Kiss, L. L., Moór, A., Szalai, T. et al. 2011, *MNRAS*, 411,878

- [Kopal 1959] Kopal, Z. 1959, in *Close Binary Stars*, The International Astrophysics Series, London: Chapman & Hall, 1959
- [2013] Kriskovics, L., Vida, K., Kovári, Zs., Garcia-Alvarez, D., & Oláh, K. 2013, *AN*, 334, 976
- [2004] Lamm, M. H., Bailer-Jones, C. A. L., Mundt, R., Herbst, W., & Scholz, A. 2004, *A&A*, 417, 557
- [2009] Lépine, S. & Simon, M. 2009, *AJ*, 137, 3632
- [2014] Malo, L., Doyon, R., Lafrenière, D. et al. 2014, *ApJ*, 762, 88
- [1995] Martin, E. L., Brandner, W. 1995, *A&A*, 294, 744
- [Masciadri et al. 2005] Masciadri, E., Mundt, R., Henning, Th., & Alvarez, C. 2005, *ApJ*, 625, 1004
- [Mason et al. 2001] Mason, B.D., Wycoff, G.L., Hartkopf, W.I., et al. 2001, *AJ*, 122, 3466
- [2008] Mentuch, E., Brandeker, A., van Kerkwijk, M.H., Jayawardhana, R., & Hauschildt, P.H. 2008, *ApJ*, 689, 1127
- [2007] Messina, S. 2007, *MmSAI*, 78, 628
- [2008] Messina, S. 2008, *A&A*, 480, 495
- [2010] Messina, S., Desidera, S., Turatto, M., Lanzafame, A.C., & Guinan, E.F. 2010, *A&A*, 520, A15
- [2011] Messina, S., Desidera, S., Lanzafame, A. C., Turatto, M., & Guinan, E. F. 2011, *A&A*, 532, A10
- [2014] Messina, S., Monard, B., Biazzo, K., Melo, C. H. F., & Frasca, A. 2014, *A&A*, 570, A19
- [Monet et al. 2013] Monet, D.G., Levine, S.E., Casian, B., et al. 2013, *AJ*, 125, 984
- [1989] Pasquini, L., Schmitt, J.H.M.M., Harnden, F.R., Tozzi, G.P., & Krautter, J. 1989, *A&A*, 218, 187
- [1991] Pasquini, L., Cutispoto, G., Gratton, R., & Mayor, M. 1991, *A&A*, 248, 72
- [2013] Pecaut, M. J. & Mamajek, E. E. 2013, *ApJS*, 208, 9
- [1997] Pojmanski, G. 1997, *AcA*, 47, 467
- [Roberts et al. 1987] Roberts, D. H., Lehar, J., & Dreher, J. W. 1987, *AJ*, 93, 968
- [Scargle 1982] Scargle, J. D. 1982, *ApJ*, 263, 835
- [2012] Schlieder, J. E., Lépine, S., & Simon, M., 2012, *Aj*, 144, 109
- [2012] Shkolnik, E. L., Anglada-Escud, G., Liu, M.C. et al. 2012, *ApJ*, 758, 56

- [2000] Siess L., Dufour E., Forestini M. 2000, A&A, 358, 593
- [1963] Stoy, R.H. 1963, Mon. Not. Astr. Soc. S. Africa, 22, 157
- [2000] Strassmeier, K.G. & Rice J.B. 2000, A&A, 360, 1019
- [Tanner et al. 2007] Tanner, A., Beichman, C., Akeson,R., et al. 2007, PASP 119, 857
- [2006] Torres, C. A. O., Quast, G. R., da Silva, L., et al. 2006, A&A, 460, 695
- [2007] van Leeuwen F. 2007, A&A, 474, 653
- [2010] Weise, P., Launhardt, R., Setiawan, J. & Henning, T. 2010, A&A, 517, 88
- [Wilson & Devinney 1971] Wilson, R.E. & Devinney, E.J. 1971, ApJ, 166, 605
- [2001] Zuckerman, B., Song, I., Bessell, M.S., & Webb, R.A. 2001, ApJ, 562, 87
- [2004] Zuckerman, B. & Song, I. 2004, ARA&A, 42, 685